

MPRA

Munich Personal RePEc Archive

Teoreme de punct fix pentru operatori multivoci contractivi in spatii metrice generalizate

Guran, Liliana

"Titu Maiorescu" University of Bucharest, Romania

26. October 2010

Online at <http://mpra.ub.uni-muenchen.de/26204/>

MPRA Paper No. 26204, posted 26. October 2010 / 08:56

FIXED POINT THEOREMS FOR MULTIVALUED CONTRACTIVE OPERATORS ON GENERALIZED METRIC SPACES

Liliana Guran

Department of Accounting and Managerial Information System,
Faculty of Economic Sciences, Titu Maiorescu University
Dâmbovnicului 22, 040441, sector 4, Bucharest, Romania.
E-mail:gliliana.math@gmail.com

Abstract

In this paper we give a fixed point results for multivalued operators on generalized metric spaces endowed with a generalized w -distance. Then we study the data dependence for this new result.

KEY WORDS AND PHRASES: multivalued weakly Picard operator, w -distance, fixed point, multivalued operator.

MATHEMATICS SUBJECT CLASSIFICATION 2000: 47H10, 54H25.

1 Introduction

It is well known that Caristi's fixed point theorem [2] is equivalent to Ekeland's variational principle [4], which is nowadays an important tool in nonlinear analysis. Most recently, many authors studied and generalized Caristi's fixed point theorem to various directions. Using the concept of Hausdorff metric, Nadler Jr. [13] has proved a multivalued version of the Banach contraction

principle which states that each closed bounded valued contraction map on a complete metric space, has a fixed point.

Recently, Bae [1] introduced a notion of multivalued weakly contractive maps and applying generalized Caristi's fixed point theorems he proved several fixed point results for such maps in the setting of metric and Banach spaces. Many authors have been using the Hausdorff metric to obtain fixed point results for multivalued maps on metric spaces, but, in fact for most cases the existence part of the results can be proved without using the concept of Hausdorff metric.

Recently, using the concept of w -distance [9], Suzuki and Takahashi [20] introduced a notion of multivalued weakly contractive in short, w -contractive maps and improved the Nadler's fixed point result without using the concept of Hausdorff metric. Most recently, Latif [10] generalized the fixed point result of Suzuki and Takahashi [[20], Theorem 1].

The concept of multivalued weakly Picard operator (briefly MWP operator) was introduced in close connection with the successive approximation method and the data dependence phenomenon for the fixed point set of multivalued operators on complete metric space, by I. A. Rus, A. Petruşel and A. Sântămărian, see [19]. In [17] is presented the theory of multivalued weakly Picard operators in L-spaces.

In 1966 A. I. Perov was introduced the concept of generalized metric space and obtained a generalization of the Banach principle for contractive operators on spaces endowed with vector-valued metrics, see [16].

The purpose of this paper is to recall the notion of generalized w -distance in a generalized metric space. Also, we present some generalizations of some fixed point results obtained in [5] with respect to a generalized w -distance and we give a data dependence result for the new theorem of fixed point.

2 Preliminaries

Let (X, d) be a metric space. We will use the following notations:

$P(X)$ - the set of all nonempty subsets of X ;

$$\mathcal{P}(X) = P(X) \cup \emptyset$$

$P_c(X)$ - the set of all nonempty closed subsets of X ;

$P_b(X)$ - the set of all nonempty bounded subsets of X ;

$P_{b,c}(X)$ - the set of all nonempty bounded and closed, subsets of X ;

For two subsets $A, B \in P_b(X)$ we recall the following functionals.

$\delta : \mathcal{P}(X) \times \mathcal{P}(X) \rightarrow \mathbb{R}_+$, $\delta(A, B) := \sup\{d(a, b) | x \in A, b \in B\}$ - the diameter functional;

$H : \mathcal{P}(X) \times \mathcal{P}(X) \rightarrow \mathbb{R}_+$, $H(A, B) := \max\{\sup_{a \in A} \inf_{b \in B} d(a, b), \sup_{b \in B} \inf_{a \in A} d(a, b)\}$ - the Pompeiu-Hausdorff functional;

If $T : X \rightarrow P(X)$ is a multivalued operator, then we denote by $FixT$ the fixed point set of T , i.e. $FixT := \{x \in X | x \in T(x)\}$.

First we define the concept of **L-space**.

Definition 2.1 Let X be a nonempty set and $s(X) := \{(x_n)_{n \in \mathbb{N}} | x_n \in X, n \in \mathbb{N}\}$. Let $c(X) \subset s(X)$ a subset of $s(X)$ and $Lim : c(X) \rightarrow X$ an operator. By definition the triple $(X, c(X), Lim)$ is called an **L-space** if the following conditions are satisfied:

- (i) If $x_n = x$, for all $n \in \mathbb{N}$, then $(x_n)_{n \in \mathbb{N}} \in c(X)$ and $Lim(x_n)_{n \in \mathbb{N}} = x$.
- (ii) If $(x_n)_{n \in \mathbb{N}} \in c(X)$ and $Lim(x_n)_{n \in \mathbb{N}} = x$, then for all subsequences, $(x_{n_i})_{i \in \mathbb{N}}$, of $(x_n)_{n \in \mathbb{N}}$ we have that $(x_{n_i})_{i \in \mathbb{N}} \in c(X)$ and $Lim(x_{n_i})_{i \in \mathbb{N}} = x$.

By the definition an element of $c(X)$ is convergent and $x := Lim(x_n)_{n \in \mathbb{N}}$ is the limit of this sequence and we can write $x_n \rightarrow x$ as $n \rightarrow \infty$.

We will denote an L-space by (X, \rightarrow) .

Let us give some examples of L-spaces, see [17].

Example 2.1 (*L-structures on Banach spaces*) Let X be a Banach space. We denote by \rightarrow the strong convergence in X and by \rightharpoonup the weak convergence in X . Then (X, \rightarrow) , (X, \rightharpoonup) are L-spaces.

Example 2.2 (*L-structures on function spaces*) let X and Y be two metric spaces. Let $\mathbb{M}(X, Y)$ the set of all operators from X to Y . We denote by \xrightarrow{p}

the point convergence on $\mathbb{M}(X, Y)$, by \xrightarrow{unif} the uniform convergence and by \xrightarrow{cont} the convergence with continuity. Then $(\mathbb{M}(X, Y), \xrightarrow{p})$, $(\mathbb{M}(X, Y), \xrightarrow{unif})$ and $(\mathbb{M}(X, Y), \xrightarrow{cont})$ are L -spaces.

Definition 2.2 Let (X, \rightarrow) be an L -space. Then $T : X \rightarrow P(X)$ is a **multi-valued weakly Picard operator** (briefly *MWP operator*) if for each $x \in X$ and each $y \in T(x)$ there exists a sequence $(x_n)_{n \in \mathbb{N}}$ in X such that:

- (i) $x_0 = x, x_1 = y$;
- (ii) $x_{n+1} \in T(x_n)$, for all $n \in \mathbb{N}$;
- (iii) the sequence $(x_n)_{n \in \mathbb{N}}$ is convergent and its limit is a fixed point of T .

Let us give some examples of MWP operators, see [17],[19].

Example 2.3 Let (X, d) be a complete metric space and $T : X \rightarrow P_d(X)$ be a Reich type multivalued operator, i.e. there exists $\alpha, \beta, \gamma \in \mathbb{R}_+$ with $\alpha + \beta + \gamma < 1$ such that

$$H(T(x), T(y)) \leq \alpha d(x, y) + \beta D(x, T(x)) + \gamma D(y, T(y)),$$

for all $x, y \in X$. Then T is a MWP operator.

Example 2.4 Let (X, d) be a complete metric space and $T : X \rightarrow P_d(X)$ be a closed multifunction for which there exists $\alpha, \beta \in \mathbb{R}_+$ with $\alpha + \beta < 1$ such that $H(T(x), T(y)) \leq \alpha d(x, y) + \beta D(y, T(y))$, for every $x \in X$ and every $y \in T(x)$. Then T is a MWP operator.

Example 2.5 Let (X, d) be a complete metric space and $T_1, T_2 : X \rightarrow P_d(X)$ for which there exists $\alpha \in]0, \frac{1}{2}[$ such that

$$H(T_1(x), T_2(y)) \leq \alpha [D(x, T_1(x)) + D(y, T_2(y))],$$

for each $x, y \in X$. Then T_1 and T_2 are a MWP operators.

The concept of w -distance was introduced by O. Kada, T. Suzuki and W. Takahashi (see[9]) as follows:

Let (X, d) be a metric space. A functional $w : X \times X \rightarrow [0, \infty)$ is called w -distance on X if the following axioms are satisfied :

1. $w(x, z) \leq w(x, y) + w(y, z)$, for any $x, y, z \in X$;
2. for any $x \in X : w(x, \cdot) : X \rightarrow [0, \infty)$ is lower semicontinuous;
3. for any $\varepsilon > 0$, exists $\delta > 0$ such that $w(z, x) \leq \delta$ and $w(z, y) \leq \delta$ implies $d(x, y) \leq \varepsilon$.

Some examples of w -distance are given in [9].

Example 2.6 *Let (X, d) be a metric space. Then the metric "d" is a w -distance on X .*

Example 2.7 *Let X be a normed linear space with norm $\|\cdot\|$. Then the function $w : X \times X \rightarrow [0, \infty)$ defined by $w(x, y) = \|x\| + \|y\|$ for every $x, y \in X$ is a w -distance.*

Example 2.8 *Let (X, d) be a metric space and let $g : X \rightarrow X$ a continuous mapping. Then the function $w : X \times Y \rightarrow [0, \infty)$ defined by:*

$$w(x, y) = \max\{d(g(x), y), d(g(x), g(y))\}$$

for every $x, y \in X$ is a w -distance.

Let us recall a crucial lemma for w -distance (see[20] for more details).

Lemma 2.1 *Let (X, d) be a metric space, and let w be a w -distance on X . Let (x_n) and (y_n) be two sequences in X , let $(\alpha_n), (\beta_n)$ be sequences in $[0, +\infty[$ converging to zero and let $x, y, z \in X$. Then the following statements hold:*

1. *If $w(x_n, y) \leq \alpha_n$ and $w(x_n, z) \leq \beta_n$ for any $n \in \mathbb{N}$, then $y = z$.*
2. *If $w(x_n, y_n) \leq \alpha_n$ and $w(x_n, z) \leq \beta_n$ for any $n \in \mathbb{N}$, then (y_n) converges to z .*
3. *If $w(x_n, x_m) \leq \alpha_n$ for any $n, m \in \mathbb{N}$ with $m > n$, then (x_n) is a Cauchy sequence.*
4. *If $w(y, x_n) \leq \alpha_n$ for any $n \in \mathbb{N}$, then (x_n) is a Cauchy sequence.*

For the rest of the paper, if $v, r \in \mathbb{R}^m$, $v := (v_1, v_2, \dots, v_m)$ and $r := (r_1, r_2, \dots, r_m)$, then by $v \leq r$ means $v_i \leq r_i$, for each $i \in \{1, 2, \dots, m\}$, while $v < r$ means $v_i < r_i$, for each $i \in \{1, 2, \dots, m\}$.

Also, $|v| := (|v_1|, |v_2|, \dots, |v_m|)$ and, if $c \in \mathbb{R}$ then $v \leq c$ means $v_i \leq c_i$, for each $i \in \{1, 2, \dots, m\}$.

If $x_0 \in X$ and $r \in \mathbb{R}_+^m$ with $r_i > 0$ for each $i \in \{1, 2, \dots, m\}$ we will denote by $B(x_0; r) := \{x \in X \mid d(x_0, x) < r\}$ the open ball centered in x_0 with radius $r := (r_1, r_2, \dots, r_m)$ and by $\tilde{B}(x_0; r) := \{x \in X \mid d(x_0, x) \leq r\}$ the closed ball centered in x_0 with radius r .

In [8] we can find the notion of generalized w -distance as follows.

Definition 2.3 *Let (X, d) a generalized metric space. The mapping $\tilde{w} : X \times X \rightarrow \mathbb{R}_+^m$ defined by $\tilde{w}(x, y) = (v_1(x, y), v_2(x, y), \dots, v_m(x, y))$ is said to be a generalized w -distance if it satisfies the following conditions:*

- (w₁) $\tilde{w}(x, y) \leq \tilde{w}(x, z) + \tilde{w}(z, y)$, for every $x, y, z \in X$;
- (w₂) $v_i : X \times X \rightarrow \mathbb{R}_+$ is lower semicontinuous, for $i \in \{1, 2, \dots, m\}$;
- (w₃) For any $\varepsilon := (\varepsilon_1, \varepsilon_2, \dots, \varepsilon_m) > 0$, for $m \in \mathbb{N}$, there exists $\delta := (\delta_1, \delta_2, \dots, \delta_m) > 0$ such that $\tilde{w}(z, x) \leq \delta$ and $\tilde{w}(z, y) \leq \delta$ implies $\tilde{d}(x, y) \leq \varepsilon$.

The notion of generalized w -distance with his properties was discussed in [8].

Let us present now an important lemma for w -distances into the terms of generalized w -distances.

Lemma 2.2 *Let (X, \tilde{d}) be a generalized metric space, and let $\tilde{w} : X \times X \rightarrow \mathbb{R}_+^m$ be a generalized w -distance on X . Let (x_n) and (y_n) be two sequences in X , let $\alpha_n = (\alpha_n^{(1)}, \alpha_n^{(2)}, \dots, \alpha_n^{(m)}) \in \mathbb{R}_+$ and $\beta_n = (\beta_n^{(1)}, \beta_n^{(2)}, \dots, \beta_n^{(m)}) \in \mathbb{R}_+$ be two sequences such that $\alpha_n^{(i)}$ and $\beta_n^{(i)}$ converge to zero for each $i \in \{1, 2, \dots, m\}$. Let $x, y, z \in X$. Then the following assertions hold:*

1. If $\tilde{w}(x_n, y) \leq \alpha_n$ and $\tilde{w}(x_n, z) \leq \beta_n$ for any $n \in \mathbb{N}$, then $y = z$.

2. If $\tilde{w}(x_n, y_n) \leq \alpha_n$ and $\tilde{w}(x_n, z) \leq \beta_n$ for any $n \in \mathbb{N}$, then (y_n) converges to z .
3. If $\tilde{w}(x_n, x_m) \leq \alpha_n$ for any $n, m \in \mathbb{N}$ with $m > n$, then (x_n) is a Cauchy sequence.
4. If $\tilde{w}(y, x_n) \leq \alpha_n$ for any $n \in \mathbb{N}$, then (x_n) is a Cauchy sequence.

Throughout this paper we will denote by $M_{m,m}(\mathbb{R}_+)$ the set of all $m \times m$ matrices with positive elements, by Θ the zero $m \times m$ matrix, by I the identity $m \times m$ matrix and by U the unity $m \times m$ matrix. If $A \in M_{m,m}(\mathbb{R}_+)$, then the symbol A^τ stands for the transpose matrix of A .

Recall that a matrix A is said to be convergent to zero if and only if $A^n \rightarrow 0$ as $n \rightarrow \infty$.

For the proof of the main result we need the following theorem, see [16].

Theorem 2.1 *Let $A \in M_{m,m}(\mathbb{R}_+)$. The following statements are equivalent:*

- (i) A is a matrix convergent to zero;
- (i) $A^n \rightarrow 0$ as $n \rightarrow \infty$;
- (ii) The eigen-values of A are in the open unit disc, i.e. $|\lambda| < 1$, for every $\lambda \in \mathbb{C}$ with $\det(A - \lambda I) = 0$;
- (iii) The matrix $I - A$ is non-singular and

$$(I - A)^{-1} = I + A + \dots + A^n + \dots;$$

(iv) The matrix $I - A$ is non-singular and $(I - A)^{-1}$ has nonnegative elements.

- (v) $A^n q \rightarrow 0$ and $qA^n \rightarrow 0$ as $n \rightarrow \infty$, for each $q \in \mathbb{R}^m$.

3 Main results

Throughout this section (X, d) is a generalized metric space in Perov's sense and w is a generalized w -distance on the generalized metric space.

Let $x_0 \in X$ and $r := (r_i)_{i=1}^n$ for each $i = \{1, 2, \dots, m\}$. Let us define:

$\tilde{B}_w(x_0; r) := \{x \in X | \tilde{w}(x_0, x) < r\}$ the open ball centered at x_0 with radius r with respect to \tilde{w} ;

$\tilde{B}_w(x_0; r) := \{x \in X | \tilde{w}(x_0, x) \leq r\}$ the closed ball centered at x_0 with radius r with respect to \tilde{w} ;

$\tilde{B}_w^d(x_0; r)$ - the closure in (X, d) of the set $\tilde{B}_w(x_0; r)$.

Theorem 3.2 *Let (X, d) be a complete generalized metric space, $x_0 \in X$, $r := (r_i)_{i=1}^n$ for each $i = \{1, 2, \dots, m\}$, $\tilde{w} : X \times X \rightarrow [0, \infty)$ a generalized w -distance on X and let $T : \tilde{B}_w(x_0; r) \rightarrow P(X)$ be a multivalued operator with the property that there exists $A = (a_{i,j})_{i,j \in \{1,2,\dots,m\}} \in \mathcal{M}_{m,m}(\mathbb{R}_+)$ a matrix convergent to zero and $B = (b_{i,j})_{i,j \in \{1,2,\dots,m\}} \in \mathcal{M}_{m,m}(\mathbb{R}_+) \setminus \{U\}$ such that, for every $x, y \in X$ and each $u \in T(x)$, there exists $v \in T(y)$ such that*

$$\tilde{w}(u, v) \leq A\tilde{w}(x, y) + BD_{\tilde{w}}(y, T(y)),$$

where $D_{\tilde{w}}(x, T(x)) := \inf\{\tilde{w}(x, y) : y \in T(x)\}$.

(This means, that for each $x, y \in Y$ and each $u \in T(x)$, there exists $v \in T(y)$ such that

$$\begin{pmatrix} w_1(u, v) \\ \dots \\ w_m(u, v) \end{pmatrix} \leq \begin{pmatrix} a_{11} & \dots & a_{1,m} \\ \dots & & \\ a_{m1} & \dots & a_{m,m} \end{pmatrix} \cdot \begin{pmatrix} w_1(x, y) \\ \dots \\ w_m(x, y) \end{pmatrix} + \begin{pmatrix} b_{11} & \dots & b_{1,m} \\ \dots & & \\ b_{m1} & \dots & b_{m,m} \end{pmatrix} \cdot \begin{pmatrix} D_{w_1}(x, y) \\ \dots \\ D_{w_m}(x, y) \end{pmatrix}$$

Suppose that:

1. $\inf\{\tilde{w}(x, y) + D_{\tilde{w}}(x, T(x))\} > 0$, for every $x, y \in X$ and $y \notin T(x)$.
2. There exists $x_1 \in T(x_0)$ such that $\tilde{w}(x_0, x_1)(I - A)^{-1} \leq r$.
3. If $u \in \mathbb{R}_+^m$ is such that $u(I - A)^{-1} \leq (I - A)^{-1}r$, then $u \leq r$.

Then $\text{Fix}T \neq \emptyset$.

Proof. Let $x_0 \in X$ and $x_1 \in T(x_0)$ such that

$$\tilde{w}(x_0, x_1)(I - A)^{-1} \leq r \leq (I - A)^{-1} \cdot r$$

Then, by (2), $x_1 \in \tilde{B}_w(x_0; r)$. For $x_1 \in T(x_0)$ there exists $x_2 \in T(x_1)$ such that

$$\begin{aligned}\tilde{w}(x_1, x_2) &\leq A\tilde{w}(x_0, x_1) + BD_{\tilde{w}}(x_1, T(x_1)) \\ &\leq A\tilde{w}(x_0, x_1) + B\tilde{w}(x_1, x_2)\end{aligned}$$

$$\text{Thus } \tilde{w}(x_1, x_2) \leq \frac{A}{U-B}\tilde{w}(x_0, x_1)$$

We denote $C := \frac{A}{U-B}$ and we observe that the matrix $C \in \mathcal{M}_{m,m}(\mathbb{R})$ is a matrix convergent to zero and satisfy the following inequalities

- $I + \frac{A}{U-B} \leq I + A + A^2 + \cdots + A^n + \cdots$, therefore $I + C \leq (I - A)^{-1}$
- $(I - C)^{-1} \leq (I - A)^{-1}$

Thus $\tilde{w}(x_1, x_2)(I - A)^{-1} \leq \frac{A}{U-B}w(x_0, x_1)(I - A)^{-1} \leq Cr$. Notice that $x_2 \in \tilde{B}_w(x_0; r)$.

Indeed, since $\tilde{w}(x_0, x_2) \leq \tilde{w}(x_0, x_1) + \tilde{w}(x_1, x_2)$ we get that $w(x_0, x_2)(I - A)^{-1} \leq \tilde{w}(x_0, x_1)(I - A)^{-1} + \tilde{w}(x_1, x_2)(I - A)^{-1} \leq Ir + Cr \leq (I - A)^{-1}r$, which immediately implies (by hypothesis (2)) that $\tilde{w}(x_0, x_2) \leq r$.

By induction, we construct the sequence $(x_n)_{n \in \mathbb{N}}$ in $\tilde{B}_w(x_0; r)$ having the properties:

- (a) $x_{n+1} \in T(x_n), n \in \mathbb{N}$;
- (b) $\tilde{w}(x_0, x_n)(I - A)^{-1} \leq (I - A)^{-1}r$, for each $n \in \mathbb{N}^*$, that means $\tilde{w}(x_0, x_n) \leq r$;
- (c) $\tilde{w}(x_n, x_{n+1})(I - A)^{-1} \leq C^n r$, for each $n \in \mathbb{N}$.

By (c), for every $m, n \in \mathbb{N}$, with $m > n$, we get that

$$\tilde{w}(x_n, x_m)(I - A)^{-1} \leq C^m(I - C)^{-1}r \leq C^n(I - A)^{-1}r.$$

By Lemma 2.2(3) we have that the sequence $(x_n)_{n \in \mathbb{N}}$ is Cauchy in the complete metric space. Denote by x^* its limit in $\tilde{B}_w^d(x_0; r)$.

Assume that $x^* \notin T(x^*)$. Fix $n \in \mathbb{N}$. Since $(x_m)_{m \in \mathbb{N}}$ is a sequence in $\tilde{B}_w(x_0; s)$ which converge to x^* and $\tilde{w}(x_n, \cdot)$ is lower semicontinuous we have

$$\tilde{w}(x_n, x^*) \leq \liminf_{m \rightarrow \infty} \tilde{w}(x_n, x_m) \leq C^n r, \text{ for every } n \in \mathbb{N}.$$

Therefore, by hypothesis (1) and using above inequality we have

$$\begin{aligned} 0 &\leq \inf\{\tilde{w}(x, x^*) + D_{\tilde{w}}(x, T(x)) : x \in X\} \\ &\leq \inf\{\tilde{w}(x_n, x^*) + \tilde{w}(x_n, x_{n+1}) : n \in \mathbb{N}\} \\ &\leq \inf\{2C^m r\} = 0. \end{aligned}$$

Which is a contradiction. Thus conclude that $x^* \in T(x^*)$.

A global version of the previous theorem is the following result.

Theorem 3.3 *Let (X, d) be a complete generalized metric space, $x_0 \in X$, $r := (r_i)_{i=1}^n$ for each $i = \{1, 2, \dots, m\}$, $\tilde{w} : X \times X \rightarrow [0, \infty)$ a generalized w -distance on X and let $T : X \rightarrow P(X)$ be a multivalued operator with the property that there exists $A \in \mathcal{M}_{m,m}(\mathbb{R}_+)$ a matrix convergent to zero and $B \in \mathcal{M}_{m,m}(\mathbb{R}_+) \setminus \{U\}$ such that, for every $x, y \in X$ and each $u \in T(x)$, there exists $v \in T(y)$ such that*

$$\tilde{w}(u, v) \leq A\tilde{w}(x, y) + BD_{\tilde{w}}(y, T(y)),$$

where $D_{\tilde{w}}(x, T(x)) := \inf\{\tilde{w}(x, y) : y \in T(x)\}$.

Suppose that $\inf\{\tilde{w}(x, y) + D_{\tilde{w}}(x, T(x))\} > 0$, for every $x, y \in X$ and $y \notin T(y)$ then

1. $FixT \neq \emptyset$.
2. The sequence $(x_n)_{n \in \mathbb{N}} \in X$ given by relation $x_{n+1} \in T(x_n)$, for all $n \in \mathbb{N}$, is convergent and its limit is a fixed point of T .
3. One has the estimation $\tilde{w}(x_n, x^*) \leq C^n \tilde{w}(x_0, x_1)$ where $C \in \mathcal{M}_{m,m}(\mathbb{R})$, $C := \frac{A}{U-B}$, and $x^* \in FixT$.

Remark 3.1 *In the condition of the previous theorem we observe that T is a MWP operator.*

4 Data dependence theorem for weakly contractive type operators in generalized metric spaces

The main result of this section is the following data dependence theorem with respect to the Theorem 3.3.

Theorem 4.4 *Let (X, d) be a complete generalized metric space, $x_0 \in X$, $r := (r_i)_{i=1}^n$ for each $i = \{1, 2, \dots, m\}$, $\tilde{w} : X \times X \rightarrow [0, \infty)$ a generalized w -distance on X and let $T_1, T_2 : X \rightarrow P(X)$ be a multivalued operator with the property that there exists $A \in \mathcal{M}_{m,m}(\mathbb{R}_+)$ a matrix convergent to zero and $B \in \mathcal{M}_{m,m}(\mathbb{R}_+) \setminus \{U\}$ such that, for every $x, y \in X$ and each $u \in T_j(x)$, for every $j \in \{1, 2\}$, there exists $v \in T_j(y)$ such that*

$$\tilde{w}(u, v) \leq A\tilde{w}(x, y) + BD_{\tilde{w}}(y, T_j(y)),$$

where $D_{\tilde{w}}(x, T_j(x)) := \inf\{\tilde{w}(x, y) : y \in T_j(x)\}$.

Suppose that the following are true:

1. $FixT_1 \neq \emptyset \neq FixT_2$.
2. We suppose that there exists $\eta := (\eta_i)_{i=1}^n$, for each $i = \{1, 2, \dots, m\}$, with $\eta > 0$, such that for every $u \in T_1(x)$ there exists $v \in T_2(x)$ such that $\tilde{w}(u, v) \leq \eta$, (respectively for every $v \in T_2(x)$ there exists $u \in T_1(x)$ such that $\tilde{w}(v, u) \leq \eta$).
3. $\inf\{\tilde{w}(x, y) + D_{\tilde{w}}(x, T_j(x))\} > 0$ for each $j \in \{1, 2\}$, for every $x, y \in X$ and $y \notin T_j(y)$.

Then for every $u^* \in FixT_1$ there exists $v^* \in FixT_2$ such that

$$\tilde{w}(u^*, v^*) \leq U(1 - C)^{-1}\eta, \text{ where } C \in \mathcal{M}_{m,m}(\mathbb{R}), C := \frac{A}{U-B};$$

(respectively for every $v^* \in FixT_2$ there exists $u^* \in FixT_1$ such that

$$\tilde{w}(v^*, u^*) \leq U(1 - C)^{-1}\eta, \text{ where } C \in \mathcal{M}_{m,m}(\mathbb{R}), C := \frac{A}{U-B}.$$

Proof. Let $u_0 \in \text{Fix}T_1$, then $u_0 \in T_1(u_0)$. Using the hypothesis (2) we have that there exists $u_1 \in T_2(u_0)$ such that $\tilde{w}(u_0, u_1) \leq \eta$.

We have that for every $u_0, u_1 \in X$ with $u_1 \in T_2(u_0)$ there exists $u_2 \in T_2(u_1)$ such that $\tilde{w}(u_1, u_2) \leq A\tilde{w}(u_0, u_1) + BD_{\tilde{w}}(u_1, T_2(u_1)) \leq A\tilde{w}(u_0, u_1) + B\tilde{w}(u_1, u_2)$.

Thus $\tilde{w}(u_1, u_2) \leq \frac{A}{U-B}\tilde{w}(u_0, u_1)$

We denote $C := \frac{A}{U-B}$ and we observe that the matrix $C \in \mathcal{M}_{m,m}(\mathbb{R})$ is a matrix convergent to zero.

Thus $\tilde{w}(u_1, u_2) \leq C\tilde{w}(u_0, u_1)$.

By induction we obtain a sequence $(u_n)_{n \in \mathbb{N}} \in X$ such that

(1) $u_{n+1} \in T_2(u_n)$, for every $n \in \mathbb{N}$;

(2) $\tilde{w}(u_n, u_{n+1}) \leq C^n \tilde{w}(u_0, u_1)$

For $n, p \in \mathbb{N}$ we have the inequality

$$\tilde{w}(u_n, u_{n+p}) \leq C^n (I - C)^{-1} \tilde{w}(u_0, u_1).$$

By the Lemma 2.2(3) we have that the sequence $(u_n)_{n \in \mathbb{N}}$ is a Cauchy sequence. Since (X, d) is a complete metric space we have that there exists $v^* \in X$ such that $u_n \xrightarrow{d} v^*$.

Assume that $v^* \notin T_2(v^*)$. Fix $n \in \mathbb{N}$. By the lower semicontinuity of $\tilde{w}(x, \cdot) : X \rightarrow [0, \infty)$ we have

$$\tilde{w}(u_n, v^*) \leq \liminf_{p \rightarrow \infty} \tilde{w}(u_n, u_{n+p}) \leq C^n (I - C)^{-1} \tilde{w}(u_0, u_1) \quad (4.1)$$

Therefore, by hypothesis (3) and using the relation 4.1 we have the inequality:

$$\begin{aligned} 0 &< \inf\{\tilde{w}(u, v^*) + D_{\tilde{w}}(u, T_2(u)) : u \in X\} \\ &\leq \inf\{\tilde{w}(u_n, v^*) + \tilde{w}(u_n, u_{n+1}) : n \in \mathbb{N}\} \\ &\leq \inf\{C^n (I - C)^{-1} \tilde{w}(u_0, u_1) + C^n \tilde{w}(u_0, u_1) : n \in \mathbb{N}\} = 0. \end{aligned}$$

Which is a contradiction. Thus we conclude that $v^* \in T_2(v^*)$.

Then, by $\tilde{w}(u_n, v^*) \leq C^n (I - C)^{-1} \tilde{w}(u_0, u_1)$, with $n \in \mathbb{N}$, for $n = 0$ we obtain $\tilde{w}(u_0, v^*) \leq U(I - C)^{-1} \tilde{w}(u_0, u_1) \leq U(I - C)^{-1} \eta$, which complete the proof. \square

References

- [1] J. S. Bae, *Fixed point theorems for weakly contractive multivalued maps*, Journal of Mathematical Analysis and Applications, vol. 284, no. 2, pp. 690697, 2003.
- [2] J. Caristi, *Fixed point theorems for mappings satisfying inwardness conditions*, Transactions of the American Mathematical Society, vol. 215, pp. 241251, 1976.
- [3] A. Granas, J. Dugundji, *Fixed Point Theory*, Berlin, Springer-Verlag, 2003.
- [4] I. Eklund, *Nonconvex minimization problems*, Bulletin of the American Mathematical Society, vol. 1, no. 3, pp. 443474, 1979.
- [5] A.-D. Filip, A. Petruşel, *Fixed Point Theorems on Spaces Endowed with Vector-Valued Metrics*, Fixed Point Theory and Applications, Volume 2010, Article ID 281381, 15 pages doi:10.1155/2010/281381.
- [6] Y. Feng, S. Liu, *Fixed point theorems for multivalued contractive mappings and multivalued Caristi type mappings*, J. Math. Anal. Appl. 317(2006), 103-112.
- [7] L.Guran, *Existence and data dependence for multivalued weakly contractive operators*, Studia Univ. Babeş-Bolyai, Mathematica, 54(2009), no. 3, 67-76.
- [8] L. Guran, *A multivalued Perov-type theorems in generalized metric spaces*, Creative Math. and Inf., 17 (2008), no. 3, 412-419;
- [9] O. Kada, T. Suzuki, W. Takahashi, *Nonconvex minimization theorems and fixed point theorems in complete metric spaces*, Math. Japonica 44(1996) 381-391.
- [10] A. Latif, *A fixed point result in complete metric spaces*, JP Journal of Fixed Point Theory and Applications, vol. 2, no. 2, pp. 169175, 2007

- [11] A. Latif, A.W. Albar , *Fixed point results for multivalued maps*, Int. J. Contemp. Math. Sci. 2, 21-24(2007), 1129-1136.
- [12] A. Latif, A. A. N. Abdou *Fixed Point Results for Generalized Contractive Multimaps in Metric Spaces*, Fixed Point Theory and Applications Volume 2009 (2009), Article ID 432130, 16 pages doi:10.1155/2009/432130.
- [13] S. B. Nadler Jr., *Multivalued contraction mappings*, Pacific Journal of Mathematics, vol. 30, pp. 475 488, 1969.
- [14] T. Lazăr, A. Petruşel, N. Shahzad, *Fixed points for non-self operators and domain invariance theorems*, Nonlinear Analysis: Theory, Methods and Applications, 70(2009), 117-125.
- [15] A.I. Perov, *On Cauchy problem for a system of ordinary differential equations*, (in Russian), Priblizhen. Metody Reshen. Differ. Uravn., 2(1964), 115-134.
- [16] A. I. Perov, A. V. Kibenko, *On a certain general method for investigation of boundary value problems* (Russian), Izv. Akad. Nauk SSSR Ser. Mat., 30(1966), 249-264.
- [17] A. Petruşel, *Multivalued weakly Picard operators and applications*, Scientiae Mathematicae Japonicae, 1(2004), 1-34.
- [18] I. A. Rus, *Generalized Contractions and Applications*, Presa Clujeană Universitară, Cluj-Napoca, 2001.
- [19] A. Petruşel, I. A. Rus, A. Sântămărian, *Data dependence of the fixed point set of multivalued weakly Picard operators*, Nonlinear Analysis, 52(2003), no. 8, 1947-1959.
- [20] T. Suzuki, W. Takahashi, *Fixed points theorems and characterizations of metric completeness*, Topological Methods in Nonlinear Analysis, Journal of Juliusz Schauder Center, 8(1996), 371-382.