

MPRA

Munich Personal RePEc Archive

Common priors for generalized type spaces

Pintér, Miklós

Corvinus University of Budapest

2011

Online at <http://mpra.ub.uni-muenchen.de/34118/>
MPRA Paper No. 34118, posted 14. October 2011 / 20:08

Common priors for generalized type spaces*

Miklós Pintér[†]

October 14, 2011

”Give me a place to stand on,
and I will move the Earth.”

Archimedes

Abstract

The notion of common prior is well-understood and widely-used in the incomplete information games literature. For ordinary type spaces the common prior is defined.

Pintér and Udvari (2011) introduce the notion of generalized type space. Generalized type spaces are models for various bounded rationality issues, for finite belief hierarchies, unawareness among others. In this paper we define the notion of common prior for generalized types spaces.

Our results are as follows: the generalization (1) suggests a new form of common prior for ordinary type spaces, (2) shows some quantum game theoretic results (Brandenburger and La Mura, 2011) in new light.

Keywords and phrases: Type spaces, Generalized type spaces, Common prior, Harsányi Doctrine, Quantum games

JEL codes: C72, D83

1 Introduction

The common prior is a central notion of games with incomplete information since the very beginnings (Harsányi, 1967-68). The intuition behind the common prior is clear, it aggregates all information of the model, it works as

*Miklós Pintér acknowledges the support by the János Bolyai Research Scholarship of the Hungarian Academy of Sciences and grant OTKA 72856.

[†]Corresponding author: Department of Mathematics, Corvinus University of Budapest, 1093 Hungary, Budapest, Fővám tér 13-15., miklos.pinter@uni-corvinus.hu.

a source of all information, that is, e.g. the players' beliefs can be deduced from it.

For ordinary type spaces (Heifetz and Samet, 1998) the common prior is known, it grabs the same intuitions as that for finite models it gives. In type spaces the players' beliefs are explicitly given by the so called type functions, so a common prior is a probability measure which is compatible with the players' beliefs. This compatibility can be well-defined by the notion of conditional probability.

Pintér and Udvari (2011) introduce the notion of generalized type space. Generalized type spaces are models for ordinary belief hierarchies and for various bounded rationality issues, e.g. for finite hierarchies of beliefs, unawareness among others. In other words, generalized type spaces are generalizations of ordinary type spaces (Heifetz and Samet, 1998), that is, every ordinary type space is a generalized type space.

In this paper we define the common prior for generalized type spaces. The intuition is the same as for ordinary type spaces, that is, a certain compatibility with the players' beliefs is required. Since in generalized type spaces the players' beliefs can be very different from those in ordinary type spaces we define the common prior as a common type function, a mapping which assign a probability measure to each state of the world, in a way such that, at each state of the world the players' beliefs (given by the type functions) are compatible with the given probability measure. However, in generalized type spaces the players might have beliefs about very few events only, so the word compatibility means less in this case than in the case of ordinary type spaces.

In the setting of generalized type spaces compatibility means at each state of the world each player cannot learn more or less from the common prior than that she learns from her own type function. In generalized type spaces, however, it can happen that each player knows nothing, in which case any mapping is a common prior. Therefore, the common priors can be very different for the very same generalized type space.

Technically, while in ordinary type spaces with common prior the type functions are conditional probabilities, this is not the case for generalized type spaces. In the generalized type space setting the type functions are less than conditional probabilities, so they can vary more.

Furthermore, in this paper we work with purely measurable generalized type spaces, so with a generalization of Heifetz and Samet (1998)'s type space. Even if most of the works in the literature use topological type spaces, see Böge and Eisele (1979); Mertens and Zamir (1985); Heifetz (1993); Brandenburger and Dekel (1993); Mertens et al (1994); Heifetz and Samet (1999); Pintér (2005) among others, the results of Heifetz and Samet (1998); Pintér

(2008, 2010) tell us to prefer the purely measurable to the topological framework.

We mention two results. First, in this paper we define the common prior as a common type function. This approach fits well to the type space setting, and makes possible for us to discuss the problem in an elegant form. However, this implies also that we have to reformulate the common prior for ordinary type spaces. This reformulation does not bring new issues for ordinary type spaces, so we can say our notion of common prior is the same as the one well-known in the literature (for ordinary type spaces).

Second, our notion of common prior can interpret the quantum correlation "device" in Brandenburger and La Mura (2011)'s paper. In other words, we can interpret the quantum correlation as an ordinary correlation for boundedly rational players, that is, as a common prior for generalized type spaces. In this sense, our result shows quantum game theoretic results in new light.

Finally, we give an explanation for the citation from Archimedes in the head of the paper. One of the main difference between ordinary and generalized type spaces is that the ordinary type spaces are Harsányi type spaces, while the the generalized type spaces are not necessarily Harsányi type spaces, that is, in ordinary type spaces the players know their own types, while it needs not to happen in generalized type spaces. That a player knows her own type is "the place to stand on". If this "place" is missing, then the players can do less, but it makes possible for external factors (common prior) to coordinate the players' beliefs more subtle ways than in models where the players know their own types. Definitely, Archimedes did not mean the same that we.

The setup of the paper is as follows. In the next section we introduce the notion of common prior for generalized type spaces in a finite, simple setting. In Section 3 we explain the introduced notion by examples. Section 4 is about applications of the previously discussed models. The last section briefly concludes. In an appendix we give the notion of common prior for the general setting.

2 Common priors for generalized type spaces

In this section we introduce the notion of common prior for generalized type spaces in the finite setting. We give the mathematically precise definitions in Appendix A.

In our model we assume that the player set N is a finite set, that is, there are finitely many players, and $N_0 = N \cup \{0\}$ is for the extended player set,

which contains the nature (Player 0) as an extra player.

In generalized type spaces the players can have very different beliefs. For instance it can happen that at a certain state of the world a player has a belief which can be represented by a probability distribution defined on all subsets of the set of the states of the world, while at an other state of the world the player has a belief which can be represented by a probability distribution defined only on some subsets of the set of the states of the world.

First, we introduce the notion of states of the world.

Definition 1. *Let the finite set Ω be the state of the states of the world and for each $i \in N_0$, let \mathcal{M}_i be a partition on Ω . Partition \mathcal{M}_i represents Player i 's information, \mathcal{M}_0 is for the information available for the nature. Let $\mathcal{M} = \vee_{i \in N_0} \mathcal{M}_i$, that is \mathcal{M} is the coarsest partition among the partitions which contain partitions \mathcal{M}_i .*

The above notion is from Aumann (1999). The states of the world are the basic elements of the world, each point of Ω is for one specific state of the world. The partitions are for modeling the information of the players. Partition \mathcal{M}_i gives which states Player i can distinguish (by observation) and which she cannot. Partition \mathcal{M}_0 is for the information the model provides, but does not belong to any player. We say this is the nature's information.

Usual in the literature that the parameters of the modeled situation, to which we also refer as the types of the nature, are given by a parameter set (Mertens and Zamir, 1985; Brandenburger and Dekel, 1993; Heifetz, 1993; Pintér, 2005). We also follow this way, and take set S as a parameter set. Moreover, we assume that S is a finite set, and each subset of S is an event.

We have already mentioned above that in generalized type spaces the players' beliefs can be truncated, that is, those need not be defined at any event; we denote the class of this type of beliefs by $\Delta(\Omega, \mathfrak{M})$. Formally, $\Delta(\Omega, \mathfrak{M})$ is for all probability measures (distributions) defined on certain subsets of Ω , $\Delta(\Omega, \mathfrak{M}) = \{\mu \text{ is a probability measure defined on the field generated by partition } \mathcal{N} : \mathcal{N} \text{ is coarser than } \mathcal{M}\}$. If at a certain state of the world a player's belief is a probability measure such that is not defined at event $A \subseteq \Omega$, then we say that the player is not aware about event A .

Definition 2. *Let $(\Omega, \mathcal{M}_i)_{i \in N_0}$ be the set of the states of the world (see Definition 2). The generalized type space based on parameter space S is a tuple $(S, \Omega, \{\mathcal{M}_i\}_{i \in N_0}, g, \{f_i\}_{i \in N})$, where*

- $g : \Omega \rightarrow S$ is such that for each $A \subseteq S$ there exists $M_1, M_2, \dots, M_n \in \mathcal{M}_0 \cup \{\emptyset\}$ such that $g^{-1}(A) = \cup_{j=1}^n M_j$,

- $f_i : \Omega \rightarrow \Delta(\Omega, \mathfrak{M})$ is such that for each $A \subseteq \Delta(\Omega, \mathfrak{M})$ there exists $M_1, M_2, \dots, M_n \in \mathcal{M}_i \cup \{\emptyset\}$ such that $f_i^{-1}(A) = \cup_{j=1}^n M_j$, $i \in N$.

Mapping g connects the parameter set to the set of the states of the world. Moreover, g preserves the information given by partition \mathcal{M}_0 , that is, the g -inverse image of any event in the parameter set is expressible by the elements of partition \mathcal{M}_0 . In other words, if two states of the world are indistinguishable by partition \mathcal{M}_0 , then these points have the same g -image.

f_i is the type function of Player i . The type function gives the player's beliefs at the states of the world. f_i also preserves Player i 's information, that is, if the two states of the world are indistinguishable by Player i , then those have the same f_i -image.

In the generalized type spaces the beliefs can vary much, the players need not be aware of all events, they might know their types or not, therefore, many things can happen. The model of generalized type spaces encompasses both Harsányi and Non-Harsányi type spaces (Heifetz and Mongin, 2001) and more.

In general it can happen that at a state of the world the players' beliefs are inconsistent, that is, the players' beliefs are very different. This can happen basically because of two things. First, when the players have different piece of information about the state of the world, so the differences in their beliefs are due to quantity of information available for the players, in this case there is no substantial disagreement about the state of the world among the players.

Second, when the differences of the players' beliefs are due to substantial disagreement, in other words, when the information available for the player differ in its quality not only in its quantity. Typically this happens when the players can agree to disagree on a commonly known event (Aumann, 1976). In the first case we say the generalized type space has (a) common prior(s), in the second we say it has not.

Definition 3 (Common prior for generalized type spaces). *Consider generalized type space $(S, \{\Omega, \mathcal{M}_i\}_{i \in N_0}, g, \{f_i\}_{i \in N})$. $f : \Omega \rightarrow \Delta(\Omega, \mathcal{M})$ is a Common Prior, if for each $\omega \in \Omega$, $i \in N$, $A \subseteq \Omega$ such that $f_i(\omega)$ is defined at event A :*

$$f(\omega)(A) = \sum_{\omega \in \Omega} f_i(\omega)(A) f(\omega) .$$

The notion of Common Prior formalizes the intuition we have discussed above. If there is a Common Prior for a generalized type space, then there is no substantial differences in the players' beliefs, that is, those are consistent.

Remark 4. We do not discuss it in details, only mention it, that Aumann (1976)'s result holds for generalized type spaces too. Formally, for any generalized type space having Common Prior, if at a state of the world event

A is commonly known, then it cannot be commonly known that the players' beliefs about event A at the state of the world do not coincide; that is, the players cannot agree to disagree.

It is also worth mentioning that the type functions f_i are not (necessarily) conditional probabilities. If type functions f_i were conditional probabilities, then those should meet the following condition, for each $\omega \in \Omega$, $i \in N$, $A, B \subseteq \Omega$ such that $f_i(\omega)$ is defined at event A , and $B \in \mathcal{M}_i$:

$$f(\omega)(A \cap B) = \sum_{\omega \in B} f_i(\omega)(A) f(\omega) . \quad (1)$$

The difference is obvious, if $A \cap B = \emptyset$, then the conditional probability must be 0 ($f(\omega)(B) > 0$, $f_i(\omega')(A)$, $\omega' \in B$ in (1)), while this is not the case in (3). In other words, since the generalized type spaces are not (necessarily) Harsányi type spaces, the type functions are not (necessarily) conditional probabilities.

3 Examples

In this section we provide two examples for illustrating the notions introduced in the previous section. The first example is about an ordinary Harsányi type space.

Example 5. Take generalized type space $(S, \{\Omega, \mathcal{M}_i\}_{i \in N_0}, g, \{f_i\}_{i \in N})$, where

- $N = \{1, 2\}$,
- $S = \{*\}$,
- $T_i = \{t_i^1, t_i^2\}$, $i \in N$,
- $\Omega = S \times T_1 \times T_2$,
- $\mathcal{M}_0 = \{\Omega\}$, $\mathcal{M}_i = \{\{*\} \times \{t_i^1\} \times \{T_{-i}\}, \{*\} \times \{t_i^2\} \times \{T_{-i}\}\}$, $i \in N$,
- for each Player $i \in N$ and state of the world $\omega \in \Omega$: $f_i(\omega)$ is defined at each subset of Ω , and $f_i(\omega)(\{*\} \times \{t_i\} \times \{t_{-i}^1\}) = f_i(\omega)(\{*\} \times \{t_i\} \times \{t_{-i}^2\}) = \frac{1}{2}$.

This type space is a Harsányi type space, that is, each player knows her own type, more precisely, each player believes with probability 1 her own type, and at each state of the world, each player's belief is defined on \mathcal{M} , that is, each player is aware, so can form beliefs, about any event.

Take state of the world $\omega = (*, t_1, t_2) \in \Omega$. If f is a common prior, then $f(\omega)(\{*\} \times \{t_1\} \times \{t_2\}) = f(\omega)(\{*\} \times \{t_1\} \times T_2 \setminus \{t_2\}) = f(\omega)(\{*\} \times T_1 \setminus \{t_1\} \times \{t_2\}) = f(\omega)(\{*\} \times T_1 \setminus \{t_1\} \times T_2 \setminus \{t_2\})$. So the unique common prior has the same value at each state of the world, see Table 1 (we do not put the only state of the nature in the model, because that does not count).

	(t_1^1, t_2^1)	(t_1^1, t_2^2)	(t_1^2, t_2^1)	(t_1^2, t_2^2)
(t_1^1, t_2^1)	1/4	1/4	1/4	1/4
(t_1^1, t_2^2)	1/4	1/4	1/4	1/4
(t_1^2, t_2^1)	1/4	1/4	1/4	1/4
(t_1^2, t_2^2)	1/4	1/4	1/4	1/4

(2)

Table 1: The common prior for the Harsányi type space of Example 5

Therefore, in this example we have got a "classical" common prior, that is, we can use a unique distribution, see Table 2.

$T_1 \setminus T_2$	t_2^1	t_2^2
t_1^1	1/4	1/4
t_1^2	1/4	1/4

Table 2: The "classical" common prior for the Harsányi type space of Example 5

In the above example we have seen that in certain cases the common prior has the same value at each state of the world, put it differently, we can say that the common prior is a probability distribution (measure) on the set of the states of the world. In the following proposition we formalize this observation, but first, we introduce a notion.

Definition 6. Consider generalized type space $(S, \{\Omega, \mathcal{M}_i\}_{i \in N_0}, g, \{f_i\}_{i \in N})$ such that it has a common prior f . For each state of the world $\omega \in \Omega$ let undirected graph (V_ω, E_ω) be defined as $V_\omega = \{B \in \cup_{i \in N} \mathcal{M}_i : f(\omega)(B) > 0\}$, and $\overline{BB'} \in E_\omega$, if there exist $A, A' \in \cup_{i \in N_0} \mathcal{M}_i$ and $B'' \in \cup_{i \in N_0} \mathcal{M}_i$ such that $A \subseteq B \cap B''$, $A' \subseteq B' \cap B''$, and $f(\omega)(A), f(\omega)(A') > 0$.

Now, we can provide the above mentioned proposition.

Proposition 7. Take generalized type space $(S, \{\Omega, \mathcal{M}_i\}_{i \in N_0}, g, \{f_i\}_{i \in N})$ such that

1. it has a common prior f ,

2. $f_i(\omega)$ is defined at each event $A \in \bigvee_{i \in N_0} \mathcal{M}_i$, $\omega \in \Omega$, $i \in N$,

3. for each $B \subseteq \mathcal{M}_i$, $\omega \in B$: $f_i(\omega)(B) = 1$, $i \in N$.

Then for all states of the world $\omega, \omega' \in \Omega$ such that there exists $B^* \in \bigwedge_{i \in N} \mathcal{M}_i$ such that

4. $\omega, \omega' \in B^*$,

5. both $(V_\omega \cap B^*, E_\omega)$ and $(V_{\omega'} \cap B^*, E_{\omega'})$ are connected undirected graphs,

$f(\omega) = f(\omega')$.

Proof. From Points 1., 2., 3. and 4. for each Player $i \in N$: $f_i(\omega)(B^*) = 1$.

Let $A \in \bigvee_{i \in N_0} \mathcal{M}_i$ be such that $A \subseteq B^*$ and $f(\omega)(A) > 0$. Then for each $A' \in \bigvee_{i \in N_0} \mathcal{M}_i$ such that there exists Player $i \in N$, $B_i \in \mathcal{M}_i$ such that $A, A' \subseteq B_i$:

$$f(\omega)(A) = \sum_{\omega \in \Omega} f_i(\omega)(A) f(\omega) = \sum_{\omega \in B_i} f_i(\omega)(A) f(\omega) = f_i(\omega_{B_i})(A) f(\omega)(B_i) ,$$

similarly,

$$f(\omega)(A') = \sum_{\omega \in \Omega} f_i(\omega)(A') f(\omega) = \sum_{\omega \in B_i} f_i(\omega)(A') f(\omega) = f_i(\omega_{B_i})(A') f(\omega)(B_i) ,$$

where $\omega_{B_i} \in B_i$.

Since $f(\omega)(A) > 0$, $f(\omega)(B_i) > 0$ and $f_i(\cdot)(A) > 0$:

$$\frac{f(\omega)(A')}{f(\omega)(A)} = \frac{f_i(\cdot)(A')}{f_i(\cdot)(A)} ,$$

that is, $\frac{f(\omega)(A')}{f(\omega)(A)}$ is given by the generalized type space (by f_i).

Since $(V_\omega \cap B^*, E_\omega)$ is connected (Point 5.), for each $A' \in \bigvee_{i \in N_0} \mathcal{M}_i$ $A' \neq A$, $\frac{f(\omega)(A')}{f(\omega)(A)}$ is given (if $A \not\subseteq B$, then $\frac{f(\omega)(A')}{f(\omega)(A)} = 0$). Furthermore, we have the equations: $f(\omega)(A') = f_i(\omega_{B_i})(A) f(\omega)(B_i)$, as many as $|\bigvee_{i \in N_0} \mathcal{M}_i| - 1$ (the number of atoms minus 1), and equation $\sum_{A' \in \bigvee_{i \in N_0} \mathcal{M}_i} f(\omega)(A') = 1$; these equations are (linearly) independent; and we have as many as $|\bigvee_{i \in N_0} \mathcal{M}_i|$ unknowns (the probabilities $f(\omega)(A')$ and $f(\omega)(A)$), so $f(\omega)$ is well-defined, that is, it is uniquely determined by the generalized type space (by the type functions).

Similarly, the above argument can be applied to $f(\omega')$, so $f(\omega')$ is well-defined too, therefore $f(\omega) = f(\omega')$. \square

In the second example the common prior is neither “classical“ nor unique.

Example 8. Take generalized type space $(S, \{\Omega, \mathcal{M}_i\}_{i \in N_0}, g, \{f_i\}_{i \in N})$, where

- $N = \{1, 2\}$,
- $S = \{*\}$,
- $T_i = \{t_i^1, t_i^2\}$, $i \in N$,
- $\Omega = S \times T_1 \times T_2$,
- $\mathcal{M}_0 = \{\Omega\}$, $\mathcal{M}_i = \{\{*\} \times \{t_i^1\} \times \{T_{-i}\}, \{*\} \times \{t_i^2\} \times \{T_{-i}\}\}$, $i \in N$,
- for each Player $i \in N$ and state of the world $\omega \in \Omega$: $f_i(\omega)$ is defined only on \mathcal{M}_i , and $f_i(\omega)(\{*\} \times \{t_i^1\} \times \{T_{-i}\}) = f_i(t_i)(\{*\} \times \{t_i^2\} \times \{T_{-i}\}) = \frac{1}{2}$, $i \in N$.

Then it is a slight calculation to see that mapping given in Table 1 is a common prior for this type space. However, so is the mapping given in Table 3 (this numerical example is from Abramsky and Brandenburger (2011)).

	(t_1^1, t_2^1)	(t_1^1, t_2^2)	(t_1^2, t_2^1)	(t_1^2, t_2^2)
(t_1^1, t_2^1)	3/8	3/8	1/8	1/8
(t_1^1, t_2^2)	1/8	1/8	3/8	3/8
(t_1^2, t_2^1)	3/8	1/8	3/8	1/8
(t_1^2, t_2^2)	1/8	3/8	1/8	3/8

(3)

It is worth noticing that the two common priors provided in Example 8 are very different. The first one (Table 1) can be interpreted as there is no relation among the states of the world, however, the second (Table 3) shows a very subtle and smart relation among the states of the world.

4 Applications

By generalized type spaces we can model finite belief hierarchies, unawareness, and other various phenomena of bounded cognitive abilities (Pintér and Udvari, 2011). In the “classical” models the common prior is a tool of coordination. By common priors the players, more precisely, the players’ beliefs, can be harmonized, so that the outcome of the game can Pareto outperform the outcomes achievable without harmonization (Aumann, 1974, 1987).

For generalized type spaces the common prior can harmonize the players’ beliefs in smarter ways than that classical common priors do. This smarter

harmonization allows outcomes to Pareto outperform those achieved by classical harmonizations.

Perhaps, the key point in general type spaces is that the players need not know their own types. From the viewpoint of mathematics, it is not difficult to formalize and handle this. However, it could be difficult to give a good decision theoretic interpretation for this type of cognitive incapability of the players.

One interpretation is Brandenburger and La Mura (2011)'s, who consider games in extensive form. If a game does not meet the property of perfect recall, then there might be decision points indistinguishable by the player, that is, the decision points are in the same information set, and the decision points are different only in the player's previous moves. This setting can be modeled as the player does not know her previous moves, that is, she does not know her own strategy, which is equivalent with that she knows her strategy, but does not her type. In other words, generalized type space can be good frameworks for this kind of problems too.

Usually, the description of incomplete information models starts with that the players learn their own types. In many cases, however, this is not realistic. For instance, many people evaluate their own physical and mental strength wrong. Many times we hear sentences like "I had thought I was strong enough, but I was not". Considering these uncertainties, and putting them into a model, we get generalized type spaces.

Filar and Beck (2007)'s model is another approach. In their model the players' incompetence makes that the players' control over their actions is not 100%, that is, even if a player intends to take a certain action, she plays a different, maybe mixed, action. Filar and Beck (2007)'s approach is similar to the trembling-hand models, but not the same. In their model the players know their incapability of playing the exact action, and they decide by using this knowledge (an other difference is that in Filar and Beck (2007) the hand can tremble in strange and not uniform ways too).

Therefore, putting Filar and Beck (2007)'s idea in the incomplete information setting, the players know their own types, but not their own strategies. However, this is equivalent with that, the players know their own strategies but not their own types. That is, we get generalized type spaces again.

5 Conclusion

In this paper we have introduced the notion of common prior for generalized type spaces. We have showed that a generalized type space can be compatible with substantially different common priors.

Moreover, for generalized type spaces the common priors can harmonize the players' beliefs on very different levels. In Example 8 one of the two common priors gives no harmonization at all (see Table 1), while the other (see Table 3) provides a very smart harmonization of the players' beliefs. This phenomenon rises substantial decision theoretical questions.

Example 8 clearly shows that the common prior can be more than the "sum" of the players' information (Pintér (2011) concludes the same on a different basis). Therefore, in certain cases generalized type spaces are not complete descriptions of decision theoretic situations. This, again, shows that it is not enough to take account of the players' beliefs (generalized type spaces), the relation of their beliefs should be also considered (common prior).

A Common prior – the abstract model

In this section we give the mathematically precise definitions of the notions used in the paper. We do not give the interpretations for the notions we discuss here, the reader can find those in the main text and in Pintér and Udvari (2011).

Notations: Let N be the set of the players, w.l.o.g. we can assume that $0 \notin N$, and let $N_0 = N \cup \{0\}$, where 0 is for the nature as a player.

For any set system $\mathcal{A} \subseteq \mathcal{P}(X)$: $\sigma(\mathcal{A})$ is the coarsest σ -field which contains \mathcal{A} .

First we introduce the notion of generalized type space. We generalize ordinary type spaces, and use terminologies, notions similar to those of Heifetz and Samet (1998).

Definition 9. *Let X be a space, \mathfrak{M} be a class of σ -fields on set X and $\Delta(X, \mathfrak{M})$ be the class of probability measures on the σ -fields of \mathfrak{M} , formally $\Delta(X, \mathfrak{M}) = \{\mu \in \Delta(X, \mathcal{M}) : \mathcal{M} \in \mathfrak{M}\}$. Then the σ -field \mathcal{A}^* on $\Delta(X, \mathfrak{M})$ is defined as follows:*

$$\mathcal{A}^* = \sigma(\{\{\mu \in \Delta(X, \mathfrak{M}) : \mu(A) \geq p\}, A \in \mathcal{M}, p \in [0, 1]\}) .$$

In other words, \mathcal{A}^ is the smallest σ -field among the σ -fields which contain the sets $\{\mu \in \Delta(X, \mathfrak{M}) : \mu(A) \geq p\}$, where $\mathcal{M} \in \mathfrak{M}$, $A \in \mathcal{M}$ and $p \in [0, 1]$ are arbitrarily chosen.*

Notice that \mathcal{A}^* is not a fixed σ -field, we mean, it depends on the measurable spaces on which the probability measures are defined.

Assumption 10. *Let the parameter space (S, \mathcal{A}) be an arbitrary measurable space.*

Henceforth we assume that (S, \mathcal{A}) is a fixed parameter space which contains all states of the nature. We can think of S as a set which encompasses all the not commonly known parameters of the considered situation.

Definition 11. *Let Ω be the space of the states of the world and for each $i \in N_0$: let \mathcal{M}_i be a σ -field on Ω . The σ -field \mathcal{M}_i represents Player i 's information, \mathcal{M}_0 is for the information available for the nature, hence it is the representative of \mathcal{A} , the σ -field of the parameter space S . Let $\mathcal{M} = \sigma(\cup_{i \in N_0} \mathcal{M}_i)$, the smallest σ -field which contains σ -fields \mathcal{M}_i .*

For the sake of brevity, henceforth – if it does not make confusion – we do not indicate the σ -fields. E.g. instead of (S, \mathcal{A}) we write S , or $\Delta(S)$ instead of $(\Delta(S, \mathcal{A}), \mathcal{A}^*)$. However, in some cases we refer to the non-written σ -field: e.g. $A \in \Delta(X, \mathcal{M})$ is a set of \mathcal{A}^* , that is, it is a measurable set in the measurable space $(\Delta(X, \mathcal{M}), \mathcal{A}^*)$, but $A \subseteq \Delta(X, \mathcal{M})$ keeps its original meaning: A is a subset of $\Delta(X, \mathcal{M})$.

Definition 12. *Let $(\Omega, \{\mathcal{M}_i\}_{i \in N_0})$ be a space of the states of the world (see Definition 11). The generalized type space based on the parameter space S is a tuple $(S, \Omega, \{\mathcal{M}_i\}_{i \in N_0}, g, \{f_i\}_{i \in N})$, where*

1. $g : \Omega \rightarrow S$ is \mathcal{M}_0 -measurable,
2. $f_i : \Omega \rightarrow \Delta(\Omega, \mathfrak{M})$ is \mathcal{M}_i -measurable, $i \in N$,

where $\mathfrak{M} = \{\mathcal{N} \text{ is a } \sigma\text{-field on } \Omega : \mathcal{N} \subseteq \mathcal{M}\}$.

The generalized type spaces are not Harsányi type spaces (Heifetz and Mongin, 2001), that is, the players do not necessarily know their own types, more precisely, they do not necessarily believe with probability 1 their own types.

Definition 13 (Common prior for generalized type spaces). *Consider generalized type space $(S, \{(\Omega, \mathcal{M}_i)\}_{i \in N_0}, g, \{f_i\}_{i \in N})$. $f : \Omega \rightarrow \Delta(\Omega, \mathcal{M})$ is a common prior, if for each $\omega \in \Omega$, $i \in N$, $A \in \mathcal{M}$ such that $f_i(\omega)$ is defined at event A :*

$$f(\omega)(A) = \int_{\Omega} f_i(\cdot)(A) \, df(\omega) .$$

The common prior is a "universal" type function, a type function which summarizes the beliefs of all the players at each state of the world. In other words, by a common prior the differences in the player's beliefs are due to the quantity of information and not to the quality of the information available to the players.

References

- Abramsky S, Brandenburger A (2011) The sheaf-theoretic structure of non-locality and contextuality, manuscript
- Aumann RJ (1974) Subjectivity and correlation in randomized strategies. *Journal of Mathematical Economics* 1:67–96
- Aumann RJ (1976) Agreeing to disagree. *The Annals of Statistics* 4(6):1236–1239
- Aumann RJ (1987) Correlated equilibrium as an expression of bayesian rationality. *Econometrica* 55(1):1–18
- Aumann RJ (1999) Interactive epistemology i: Knowledge. *International Journal of Game Theory* 28:263–300
- Böge W, Eisele T (1979) On solutions of bayesian games. *International Journal of Game Theory* 8(4):193–215
- Brandenburger A, Dekel E (1993) Hierarchies of beliefs and common knowledge. *Journal of Economic Theory* 59:189–198
- Brandenburger A, La Mura P (2011) Quantum decision theory, manuscript
- Filar J, Beck J (2007) Games, Incompetence and Training, *Annals of the International Society of Dynamic Games*, vol 9, Springer, pp 93–110
- Harsányi J (1967-68) Games with incomplete information played by bayesian players part i., ii., iii. *Management Science* 14:159–182, 320–334, 486–502
- Heifetz A (1993) The bayesian formulation of incomplete information - the non-compact case. *International Journal of Game Theory* 21:329–338
- Heifetz A, Mongin P (2001) Probability logic for type spaces. *Games and Economic Behavior* 35(1-2):31–53
- Heifetz A, Samet D (1998) Topology-free typology of beliefs. *Journal of Economic Theory* 82:324–341
- Heifetz A, Samet D (1999) Coherent beliefs are not always types. *Journal of Mathematical Economics* 32:475–488
- Mertens JF, Zamir S (1985) Formulation of bayesian analysis for games with incomplete information. *International Journal of Game Theory* 14:1–29

- Mertens JF, Sorin S, Zamir S (1994) Repeated games part a. CORE Discussion Paper No 9420
- Pintér M (2005) Type space on a purely measurable parameter space. *Economic Theory* 26:129–139
- Pintér M (2008) Every hierarchy of beliefs is a type. CoRR abs/0805.4007, URL <http://arxiv.org/abs/0805.4007>
- Pintér M (2010) The non-existence of a universal topological type space. *Journal of Mathematical Economics* 46:223–229
- Pintér M (2011) A note on the common prior, manuscript
- Pintér M, Udvari Z (2011) Generalized type spaces, manuscript