

CBM

CBM  
R

7626  
1989  
401


UNIVERSITY

UNIVERSITEIT  
BRABANT

POSTBOX 90153  
5000 LE TILBURG  
THE NETHERLANDS



DEPARTMENT OF ECONOMICS  
RESEARCH MEMORANDUM



REGULAR TWO-GRAPHS AND EXTENSIONS OF  
PARTIAL GEOMETRIES

W. R. HAEMERLE

ABSTRACT

1965

1965

REGULAR TWO-GRAPHS AND EXTENSIONS OF PARTIAL GEOMETRIES

by

W.H. Haemers

Tilburg University,  
Tilburg, The Netherlands.

ABSTRACT.

We study so called two-graph geometries. These are geometries that carry a regular two-graph, but also constitute one-point extensions of partial geometries. First we develop some theory, then we go through lists of known regular two-graphs and partial geometries in order to find examples. Some are found, including one that extends the partial geometry with parameters  $s=4$ ,  $t=17$ ,  $\alpha=2$ .

## 1. INTRODUCTION.

Extensions of  $t$ -designs, especially one-point extensions, have been studied a lot in the past. More recently, people became interested in extensions of finite geometries, such as generalised quadrangles (there exist various papers on this subject; Cameron [6] gives a survey) or, more generally, partial geometries (see Hobart & Hughes [17]). The points of a partial geometry carry a strongly regular graph, whilst regular two-graphs are, in a certain sense, extensions of strongly regular graphs by one point. For this reason it seemed worthwhile to investigate a combination of these objects, being one-point extensions of partial geometries with the structure of a regular two-graph on the points. We call such structures two-graph geometries. The present paper is a first attempt to study these geometries.

The reader is assumed to be familiar with the theory of designs and strongly regular graphs (see for instance Cameron & Van Lint [7]). We shall briefly survey the relevant results on two-graphs and partial geometries. A *two-graph* consists of a finite set  $\Omega$  together with a set  $\Delta$  of triples (called *coherent* triples) from  $\Omega$ , such that every 4-subset of  $\Omega$  contains an even number of coherent triples. Let  $\nabla$  denote the set of non-coherent triples. Then  $(\Omega, \nabla)$  is also a two-graph, called the *complement* of  $(\Omega, \Delta)$ . The two-graph  $(\Omega, \Delta)$  is *empty [complete]* if  $\Delta [\nabla]$  is empty;  $(\Omega, \Delta)$  is *regular* if every pair of points from  $\Omega$  is contained in a constant number  $a$  of coherent triples. For any point  $\omega \in \Omega$  of a regular two-graph  $(\Omega, \Delta)$ , the matrix  $A_\omega$ , defined by

$$(A_\omega)_{\beta\gamma} = \begin{cases} -1 & \text{if } \{\beta, \gamma, \omega\} \in \Delta, \text{ or } \omega \in \{\beta, \gamma\}, \\ 0 & \text{if } \beta = \gamma, \\ 1 & \text{otherwise,} \end{cases}$$

has just two eigenvalues  $\rho_1$  and  $\rho_2$  ( $\rho_1 > \rho_2$ ). These eigenvalues have opposite sign and are odd integers if  $\rho_1 \neq -\rho_2$ , furthermore

$$|\Omega| = 1 - \rho_1 \rho_2, \quad a = -(\rho_1 + 1)(\rho_2 + 1)/2.$$

The *derived graph*  $\Gamma_\omega$  of  $(\Omega, \Delta)$  with respect to  $\omega$  has vertex set  $\Omega \setminus \{\omega\}$ , two vertices  $\beta$  and  $\gamma$  being adjacent if  $\{\beta, \gamma, \omega\} \in \Delta$ . So by deleting row and column  $\omega$  from  $A_\omega$ , we obtain the  $(-1, 1, 0)$  adjacency matrix of  $\Gamma_\omega$ . For any  $\omega$ , the derived graph of a regular two-graph (not complete or empty) is strongly regular with parameters  $(V, K, \lambda, M)$ , where

$$V = |\Omega| - 1 = -\rho_1 \rho_2, \quad K = \alpha = -(\rho_1 + 1)(\rho_2 + 1)/2,$$

$$(1) \quad \lambda = 1 - (\rho_1 + 3)(\rho_2 + 3)/4, \quad M = K/2 = -(\rho_1 + 1)(\rho_2 + 1)/4.$$

Conversely, to a strongly regular graph with parameters  $(V, K, \lambda, K/2)$  there corresponds a regular two-graph. A clique (or coherent set) of  $(\Omega, \Delta)$  is a subset  $c$  of  $\Omega$ , such that every triple from  $c$  is coherent. Clearly, for  $\omega \in c$ ,  $c \setminus \{\omega\}$  is a clique (in the normal sense) in  $\Gamma_\omega$ . A clique  $c$  of  $(\Omega, \Delta)$  satisfies:

$$(2) \quad |c| \leq 1 - \rho_2.$$

Two-graphs have been introduced by G. Higman, and were studied mainly by Seidel and Taylor [24] [25] [28].

A design is denoted by the pair  $(\Phi, B)$ , where  $\Phi$  is the set of points and  $B$  is the set of blocks. An *anti-flag* of a design  $(\Phi, B)$  is a pair  $(\varphi, b)$  with  $\varphi \in \Phi$ ,  $b \in B$  and  $\varphi \notin b$ .

A *partial geometry*  $pg(s, t, \alpha)$  is a  $1-(V, s+1, t+1)$  design  $(\Phi, B)$ , where any two distinct lines (= blocks) meet in at most one point, such that for every anti-flag  $(\varphi, b)$  there are precisely  $\alpha$  points on  $b$  collinear with  $\varphi$ . It follows that  $V = |\Phi| = (s+1)(st+\alpha)/\alpha$ ,  $|B| = (t+1)(st+\alpha)/\alpha$ . If we interchange the roles of points and lines we obtain the *dual* partial geometry  $pg(t, s, \alpha)$ . The *point graph* of a  $pg(s, t, \alpha)$  has vertex set  $\Phi$ ; two vertices are adjacent if they are collinear. The point graph of a  $pg(s, t, \alpha)$  is strongly regular with parameters  $(V, K, \lambda, M) = (V, s(t+1), s-1+t(\alpha-1), \alpha(t+1))$ . A *one-point extension* of  $pg(s, t, \alpha)$  is a design for which the derived design with respect to any point is a  $pg(s, t, \alpha)$ .



Partial geometries were introduced by Bose [1] and have been studied a lot. Some general references are: Brouwer & Van Lint [2] and De Clerck [3].

## 2. TWO-GRAPH GEOMETRIES.

1. DEFINITION. A *two-graph geometry* is a  $2-(v,k,\lambda)$  design  $(\Omega, C)$  satisfying the following properties:

- i. two distinct blocks of  $C$  have at most two points in common (therefore blocks are called circles),
- ii. any set of four points contains an even number of cocircular triples,
- iii.  $v = 1 + (k-1)(2\lambda-1)$ .

2. PROPOSITION. Let  $\Delta$  be the set of cocircular triples of a two-graph geometry  $(\Omega, C)$ . Then  $(\Omega, \Delta)$  is a regular two-graph with eigenvalues

$$\rho_1 = 2\lambda-1, \rho_2 = 1-k.$$

Proof. By i and ii  $(\Omega, \Delta)$  is a two graph. Since  $(\Omega, C)$  is a  $2-(v,k,\lambda)$  design,  $(\Omega, \Delta)$  is regular with  $\alpha = -(\rho_1+1)(\rho_2+1)/2 = \lambda(k-2)$ . Using  $v = |\Omega| = 1+(k-1)(2\lambda-1) = 1-\rho_1\rho_2$  and  $\rho_1 > 0 > \rho_2$  the values of  $\rho_1$  and  $\rho_2$  follow.

Note that we did not use property iii to prove that  $(\Omega, \Delta)$  is a regular two-graph, it is only used to compute  $\rho_1$  and  $\rho_2$ . In fact, once  $(\Omega, \Delta)$  is defined, property iii can be replaced by:

iii'. the circles of  $C$  are maximal cliques of  $(\Omega, \Delta)$ .

Herein maximal means that the bound  $-\rho_2+1$  given in (2) is met. As usual, the number of circles is denoted by  $b$  and the number of circles through a fixed point by  $r$ . Then  $\lambda(v-1) = r(k-1)$  and  $bk = vr$  yield

$$r = \lambda(2\lambda-1) = \rho_1(\rho_1+1)/2,$$

$$b = \lambda(2\lambda-1)^2 - 2\lambda(2\lambda-1)(\lambda-1)/k = \rho_1^2(\rho_1+1)/2 + \rho_1(\rho_1^2-1)/2(\rho_2-1).$$

We call a regular two-graph *geometric* if it corresponds to a two-graph geometry. Clearly, for a two-graph to be geometric the following divisibility condition must be satisfied:

$$(3) \quad 2(-\rho_2 + 1) \mid \rho_1(\rho_1^2 - 1).$$

The following result is straight forward.

3. PROPOSITION. A regular two-graph with eigenvalues  $\rho_1$  and  $\rho_2$  is geometric if and only if there exists a set  $C$  of cliques of size  $(1-\rho_2)$ , such that every coherent triple is covered by a unique clique of  $C$ .

A regular two-graph with  $\rho_2 = -1$  is empty (no triple is coherent). Also for the next case,  $\rho_2 = -3$ , two-graph geometries are nothing special, because of the following result.

4. PROPOSITION. Let  $C$  be the set of all 4-cliques of a regular two-graph  $(\Omega, \Delta)$  with  $\rho_2 = -3$ . Then  $(\Omega, C)$  is the unique two-graph geometry corresponding to  $(\Omega, \Delta)$ .

Proof. In a regular two-graph with eigenvalues  $\rho_1$  and  $\rho_2$  each coherent triple is contained in exactly  $\wedge = 1 - (\rho_1 + 3)(\rho_2 + 3)/4$  cliques of size 4, by use of (1). This number  $\wedge$  equals 1 if  $\rho_2 = -3$ , hence Proposition 3. gives the result.

Seidel's [22] determination of all regular two-graphs with  $\rho_2 = -3$  leads to

5. COROLLARY. Two-graph geometries with  $\rho_2 = -3$  (i.e.  $k = 4$ ) exist if and only if  $\rho_1 = 1, 3, 5$  or  $9$  (i.e.  $\lambda = 1, 2, 3$  or  $5$ ) and are unique

Note that two-graph geometries with  $\rho_1 = 1$  are degenerate: there is just one circle of size  $k = v$ , and the two-graphs are complete.

Let  $(\omega, c)$  be an anti-flag of a design  $(\Omega, C)$ . The *anti-flag graph*  $\Gamma_{\omega, c}$  has vertex set  $c$ , two vertices  $\beta$  and  $\gamma$  ( $\beta \neq \gamma$ ) are adjacent whenever  $\omega, \beta$  and  $\gamma$  are covered by a block of  $C$ .



6. PROPOSITION. A  $2-(v,k,\lambda)$  design  $(\Omega, C)$  with block intersection sizes at most 2, is a two-graph geometry if and only if each anti-flag graph is the disjoint union of two complete graphs of size  $k/2$ .

Proof. Suppose  $(\Omega, C)$  is a two-graph geometry. Let  $(\omega, c)$  be an anti-flag of  $(\Omega, C)$  and let  $\beta, \gamma$  and  $\delta$  be three distinct points of  $c$ . Since  $\{\beta, \gamma, \delta, \omega\}$  contains an even number of cocircular triples, the subgraph of  $\Gamma_{\omega, c}$  induced by  $\beta, \gamma$  and  $\delta$  is either a triangle or has just one edge. Thus  $\Gamma_{\omega, c}$  is the complete graph or the disjoint union of two complete graphs. Conversely, it is easily seen that any 4-set contains 0, 2 or 4 cocircular triples if each anti-flag graph is the disjoint union of two complete graphs.

Next fix  $c \in C$ . For  $\omega \in \Omega \setminus c$ , let  $m_\omega$  denote the size of a component of  $\Gamma_{\omega, c}$ . Counting in two ways the total number of triples  $(\omega, \beta, \gamma)$  with  $\omega \in \Omega \setminus c$ , and  $\beta, \gamma$  adjacent vertices in  $\Gamma_{\omega, c}$  gives

$$\sum_{\omega \notin c} (m_\omega(m_\omega - 1) + (k - m_\omega)(k - m_\omega - 1)) = k(k-1)(\lambda-1)(k-2)$$

The left hand side is at least  $\lambda = (v-k)k(\frac{1}{2}k-1)$  with equality if and only if  $m_\omega = k/2$  for all  $\omega \notin c$ . This proves the result, because  $\lambda$  equals the right hand side, precisely when  $v = 1 + (k-1)(2\lambda-1)$ .

By definition, a 2-design is a one-point extension of a partial geometry  $pg(s, t, \alpha)$  if and only if any two distinct blocks meet in at most 2 points and each anti-flag graph is regular of degree  $\alpha$ . Therefore we have:

7. THEOREM. A two-graph geometry  $(\Omega, C)$  with eigenvalues  $\rho_1$  and  $\rho_2$  is a one point extension of a partial geometry with parameters

$$s = -\rho_2 - 1, \quad t = (\rho_1 - 1)/2, \quad \alpha = (-\rho_2 - 1)/2.$$

So, only partial geometries with  $s = 2\alpha$  occur. Clearly, the point graph of the partial geometry with respect to  $\omega \in \Omega$  (say) is the derived graph  $\Gamma_\omega$  of  $(\Omega, \Delta)$ . Such strongly regular graphs satisfy  $K = 2\lambda$  ( $(V, K, \lambda, M)$  is the set of parameters), which is equivalent to  $s = 2\alpha$ .

Clearly, the anti-flag graph of a one-point extension of a  $pg(s,t,1)$  (i.e. a generalised quadrangle) consists of disjoint edges. So by Proposition 6 we have:

8. PROPOSITION. A one-point extension of a  $pg(2,t,1)$  is a two graph geometry

This result needs not to be true for  $\alpha > 1$ . For instance, there exists one-point extensions of  $pg(4,1,2)$  for which some anti-flag graphs are hexagons. So they are not two-graph geometries. But no other exceptions are known (to us). The two-graph geometries corresponding to the above proposition have  $\rho_2 = -3$ . So by Corollary 5 we have the following result due to Beukenhout [3] (see also Thas [29]).

9. COROLLARY. One-point extensions of  $pg(2,t,1)$  exist and are unique.

### 3. REGULAR TWO-GRAPHS AND PARTIAL GEOMETRIES.

Next we investigate known or feasible regular two-graphs and partial geometries with  $s = 2\alpha$ . We more or less follow the surveys of Seidel [24] and De Clerck [10]. Since the point graph of a partial geometry with  $s = 2\alpha$  is strongly regular graph with  $K = 2M$ , the regular two-graph exists if the partial geometry exists. The converse, however, is not true.

Case 1:  $\rho_1 = -\rho_2 - 2$ , or equivalently,  $t = \alpha - 1$ .

The regular two-graph corresponds to a regular symmetric Hadamard matrix with constant diagonal. The corresponding partial geometries are duals of block designs with  $\lambda = 1$ . In such a partial geometry any two lines meet. This implies that two circles of the two-graph geometry can only have no or two points in common. The two-graph geometry is therefore a quasi-symmetric block design. The divisibility condition (3) leads to  $\rho_1 = 1, 3$  or  $9$ . The case  $\rho_1 = 1, \rho_2 = -3$  is treated in Corollary 5. For the other two cases the parameters  $(\rho_1, \rho_2, v, b, k, r, \lambda, s, t, \alpha)$  are  $(3, -5, 16, 16, 6, 6, 2, 4, 1, 2)$  and  $(9, -11, 100, 375, 12, 45, 5, 10, 4, 5)$ . The first one is a  $2-(16, 6, 2)$  design. There exist precisely three such designs, but only one satisfies condition

2.1.ii, viz. the unique 2-(16,6,2) design with characteristic 3, see Cameron [5]. Nothing is known about the second case. Mavron & Shrikhande [21] also found the mentioned possibilities in their classification of quasi-symmetric block designs with block intersection sizes 0 and 2 and an additional requirement, a little weaker than condition 2.1.ii.

Case 2:  $\rho_1 = -\rho_2$ , or equivalently,  $t = \alpha$ .

The regular two-graphs are the ones associated to conference matrices with integral eigenvalues. The partial geometries  $pg(s, s/2, s/2)$  are dual nets; they corresponding to  $(s-2)/2$  mutually orthogonal latin squares of order  $s+1$ . The parameters are:

$$v = \rho_1^2 + 1, \quad b = \rho_1(\rho_1^2 + 1)/2, \quad k = \rho_1 + 1, \quad r = \rho_1(\rho_1 + 1)/2,$$

$$\lambda = (\rho_1 + 1)/2, \quad s = \rho_1 - 1, \quad t = \alpha = (\rho_1 - 1)/2.$$

It was observed by Fisher [14] that such two-graph geometries can be constructed from the inversive plane over the field with  $\rho_1$  elements. So they exist whenever  $\rho_1$  is an odd prime power. The required set of circles is just one orbit of the group generated by the inversions acting on the blocks (circles) of the inversive plane. Wilbrink [32] proved that the corresponding two-graphs are the Paley two-graphs. The derived partial geometry is the corresponding half of the affine plane derived from the original inversive plane. The lines of this partial geometry are the only maximal cliques in the Paley graph (the Paley graph is the derived graph of the Paley two-graph with respect to any point), see Blokhuis [0]. This implies that the Paley two-graph is geometric in a unique way; the circles are just all maximal cliques.

The above cases together with Corollary 5 cover all two-graph designs whose corresponding partial geometry is improper (i.e.  $\alpha = 1$ ,  $t$  or  $t+1$ ). For the remaining cases  $\alpha < t$  holds, which implies  $\rho_1 > -\rho_2$ .

Case 3:  $\rho_1 = -\rho_2 + 2$ , or equivalently,  $t = \alpha + 1$ .



The regular two-graphs are the complements of the ones considered in Case 1. Partial geometries with  $t = \alpha + 1$  are classified by De Clerck [9]. The parameters for this case are:

$$v = (\rho_1 - 1)^2, \quad b = \rho_1(\rho_1^2 - 1)/2, \quad k = \rho_1 - 1, \quad r = \rho_1(\rho_1 + 1)/2,$$

$$\lambda = (\rho_1 + 1)/2, \quad s = \rho_1 - 3, \quad t = (\rho_1 - 1)/2, \quad \alpha = (\rho_1 - 3)/2.$$

A  $\text{pg}(2\alpha, \alpha + 1, \alpha)$  can be constructed from a projective plane of order  $2\alpha + 2$  possessing a hyperoval. Such planes do exist if the order is a power of 2. If  $\alpha = 1$  the two-graph geometry exists (Corollary 5). If  $\alpha = 2$  or 4 the partial geometry does not exist (by De Clerck [9] and Lam et al [19] respectively). The regular two-graph is known for many more values of  $\rho_1 = 2\alpha + 3$  than the corresponding partial geometry, including for  $\alpha = 2$  and  $\alpha = 4$ . The symplectic two-graphs, for instance, belong to this case. By De Clerck, Gevaert & Thas [12] they are not geometric if  $\rho_1 = 9$  or 17. In fact, for no other value of  $\alpha$  existence of such a two-graph geometry is settled. The smallest candidate has parameters  $\rho_1 = 9$ ,  $\rho_2 = -7$ ,  $v = 64$ ,  $b = 360$ ,  $k = 8$ ,  $r = 45$ ,  $\lambda = 5$ ,  $s = 6$ ,  $t = 4$ ,  $\alpha = 3$ . By Mathon [20], there are just two such partial geometries. Storme [27] and Tonchev [31] proved by computer that one of the two (the one corresponding to the hyperoval in the plane of order 8) cannot be extended to a two-graph geometry.

Case 4:  $\rho_1 = 2^m - 1$ ,  $\rho_2 = -2^{m-1} - 1$ , or equivalently  $\alpha = 2^{m-2}$ ,  $t = 2^{m-1} - 1$ .

By De Clerck, Dye & Thas [11] partial geometries and (hence) regular two-graphs with these parameters are known if  $m$  is even. The two-graphs exist for all  $m > 1$ , they are the complements of the orthogonal two-graphs  $\Omega^+(2m, 2)$ , see Seidel [23]. By condition (3) only  $\rho_1 = 3, 7$  and 15 are possible. The first possibility exists (Corollary 5), the second one doesn't (see Case 3) and the remaining one has parameters:

$$\rho_2 = -9, \quad v = 136, \quad k = 10, \quad b = 1632, \quad r = 120, \quad \lambda = 8, \quad s = 8, \quad t = 7, \quad \alpha = 4.$$

Tonchev [31] showed by computer that the known  $\text{pg}(8, 7, 4)$  (see Cohen [8] and Haemers & Van Lint [16] for other ways to construct this partial geometry) does not extend to a two-graph geometry. Since  $\text{pg}(8, 7, 4)$  is conjectured to be unique, the two-graph geometry probably does not exist.

Case 5.  $\rho_1 = 2^m + 1$ ,  $\rho_2 = -2^{m-1} + 1$ , or equivalently,  $t = 2^{m-1}$ ,  $\alpha = 2^{m-2} - 1$ .

For the other parameters we find

$$v = 2^{2m-1} - 2^{m-1}, \quad b = (2^{2m-1})(2^m + 1), \quad k = 2^{m-1},$$

$$r = (2^m + 1)(2^{m-1} + 1), \quad \lambda = 2^{m-1} + 1, \quad s = 2^{m-1} - 2.$$

The corresponding regular two-graphs are the orthogonal two-graphs  $\Omega^-(2m, 2)$ , see Seidel [23]. Corollary 5 takes care of  $m = 3$ . For  $m > 3$  existence of the partial geometry is still open. De Clerck & Tonchev [13] showed that for  $m = 4$  the corresponding  $pg(6, 8, 3)$  can only have automorphisms of order 2 and 3, leaving not much hope for finding the two-graph geometry.

Case 6:  $\rho_1 = \rho_2^2$ , or equivalently,  $t = 2\alpha(\alpha + 1)$ .

Regular two-graphs with these eigenvalues were constructed by Taylor [28] whenever  $\sqrt{\rho_1}$  is an odd prime power. The remaining parameters are:

$$v = \rho_1 \sqrt{\rho_1} + 1, \quad b = \rho_1(\rho_1 + 1)(\rho_1 - \sqrt{\rho_1} + 1)/2, \quad k = \sqrt{\rho_1} + 1,$$

$$r = \rho_1(\rho_1 + 1)/2, \quad \lambda = (\rho_1 + 1)/2, \quad s = 2\alpha = \sqrt{\rho_1} - 1, \quad t = (\rho_1 - 1)/2.$$

Again the first one exists ( $\rho_1 = 9, \rho_2 = -3$ ) by Corollary 5. For  $\rho_2 = -5$  and  $\rho_2 = -7$  Spence [26] proved that the derived strongly regular graph is not geometric. For  $-\rho_2 > 7$  nothing is known. Also the Ree groups provide regular two-graphs with these eigenvalues whenever  $-\rho_2$  is an odd power of 3. We have no idea whether these two-graphs can be geometric.

Case 7.  $\rho_2 = -5$ , or equivalently,  $\alpha = 2$ .

Then  $k = 6, s = 4, \alpha = 2$ . For  $\rho_1, 15, 19, 35$  and  $55$  are the only possible values that have not been considered before. If  $\rho_1 = 15, 19$  the regular two-graph nor the partial geometry is known to exist. For  $\rho_1 = 35$  a two-graph geometry is realised in the next section. For  $\rho_1 = 55$  there is a unique regular two-graph and a unique derived strongly regular graph. Nevertheless, existence of the partial geometry and the two-graph geometry is yet unsolved.

#### 4. AN EXCEPTIONAL TWO-GRAPH GEOMETRY.

In this section we construct a two graph geometry  $(\Omega, C)$  with parameters:  $\rho_1 = 35$ ,  $\rho_2 = -5$ ,  $v = 176$ ,  $b = 18480$ ,  $k = 6$ ,  $r = 630$ ,  $\lambda = 18$ ,  $s = 4$ ,  $t = 17$ ,  $\alpha = 2$ . The regular two graph  $(\Omega, \Delta)$  is the one having the Higman-Sims group HS acting on  $\Omega$  as a 2-transitive automorphism group, see Taylor [28]. The partial geometry (with respect to any point) is the one constructed by the author [15]. The group of the two-graph geometry will be the Mathieu group  $M_{22}$ , which is a subgroup of HS and the corresponding action on  $\Omega$  is rank 3. For the construction we need some properties of this action.

10. LEMMA The action of  $M_{22}$  on  $(\Omega, \Delta)$  satisfies:

- i. There exists an orbit  $C$  of size 18480 on the 6-cliques of  $(\Omega, \Delta)$ .
- ii. Every triple from  $\Delta$  is contained in a 6-clique of  $C$ .

Proof. Fix a point  $\omega \in \Omega$ . The subgroup of  $M_{22}$  stabilizing  $\omega$  is  $A_7$ . It is an automorphism group of  $\Gamma_\omega$  (the full automorphism group of  $\Gamma_\omega$  is  $\text{PEU}(5,2)$ ). We can define  $\Gamma_\omega$  on the edges of the Hoffman-Singleton graph (for short HoSi), where two edges are adjacent whenever they are disjoint and possess an interconnecting edge (see Hubaut [18]). The group of automorphisms of HoSi, that fixes (setwise) a distinguished 15-coclique is  $A_7$ . Its action on the edges is the action on  $\Gamma_\omega$ , just mentioned. This description is worked out in some detail in [15], in order to construct  $\text{pg}(4,17,2)$ . Using this description, the following facts are straightforward:

- The 5-cliques of  $\Gamma_\omega$  are one-factors in Petersen subgraphs of HoSi.
- The group  $A_7$  has two orbits on the Petersen subgraphs of HoSi. The sizes are 105 (the 'special Petersen graphs' in [15]) and 420, respectively.
- The subgroup of  $A_7$  that stabilizes any Petersen subgraph  $P$  of HoSi acts transitively on the one-factors of  $P$ .

So  $A_7$  has two orbits on the 5-cliques of  $\Gamma_\omega$ ; one of size 630 (the lines of  $\text{pg}(4,17,2)$ ), and one of size 2520. Any such 5-clique is, together with  $\omega$ , a 6-clique of  $(\Omega, \Delta)$ . Thus (remember that  $M_{22}$  acts transitively on  $\Omega$ ) there are  $176 \times 3150 / 6 = 92400$  6-cliques, and  $M_{22}$  acts either transitively on  $\Omega$ , or has two orbits; one of size 18480 and one of size 73920, respectively. The first option, however, cannot occur, since 92400 doesn't divide the order



of  $M_{22}$  (= 443520). This proves i. Next let  $\{\omega, \beta, \gamma\} \in \Delta$ . Then  $\{\beta, \gamma\}$  is an edge of  $\Gamma_\omega$ , which is contained in a (unique) 5-clique of the smaller orbit (i.e. a line of  $\text{pg}(4,17,2)$ ). Hence  $\{\omega, \beta, \gamma\}$  is contained in a 6-clique of  $C$ , proving ii.

11. THEOREM. With  $C$  as in Lemma 10,  $(\Omega, C)$  is a two-graph geometry.

Proof. By i of the lemma, the cliques of  $C$  cover at most  $18480 \cdot 20 = 369600$  coherent triples. This, however, is precisely the total number of coherent triples. Therefore by ii every triple is covered exactly ones by a clique of  $C$ , so  $(\Omega, C)$  is a two-graph geometry by Proposition 3.

It is clear that the parameters of this two-graph geometry are the ones mentioned above. There must be several ways to prove Lemma 10. For instance, another proof could go along the lines of the construction by Calderbank & Wales [4] of  $\text{pg}(4,17,2)$ ; instead of HoSi, they start from the Steiner system  $S(5,8,24)$ . An approach that does not use one of these two construction methods of  $\text{pg}(4,17,2)$ , would give a new way to describe this partial geometry.

It has been checked (using a computer) that there is a unique way to extend the  $\text{pg}(4,17,2)$  to a two-graph geometry such that all automorphisms  $(A_7)$  of the partial geometry are preserved.

## 5. CONCLUDING REMARKS.

Apart from the existence questions mentioned in Section 3, there are several other problems that look interesting. We mention a few.

We don't know examples of non-isomorphic two-graph geometries with the same parameters. For  $\rho_2 = -3$  they are unique (Corollary 5). Is it premature to conjecture that two-graph geometries are necessarily unique, but it seems save to do so for the sporadic one of the previous section, because of the remark at the end (it is even conceivable that the regular two-graph and the partial geometry are unique).

The relation between two-graphs and Seidel switching leads to a class of  $(-1,1,0)$  incidence matrices of a two graph geometry in the following way. Consider the incidence matrix  $N$  of  $(Q,C)$ . Let  $\Gamma$  be a (strong) graph in the switching class of  $(Q,\Delta)$ . Each circle of  $C$  corresponds to a disjoint union of two complete graphs in  $\Gamma$ . Sign the non-zero entries of each column of  $N$  with + and - according to the partition of the corresponding circle, just described. The matrix obtained in this manner has some interesting properties; for instance, its rank equals the multiplicity of  $\rho_2$ . It is not clear how this can be explored.

If for a two-graph geometry,  $2(-\rho_2+1)$  divides  $\rho_1^2-1$  (compare with (3)), then it is feasible that a subset of the circles forms a partial geometry  $pg(-\rho_2, (\rho_1-1)/2, (-\rho_2+1)/2)$  ( $= pg(s+1, t, \alpha+1)$ ). For Fisher's two-graph geometries (Section 3, Case 2) it would mean that a subset of the circles is the dual of a  $2-((\rho_1^2+1)/2, (\rho_1+1)/2, 1)$  design. If  $\rho_1 = 3$ , this is possible, however, Thas [30] proved that it is impossible for  $\rho_1 > 3$ . An affirmative answer for the sporadic example of Section 4 would give a new partial geometry  $pg(5,17,3)$ .

ACKNOWLEDGEMENT. We thank Frank De Clerck and Hennie Wilbrink for several useful discussions.

## REFERENCES.

- [0] A. Blokhuis, *On subsets of  $GF(q^2)$  with square differences*, Proceedings KNAW A87, (1984).
- [1] R.C. Bose, *Strongly regular graphs, partial geometries and partially balanced designs*, Pacific J. Math. 13 (1963) 389-419.
- [2] A.E. Brouwer & J.H. van Lint, *Strongly regular graphs and partial geometries*, in: Enumeration and Design (eds. M. Jackson and S. Vanstone), Acad. Press (1984) 85-122.
- [3] F. Buekenhout, *Extensions of polar spaces and the doubly transitive symplectic groups*, Geom. Ded. 6 (1977) 13-21.
- [4] R. Calderbank & D.B. Wales, *The Haemers partial geometry and the Steiner system  $S(5,8,24)$* , Discrete Math. 51 (1984) 125-136.
- [5] P.J. Cameron, *Biplanes*, Math. Z. 131, 85-101.
- [6] P.J. Cameron, *Extended generalised quadrangles - a survey*, Proceedings Bose Memorial Conference, Calcutta 1988 (to appear).
- [7] P.J. Cameron & J.H. van Lint, *Graphs, codes and designs*, London Math. Soc. Lect. Notes Ser. 43, Cambridge 1980.
- [8] A.M. Cohen, *A new partial geometry with parameters  $(s,t,\alpha) = (7,8,4)$* , J. of Geometry 16 (1981) 181-186.
- [9] F. De Clerck, *The pseudo-geometric  $(t,s,s-1)$ -graphs*, Simon Stevin Vol. 53 no. 4 (1979) 301-317.
- [10] F. De Clerck, *Strongly regular graphs and partial geometries*, Preprint 37 (1985), Dept. of Math., Univ. of Napels, Italy.
- [11] F. De Clerck, R.H. Dye & J.A. Thas, *An infinite class of partial geometries associated with the hyperbolic quadric in  $pg(4n-1,2)$* , Europ. J. Combinatorics 1 (1980) 323-326.
- [12] F. De Clerck, H. Gevaert & J.A. Thas, *Partial geometries and copolar spaces*, Proceedings Combinatorics '88, Ravello 1988 (to appear).
- [13] F. De Clerck & V.D. Tonchev, *Partial geometries and quadratics* (preprint).
- [14] J.C. Fisher, *Geometry according to Euclid*, Preprint 11 (1977), Dept. of Math., Univ. of Regina, Canada.
- [15] W.H. Haemers, *A new partial geometry constructed from the Hoffman-Singleton graph*, In: Finite Geometries and Designs (eds. P.J. Cameron,

- J.W.P. Hirschfeld and D.R. Hughes), London Math. Soc. Lect. Notes Ser. 49, Cambridge (1981) 119-127.
- [16] W.H. Haemers & J.H. van Lint, *A partial geometry  $pg(9,8,4)$* , Annals of Discrete Math. 15 (1982) 205-212.
- [17] S.A. Hobart & D.R. Hughes, *Extended partial geometries: nets and dual nets* (preprint).
- [18] X.L. Hubaut, *Strongly regular graphs*, Discrete Math. 13 (1975) 357-381.
- [19] C.W.H. Lam, L. Thiel, S. Swiercz & J. McKay, *The nonexistence of ovals in a projective plane of order 10*, Discrete Math. 45 (1983) 319-321.
- [20] R. Mathon, *The partial geometries  $pg(5,7,3)$* , Congressus Numerantium 31 (1981) 129-139.
- [21] V.C. Mavron & M.S. Shrikhande, *On designs with intersection numbers 0 and 2*, Archiv der Math. (to appear).
- [22] J.J. Seidel, *Strongly regular graphs with  $(-1,1,0)$  adjacency matrix having eigenvalue 3*, Linear Algebra and Appl. 1 (1968) 281-298.
- [23] J.J. Seidel, *On two-graphs and Shult's characterization of symplectic and orthogonal geometries over  $GF(2)$* , Report 73-WSK-02, Eindhoven Univ. of Technology, The Netherlands, 1973.
- [24] J.J. Seidel, *A survey of two-graphs*, in: Proc. Intern. Colloq. Theorie Combinatorie (Roma 1973), tomo I, Accad. Naz. Lincei (1976) 481-511.
- [25] J.J. Seidel & D.E. Taylor, *Two-graphs, a second survey*, in: Proc. Intern. Colloq. Algebraic Methods in Graph Theory, Szeged 1978, Coll. Math. Soc. Bolyai 25, 689-711.
- [26] E. Spence, *Is Taylor's graph geometric?* (in preparation).
- [27] L. Storme (personal communication).
- [28] D.E. Taylor, *Regular 2-graphs*, Proc. London Math. Soc. 35 (1977) 257-274.
- [29] J.A. Thas, *Extensions of finite generalised quadrangles*, Symp. Math. 28 (1986) 127-143.
- [30] J.A. Thas (personal communication).
- [31] V.D. Tonchev (personal communication).
- [32] H.A. Wilbrink, *Two-graphs and geometries* (manuscript).



## IN 1988 REEDS VERSCHENEN

- 297 Bert Bettonvil  
Factor screening by sequential bifurcation
- 298 Robert P. Gilles  
On perfect competition in an economy with a coalitional structure
- 299 Willem Selen, Ruud M. Heuts  
Capacitated Lot-Size Production Planning in Process Industry
- 300 J. Kriens, J.Th. van Lieshout  
Notes on the Markowitz portfolio selection method
- 301 Bert Bettonvil, Jack P.C. Kleijnen  
Measurement scales and resolution IV designs: a note
- 302 Theo Nijman, Marno Verbeek  
Estimation of time dependent parameters in linear models  
using cross sections, panels or both
- 303 Raymond H.J.M. Gradus  
A differential game between government and firms: a non-cooperative  
approach
- 304 Leo W.G. Strijbosch, Ronald J.M.M. Does  
Comparison of bias-reducing methods for estimating the parameter in  
dilution series
- 305 Drs. W.J. Reijnders, Drs. W.F. Verstappen  
Strategische bespiegelingen betreffende het Nederlandse kwaliteits-  
concept
- 306 J.P.C. Kleijnen, J. Kriens, H. Timmermans and H. Van den Wildenberg  
Regression sampling in statistical auditing
- 307 Isolde Woittiez, Arie Kapteyn  
A Model of Job Choice, Labour Supply and Wages
- 308 Jack P.C. Kleijnen  
Simulation and optimization in production planning: A case study
- 309 Robert P. Gilles and Pieter H.M. Ruys  
Relational constraints in coalition formation
- 310 Drs. H. Leo Theuns  
Determinanten van de vraag naar vakantiereizen: een verkenning van  
materiële en immateriële factoren
- 311 Peter M. Kort  
Dynamic Firm Behaviour within an Uncertain Environment
- 312 J.P.C. Blanc  
A numerical approach to cyclic-service queueing models

- 313 Drs. N.J. de Beer, Drs. A.M. van Nunen, Drs. M.O. Nijkamp  
Does Morkmon Matter?
- 314 Th. van de Klundert  
Wage differentials and employment in a two-sector model with a dual  
labour market
- 315 Aart de Zeeuw, Fons Groot, Cees Withagen  
On Credible Optimal Tax Rate Policies
- 316 Christian B. Mulder  
Wage moderating effects of corporatism  
Decentralized versus centralized wage setting in a union, firm,  
government context
- 317 Jörg Glombowski, Michael Krüger  
A short-period Goodwin growth cycle
- 318 Theo Nijman, Marno Verbeek, Arthur van Soest  
The optimal design of rotating panels in a simple analysis of  
variance model
- 319 Drs. S.V. Hannema, Drs. P.A.M. Versteijne  
De toepassing en toekomst van public private partnership's bij de  
grote en middelgrote Nederlandse gemeenten
- 320 Th. van de Klundert  
Wage Rigidity, Capital Accumulation and Unemployment in a Small Open  
Economy
- 321 M.H.C. Paardekooper  
An upper and a lower bound for the distance of a manifold to a nearby  
point
- 322 Th. ten Raa, F. van der Ploeg  
A statistical approach to the problem of negatives in input-output  
analysis
- 323 P. Kooreman  
Household Labor Force Participation as a Cooperative Game; an Empiri-  
cal Model
- 324 A.B.T.M. van Schaik  
Persistent Unemployment and Long Run Growth
- 325 Dr. F.W.M. Boekema, Drs. L.A.G. Oerlemans  
De lokale produktiestructuur doorgelicht.  
Bedrijfstakverkenningen ten behoeve van regionaal-economisch onder-  
zoek
- 326 J.P.C. Kleijnen, J. Kriens, M.C.H.M. Lafleur, J.H.F. Pardoel  
Sampling for quality inspection and correction: AOQL performance  
criteria



- 327 Theo E. Nijman, Mark F.J. Steel  
Exclusion restrictions in instrumental variables equations
- 328 B.B. van der Genugten  
Estimation in linear regression under the presence of heteroskedasticity of a completely unknown form
- 329 Raymond H.J.M. Gradus  
The employment policy of government: to create jobs or to let them create?
- 330 Hans Kremers, Dolf Talman  
Solving the nonlinear complementarity problem with lower and upper bounds
- 331 Antoon van den Elzen  
Interpretation and generalization of the Lemke-Howson algorithm
- 332 Jack P.C. Kleijnen  
Analyzing simulation experiments with common random numbers, part II: Rao's approach
- 333 Jacek Osiewalski  
Posterior and Predictive Densities for Nonlinear Regression. A Partly Linear Model Case
- 334 A.H. van den Elzen, A.J.J. Talman  
A procedure for finding Nash equilibria in bi-matrix games
- 335 Arthur van Soest  
Minimum wage rates and unemployment in The Netherlands
- 336 Arthur van Soest, Peter Kooreman, Arie Kapteyn  
Coherent specification of demand systems with corner solutions and endogenous regimes
- 337 Dr. F.W.M. Boekema, Drs. L.A.G. Oerlemans  
De lokale produktiestructuur doorgelicht II. Bedrijfstakverkenningen ten behoeve van regionaal-economisch onderzoek. De zeescheepsnieuwbouwindustrie
- 338 Gerard J. van den Berg  
Search behaviour, transitions to nonparticipation and the duration of unemployment
- 339 W.J.H. Groenendaal and J.W.A. Vingerhoets  
The new cocoa-agreement analysed
- 340 Drs. F.G. van den Heuvel, Drs. M.P.H. de Vor  
Kwantificering van ombuigen en bezuinigen op collectieve uitgaven 1977-1990
- 341 Pieter J.F.G. Meulendijks  
An exercise in welfare economics (III)

- 342 W.J. Selen and R.M. Heuts  
A modified priority index for Günther's lot-sizing heuristic under capacitated single stage production
- 343 Linda J. Mittermaier, Willem J. Selen, Jeri B. Waggoner, Wallace R. Wood  
Accounting estimates as cost inputs to logistics models
- 344 Remy L. de Jong, Rashid I. Al Layla, Willem J. Selen  
Alternative water management scenarios for Saudi Arabia
- 345 W.J. Selen and R.M. Heuts  
Capacitated Single Stage Production Planning with Storage Constraints and Sequence-Dependent Setup Times
- 346 Peter Kort  
The Flexible Accelerator Mechanism in a Financial Adjustment Cost Model
- 347 W.J. Reijnders en W.F. Verstappen  
De toenemende importantie van het verticale marketing systeem
- 348 P.C. van Batenburg en J. Kriens  
E.O.Q.L. - A revised and improved version of A.O.Q.L.
- 349 Drs. W.P.C. van den Nieuwenhof  
Multinationalisatie en coördinatie  
De internationale strategie van Nederlandse ondernemingen nader beschouwd
- 350 K.A. Bubshait, W.J. Selen  
Estimation of the relationship between project attributes and the implementation of engineering management tools
- 351 M.P. Tummers, I. Woittiez  
A simultaneous wage and labour supply model with hours restrictions
- 352 Marco Versteijne  
Measuring the effectiveness of advertising in a positioning context with multi dimensional scaling techniques
- 353 Dr. F. Boekema, Drs. L. Oerlemans  
Innovatie en stedelijke economische ontwikkeling
- 354 J.M. Schumacher  
Discrete events: perspectives from system theory
- 355 F.C. Bussemaker, W.H. Haemers, R. Mathon and H.A. Wilbrink  
A (49,16,3,6) strongly regular graph does not exist
- 356 Drs. J.C. Caanen  
Tien jaar inflatieneutrale belastingheffing door middel van vermogensaftrek en voorraadaf trek: een kwantitatieve benadering

- 357 R.M. Heuts, M. Bronckers  
A modified coordinated reorder procedure under aggregate investment  
and service constraints using optimal policy surfaces
- 358 B.B. van der Genugten  
Linear time-invariant filters of infinite order for non-stationary  
processes
- 359 J.C. Engwerda  
LQ-problem: the discrete-time time-varying case
- 360 Shan-Hwei Nienhuys-Cheng  
Constraints in binary semantical networks
- 361 A.B.T.M. van Schaik  
Interregional Propagation of Inflationary Shocks
- 362 F.C. Drost  
How to define UMVU
- 363 Rommert J. Casimir  
Infogame users manual  
Rev 1.2 December 1988
- 364 M.H.C. Paardekooper  
A quadratically convergent parallel Jacobi-process for diagonal  
dominant matrices with nondistinct eigenvalues
- 365 Robert P. Gilles, Pieter H.M. Ruys  
Characterization of Economic Agents in Arbitrary Communication  
Structures
- 366 Harry H. Tigelaar  
Informative sampling in a multivariate linear system disturbed by  
moving average noise
- 367 Jörg Glombowski  
Cyclical interactions of politics and economics in an abstract  
capitalist economy

## IN 1989 REEDS VERSCHENEN

- 368 Ed Nijssen, Will Reijnders  
"Macht als strategisch en tactisch marketinginstrument binnen de distributieketen"
- 369 Raymond Gradus  
Optimal dynamic taxation with respect to firms
- 370 Theo Nijman  
The optimal choice of controls and pre-experimental observations
- 371 Robert P. Gilles, Pieter H.M. Ruys  
Relational constraints in coalition formation
- 372 F.A. van der Duyn Schouten, S.G. Vanneste  
Analysis and computation of  $(n,N)$ -strategies for maintenance of a two-component system
- 373 Drs. R. Hamers, Drs. P. Verstappen  
Het company ranking model: a means for evaluating the competition
- 374 Rommert J. Casimir  
Infogame Final Report
- 375 Christian B. Mulder  
Efficient and inefficient institutional arrangements between governments and trade unions; an explanation of high unemployment, corporatism and union bashing
- 376 Marno Verbeek  
On the estimation of a fixed effects model with selective non-response
- 377 J. Engwerda  
Admissible target paths in economic models
- 378 Jack P.C. Kleijnen and Nabil Adams  
Pseudorandom number generation on supercomputers
- 379 J.P.C. Blanc  
The power-series algorithm applied to the shortest-queue model
- 380 Prof. Dr. Robert Bannink  
Management's information needs and the definition of costs, with special regard to the cost of interest
- 381 Bert Bettonvil  
Sequential bifurcation: the design of a factor screening method
- 382 Bert Bettonvil  
Sequential bifurcation for observations with random errors



- 383 Harold Houba and Hans Kremers  
Correction of the material balance equation in dynamic input-output models
- 384 T.M. Doup, A.H. van den Elzen, A.J.J. Talman  
Homotopy interpretation of price adjustment processes
- 385 Drs. R.T. Frambach, Prof. Dr. W.H.J. de Freytas  
Technologische ontwikkeling en marketing. Een oriënterende beschouwing
- 386 A.L.P.M. Hendriks, R.M.J. Heuts, L.G. Hoving  
Comparison of automatic monitoring systems in automatic forecasting
- 387 Drs. J.G.L.M. Willems  
Enkele opmerkingen over het inversificerend gedrag van multinationale ondernemingen
- 388 Jack P.C. Kleijnen and Ben Annink  
Pseudorandom number generators revisited
- 389 Dr. G.W.J. Hendrikse  
Speltheorie en strategisch management
- 390 Dr. A.W.A. Boot en Dr. M.F.C.M. Wijn  
Liquiditeit, insolventie en vermogensstructuur
- 391 Antoon van den Elzen, Gerard van der Laan  
Price adjustment in a two-country model
- 392 Martin F.C.M. Wijn, Emanuel J. Bijnen  
Prediction of failure in industry  
An analysis of income statements
- 393 Dr. S.C.W. Eijffinger and Drs. A.P.D. Gruijters  
On the short term objectives of daily intervention by the Deutsche Bundesbank and the Federal Reserve System in the U.S. Dollar - Deutsche Mark exchange market
- 394 Dr. S.C.W. Eijffinger and Drs. A.P.D. Gruijters  
On the effectiveness of daily interventions by the Deutsche Bundesbank and the Federal Reserve System in the U.S. Dollar - Deutsche Mark exchange market
- 395 A.E.M. Meijer and J.W.A. Vingerhoets  
Structural adjustment and diversification in mineral exporting developing countries
- 396 R. Gradus  
About Tobin's marginal and average  $q$   
A Note
- 397 Jacob C. Engwerda  
On the existence of a positive definite solution of the matrix equation  $X + A^T X^{-1} A = I$

- 398 Paul C. van Batenburg and J. Kriens  
Bayesian discovery sampling: a simple model of Bayesian inference in auditing
- 399 Hans Kremers and Dolf Talman  
Solving the nonlinear complementarity problem
- 400 Raymond Gradus  
Optimal dynamic taxation, savings and investment



**Bibliotheek K. U. Brabant**



17 000 01086030 3