



*Demographic Research* a free, expedited, online journal  
of peer-reviewed research and commentary  
in the population sciences published by the  
Max Planck Institute for Demographic Research  
Konrad-Zuse Str. 1, D-18057 Rostock · GERMANY  
[www.demographic-research.org](http://www.demographic-research.org)

---

## ***DEMOGRAPHIC RESEARCH***

**VOLUME 20, ARTICLE 2, PAGES 3-6  
PUBLISHED 20 JANUARY 2009**

<http://www.demographic-research.org/Volumes/Vol20/2/>

DOI: 10.4054/DemRes.2009.20.2

*Formal Relationships 1*

**Life lived equals life left in  
stationary populations**

**Joshua R. Goldstein**

This article is part of the Special Collection "Formal Relationships".  
Guest Editors are Joshua R. Goldstein and James W. Vaupel.

© 2009 Joshua R. Goldstein.

*This open-access work is published under the terms of the Creative Commons Attribution NonCommercial License 2.0 Germany, which permits use, reproduction & distribution in any medium for non-commercial purposes, provided the original author(s) and source are given credit. See <http://creativecommons.org/licenses/by-nc/2.0/de/>*

## **Table of Contents**

1	Relationship	3
2	Proof	4
3	History and related results	4
4	Applications	4
	References	6

## Life lived equals life left in stationary populations

Joshua R. Goldstein<sup>1, 2</sup>

### Abstract

The average age of the individuals in a population is equal to the average remaining life expectancy when the population is stationary.

### 1. Relationship

The mean age of a stationary population is equal to the average remaining life expectancy. Let  $a$  denote age,  $e(a)$  be life expectancy at age  $a$ , and  $\ell(a)$  be the proportion surviving to age  $a$ . In a stationary population, the density of individuals aged  $a$  is  $\ell(a)/e(0)$ .

If we define the mean life lived and the mean life left as, respectively,

$$(1) \quad \bar{a} = \frac{\int_0^{\omega} a \ell(a) da}{e(0)}$$

and

$$(2) \quad \bar{e} = \frac{\int_0^{\omega} e(a) \ell(a) da}{e(0)},$$

then

$$(3) \quad \bar{a} = \bar{e}.$$

---

<sup>1</sup>Max-Planck-Institute for Demographic Research. E-mail: goldstein@demogr.mpg.de

<sup>2</sup>I thank James W. Vaupel and an anonymous reviewer.

## 2. Proof

Since life expectancy at age  $a$  is  $e(a) = \int_a^\omega \ell(x) dx / \ell(a)$ ,

$$(4) \quad \bar{e} = \frac{\int_0^\omega \ell(a) \left( \int_a^\omega \ell(x) / \ell(a) dx \right) da}{e(0)} = \frac{\int_0^\omega \int_a^\omega \ell(x) dx da}{e(0)}.$$

Reversing the order of integration, with the corresponding change in the limits of integration, gives

$$(5) \quad \bar{e} = \frac{\int_0^\omega \left( \int_a^\omega \ell(x) dx \right) da}{e(0)} = \frac{\int_0^\omega \ell(x) \left( \int_0^x 1 da \right) dx}{e(0)} = \frac{\int_0^\omega x \ell(x) dx}{e(0)} = \bar{a}.$$

Q.E.D.

## 3. History and related results

Kim and Aron (1989) gave the same result using integration by parts. A more general form of the result is known from the distributional equality of backward and forward recurrences in renewal theory: see Cox (1962). Keyfitz (1985:74) calls  $\bar{e}$  the average expectation of life, denoting it  $E$ .

It is also possible to show that in a growing stable population  $\bar{e} > \bar{a}$ , and in a shrinking stable population  $\bar{e} < \bar{a}$ .

The mean age of a stationary population can be expressed in terms of the mean age of death and the coefficient of variation (CV) of age of death,

$$(6) \quad \bar{a} = \frac{e(0) + \frac{\sigma^2}{e(0)}}{2} = \frac{e(0)}{2} (1 + CV^2),$$

where  $\sigma^2$  is the variance of age of death and  $CV$  is  $\sigma / e(0)$ .

Thus, average remaining life expectancy  $\bar{e}$  can also be expressed this way, showing the dependence on the mean and variability of life's length.

## 4. Applications

Time lived and time left are central to debates about population aging. Increases in  $\bar{a}$  are seen as making the population "older"; but increasing distance from death  $\bar{e}$  is seen as making the population effectively younger (Ryder 1975; Miller 2001). The equality of  $\bar{a}$  and  $\bar{e}$  in stationary populations means that mortality decline<sup>3</sup> increases both by the

<sup>3</sup>More precisely, mortality decline that does not influence the birth rate.

same amount. As illustrated by Sanderson and Scherbov (2005), populations can even be getting older and younger at the same time, depending on one's point of view. The quantity  $\bar{a}/(\bar{a} + \bar{e})$  can be used as a measure of relative aging, varying between 0 and 1, taking the value  $\frac{1}{2}$  in stationary populations.

Applications of equality (2) include estimation of  $\bar{a}$  when  $\bar{e}$  is more easily observed and vice versa. Kim and Aron give an example in epidemiological modeling. A potential application for household demography would be to infer average time remaining in the household from retrospective estimates of average time in the household.

Note: On Tuesday, March 3, 2009, three small typos were corrected in this PDF, on pages 3 and 4.

## References

- Cox, D. R. (1962). *Renewal Theory*. London: Methuen and Co.
- Keyfitz, N. (1985). *Applied Mathematical Demography*. New York: Springer, second edition.
- Kim, Y. J. and Aron, J. L. (1989). On the equality of average age and average expectation of remaining life in a stationary population. *SIAM Review* 31(1): 110–113. doi:[10.1137/1031005](https://doi.org/10.1137/1031005).
- Miller, T. (2001). Increasing longevity and medicare expenditures. *Demography* 38(2): 215–226. doi:[10.1353/dem.2001.0018](https://doi.org/10.1353/dem.2001.0018).
- Ryder, N. B. (1975). Notes on stationary populations. *Population Index* 41(1): 3–28. doi:[10.2307/2734140](https://doi.org/10.2307/2734140).
- Sanderson, W. C. and Scherbov, S. (2005). Average remaining lifetimes can increase as human populations age. *Nature* 435: 811–813. doi:[10.1038/nature03593](https://doi.org/10.1038/nature03593).