

CentER



Discussion Paper

No. 2003–66

CONDITIONS FOR SINGULAR INCIDENCE MATRICES

By W.H. Haemers

July 2003

ISSN 0924-7815

Conditions for singular incidence matrices

Willem H. Haemers

Tilburg University, Dept. of Econometrics and O.R.,
Tilburg, The Netherlands, e-mail: haemers@uvt.nl

Abstract

Suppose one looks for a square integral matrix N , for which NN^\top has a prescribed form. Then the Hasse-Minkowski invariants and the determinant of NN^\top lead to necessary conditions for existence. The Bruck-Ryser-Chowla theorem gives a famous example of such conditions in case N is the incidence matrix of a square block design. This approach fails when N is singular. In this paper it is shown that in some cases conditions can still be obtained if the kernels of N and N^\top are known, or known to be rationally equivalent. This leads for example to non-existence conditions for self-dual generalised polygons, semi-regular square divisible designs and distance-regular graphs.

1 Introduction

Consider a square $2-(v, k, \lambda)$ design with incidence matrix N . (We prefer the name ‘square’ above ‘symmetric’.) Then $NN^\top = \lambda J_v + (k - \lambda)I_v$, where J_v is the $v \times v$ all-ones matrix and I_v is the identity matrix of size v . The Bruck-Ryser-Chowla theorem is based on two observations (see for example [5]). The first one is that $\det N = \det N^\top$ is an integer. Therefore $\det(\lambda J_v + (k - \lambda)I_v)$ is an integral square, hence $k - \lambda$ is a square if v is even. The other observation is that, since N is a non-singular rational matrix, $\lambda J_v + (k - \lambda)I_v$ is rationally congruent to I_v , and therefore these two matrices have the same Hasse-Minkowski invariants. These invariants can be expressed in terms of v , k and λ from which it follows that for odd v the Diophantine equation $(k - \lambda)X^2 + (-1)^{(v-1)/2}\lambda Y^2 = Z^2$ has an integral solution different from $X = Y = Z = 0$. Similar approaches work for other square incidence structures for which the determinant or the Hasse-Minkowski invariants of NN^\top are known. See for example [5], Chapter 12. It is clear that this approach gives no conditions if N is singular. In the present paper we modify the mentioned approach such that we still find conditions for singular N . The key lemma is a simple trick that changes a singular N into a non-singular matrix M in such a way that for some types of designs it is still possible to compute the Hasse-Minkowski invariants or the (square free part of the) determinant of MM^\top .

Lemma 1 Suppose N is a rational $v \times v$ matrix of rank $v - m$. Let Z be a rational $v \times v$ matrix of rank m , such that $N^\top Z = NZ^\top = O$. Define $M = N + Z$, then

i. $MM^\top = NN^\top + ZZ^\top$,

ii. the eigenvalues of MM^\top are the positive eigenvalues of NN^\top together with the positive eigenvalues of ZZ^\top ,

iii. MM^\top is non-singular.

Proof. Part *i* is straightforward. To prove *ii*, first notice that NN^\top and ZZ^\top commute, so they have a common orthogonal basis of eigenvectors. Suppose \mathbf{v} is such an eigenvector that corresponds to a positive eigenvalue of NN^\top . Then \mathbf{v} is orthogonal to the kernel of NN^\top , which is the span of the columns of Z . Hence $Z^\top \mathbf{v} = \mathbf{0}$, so the corresponding eigenvalue of ZZ^\top equals 0. Similarly, a positive eigenvalue of ZZ^\top corresponds to an eigenvalue 0 of NN^\top . This proves *ii*, since NN^\top has $v - m$ positive eigenvalues, and ZZ^\top has m positive eigenvalues. Statement *iii* follows because MM^\top has only positive eigenvalues. \square

For a given N , a matrix Z with the required properties always exists. One way to make such a Z is the following. Take rational $v \times m$ matrices L and R , whose columns form a basis for the left and the right kernel of N , respectively. Then $\text{rank } L = \text{rank } R = m$ and $N^\top L = NR = O$. Therefore $Z = LR^\top$ has the desired properties.

In the coming sections we will consider two kinds of square designs for which something new can be said: Self-dual designs and semi-regular square divisible designs.

2 Self-dual designs

Consider two m -dimensional subspaces V and W of the vectorspace \mathbb{Q}^v . Let L and R be rational $v \times m$ matrices whose columns span V and W , respectively. We call the subspaces V and W *rationally equivalent* if $L^\top L$ and $R^\top R$ are rationally congruent matrices, which means that $S^\top L^\top L S = R^\top R$ for some non-singular rational matrix S . Note that rational equivalence of vectorspaces does not depend on the choice of L and R indeed.

Lemma 2 Let N be a rational $v \times v$ matrix. If the left kernel and the right kernel of N are rationally equivalent then the product of the non-zero eigenvalues of NN^\top is a rational square.

Proof. Let L and R be $v \times m$ matrices whose columns form a basis for the left and the right kernel of N , respectively. Put $Z = LR^\top$. Then $ZZ^\top = LR^\top RL^\top = LS^\top L^\top L SL^\top$ (with S as above). The non-zero eigenvalues of $L(S^\top L^\top L SL^\top)$ coincide with the non-zero eigenvalues of $(S^\top L^\top L SL^\top)L$. But $\det(S^\top L^\top L SL^\top L) = (\det S)^2 (\det L^\top L)^2$ which is a non-zero rational square. Thus we have that the product of the non-zero eigenvalues of ZZ^\top is a square, and

Lemma 1 finishes the proof. □

If N is the incidence matrix of a self-dual design (that is, N and N^\top are isomorphic), then left and right kernel of N are obviously rationally equivalent and Lemma 2 gives:

Theorem 1 *If N is the incidence matrix of a self-dual design, then the product of the positive eigenvalues of NN^\top is an integral square.*

For example if N is the incidence matrix of a self-dual partial geometry with parameters s ($= t$) and α (see [4]), the non-zero eigenvalues of NN^\top are $(s + 1)^2$ of multiplicity 1, and $2s + 1 - \alpha$ of multiplicity $s^2(s + 1)^2/\alpha(2s + 1 - \alpha)$. So if the latter multiplicity is odd, $2s + 1 - \alpha$ is a square. In particular if $\alpha = 1$, the partial geometry is a generalised quadrangle of order s (denoted by $GQ(s)$) and we find:

Corollary 1 *There exists no self-dual $GQ(s)$ if $s \equiv 2 \pmod{4}$ and $2s$ is not a square.*

For example no $GQ(6)$ is self-dual. Similarly, if N is the incidence matrix of a generalised hexagon of order s (denoted by $GH(s)$), the non-zero eigenvalues of NN^\top are $(s + 1)^2$, s and $3s$ of multiplicity 1, $s(1 + s)^2(1 - s + s^2)/2$ and $s(1 + s)^2(1 + s + s^2)/6$, respectively (see for example [2] p.203). Thus we find:

Corollary 2 *There exists no self-dual $GH(s)$ if $s \equiv 2 \pmod{4}$.*

Stronger condition are known if the incidence matrix of a $GQ(s)$ or $GH(s)$ is symmetric (see [7] p.309). A symmetric incidence matrix clearly implies that the structure is self-dual, but the converse is not true in general.

3 Square divisible designs

Another case when Lemma 1 can be applied is when the left and right kernel of N are determined by the design requirements. Note that the left kernel of N is the kernel of NN^\top , and similarly, the right kernel of N is the kernel of $N^\top N$. So the lemma applies for square incidence matrices N for which NN^\top and $N^\top N$ are prescribed. For example, consider a 2 -(v, k, λ) design with a $v \times b$ incidence matrix where $b > v$. Extend the $v \times b$ incidence matrix with $b - v$ zero rows. For the $b \times b$ matrix N thus obtained NN^\top is known, and so is its left kernel. The right kernel of N is in general not known, but there are some types of designs for which $N^\top N$ is prescribed. These include strongly resolvable designs and triangular designs. For these designs Bruck-Ryser-Chowla type conditions have been worked out; see [6], [5] and [3], so we will not do it again.

In this section we consider semi-regular square divisible designs. A divisible design (also called group-divisible design) with parameters k, g, n, λ_1 and λ_2 , is an incidence structure,

denoted by $GD(k, g, n, \lambda_1, \lambda_2)$, for which the points can be ordered such that the incidence matrix N satisfies

$$NN^\top = \lambda_2 J_v + (\lambda_1 - \lambda_2)K_{n,g} + (r - \lambda_1)I_v, \quad \text{and} \quad N^\top J_v = kJ_v,$$

where $K_{n,g}$ is the block diagonal matrix $I_n \otimes J_g$, $v = ng$ is the number of points and $r = ((n-1)g\lambda_2 + (g-1)\lambda_1)/(k-1)$ is the replication number. The eigenvalues of NN^\top are easily seen to be kr , $r - \lambda_1$, and $g(\lambda_1 - \lambda_2) + r - \lambda_1$ with multiplicities 1, $n(g-1)$ and $n-1$, respectively. Assume that N is a square matrix. Then $r = k$, and the eigenvalues of NN^\top become k^2 , $k - \lambda_1$ and $k^2 - gn\lambda_2$. If N is non-singular, the divisible design is called regular, and necessary conditions for existence have been known for a long time, see [1], [5] p.228, or [2] p.23. If N is singular, either $k = \lambda_1$ and $N = N' \otimes J_n$, where N' is the incidence matrix of a square block design (then the divisible design is called singular), or $k^2 = ng\lambda_2$ and the divisible design is called semi-regular.

Theorem 2 *Let D be a design with the property that both D and its dual are a semi-regular $GD(k, g, n, \lambda_1, \lambda_2)$. Then*

- i. if g is even and n is odd, $k - \lambda_1$ is an integral square,*
- ii. if g is even and $n \equiv 2 \pmod{4}$ then $k - \lambda_1$ is the sum of two integral squares,*
- iii. if g and n are odd, the equation $(k - \lambda_1)X^2 + (-1)^{(g-1)/2}gY^2 = Z^2$ has an integral solution different from $X = Y = Z = 0$.*

Proof. Suppose N is the incidence matrix of D . We may assume that $NN^\top = N^\top N$, which implies that N^\top and N have the same kernel, so by Lemma 2 the product of the non-zero eigenvalues of NN^\top is a square, which proves *i*. Define $Z = (J_n - nI_n) \otimes J_g$. Then $\text{rank } Z = n - 1$, and $NN^\top Z = N^\top NZ = O$, so Z satisfies the requirement for Lemma 1. Hence

$$MM^\top = NN^\top + ZZ^\top = (\lambda_2 - gn)J_v + (\lambda_1 - \lambda_2 + gn^2)K_{n,g} + (k - \lambda_1)I_v.$$

has eigenvalues k^2 , $\rho = k - \lambda_1$ and $\sigma = g^2n^2$ of multiplicity 1, $n(g-1)$ and $n-1$ respectively. The Hasse-Minkowski invariant $C_p(MM^\top)$ with respect to the odd prime p of a matrix MM^\top of the above form is known, see for example [1].

$$C_p(MM^\top) = (\rho, -1)_p^{n(g-1)(n+g-1)/2} (\sigma, -1)_p^{n(n-1)/2} (\sigma, g)_p^n (\rho, g)_p^n (\sigma, \lambda_2 - gn)_p =$$

$$(\rho, -1)_p^{n(g-1)(n+g-1)/2} (\rho, g)_p^n,$$

where $(a, b)_p$ is the Hilbert norm residue symbol, defined by $(a, b)_p = 1$ if for all t the congruence $aX^2 + bY^2 \equiv 1 \pmod{p^t}$ has a rational solution, and $(a, b)_p = -1$ otherwise. Since M is a non-singular rational matrix, $C_p(MM^\top) = C_p(I_v) = 1$ for every odd prime p , and the conditions *ii* and *iii* follow. \square

For example there exists no $GD(18, 4, 9, 6, 9)$ for which the dual is also such a design. Note that in case $n = 1$, D is a square block design and the conditions are those of Bruck, Ryser and Chowla. The above theorem also has consequences for distance-regular graphs. Some putative distance-regular graphs imply the existence of square divisible designs (see [2] p.22), and in case these divisible designs are semi-regular we obtain new conditions.

Corollary 3 *Suppose there exists a distance-regular graph of diameter 4 with $2g^2\mu$ vertices and intersection array $\{g\mu, g\mu - 1, (g - 1)\mu, 1; 1, \mu, g\mu - 1, g\mu\}$. Then*

i. If μ is odd and $g \equiv 2 \pmod{4}$ then $g\mu$ is the sum of two integral squares.

ii. If μ and g are odd, then the equation $\mu X^2 + (-1)^{(g-1)/2} Y^2 = gZ^2$ has an integral solution different from $X = Y = Z = 0$.

Proof. Such a distance-regular graph is the incidence graph of a $GD(g\mu, g, g\mu, 0, \mu)$ for which the dual is also such a design. \square

For example a distance-regular graph with intersection array $\{15, 14, 12, 1; 1, 3, 14, 15\}$ does not exist. Note that a distance-regular graph with intersection array $\{g\mu - 1, (g - 1)\mu, 1; 1, \mu, g\mu - 1\}$ also gives rise to a semi-regular square divisible design; see [2], p.24. But here we find no new restrictions.

Acknowledgement. I thank Edwin van Dam for many relevant conversations.

References

- [1] R.C. Bose and W.S. Connor, *Combinatorial properties of group divisible incomplete block designs*, Ann. Math. Statist. 23 (1952) 367-383.
- [2] A.E. Brouwer, A.M. Cohen and A. Neumaier, *Distance-Regular Graphs*, Springer, Heidelberg, 1989.
- [3] M.J. Coster and W.H. Haemers, *Quasi-symmetric designs related to the triangular graph*, Designs, Codes and Cryptography 5 (1995), 27-42.
- [4] F. De Clerck and H. Van Maldeghem, *Some classes of rank 2 geometries*, in: Handbook of Incidence Geometry (F. Buekenhout ed.), Elsevier Science B.V., 1995, pp. 433-475.
- [5] D. Raghavarao, *Constructions and combinatorial problems in designs of experiments*, John Wiley & Sons, Inc., 1971.
- [6] S.S. Shrikhande, D. Raghavarao and S.K. Tharthare, *Non-existence of some unsymmetrical partially balanced incomplete block designs*, Canad. J. Math. 15 (1963), 686-701.
- [7] H. Van Maldeghem, *Generalized polygons*, Birkhäuser, 1991.