

Der Open-Access-Publikationsserver der ZBW – Leibniz-Informationzentrum Wirtschaft
The Open Access Publication Server of the ZBW – Leibniz Information Centre for Economics

Klein, Ingo; Fischer, Matthias J.

Working Paper

Kurtosis ordering of the generalized secant hyperbolic distribution: a technical note

Diskussionspapiere // Friedrich-Alexander-Universität Erlangen-Nürnberg, Lehrstuhl für Statistik und Ökonometrie, No. 54/2003

Provided in cooperation with:

Friedrich-Alexander-Universität Erlangen-Nürnberg (FAU)

Suggested citation: Klein, Ingo; Fischer, Matthias J. (2003) : Kurtosis ordering of the generalized secant hyperbolic distribution: a technical note, Diskussionspapiere // Friedrich-Alexander-Universität Erlangen-Nürnberg, Lehrstuhl für Statistik und Ökonometrie, No. 54/2003, <http://hdl.handle.net/10419/29581>

Nutzungsbedingungen:

Die ZBW räumt Ihnen als Nutzerin/Nutzer das unentgeltliche, räumlich unbeschränkte und zeitlich auf die Dauer des Schutzrechts beschränkte einfache Recht ein, das ausgewählte Werk im Rahmen der unter

→ <http://www.econstor.eu/dspace/Nutzungsbedingungen> nachzulesenden vollständigen Nutzungsbedingungen zu vervielfältigen, mit denen die Nutzerin/der Nutzer sich durch die erste Nutzung einverstanden erklärt.

Terms of use:

The ZBW grants you, the user, the non-exclusive right to use the selected work free of charge, territorially unrestricted and within the time limit of the term of the property rights according to the terms specified at

→ <http://www.econstor.eu/dspace/Nutzungsbedingungen>
By the first use of the selected work the user agrees and declares to comply with these terms of use.

**Friedrich-Alexander-Universität
Erlangen-Nürnberg**
Wirtschafts-und Sozialwissenschaftliche Fakultät

Diskussionspapier

54 / 2003

**Kurtosis ordering of the generalized secant hyperbolic
distribution — A technical note**

Ingo Klein and Matthias Fischer



Lehrstuhl für Statistik und Ökonometrie
Lehrstuhl für Statistik und empirische Wirtschaftsforschung
Lange Gasse 20 · D-90403 Nürnberg

Kurtosis ordering of the generalized secant hyperbolic distribution – A technical note

Ingo Klein and Matthias Fischer
Department of Statistics and Econometrics,
University of Erlangen-Nuremberg

Abstract: Two major generalizations of the hyperbolic secant distribution have been proposed in the statistical literature which both introduce an additional parameter that governs the kurtosis of the generalized distribution. The generalized hyperbolic secant (GHS) distribution was introduced by Harkness and Harkness (1968) who considered the p -th convolution of a hyperbolic secant distribution. Another generalization, the so-called generalized secant hyperbolic (GSH) distribution was recently suggested by Vaughan (2002). In contrast to the GHS distribution, the cumulative and inverse cumulative distribution function of the GSH distribution are available in closed-form expressions. We use this property to proof that the additional shape parameter of the GSH distribution is actually a kurtosis parameter in the sense of van Zwet (1964).

Keywords: kurtosis ordering; hyperbolic secant distribution.

1 Introduction

The hyperbolic secant distribution — which was studied by Baten (1934) and Talacko (1956) — has not received sufficient attention in the literature, although it has a lot of nice properties: All moments and the moment-generating function exist, it is reproductive (i.e. the class is preserved under convolution) and both the cumulative and the inverse cumulative distribution function admit a closed form. In addition, the hyperbolic secant distribution exhibits more leptokurtosis than the normal and even more than the logistic distribution. Nevertheless, generalizations have been proposed that introduce an additional parameter which increase the "kurtosis" of the hyperbolic secant distribution. At first, Baten (1934) discussed the sum of n independent random variables, each having the same hyperbolic secant distribution. More general, Harkness and Harkness (1968) investigated the p -th convolution of hyperbolic secant variables for arbitrary positive $p > 0$ which, unfortunately, doesn't admit a closed-form representation for the cumulative and the inverse cumulative distribution function. Recently, Vaughan (2002) suggested a family of symmetric distributions, the so-called generalized secant hyperbolic (GSH) distribution. This family includes both the hyperbolic secant and the logistic distribution. Moreover, it closely approximates the Student t -distribution with corresponding kurtosis. In addition, the moment-generating function and all moments exist, and the cumulative and inverse cumulative distribution are again available in closed form. The range of "kurtosis" — measured by the fourth standardized moments — varies from 1.8 to infinity. Within this note we proof that the additional parameter of the GSH distribution is indeed a kurtosis parameter which preserves the kurtosis ordering of van Zwet (1964).

2 The generalized secant hyperbolic (GSH) distribution

The above-mentioned standard *generalized secant hyperbolic* (GSH) distribution – which is able to model both thin and fat tails – was introduced by Vaughan (2002) and has density

$$f_{GSH}(x; t) = c_1(t) \cdot \frac{\exp(c_2(t)x)}{\exp(2c_2(t)x) + 2a(t)\exp(c_2(t)x) + 1}, \quad x \in \mathbb{R} \quad (1)$$

with

$$\begin{aligned} a(t) &= \cos(t), \quad c_2(t) = \sqrt{\frac{\pi^2 - t^2}{3}}, \quad c_1(t) = \frac{\sin(t)}{t} \cdot c_2(t), \quad \text{for } -\pi < t \leq 0, \\ a(t) &= \cosh(t), \quad c_2(t) = \sqrt{\frac{\pi^2 + t^2}{3}}, \quad c_1(t) = \frac{\sinh(t)}{t} \cdot c_2(t), \quad \text{for } t > 0 \end{aligned}$$

The density from (1) is chosen so that $X \sim f_{GSH}(x)$ has zero mean and unit variance, the range of the "kurtosis parameter" $t \in (-\pi, \infty)$. The GSH distribution includes the logistic distribution ($t = 0$) and the hyperbolic secant distribution ($t = -\pi/2$) as special cases and the uniform distribution on $(-\sqrt{3}, \sqrt{3})$ as limiting case for $t \rightarrow \infty$. Vaughan (2002) derived, amongst other properties, the cumulative distribution function, depending on the parameter t , given by

$$F_{GSH}(x; t) = \begin{cases} 1 + \frac{1}{t} \operatorname{arccot} \left(-\frac{\exp(c_2(t)x) + \cos(t)}{\sin(t)} \right) & \text{for } t \in (-\pi, 0), \\ \frac{\exp(\pi x / \sqrt{3})}{1 + \exp(\pi x / \sqrt{3})} & \text{for } t = 0, \\ 1 - \frac{1}{t} \operatorname{arccoth} \left(\frac{\exp(c_2(t)x) + \cosh(t)}{\sinh(t)} \right) & \text{for } t > 0. \end{cases}$$

the inverse distribution function, given by

$$F_{GSH}^{-1}(u; t) = \begin{cases} \frac{1}{c_2(t)} \ln \left(\frac{\sin(tu)}{\sin(t(1-u))} \right) & \text{für } t \in (-\pi, 0), \\ \frac{\sqrt{3}}{\pi} \ln \left(\frac{u}{1-u} \right) & \text{für } t = 0, \\ \frac{1}{c_2(t)} \ln \left(\frac{\sinh(tu)}{\sinh(t(1-u))} \right) & \text{für } t > 0. \end{cases}$$

However, it was not proved that the "kurtosis parameter" t is actually a kurtosis parameter in the sense of van Zwet(1964). This will be done in the next section.

3 GSH distribution and kurtosis ordering

Van Zwet (1964) introduced a partial ordering of kurtosis \preceq_S on the set of symmetric distribution functions \mathcal{F}^s . Let $F, G \in \mathcal{F}^s$ and μ_F denote the location of symmetry of F , then \preceq_S is defined by

$$(A) \quad F \preceq_S G : \iff G^{-1}(F(x)) \text{ is convex for } x > \mu_F$$

and means that G has higher kurtosis than F . Balanda and MacGillivray (1990) generalized this partial ordering of van Zwet by using so-called spread functions defined as symmetric differences of quantiles:

$$S_F(u) = F^{-1}(u) - F^{-1}(1-u), \quad u \geq 0.5.$$

In the sense of Balanda and MacGillivray (1990), an arbitrary continuous, monotone increasing distribution function F has less kurtosis than an equally distribution function G if

$$(B) \quad F \preceq_S G : \iff S_G(S_F^{-1}(x)) \quad \text{is convex for } x > F^{-1}(0.5).$$

If F is symmetric, $F^{-1}(u) = -F^{-1}(1 - u)$ for $u > 0.5$, so that $S_F(u) = 2F^{-1}(u) \quad u \geq 0.5$. This means that the spread function essentially coincides with the quantile function. It can be shown that (A) and (B) coincide in this case. The aim of this note is to prove that the parameter t from (1) is a kurtosis parameter in the sense of van Zwet (1964).

Proposition 3.1 (Kurtosis ordering) *Assume that X_1 (X_2) follows a generalized secant hyperbolic distribution with parameter t_1 (t_2) and corresponding cumulative distribution functions F_1 (F_2). If $t_1 < t_2$, then $F_2 \preceq_S F_1$, i.e. F_1 has higher kurtosis than F_2 (in the sense of van Zwet).*

Proof: To prove this result, we distinguish between 4 cases:

$$\text{Case 1: } -\pi \leq t_1 < t_2 < 0,$$

$$\text{Case 2: } -\pi \leq t_1, t_2 = 0,$$

$$\text{Case 3: } t_1 = 0, t_2 > 0,$$

$$\text{Case 4: } 0 < t_1 < t_2$$

and refer to transitivity which was shown by Oja (1981, Theorem 5.1). According to equation (A), we have to show that

$$F_1^{-1}(F_2(u)) \quad \text{is convex for } 1/2 \leq u < 1$$

or, equivalently,

$$A(u) \equiv \frac{\partial F_1^{-1}(u)/\partial u}{\partial F_2^{-1}(u)/\partial u} \quad \text{is strictly monotone increasing for } 1/2 \leq u < 1.$$

Case 1: Assume $-\pi \leq t_1 < t_2 < 0$.

Preliminary remarks: If $-\pi \leq t_1 < t_2 < 0$ and $1/2 \leq u < 1$, then $t_1(u - 1/2) \in [-\pi/2, 0]$ and $t_2(u - 1/2) \in [-\pi/2, 0]$. Moreover, both $\sin(x)$ and $\cos(x)$ are strictly monotone increasing on $[-\pi/2, 0]$.

Part 1: Paraphrasing $A(u)$. Firstly,

$$\frac{\partial F_i^{-1}(u)}{\partial u} = t_i/c_2(t_i) \left[\cot(t_i u) + \cot(t_i(1 - u)) \right] \quad \text{for } 0 < u < 1, i = 1, 2.$$

Consequently,

$$A(u) = \frac{t_1/c_2(t_1)}{t_2/c_2(t_2)} \cdot \frac{\cot(t_1 u) + \cot(t_1(1 - u))}{\cot(t_2 u) + \cot(t_2(1 - u))}.$$

Using $\cot(\alpha) + \cot(\beta) = \frac{\sin(\alpha+\beta)}{\sin(\alpha)\sin(\beta)}$ (see Bronstein and Semendjajew, [2.5.2.1.1]),

$$A(u) = \frac{t_1/c_2(t_1)}{t_2/c_2(t_2)} \cdot \frac{\frac{\sin(t_1)}{\sin(t_1u)\sin(t_1(1-u))}}{\frac{\sin(t_2)}{\sin(t_2u)\sin(t_2(1-u))}} = \frac{t_1/c_2(t_1)\sin(t_1)}{t_2/c_2(t_2)\sin(t_2)} \cdot \frac{\sin(t_2u)\sin(t_2(1-u))}{\sin(t_1u)\sin(t_1(1-u))}.$$

Now, because $\pi \leq t_1 < t_2 < 0$, $\sin(t_1) < 0$ and $\sin(t_2) < 0$. Hence,

$$K(t_1, t_2) \equiv \frac{t_1/c_2(t_1)\sin(t_1)}{t_2/c_2(t_2)\sin(t_2)} > 0.$$

Finally, using $\sin(\alpha)\sin(\beta) = 1/2(\cos(\alpha-\beta) - \cos(\alpha+\beta))$ (see Bronstein and Semendjajew, [2.5.2.1.1]),

$$A(u) = K(t_1, t_2) \cdot \frac{\cos(2t_2(u-1/2)) - \cos(t_2)}{\cos(2t_1(u-1/2)) - \cos(t_1)} > 0,$$

because $\cos(x)$ is strictly monotone increasing for $x \in [-\pi, 0)$.

Part 2: Proof of the convexity. We have to show that $A(u)$ is strictly monotone increasing on $[1/2, 1)$. This is true, if $A'(u) > 0$ for $[1/2, 1)$:

$$A'(u) = \frac{K(t_1, t_2)}{N} \left\{ \left(-\sin(2t_2(u-1/2))2t_2 \right) \cdot \left(\cos(2t_1(u-1/2)) - \cos(t_1) \right) - \left(-\sin(2t_1(u-1/2))2t_1 \right) \cdot \left(\cos(2t_2(u-1/2)) - \cos(t_2) \right) \right\}$$

with $N \equiv [\cos(2t_1(u-1/2)) - \cos(t_1)]^2 > 0$. Using the monotony of the cosinus on $[-\pi, 0)$,

$$-2t_2(\cos(2t_1(u-1/2)) - \cos(t_1)) > 0 \quad \text{and} \quad -2t_1(\cos(2t_2(u-1/2)) - \cos(t_2)) > 0$$

for $1/2 \leq u < 1$ and $-\pi \leq t_1 < t_2 < 0$. Defining

$$K^*(t_1, t_2, u) \equiv \min \left\{ -2t_2(\cos(2t_1(u-0.5)) - \cos(t_1)); -2t_1(\cos(2t_2(u-0.5)) - \cos(t_2)) \right\},$$

we get

$$\begin{aligned} A'(u) &\geq \frac{K(t_1, t_2)K^*(t_1, t_2, u)}{N} \left(\sin(2t_2(u-1/2)) - \sin(2t_1(u-1/2)) \right) \\ &= \frac{K(t_1, t_2)K^*(t_1, t_2, u)}{N} \left(2\sin(t_2(u-1/2))\cos(t_2(u-1/2)) \right. \\ &\quad \left. - 2\sin(t_1(u-1/2))\cos(t_1(u-1/2)) \right), \end{aligned}$$

where we used $\sin(2\alpha) = 2\sin(\alpha)\cos(\alpha)$ (see Gradshteyn and Ryzhik, [1.333.1]). According to the preliminary remarks,

$$\sin(t_2(u-1/2))\cos(t_2(u-1/2)) - \sin(t_1(u-1/2))\cos(t_1(u-1/2)) > 0$$

for $-\pi \leq t_1 < t_2 < 0$ and $1/2 \leq u < 1$ implying that $A'(u) > 0$ for $1/2 \leq u < 1$.

Case 2: Assume $t_1 \in [-\pi, 0)$ and $t_2 = 0$.

The inverse distribution function of a GSH variable with $t_2 = 0$ is given by

$$F_2^{-1}(u) = \frac{\sqrt{3}}{\pi} \ln \left(\frac{u}{1-u} \right), \quad 0 < u < 1.$$

Consequently, for $0 < u < 1$,

$$\frac{\partial F_2^{-1}(u)}{\partial u} = \frac{\sqrt{3}}{\pi} \left(\frac{1}{u} + \frac{1}{1-u} \right) = \frac{\sqrt{3}}{\pi} \left(\frac{1}{u(1-u)} \right).$$

Thus, for $-\pi < t_2 < 0$ and $0 < u < 1$,

$$\begin{aligned} A(u) &= \frac{\partial F_1^{-1}(u)/\partial u}{\partial F_2^{-1}(u)/\partial u} = \frac{t_1/c_2(t_1)[\cot(t_1 u) + \cot(t_1(1-u))]}{\sqrt{3}/\pi \cdot 1/(u(1-u))} \\ &= \frac{t_1/c_2(t_1)}{\sqrt{3}/\pi} \frac{\sin(t_1)(u(1-u))}{\sin(t_1 u) \sin(t_1(1-u))} = \frac{\pi \sin(t_1)}{c_2(t_1)t_1\sqrt{3}} \cdot \frac{(t_1 u)(t_1(1-u))}{\sin(t_1 u) \sin(t_1(1-u))}. \end{aligned}$$

The first derivation is given by

$$\begin{aligned} A'(u) &= K(t_1) \left(\frac{t_1 \sin(t_1 u) - (t_1)^2 u \cos(t_1 u)}{\sin(t_1 u)^2} \cdot \frac{t_1(1-u)}{\sin(t_1(1-u))} \right. \\ &\quad \left. + \frac{-t_1 \sin(t_1(1-u)) + (t_1)^2(1-u) \cos(t_1(1-u))}{\sin(t_1(1-u))^2} \cdot \frac{t_1 u}{\sin(t_1 u)} \right) \\ &= \frac{K(t_1)t_1 t_1 u t_1(1-u)}{\sin(t_1 u) \sin(t_1(1-u))} \left(\left[\frac{1}{t_1 u} - \cot(t_1 u) \right] - \left[\frac{1}{t_1(1-u)} - \cot(t_1(1-u)) \right] \right) \end{aligned}$$

for $-\pi < t_1 < 0$ and $0 < u < 1$ with

$$K(t_1) \equiv \frac{\pi \sin(t_1)}{c_2(t_1)t_1\sqrt{3}} > 0.$$

Note that a series expansion to the $\cot(x)$ for $0 < |x| < \pi$ (see Gradshteyn and Ryzhik, [1.441.7]) is given by

$$\cot(x) = \frac{1}{x} - \sum_{i=1}^{\infty} \frac{2^{2i}|B_{2i}|}{(2i)!} x^{2i-1}, \quad (2)$$

where B_i , $i = 1, 2, \dots$, denotes the numbers of Bernoulli. Applying (2) for $x = t_1 u$ and $x = t_1(1-u)$,

$$A'(u) = K(t_1)t_1 \frac{t_1 u t_1(1-u)}{\sin(t_1 u) \sin(t_1(1-u))} \left(\sum_{i=1}^{\infty} \frac{2^{2i}|B_{2i}|}{(2i)!} (t_1 u)^{2i-1} - \sum_{i=1}^{\infty} \frac{2^{2i}|B_{2i}|}{(2i)!} (t_1(1-u))^{2i-1} \right).$$

For $-\pi < t_1 < 0$,

$$K(t_1)t_1 \frac{t_1 u t_1(1-u)}{\sin(t_1 u) \sin(t_1(1-u))} < 0.$$

The term in brackets is negative because $t_1(1-u) > t_1u$ and $(t_1(1-u))^{2i-1} > (t_1u)^{2i-1}$, $i = 1, 2, \dots$ for $1/2 < u < 1$ and $t_1 < 0$. Combining both results, $A'(u) > 0$ for $-\pi < t_1 < 0$ and $1/2 < u < 1$. This completes the proof of case 2.

Case 3: Assume $t_1 = 0$ and $t_2 > t_1$.

As calculated above,

$$\frac{\partial F_2^{-1}(u)}{\partial u} = \frac{t_2}{c_2(t_2)} \cdot \left[\coth(t_2u) + \coth(t_2(1-u)) \right], \quad 0 < u < 1,$$

with $c_2(t_2) = \sqrt{\frac{\pi^2 + t_2^2}{3}} > 0$ for $t_2 > 0$. It now follows that

$$\begin{aligned} A(u) &= \frac{\partial F_1^{-1}(u)/\partial u}{\partial F_2^{-1}(u)/\partial u} = \frac{\sqrt{3}/\pi}{t_2/c_2(t_2)} \cdot \frac{1/(u(1-u))}{\coth(t_2u) + \coth(t_2(1-u))} \\ &= \frac{\sqrt{3}t_2/\pi}{\sinh(t_2)/c_2(t_2)} \cdot \frac{\sinh(t_2u) \sinh(t_2(1-u))}{t_2ut_2(1-u)}. \end{aligned}$$

Defining

$$K(t_2) \equiv \frac{\sqrt{3}t_2/\pi}{\sinh(t_2)/c_2(t_2)} > 0 \quad \text{for } t_2 > 0,$$

then – analogue to case 2, but now using hyperbolic functions – for $t_2 > 0$,

$$\begin{aligned} A'(u) &= K(t_2)t_2 \cdot \frac{\sinh(t_2u) \sinh(t_2(1-u))}{t_2ut_2(1-u)} \cdot \left(\left[\coth(t_2u) - 1/(t_2u) \right] \right. \\ &\quad \left. - \left[\coth(t_2(1-u)) - 1/(t_2(1-u)) \right] \right) \end{aligned}$$

Now define $z(x) \equiv \coth(x) - 1/x$ which is strictly monotone increasing for $x > 0$, because by means of $\sinh(x) > x$ for $x > 0$,

$$z'(x) = -\frac{1}{(\sinh(x))^2} + \frac{1}{x^2} > 0.$$

Hence, from $t_2(1-u) < t_2u$ for $1/2 < u < 1$ and $t_2 > 0$ follows

$$\left[\coth(t_2u) - 1/(t_2u) \right] - \left[\coth(t_2(1-u)) - 1/(t_2(1-u)) \right] > 0,$$

This completes the proof of case 3.

Case 4: Assume $t_1 > 0$ and $t_2 > t_1$.

Similar to case 1 we have for $0 < u < 1$

$$A(u) = \frac{t_1/c_2(t_1)}{t_2/c_2(t_2)} \cdot \frac{\frac{\sinh(t_1)}{\sinh(t_1u) \sinh(t_1(1-u))}}{\frac{\sinh(t_2)}{\sinh(t_2u) \sinh(t_2(1-u))}} = \frac{t_1/c_2(t_1) \sinh(t_1)}{t_2/c_2(t_2) \sinh(t_2)} \cdot \frac{\sinh(t_2u) \sinh(t_2(1-u))}{\sinh(t_1u) \sinh(t_1(1-u))}.$$

Defining

$$K(t_1, t_2) \equiv \frac{t_1/c_2(t_1) \sinh(t_1)}{t_2/c_2(t_2) \sinh(t_2)} > 0,$$

the first derivative of A is given by

$$A'(u) = K(t_1, t_2) \cdot \frac{\sinh(t_2u) \sinh(t_2(1-u))}{\sinh(t_1u) \sinh(t_1(1-u))} \cdot \left[t_2(\coth(t_2u) - \coth(t_2(1-u))) - t_1(\coth(t_1u) - \coth(t_1(1-u))) \right].$$

Now,

$$z(t) = \coth(tu) - \coth(t(1-u)) \text{ is strictly monotone increasing for } t > 0, \quad (3)$$

if the first derivative $z'(t)$ is positive: For $t > 0$, $1/2 < u < 1$,

$$\begin{aligned} \frac{1-u}{[\sinh(t(1-u))]^2} - \frac{u}{[\sinh(tu)]^2} &> 0 && \iff \\ \frac{t(1-u)}{[\sinh(t(1-u))]^2} - \frac{tu}{[\sinh(tu)]^2} &> 0 && \iff \\ 1 - \frac{tu}{t(1-u)} \frac{[\sinh(t(1-u))]^2}{[\sinh(tu)]^2} &> 0 && \iff \frac{\frac{[\sinh(t(1-u))]^2}{t(1-u)}}{\frac{[\sinh(tu)]^2}{tu}} < 1. \end{aligned}$$

To prove the last inequality, we show that $f(x) = \frac{[\sinh(x)]^2}{x}$ is monotone increasing for $x > 0$. Using $x \geq \tanh(x)$,

$$\begin{aligned} f'(x) &= \frac{2 \sinh(x) \cosh(x)x - [\sinh(x)]^2}{x^2} = \frac{[\cosh(x)]^2}{x^2} (2 \tanh(x)x - [\tanh(x)]^2) \\ &\geq \frac{[\cosh(x)]^2}{x^2} \tanh(x)x \geq 0. \end{aligned}$$

From equation (3) follows that

$$\coth(t_2u) - \coth(t_2(1-u)) > \coth(t_1u) - \coth(t_1(1-u))$$

and

$$t_2 \left[\coth(t_2u) - \coth(t_2(1-u)) \right] - t_1 \left[\coth(t_1u) - \coth(t_1(1-u)) \right] > 0.$$

Consequently, $A'(u) > 0$ for $1/2 < u < 1$. \square

References

- [1] Balanda, K. P. and H. L. MacGillivray: *Kurtosis and spread*. The Canadian Journal of Statistics, **18**(1):17-30, 1990.
- [2] Baten, W. D.: *The Probability Law for the Sum of n Independent Variables, each Subject to the Law $(1/2h)\text{sech}(\pi x/2h)$* . Bulletin of the American Mathematical Society, **40**:284-290, 1934.
- [3] Bronstein, I. N. and K. A. Semendjajew: *Taschenbuch der Mathematik*. Verlag Harri Deutsch, Frankfurt(Main), 1980.
- [4] Gradshteyn, I. S. and I. M. Ryzhik: *Table of Integrals, Series, and Products*. Academic Press, New York, 2000.
- [5] Harkness, W. L. and M. L. Harkness: *Generalized Hyperbolic Secant Distributions*. Journal of the American Statistical Association, **63**:329-337, 1968.
- [6] Oja, H.: *On location, scale, skewness and kurtosis of univariate distributions*. Scandinavian Journal of Statistics **8**, 154–168.
- [7] Van Zwet, W. R.: *Convex Transformations of Random Variables*. Mathematical Centre Tracts No. 7. Mathematical Centre, Amsterdam, 1964.
- [8] Vaughan, D. C.: *The Generalized Hyperbolic Secant distribution and its Application*. Communications in Statistics – Theory and Methods, **31**(2):219-238, 2002.

Adress of the authors:

Prof. Dr. Ingo Klein
Lehrstuhl für Statistik und Ökonometrie
Universität Erlangen-Nürnberg
Lange Gasse 20
D-90403 Nürnberg
Tel. +60 911 5320271
Fax +60 911 5320277
Elec. Mail: Ingo.Klein@wiso.uni-erlangen.de
<http://www.statistik.wiso.uni-erlangen.de>

Dr. Matthias Fischer
Lehrstuhl für Statistik und Ökonometrie
Universität Erlangen-Nürnberg
Lange Gasse 20
D-90403 Nürnberg
Tel. +60 911 5320271
Fax +60 911 5320277
Elec. Mail: Matthias.Fischer@wiso.uni-erlangen.de
<http://www.statistik.wiso.uni-erlangen.de>