

ENHANCING PRODUCTIVITY OF WATER UNDER VARIABLE CLIMATE

K.P.C. Rao ICRISAT – Nairobi, P O Box 39063, Nairobi
Tel (Off) 020 524192
Mobile: 0721409239
Email: k.p.rao@cgiar.org

G.E. Okwach, KARI Katumani Research Centre, P.O. Box 340 Machakos
Phone / Fax: 044 21122
Mobile: 0733787401
Email: geokwach@africaonline.co.ke

Abstract

efficient use deserves special attention in our efforts to increase the productivity and profitability of agriculture in these areas. Rainfall in these regions is highly variable and the risk associated with such variable weather acts as a major deterrent for the farmers to invest in expensive inputs such as fertilizers and improved seeds required to achieve higher productivity. Availability of skilful forecasts from the growing understanding and advances in modeling the global climate system have opened up new opportunities for farmers to consider a number of adjustments to the management practices based on seasonal conditions predicted for the forth-coming season.

Machakos district in Kenya is characterized as hot and dry with bimodal distribution of rainfall. Annual rainfall at Katumani ranged between 330 and 1260 mm with a coefficient of variation of 28%. Average seasonal rainfall is less than 300 mm with more than 40% of the seasons receiving less than 250 mm. Maize is the main food crop in the district with an average productivity of 0.8 t ha⁻¹. However, average maize yields have declined by nearly 50% to 0.4 t/ha during the decade 1993-2002 mainly due to the adoption of low input management techniques. Further analysis of yield trends confirmed that the farmers' strategy is well suited for the below normal seasons but failed to capitalize on the good seasonal conditions during normal and above normal seasons.

The reliability of hindcasts generated by IRI for 43 SR seasons starting from 1961 was evaluated and their potential value in reducing risk and improving productivity and profitability was assessed for Katumani situation using crop simulation model APSIM. Though the available skill in forecasts is not sufficient to predict accurately the amount of rainfall or its distribution in advance, it is possible to predict with some certainty whether the coming season is going to be below normal or not. Model simulations indicated significant gains in productivity and profitability with simple adjustments identified by the farmer such as application of recommended dose of fertilizer and high plant population in the seasons forecasted as normal to above, and low risk farmer strategies during the years forecasted as dry years.

Introduction

Water is the most limiting factor for crop production in the semi-arid tropics (SAT) and its efficient use deserves special attention in our efforts to increase the productivity and profitability of agriculture in these areas. Rainfall, the only source of water shows high temporal and spatial variability and the risk associated with such variable weather acts as a major deterrent for the farmers to invest in expensive inputs such as fertilizers and improved seeds required to achieve higher productivity. Farmers, particularly smallholders in developing countries, show risk-averse behavior (Binswanger, 1980) and adopt conservative management strategies that reduce negative impacts in poor years, but at the expense of reduced average productivity and profitability (Rosenzweig and Binswanger, 1993; Zimmerman and Carter, 2003). According to IPCC (2001), global changes in climate are

expected to further exacerbate the variability. This means that farmers will have to deal with more uncertain weather, with extreme events occurring more frequently. If deliberate attention is not directed to managing impacts of climate variability, majority poor farmers especially in semi-arid areas will face higher insecurity in food and incomes.

Much of the past research on managing climate variability has been devoted to the analysis and understanding of the complexities associated with the variability and distribution of rainfall (Sivakumar, et al. 1983, Janowiak 1988, and Hulme, 1992). However, many critical agricultural decisions must be made several months before impacts of climate are realized, making it difficult to tailor the management to the seasons potential. The Response Farming technique tried in Kenya in the late 80's was an attempt to predict the rainy season potential and adjust farming practices to the prevailing environmental conditions (Stewart and Faught, 1984), but met with limited success due to difficulties in assessing the season's potential. Other risk management strategies developed include maintaining storage reserves, diversifying production, insurance, forward selling, futures trading, government subsidies and taxation incentives have been developed (Kurukulasuriya and Rosenthal, 2003). However, adoption of these interventions requires good institutional and policy support which is limiting in many developing countries in general and in Africa in particular.

Recent developments in the understanding of interactions between the atmosphere, sea and land surfaces, and in modeling the global climate system made it possible to predict climatic conditions months in advance in many parts of the world (Goddard et al., 2001, Mutai et al., 1998; Indeje et al., 2000; Hansen and Indeje, 2003). Some efforts were also made to use seasonal climate forecasts in disaster preparedness by agencies such as FEWSNET but use of seasonal climate forecasts in farm level decision making is minimal in the region.

In this paper we present the results of a case study conducted to assess the potential value of seasonal climate forecasts in reducing risk and improving productivity and profitability of small holder farms in Machakos district, Kenya. The case study is based on Machakos district crop production data, results of a farmer survey conducted in Mwala division of Machakos district, data from a long-term trial conducted at Katumani research station and results of system simulation analysis using crop simulation model APSIM (McCown et al., 1996).

Climate variability and crop production in Machakos

The district Machakos is generally characterized as hot and dry with bimodal distribution of rainfall. Throughout the district rainfall is subject to pronounced variability from year to year and breaks in rain occur often and any time during the rainy season. Long-term rainfall data for the period 1957-2003 recorded at Katumani research station was analyzed to get a good understanding of the variability in frequency and distribution of seasonal rainfall. Annual rainfall at Katumani ranged between 330 and 1260 mm with a coefficient of variation of 28%. Nearly 85% of the average annual rainfall is received during the two cropping seasons, long rains (LR) between March and May and short rains (SR) between October and December. Though both SR and LR seasons receive similar amounts of rainfall, SR seasons are more reliable than the LR seasons and therefore more important for crop production. With an average seasonal rainfall of less than 300 mm and a coefficient of variation more than 40% the district is considered as marginal area for maize production (Dowker, 1961). About 40% of all seasons received less than 250 mm rainfall while 27% recorded rainfall in excess of 350 mm (Table 1). The average seasonal rainfall of below normal SR and LR seasons is about a third of that received during the above normal seasons. The big difference in the seasonal rainfall presents different opportunities and challenges for the management to tailor crop mix and/or management practices such that the seasonal potentials are realized and risks are minimized.

Since agriculture in the district is predominantly rainfed, maize yield trends are closely related to trends in rainfall (Figure 1). Long-term average yield of maize in the district is 0.8 t/ha. However, since 1990 a strong declining trend was observed in the maize productivity resulting in steep fall in maize yields. Average maize yields declined by nearly 50% to 0.4

t/ha during the decade 1993-2002. Similar declining trend in maize yields during the same period was also observed at the national level. Further analysis of district level information indicated that much of this decline is coming from the districts having high percentage of medium and low potential areas primarily located in semi-arid and arid environments. The two major factors contributing to the observed decline in yields could be declining soil fertility as a result of non-application of fertilizers and extension of agriculture into more marginal areas. Because fertilizers are expensive and the risk of losing on investment is very high, farmers in these environments tend not to apply fertilizers. At the same time increasing population and limited availability of good agricultural land is pushing agriculture into more marginal lands and environments where the need for external inputs and risks of crop failure are high.

District level production data for maize was also analyzed for trends in crop productivity during various seasons classified as below normal (< 250 mm), normal (250 – 350 mm), and above normal (> 350 mm) and results are presented in Figure 2. It is interesting to note that maize yields during the years in which both LR and SR seasons received above normal rainfall are lower than the yields recorded during the years in which both seasons received below normal rainfall. Productivity of maize per mm of rainfall followed a similar trend except that the productivity when both seasons were below normal is higher than that during any other year (Figure 3). The average productivity achieved is 2.9 kg maize grain mm⁻¹ of rain. The productivity during wettest years is about 1.2 kg maize grain mm⁻¹ while during the normal years it is about 3 kg maize grain mm⁻¹. Since loss of rain water through runoff and erosion is high during the wet years, we tried to estimate the productivity using effective rainfall (rainfall-runoff-drainage). We have estimated the effective rainfall using system simulation model APSIM which was earlier calibrated and validated for the Katumani location by (Okwach and Simiyu, 1999; Okwach, 2002). Productivity of rainwater when based on effective rainfall increased to 4.2 kg maize grain mm⁻¹ which is nearly 45% higher than that observed with total seasonal rainfall. However, there is no change in the observed trend.

The observed productivity of rain water is very similar to that recorded under low input system with 22,000 plants ha⁻¹ and no fertilizer application in a long-term trial conducted at Katumani research station between 1990 and 1999. This treatment is very similar to what farmers normally do on their farms. In the same trial, average productivity of effective rainfall is more than doubled when plant population is increased to 53,000 maize plants ha⁻¹ and urea fertilizer equivalent to 70 kg N ha⁻¹ was applied. Data from a long-term trial was also analyzed to identify trends in maize yields in different seasons. Of the total 20 seasons over which the trial was conducted, eight seasons were below normal, five were normal and the remaining six seasons were above normal rainfall. The productivity during above normal years with or without moisture conservation through application of mulch is less than that during normal years but higher than that in the below normal seasons when no fertilizer was applied. Fertilizer application increased the yields significantly in all seasons but the increase is more in normal and above normal seasons (Figure 4). During the normal and above normal seasons application of fertilizer resulted in a gain of 1 t maize grain ha⁻¹. However, application of fertilizer is profitable in only three of the eight below normal seasons. Crop completely failed during the LR season of 1993 and no increase in yield due to fertilizer application was observed during 1998 LR season and 1996 SR season. This risk of losing under unfavorable seasonal conditions is the major constraint in farmers using fertilizers.

Seasonal climate forecasts and their reliability

Farmers would be able to consider a number of adjustments in the management practices used if they had prior knowledge of what the rainfall conditions are going to be during the forthcoming season. One way of having advance information about the forthcoming season is through use of long-term/seasonal climate forecasts made by institutions such as International Research Institute for Climate Prediction (IRI) and ICPAC (IGAD Climate Prediction and Application Centre formerly Drought Monitoring Centre). Since 1998, ICPAC in collaboration with several international climate centers is providing seasonal climate

outlooks through its regional climate outlook forums and IRI has the capability to develop hindcasts using GCM SST data.

The reliability of hindcasts generated by IRI for 43 SR seasons starting from 1961 was evaluated by comparing the predicted with the observed seasonal conditions (Table 2). The predicted and observed rainfall amounts correlated poorly with a coefficient of determination (R^2) of 0.336 which shows that existing skill in predicting the amount of rainfall is not very high. We then looked into the type of season by classifying the season using the criteria described earlier and amount of rainfall hindcasted. While a total of thirteen seasons were predicted to receive below normal rainfall, the prediction was turned out to be true in ten seasons or in 77% of the instances. The predictability of normal seasons is better with an accuracy of 84%. The predictability of above normal seasons is least amongst the three groups with prediction coming true in 55% years. However, none of the seasons predicted to receive normal or above normal rainfall were turned out to be below normal rainfall.

During a one day workshop with farmers at Mwala, farmers were asked to assess the reliability of these hindcasts by comparing them with actual rainfall recorded at Katumani research station and their own experiences. According to the farmer assessment, 32 of the 43 predictions (about 74 %) are extremely good and use of these forecasts in farm management can result in substantial productivity gains during wet years and in minimizing losses during dry years. Farmers ranked eight predictions as good during which the gap between predicted and observed rainfall amounts is high but both observed and predicted rainfall amounts are more than that required for harvesting a good crop.

Management decisions that can be influenced by forecasts and potential benefits

During the workshop, farmers were also asked to identify how this information would benefit them and what adjustments they would like to make using the existing skill in the forecasts. Farmers felt that the forecasts are extremely good in identifying whether the forthcoming season is going to be below normal or normal to above normal. It is only in three out of the 43 seasons that the predictions went wrong and all three of them are under predictions. According to the farmers, the possibility of making loss from under predictions is less than that from over predictions. Hence, they did not consider this as a constraint.

Farmers then identified a number of management decisions that can be made using the existing skill in the forecasts (Table 3). Farmer response to forecast based decision-making indicates that they clearly understood the variability in seasonal rainfall and the potential role forecasts can play in improving management of their farms. The management practices identified, for example, the need to plant drought tolerant or drought escaping crop varieties if the forecast indicate a dry season and increasing manure and fertilizer inputs when the forecast is wet season is a clear demonstration that small holder farmer can make tactical decisions if the required information is made available.

Potential benefits from the changed decisions

Using the system simulation model APSIM, a scenario analysis was conducted to estimate the potential benefit from the adjustments identified by the farmers based on the hindcasted seasonal conditions. The management options simulated include application of 30, 40, and 60 kg nitrogen ha^{-1} with a density of 35,000 maize plants ha^{-1} during normal to above normal seasons; and no fertilizer and 22,000 maize plants ha^{-1} during below normal seasons. Though the adjustments were made only to the SR seasons that were predicted to be normal or above normal, significant gains were also observed during the below normal years when low input farmer practice is used. This is the spill over benefit coming from the residual effect of the adjustments made during the other seasons. The simulation analysis has clearly indicated that the forecast based farming can result in an overall average gain ranging from 139-251% when adjustments were made only to those years predicted to be either normal or above normal (Table 4).

Summary and conclusions

Given the high variation in seasonal rainfall and the need to plan farm operations without knowing the seasonal conditions, farmers in semi-arid regions generally favor using low risk conservative management strategies which do not capitalize on the opportunities created by the better seasonal conditions during the normal and above normal seasons. Farmers would be able to consider a number of adjustments to the management practices used if they had prior knowledge of what the rainfall conditions are going to be during the forth-coming season. This study has highlighted the benefits that can be derived by these adjustments.

For use in farm level decision making, the forecasts should preferably give information about the expected amount and distribution of rainfall. However, with the current understanding of climatic anomalies and the factors contributing to them it is still not possible to predict accurately the amount of rainfall or its distribution in advance. But the existing skill is good enough to predict with some certainty whether the coming season is going to be below normal or not. This in itself is an important piece of information from which significant benefits can be derived. As indicated by the simulation analysis, there is a potential to increase the yields by 2 to 3 times through adoption of simple adjustments to the management involving very low levels of risk under variable climatic conditions.

Use of climate information and seasonal climate forecasts are not systematically explored. While the ability to forecast weather events has increased and is expected improve further, our ability to transmit this information to the end user in the form that can be utilized by them is not yet developed. The approach presented in this paper not only helps in coping with current climate variability, but has the potential to serve as an adaptation strategy to long term climate change.

Table 1: Average seasonal rainfall (mm) recorded at Katumani (1957-2003) during short and long rain seasons

Seasons with Rainfall	Short Rains (Oct-Dec)			Long Rains (Mar-May)		
	Average rain (mm)	No of years	CV (%)	Average rain (mm)	No of years	CV (%)
<250 mm	190	22	20.5	151	17	32.8
250-350 mm	300	15	10.9	293	14	10.9
>350 mm	507	10	29.3	415	16	12.9
All years	292	47	48.8	283	47	42.6

Table 2: Observed and predicted short rain season types at Katumani

Rainfall class	Predicted	Observed BN	Observed N	Observed AN	Avg. RF Predicted	Avg. RF Observed
Below normal (BN) <250 mm	13	10	2	1	207	221
Normal (N) 250-300 mm	19	0	16	3	299	287
Above normal (AN) >350 mm	11	0	5	6	437	437

Table 3: Some farmer identified management options for below normal and normal to above normal seasons.

Management decisions	
Dry season	Normal to wet season
1. Use low plant density (2.2 plants/m ²)	1. Use higher plant density (3.5 to 4.5 plants/m ²)
2. Reduce labor and other input use	2. Apply fertilizer
3. Increased use of drought tolerant crops such sorghum, millet, green grams, and cassava	3. Plant hybrid maize varieties such as pioneer
4. Plough and plant early before the start of the rain	4. Adopt intercropping
5. Adopt water conservation measures	5. Strengthen terraces
6. Reduce area under cultivation	6. Increase area under cultivation

Table 4: Expected gain in maize yield (kg/ha) with forecast based adjustments to SR seasons predicted to receive normal to above normal rainfall

Type of season	Farmer practice	Forecast based farming with 35,000 plants ha ⁻¹ and		
		30 kg N ha ⁻¹	40 kg N ha ⁻¹	60 kg N ha ⁻¹
Dry	555	951 (71)	1052 (90)	1206 (117)
Normal to wet	666	1879 (182)	2286 (243)	2822 (323)
All	613	1467 (139)	1747 (185)	2151 (251)

(Figures in parenthesis indicate percent gain)




Figure 1: Rainfall and maize yields in Machakos district




Figure 2: Productivity of Maize (t ha⁻¹) during below normal (<250 mm), normal (250-350 mm), and above normal (>350 mm) short and long rains seasons for the period 1970-2002




Figure 3. Productivity of total and effective rainfall (kg maize grain/ mm rain) during below normal (<250 mm), normal (250-350 mm) and above normal (>350 mm) short and long rains seasons for the period 1970-2002




Figure 4. Average maize yields (kg/ha) recorded under farmer practice, water conservation by mulching (WC) and with WC and application of 70 Kg N/ha treatments during below normal (<250 mm), normal (250-350 mm) and above normal (>350 mm) crop seasons between 1990 and 1999

Literature cited

- Binswanger, H., (1980). Attitudes towards risk: experimental measures in rural India. *Am. J. Agric. Econ.* 62:395-470.
- Dowkwer, B.D. (1961). Rainfall variability and maize yield in Machakos. *E Afr. Agric. For. J.* 28:134-138.
- Goddard, L., Mason, S.J., Zebiak, S.E., Ropelewski, C.F., Basher, R., Cane, M.A. (2001). Current Approaches to Seasonal to Interannual Climate Predictions. *International Journal of Climatology.* 21: 1111-1152
- Hansen, J.W., and Indeje, M. (2003). Linking dynamic seasonal climate forecasts with crop simulation for maize yield prediction in semi-arid Kenya. *Agricultural and Forest meteorology.*
- Hulme, M. (1992). Rainfall changes in Africa: 1931-60 to 1961-90 *Int. J. Climatol.* 12: 685-699
- Indeje, M., Semazzi, F.H.M., Ogallo, L.J. (2000). ENSO signals in East African rainfall and their prediction potentials. *Int. J. Climatol.* 20:19-46.
- IPCC (2001). *Climate Change 2001: Impacts, Adaptations and Vulnerability. Report of working group II.*
- Janowiak, J.E. (1988). An investigation of interannual rainfall variability in Africa *J. Climate.* 1:240-255.
- Kurukulasuriya, P. and Rosenthal, S. (2003). *Climate Change and Agriculture: A Review of Impacts and Adaptations World bank Climate change series Paper no 91. Paper prepared and published for the Rural Development Group and Environment Department of the World Bank. Pp106.*
- McCown, R. L., Hammer, G. L., Hargreaves, J. N. G., Holzworth, D. and Freebairn, D. M. (1996). APSIM: A novel software system for model development, model testing, and simulation in agricultural systems research. *Agricultural Systems* 50: 255-271.
- Mutai, C.C., Ward, M.N., Coleman, A.W. (1998). Towards the prediction of the East Africa short rains based on sea-surface temperature-atmosphere coupling. *Int. J. Climatol.* 18: 975-997.
- Okwach, G.E. (2002). Developing an appropriate tool for managing runoff, erosion, and crop production in semi-arid eastern Kenya: A Case for APSIM model. CARMASAK technical report No. 3. Kenya Agricultural Research Institute.
- Okwach, G.E. and Simiyu, C.S. (1999). *Effects of Land Management on Runoff, Erosion, and Crop Production in a semi-arid area of Kenya. East African Agricultural and Forestry Journal Special Edition* 65: 125-142.
- Rosenzweig, M.R., and Binswanger, H.P., (1993). Wealth, weather risk and the composition and profitability of agricultural investments. *The Economic Journal* 103:56-78.
- Sivakumar, M. V. K., Singh, P. and Williams, J. H. (1983). Agroclimatic aspects in planning for improved productivity of Alfisols. In: *Alfisols in the semi-arid tropics: a consultants' workshop, 1-3 Dec. 1983, ICRISAT Centre, India, pp 15-30.*

Stewart, J.I and Faught, W.A., (1984). Response farming of maize and beans at Katumani, Machakos district, Kenya: Recommendations, Yield expectations, and economic benefits. *East African Agricultural and Forestry Journal*. 44:29-51

Zimmerman, F.J., and Carter, M.R., (2003). Asset smoothing, consumption smoothing and the reproduction of inequality under risk and subsistence constraints. *J. Development Economics* 71:233-260.