

Der Open-Access-Publikationsserver der ZBW – Leibniz-Informationzentrum Wirtschaft  
*The Open Access Publication Server of the ZBW – Leibniz Information Centre for Economics*

Dette, Holger; Melas, Viatcheslav B.

Working Paper

## A note on some extremal problems for trigonometric polynomials

Technical Report / Universität Dortmund, SFB 475 Komplexitätsreduktion in Multivariaten Datenstrukturen, No. 2005,20

**Provided in cooperation with:**  
Technische Universität Dortmund

Suggested citation: Dette, Holger; Melas, Viatcheslav B. (2005) : A note on some extremal problems for trigonometric polynomials, Technical Report / Universität Dortmund, SFB 475 Komplexitätsreduktion in Multivariaten Datenstrukturen, No. 2005,20, <http://hdl.handle.net/10419/22611>

**Nutzungsbedingungen:**

Die ZBW räumt Ihnen als Nutzerin/Nutzer das unentgeltliche, räumlich unbeschränkte und zeitlich auf die Dauer des Schutzrechts beschränkte einfache Recht ein, das ausgewählte Werk im Rahmen der unter

→ <http://www.econstor.eu/dspace/Nutzungsbedingungen> nachzulesenden vollständigen Nutzungsbedingungen zu vervielfältigen, mit denen die Nutzerin/der Nutzer sich durch die erste Nutzung einverstanden erklärt.

**Terms of use:**

*The ZBW grants you, the user, the non-exclusive right to use the selected work free of charge, territorially unrestricted and within the time limit of the term of the property rights according to the terms specified at*

→ <http://www.econstor.eu/dspace/Nutzungsbedingungen>  
*By the first use of the selected work the user agrees and declares to comply with these terms of use.*

# A note on some extremal problems for trigonometric polynomials

Holger Dette

Ruhr-Universität Bochum  
Fakultät für Mathematik  
44780 Bochum, Germany  
e-mail: holger.dette@rub.de  
FAX: +49 234 3214 559

Viatcheslav B. Melas

St. Petersburg State University  
Department of Mathematics  
St. Petersburg  
Russia  
email: v.melas@pobox.spbu.ru

July 6, 2005

## Abstract

We consider the problem of finding the trigonometric polynomial

$$\vartheta_0 + \sum_{j=1}^m \vartheta_{2j-1} \sin(jx) + \vartheta_{2j} \cos(jx)$$

with minimal sup-norm on the interval  $[-\pi, \pi]$ , where one coefficient in the polynomial, say  $\vartheta_k$  ( $0 \leq k \leq 2m$ ), has been fixed. A complete solution of this problem is given, which depends sensitively on the ratio  $k/m$ , and the problem of uniqueness is discussed. Several examples are presented to illustrate the main properties of the extremal polynomials.

AMS Subject Classification: 42A10, 42A05

Keywords and Phrases: Trigonometric polynomial, extremal polynomial, Chebyshev system, extremal problem

## 1 Introduction

Consider the class

$$(1.1) \mathcal{T} = \left\{ \varphi_m(x) = \vartheta_0 + \sum_{j=1}^m \vartheta_{2j-1} \sin(jx) + \vartheta_{2j} \cos(jx) \mid x \in [-\pi, \pi]; \vartheta_i \in \mathbb{R}, \quad i = 0, \dots, 2m \right\}$$

of all trigonometric polynomials of degree  $m$ , and let

$$\|f\|_\infty = \max_{|x| \leq \pi} |f(x)|$$

denote the sup-norm of a continuous function on the interval  $[-\pi, \pi]$ . In the present paper we derive an analytic solution of the extremal problem

$$(1.2) \quad \min\{\|\varphi_m\|_\infty \mid \varphi_m \in \mathcal{T}, |\vartheta_k| = 1\};$$

where  $k \in \{0, \dots, 2m\}$  is arbitrary but fixed. Extremal problems of this type have been originally studied for ordinary polynomials by Chebyshev (1859), who determined the minimal sup-norm of all polynomials with a fixed degree and leading coefficient 1 [see e.g. Achieser (1956) or Natanson (1955) among others]. In the context of algebraic and trigonometric polynomials numerous extremal problems have been studied in the literature [see Rivlin (1974) or Milavonović, Mitrinović and Rassias (1994) among many others] but to the knowledge of the authors a solution of the above problem is not available. Note that the extremal problem (1.2) can be reformulated to the problem of finding the best constant, say  $c_{m,\infty}$ , such that the inequality

$$(1.3) \quad 1 = |\vartheta_k| \leq c_{k,m,\infty} \|\varphi_m\|_\infty \quad \forall \varphi_m \in \mathcal{T}$$

holds. This problem is related to the problem of determining the best constant  $c_{m,p}$  such that

$$(1.4) \quad \max_{j=0}^{2m} |\vartheta_j| \leq c_{m,p} \|\vartheta_m\|_p \quad \forall \varphi_m \in \mathcal{T},$$

(here  $\|f\|_p = (\int_{-\pi}^{\pi} |f(x)|^p dx)^{1/p}$  denotes the  $L^p$ -norm), which was recently solved by Marshall and Ganzburg (1999). Similar problems for the  $L_1$ -norm of nonnegative trigonometric polynomials were studied by Peherstorfer (1982). We finally note that the extremal problem (1.2) is also equivalent to the problem

$$(1.5) \quad \max\{|\vartheta_k| \mid \varphi_m \in \mathcal{T}, \|\varphi_m\|_\infty \leq 1\}$$

in the following sense. If  $\varphi_m$  is a solution of the extremal problem (1.2), then  $\varphi_m/\|\varphi_m\|_\infty$  is a solution of the extremal problem (1.5) and vice versa. It is also easy to see that  $-\varphi_m$  is a solution of the extremal problem (1.2) [or equivalently (1.3) or (1.5)] if and only if  $\varphi_m$  is a solution, and throughout this paper we do not distinguish these two solutions.

It is the purpose of the present note to derive an explicit solution of the extremal problem (1.2), [or equivalently (1.3), (1.5)]. In Section 2 we provide a necessary and sufficient condition for a trigonometric polynomial to be a solution of the extremal problem. It turns out that this solution depends sensitively on the value of  $k \in \{0, \dots, 2m\}$ , and two different scenarios are identified in Section 3 and Section 4 corresponding to the cases  $\lfloor \frac{k+1}{2} \rfloor \leq m/3$  and  $\lfloor \frac{k+1}{2} \rfloor > m/3$ , respectively. Finally, in Section 5 the problem of uniqueness is discussed and it is demonstrated that the extremal problem (1.2) can have uncountable many solutions. Moreover, several examples are presented, which illustrate some peculiar properties of the extremal polynomials in the case  $\lfloor \frac{k+1}{2} \rfloor < m/3$ .

## 2 A necessary and sufficient condition for the solution of the extremal problem

For the sake of a transparent notation we introduce the functions

$$(2.1) \quad f_j(x) = \begin{cases} 1 & \text{if } j = 0 \\ \sin(\ell x) & \text{if } j = 2\ell - 1; \quad \ell = 1, \dots, m \\ \cos(\ell x) & \text{if } j = 2\ell; \quad \ell = 1, \dots, m \end{cases}$$

and rewrite a trigonometric polynomial  $\varphi_m \in \mathcal{T}$  as

$$(2.2) \quad \varphi_m(x) = \sum_{j=0}^{2m} \vartheta_j f_j(x).$$

The following result characterizes the solution of the extremal problem (1.5).

**Theorem 2.1.** *A trigonometric polynomial of the form (2.2) with  $\|\varphi_m\|_\infty \leq 1$  is a solution of the extremal problem (1.5) if and only if there exist points  $x_1, \dots, x_N \in [-\pi, \pi]$  and real constants  $A_1, \dots, A_N$  ( $N \in \mathbb{N}$ ) such that  $N \leq 2m$  and*

$$(i) \quad \text{sign}(A_j)\varphi_m(x_j) = 1 \quad \text{whenever } A_j \neq 0 \quad (j = 1, \dots, N);$$

$$(ii) \quad \sum_{j=1}^N A_j f_i(x_j) = 0; \quad \text{for all } i \in \{0, 1, \dots, 2m\} \setminus \{k\};$$

$$(iii) \quad \sum_{j=1}^N A_j f_k(x_j) = 1.$$

Moreover, if  $\hat{\varphi}_m = \sum_{j=0}^{2m} \hat{\vartheta}_j f_j$  is a solution of the extremal problem (1.5) the coefficient  $\hat{\vartheta}_k$  corresponding to the function  $f_k$  in the polynomial  $\hat{\varphi}_m$  is given by

$$(2.3) \quad \hat{\vartheta}_k = \sum_{j=1}^N |A_j|.$$

**Proof.** Using Theorem 1.3 in Singer (1970) it follows that a trigonometric polynomial  $\tilde{\varphi}_m$  is a solution of the extremal problem (1.2) if and only if there exist  $N \leq 2m$  points  $x_1, \dots, x_N$  and  $N$  scalars  $\mu_1, \dots, \mu_N$  with  $\sum_{i=1}^N |\mu_i| = 1$  such that

$$(2.4) \quad \sum_{j=1}^N \mu_j f_i(x_j) = 0 \quad i \in \{0, \dots, 2m\} \setminus \{k\}$$

$$(2.5) \quad \sum_{j=1}^N \mu_j \tilde{\varphi}_m(x_j) = \|\tilde{\varphi}_m\|_\infty$$

$$(2.6) \quad \tilde{\varphi}_m(x_j) \text{sign}(\mu_j) = \|\tilde{\varphi}_m\|_\infty \quad j \in \{0, \dots, N\}$$

Define  $A_j = \mu_j / \|\tilde{\varphi}_m\|_\infty$ , note that  $\tilde{\varphi}_m(x) = f_k(x) + \sum_{i \neq k} \vartheta_i f_i(x)$  and define  $\hat{\varphi}_m(x) = \tilde{\varphi}_m(x) / \|\tilde{\varphi}_m\|_\infty$ . Because the extremal problems (1.2) and (1.5) are equivalent, it is now easy to see that the trigonometric polynomial  $\hat{\varphi}_m$  is a solution of the extremal problem (1.5), if and only if there exist points  $x_1, \dots, x_N$  and constants  $A_1, \dots, A_N$  such that the conditions (i) – (iii) of Theorem 2.1 are satisfied. Finally, we obtain from the conditions (i) – (iii) the representation

$$1 = \sum_{j=1}^N A_j f_k(x_j) = \frac{1}{\hat{\vartheta}_k} \sum_{j=1}^N A_j \hat{\varphi}_m(x_j) = \frac{1}{\hat{\vartheta}_k} \sum_{j=1}^N |A_j|,$$

which proves the remaining assertion (2.3) of Theorem 2.1.  $\square$

### 3 Extremal polynomials in the case $k = 0, m/3 < \lfloor \frac{k+1}{2} \rfloor \leq m$

In this section we provide a solution of the extremal problem (1.2) for the case  $k = 0$  and “large” values of  $k$ . It turns out that in this case the solution is very transparent.

**Theorem 3.1.** *Assume that  $k = 0$  or  $m/3 < \lfloor \frac{k+1}{2} \rfloor \leq m$ , then the trigonometric polynomial  $\hat{\varphi}_m(x) = f_k(x)$  is a solution of the extremal problem (1.2). In particular the constant  $c_{m,\infty}$  in (1.3) is given by  $c_{m,\infty} = 1$  if  $k = 0$  or  $m/3 < \lfloor \frac{k+1}{2} \rfloor \leq m$ .*

**Proof.** Consider the case  $k = 2\ell$  with  $\ell > m/3$ , for which we have to show that the solution of the extremal problem is given by  $\hat{\varphi}_m(x) = f_k(x) = \cos(\ell x)$ . Obviously, we have  $\|\hat{\varphi}_m\|_\infty = 1$ . Define

$$x_j = \frac{j-1}{\ell} \pi - \pi; \quad j = 1, \dots, 2\ell, \quad (3.1)$$

$$A_j = \frac{1}{2\ell} \cos(\ell x_j) = \frac{(-1)^{\ell-j+1}}{2\ell}; \quad j = 1, \dots, 2\ell,$$

and observe the identities

$$\sum_{j=1}^q \cos\left(s \left(\frac{j-1}{q} \pi - \pi\right)\right) \cos\left(i \frac{j-1}{q} \pi - \pi\right) = 0 \quad (3.2)$$

for all  $i \in \{0, \dots, s-1, s+1, \dots, m\}$  such that  $s+i$  and  $|s-i|$  are not multipliers of  $2q$  [see e.g. Rivlin (1974), Exercise 1.5.28]. Using these equalities it is easy to see that

$$\sum_{j=1}^{2\ell} A_j f_{2i}(x_j) = \frac{1}{2\ell} \sum_{j=1}^{2\ell} \cos(\ell x_j) \cos(ix_j) = 0, \quad (3.3)$$

whenever  $i \in \{1, \dots, m\} \setminus \{\ell\}$  [note that we require the condition  $\ell > m/3$  for the application of formula (3.2)]. On the other hand it follows from the definition of the points  $x_j$  in (3.1) that

$$x_{2\ell-j} + x_{j+2} = 0, \quad j = 0, \dots, \ell-1; \quad x_1 = -\pi; \quad x_{\ell+1} = 0,$$

and a direct calculation shows

$$\sum_{j=1}^{2\ell} A_j f_{2i-1}(x_j) = \frac{1}{2\ell} \sum_{j=1}^{2\ell} \cos(\ell x_j) \sin(i x_j) = 0,$$

which proves the conditions (i) and (ii) of Theorem 2.1. Finally, the remaining condition (iii) of this Lemma is obvious from

$$\sum_{j=1}^{2\ell} A_j f_{2\ell}(x_j) = \frac{1}{2\ell} \sum_{j=1}^{2\ell} \cos^2((j-1-\ell)\pi) = 1,$$

and by Theorem 2.1 the function  $\varphi_m(x) = f_{2\ell}(x) = \cos(\ell x)$  is a solution of the extremal problem (1.5) which is equivalent to the extremal problem (1.2).

The case  $k = 0$  is obtained exactly in the same way. Finally, the remaining case  $k = 2\ell - 1$ ,  $\ell > m/3$  is treated by similar arguments using the the points

$$x_j = \frac{2j-1}{2\ell}\pi - \pi, \quad j = 1, \dots, 2\ell,$$

the weights

$$A_j = \frac{1}{2\ell} \sin(\ell x_j), \quad j = 1, \dots, 2\ell$$

in (3.1) and observing the identity

$$\sum_{j=1}^q \sin\left(s\left(\frac{2j-1}{2q}\pi - \pi\right)\right) \sin\left(i\left(\frac{2j-1}{2q}\pi - \pi\right)\right) = 0$$

for all  $i \in \{0, \dots, s-1, s+1, \dots, m\}$  such that  $s+i$  and  $|s-i|$  are not multipliers of  $2q$  [see Rivlin (1974)]. The details are omitted for the sake of brevity. □

## 4 Extremal polynomials in the case $1 \leq \lfloor \frac{k+1}{2} \rfloor \leq m/3$

It will be demonstrated in the present section that the solution of the extremal problem (1.2) in the case  $1 \leq \lfloor \frac{k+1}{2} \rfloor \leq m/3$  is substantially more complicated. Throughout this section we define the index

$$(4.1) \quad p = \lfloor \frac{m+\ell}{2\ell} \rfloor + 1,$$

which will be an important quantity for the classification of the solution of the extremal problem. In particular we have to distinguish the following three cases

$$(4.2) \quad k = 2\ell, \quad p \text{ is odd}$$

$$(4.3) \quad k = 2\ell - 1$$

$$(4.4) \quad k = 2\ell, \quad p \text{ is even}$$

Trigonometric polynomials of the form

$$(4.5) \quad \varphi(x, \vartheta) = \begin{cases} \sum_{j=1}^{p-1} \vartheta_{2\ell(2j-1)} f_{2\ell(2j-1)}(x) = \sum_{j=1}^{p-1} \vartheta_{2\ell(2j-1)} \cos(\ell(2j-1)x) & \text{if } k = 2\ell \\ \sum_{j=1}^{p-1} \vartheta_{2\ell(2j-1)-1} f_{2\ell(2j-1)-1}(x) = \sum_{j=1}^{p-1} \vartheta_{2\ell(2j-1)-1} \sin(\ell(2j-1)x) & \text{if } k = 2\ell - 1 \end{cases}$$

will serve as appropriate candidates for the extremal polynomial [recall the definition of the function  $f_k(x)$  in (2.1)]. The following result is the basic tool for the solution of the extremal problem (1.2).

**Lemma 4.1.** *Assume that  $1 \leq \lfloor \frac{k+1}{2} \rfloor \leq \frac{m}{3}$ .*

(a) *If condition (4.2) is satisfied, define*

$$(4.6) \quad x_i = \left(2i - 1 + 2\left\lfloor \frac{i-1}{p-1} + \frac{1}{2} \right\rfloor\right) \frac{\pi}{2p\ell}; \quad i = 1, \dots, p-1,$$

*then there exists a unique trigonometric polynomial of the form (4.5), which is determined by of the system of equations*

$$(4.7) \quad \varphi(x_i, \vartheta) = \frac{f_k(x_i)}{|f_k(x_i)|}; \quad i = 1, \dots, p-1,$$

$$(4.8) \quad \varphi'(x_i, \vartheta) = 0; \quad i = 1, \dots, p-1.$$

(b) *If condition (4.3) is satisfied, define*

$$(4.9) \quad x_i = \left(i + \left\lfloor \frac{i-1}{p-1} \right\rfloor\right) \frac{\pi}{p\ell}; \quad i = 1, \dots, p-1,$$

*then there exists a unique trigonometric polynomial of the form (4.5) which is determined by the equations (4.7) and (4.8) .*

(c) *If condition (4.4) is satisfied, define*

$$(4.10) \quad x_i = \left(i - 1 + \left\lfloor \frac{i-1}{p-1} + \frac{1}{2} \right\rfloor\right) \frac{\pi}{p\ell}; \quad i = 1, \dots, p,$$

*then there exists a unique trigonometric polynomial of the form (4.5) which is determined by (4.8) and*

$$(4.11) \quad \varphi(x_i, \vartheta) = \frac{f_k(x_i)}{|f_k(x_i)|}; \quad i = 1, \dots, p.$$

**Proof.** We only prove case (a), the other cases are shown exactly in the same way. Recall that in this case  $k = 2\ell$ ,  $\ell = \lfloor \frac{k+1}{2} \rfloor$  and  $p$  is odd which obviously implies  $p \geq 3$ . We define the  $(p-1) \times (p-1)$ -matrix

$$(4.12) \quad B = \begin{pmatrix} \cos(\ell x_1) & \cos(3\ell x_1) & \dots & \cos((2p-3)\ell x_1) \\ \vdots & \vdots & \vdots & \vdots \\ \cos(\ell x_{\frac{p-1}{2}}) & \cos(3\ell x_{\frac{p-1}{2}}) & \dots & \cos((2p-3)\ell x_{\frac{p-1}{2}}) \\ -\ell \sin(\ell x_1) & -3\ell \sin(3\ell x_1) & \dots & -(2p-3)\ell \sin((2p-3)\ell x_1) \\ \vdots & \vdots & \vdots & \vdots \\ -\ell \sin(\ell x_{\frac{p-1}{2}}) & -3\ell \sin(3\ell x_{\frac{p-1}{2}}) & \dots & -(2p-3)\ell \sin((2p-3)\ell x_{\frac{p-1}{2}}) \end{pmatrix}$$

and the vector

$$(4.13) \quad e = \left( \frac{\cos(\ell x_1)}{|\cos(\ell x_1)|}, \dots, \frac{\cos(\ell x_{\frac{p-1}{2}})}{|\cos(\ell x_{\frac{p-1}{2}})|}, 0, \dots, 0 \right)^T \in \mathbb{R}^{p-1}.$$

Observing the fact that

$$x_i + x_{p-i} = \frac{\pi}{\ell} \quad i = 1, \dots, \frac{p-1}{2}$$

[by definition (4.6)] and the identities

$$\begin{aligned} \cos((2j-1)\ell(x - \frac{\pi}{\ell})) &= -\cos((2j-1)\ell x), \\ \sin((2j-1)\ell(x - \frac{\pi}{\ell})) &= -\sin((2j-1)\ell x), \end{aligned}$$

it follows that the system of equations in (4.7) and (4.8) for an arbitrary trigonometric polynomial of the form

$$\varphi(x, \vartheta) = \sum_{j=1}^{p-1} \vartheta_{2\ell(2j-1)} f_{2\ell(2j-1)}(x) = \sum_{j=1}^{p-1} \vartheta_{2\ell(2j-1)} \cos((2j-1)\ell x)$$

is equivalent to the system

$$(4.14) \quad B\vartheta = e.$$

Therefore the assertion in part (a) of Lemma 4.1 follows, if it can be shown that the matrix  $B$  is non-singular. For this purpose consider for  $s \in \mathbb{N}$  the matrix

$$A = \begin{pmatrix} \cos x_1 & \cos(3x_1) & \dots & \cos((4s-1)x_1) \\ \cos x_2 & \cos(3x_2) & \dots & \cos((4s-1)x_2) \\ \vdots & \vdots & \vdots & \vdots \\ \cos x_{2s} & \cos(3x_{2s}) & \dots & \cos((4s-1)x_{2s}) \end{pmatrix} = (u_1, \dots, u_{2s}) \in \mathbb{R}^{2s \times 2s},$$

where  $x_1 < x_2 < \dots < x_{2s}$  are arbitrary numbers and the vector  $u_\ell$  is defined by

$$u_\ell = (\cos((2\ell-1)x_1), \dots, \cos((2\ell-1)x_{2s}))^T; \quad \ell = 1, \dots, 2s.$$



Let  $U_{2\ell-1}(x) = \sin(2\ell \arccos x) / \sin(\arccos x)$  denote the  $(2\ell - 1)$ th Chebyshev polynomial of the second kind [see Szegö (1975)]. Observing the identity

$$\sum_{j=1}^{\ell} \cos((2j-1)x) = \frac{1}{2} \frac{\sin(2\ell x)}{\sin x} = \frac{1}{2} U_{2\ell-1}(z)$$

with  $z = \cos x$  [see Jolley (1961), formula 420] it follows that

$$\begin{aligned} \det A &= \det(u_1, u_1 + u_2, u_1 + u_2 + u_3, \dots, u_1 + \dots + u_{2s}) \\ &= \left(\frac{1}{2}\right)^{2s} \det\left(U_{2j-1}(z_i)\right)_{i,j=1}^{2s} \end{aligned}$$

where  $z_i = \cos x_i$  ( $i = 1, \dots, 2s$ ). It is well known that the leading coefficient of  $U_j(x)$  is  $2^j$  [see Szegö (1975)], and we obtain (using the Vandermonde determinant formula)

$$\begin{aligned} \det A &= \left(\frac{1}{2}\right)^{2s} 2^{\sum_{j=1}^{2s} (2j-1)} \det(z_i^{2j-1})_{i,j=1}^{2s} \\ &= 2^{2s(2s-1)} \prod_{i=1}^{2s} \cos x_i \prod_{1 \leq i < j \leq 2s} (\cos^2 x_j - \cos^2 x_i). \end{aligned}$$

Subtracting the  $i$ th row of the matrix  $A$  from the  $(i+1)$ th row ( $i = 1, \dots, s$ ) it follows that the determinant of the matrix  $B$  defined by (4.12) (with  $s = \frac{p-1}{2}$ ) can be obtained as

$$\begin{aligned} \det B &= \lim_{x_{i+s} \rightarrow x_i; i=1, \dots, s} \frac{\det A}{\prod_{i=1}^s (x_{i+s} - x_i)} \\ &= -2^{2s(2s-1)} \prod_{i=1}^s (\cos^2 x_i \sin 2x_i) \prod_{1 \leq i < j \leq s} (\cos^2 x_i - \cos^2 x_j)^4. \end{aligned}$$

Finally, if  $s = \frac{p-1}{2}$  and the points  $x_1, \dots, x_{\frac{p-1}{2}}$  are given by (4.6), then it follows that  $x_j \in (0, \frac{\pi}{2})$  ( $j = 1, \dots, \frac{p-1}{2}$ ), which implies  $\det B \neq 0$ . This proves the assertion of Lemma 4.1.  $\square$

**Theorem 4.2.** *Assume that  $m > 2$ . For any  $k$  such that  $1 \leq \lfloor \frac{k+1}{2} \rfloor \leq \frac{m}{3}$  the trigonometric polynomial defined by Lemma 4.1 is a solution of the extremal problem (1.2). Moreover, the constant  $c_{m,\infty}$  in the equivalent problem (1.3) depends only on  $p$  and is given by*

$$c_{m,\infty} = \left\{ \frac{2}{p} \cot\left(\frac{\pi}{2p}\right) \right\}^{-1},$$

where  $p$  is defined by (4.1).

**Proof.** Again we restrict ourselves to the case (4.2) where  $k = 2\ell$  and  $p$  is odd. We consider the trigonometric polynomial  $\hat{\varphi}(x) = \varphi(x, \hat{\vartheta})$  defined by the first part of Lemma 4.1 and show that

$$(4.15) \quad |\varphi(x, \hat{\vartheta})| \leq 1 \quad \forall x \in [-1, 1].$$




Figure 4.1: The function  $\bar{Q}$  defined in (4.17) for  $p = 5$  (left panel) and  $p = 7$  (right panel), which is considered in the proof of Theorem 4.2.

For this we introduce the function

$$(4.16) \quad Q(x) = \varphi(x, \hat{v}) + \cos(2plx),$$

and note that it follows from the representation (4.5) that

$$(4.17) \quad Q(x) = Q(\ell u) = \bar{\varphi}(u) + \cos(2pu) =: \bar{Q}(u)$$

where  $u = \ell x$  and the function  $\bar{\varphi}$  is defined by

$$\bar{\varphi}(u) = \sum_{j=1}^{p-1} \vartheta_{2\ell(2j-1)} \cos((2j-1)u).$$

Similarly, define  $u_i = \ell x_i$  ( $i = 1, \dots, p-1$ ), where  $x_i$  is given by (4.6), note that

$$u_i = \frac{2i-1}{2p}\pi; \quad i = 1, \dots, \frac{p-1}{2},$$

and that  $u_1, \dots, u_{\frac{p-1}{2}} \in (0, \frac{\pi}{2})$  are roots of trigonometric polynomial  $\bar{Q}(u)$  with at least multiplicity two, by Lemma 4.1. Moreover,  $\bar{\varphi}(u)$  changes in the interval  $[u_{\frac{p-1}{2}}, \frac{\pi}{2}]$  from 1 to 0, while  $-\cos(2pu)$  changes in the interval  $[u_{p-1}, \frac{p-1}{p}\pi] = [\frac{2p-3}{2p}\pi, \frac{p-1}{p}\pi]$  from 1 to  $-1$  and in the interval  $[\frac{p-1}{2p}\pi, \frac{\pi}{2}]$  from  $-1$  to 1 (note that  $p$  is odd). Therefore there exists at least one more root of the trigonometric polynomial  $\bar{Q}(u)$  in the interval  $(u_{\frac{p-1}{2}}, \frac{\pi}{2})$ . A similar argument shows that  $\bar{Q}(u)$  has at least  $p$  roots in the interval  $(\frac{\pi}{2}, \pi)$  (counted with multiplicities). Therefore the polynomial  $\bar{Q}(u)$  has at least  $2p$  roots in the interval  $(0, \pi)$ . A typical behaviour of  $\bar{Q}(u)$  is depicted in Figure 4.1 for the case  $p = 5$  and  $p = 7$ . As a consequence we obtain that  $|\bar{\varphi}(u)| \leq 1 \quad \forall u \in [0, \pi]$  because otherwise the polynomial  $\bar{Q}(u)$  would have at least  $2p + 1$  roots which is impossible. By symmetry it follows that  $\|\bar{\varphi}\|_{\infty} \leq 1$  and we also have  $\|\hat{\varphi}\|_{\infty} \leq 1$ .

We now use Theorem 2.1 to prove that the trigonometric polynomial  $\hat{\varphi}$  is a solution of the extremal problem (1.5). For this we consider the points

$$x_i = \left(2i - 1 + 2\left\lfloor \frac{i-1}{p-1} + \frac{1}{2} \right\rfloor\right) \frac{\pi}{2p\ell} \quad i = 1, \dots, n = \ell(p-1)$$

where  $n = \ell(p-1)$ , and

$$x_{n+j} = -x_j \quad j = 1, \dots, n.$$

Define  $N = 2n$  and the weights  $A_1, \dots, A_N$  by

$$A_j = \frac{1}{p\ell} \cos(\ell x_j) \quad j = 1, \dots, N.$$

With these notations it follows for  $i = 2s, s \in \{0, \dots, m\} \setminus \{\ell\}$  that

$$\begin{aligned} \sum_{j=1}^N A_j f_{2s}(x_j) &= \frac{2}{p\ell} \sum_{j=1}^{(p-1)\ell} \cos(\ell x_j) \cos(s x_j) \\ &= \frac{2}{p\ell} \sum_{j=1}^{p\ell} \cos\left(\ell \frac{2j-1}{2p\ell} \pi\right) \cos\left(s \frac{2j-1}{2p\ell} \pi\right) = 0, \end{aligned}$$

where we have used the identity

$$\sum_{j=1}^q \cos\left(\ell \frac{2j-1}{2q} \pi\right) \cos\left(s \frac{2j-1}{2q} \pi\right) = 0 \quad \text{if } s \neq \ell$$

[see Rivlin (1974), Exercise 1.5.26]. On the other hand we obtain

$$\begin{aligned} \sum_{j=1}^N A_j f_{2\ell}(t_j) &= \frac{2}{p\ell} \sum_{j=1}^{p\ell} \cos^2\left(\ell \frac{2j-1}{2p\ell} \pi\right) = 1 + \frac{1}{p\ell} \sum_{j=1}^{p\ell} \cos\left(\ell \frac{2j-1}{p\ell} \pi\right) \\ &= 1 + \frac{1 \sin(2\ell\pi)}{2 \sin(\pi/p)} = 1, \end{aligned}$$

[see Jolley (1961), formula 420] and by symmetry (using the fact that  $A_{j+n} = A_j$ )

$$\sum_{j=1}^N A_j f_{2s-1}(x_j) = \sum_{j=1}^N A_j \sin(s x_j) = 0, \quad s \in \{1, \dots, m\}.$$

This shows that conditions (i) - (iii) of Theorem 2.1 are satisfied and consequently the trigonometric polynomial  $\hat{\varphi}(x) = \varphi(x, \hat{\vartheta})$  is a solution of the extremal problem (1.5). For the calculation of the value  $\hat{\vartheta}_{2\ell}$  corresponding to function  $\cos(\ell x)$  in the trigonometric polynomial  $\hat{\varphi}(x)$  we use the representation (2.3) in Theorem 2.1 and obtain

$$\begin{aligned} \hat{\vartheta}_{2\ell} &= \sum_{j=1}^N |A_j| = \frac{1}{p\ell} \sum_{j=1}^N |\cos \ell x_j| = \frac{2}{p\ell} \sum_{j=1}^{\ell(p-1)} |\cos \ell x_j| \\ &= \frac{2}{p\ell} \sum_{j=1}^{p\ell} \left| \cos\left(\frac{2j-1}{2p} \pi\right) \right| = \frac{4}{p} \sum_{j=1}^{\frac{p-1}{2}} \cos\left(\frac{2j-1}{2p} \pi\right) \\ &= \frac{2}{p} \cot\left(\frac{\pi}{2p}\right), \end{aligned}$$

where the last line follows again from formula 420 in Jolley (1961). This completes the proof of Theorem 4.2.  $\square$

**Remark 4.3.** Note that it follows from the proof of Theorem 4.2 that the extremal polynomial defined by (4.5) has  $2\ell(p-1)$  extremal points, if condition (4.2) or (4.3) is satisfied, while there are  $2\ell(p-1) + 3$  extremal points in the case (4.4). Moreover, it follows from the representation (4.10) that the points  $-\pi$  and  $\pi$  are extremal points if condition (4.4) is satisfied. In the two other cases the boundary points of the interval  $[-\pi, \pi]$  are not extremal points.

**Remark 4.4.** It is interesting to note that Theorem 3.1 is contained in Theorem 4.2 (note that our proof does not require the condition  $1 \leq \lfloor \frac{k+1}{2} \rfloor \leq \frac{m}{3}$ . To see this note in the case  $\ell > \frac{m}{3}$  we have  $p = \lfloor \frac{m+3\ell}{2\ell} \rfloor = 2$ , by an elementary calculation, and therefore the polynomial in (4.5) reduces to  $\cos(\ell t)$  or  $\sin(\ell t)$  corresponding to the case  $k = 2\ell$  and  $k = 2\ell - 1$ , respectively. In order to emphasize the different structure of the solution of the extremal problem (1.2) in the cases  $1 \leq \lfloor \frac{k+1}{2} \rfloor \leq \frac{m}{3}$  and  $\frac{m}{3} < \lfloor \frac{k+1}{2} \rfloor \leq m$  both scenarios are presented in different sections.

**Remark 4.5.** The extremal polynomial in Theorem 4.2 can be represented explicitly as

$$(4.18) \quad \hat{\varphi}(x) = \sum_{j=1}^n \text{sign}(f_k(x_j)) \ell_{jk}(x),$$

where the functions  $\ell_{jk}(t)$  are defined as

$$\ell_{jk}(x) = \prod_{i \neq j} \left( \frac{f_k(x) - f_k(x_i)}{f_k(x_j) - f_k(x_i)} \right)^{s_i} \{a_j(f_k(x) - f_k(x_j)) + 1\}$$

with

$$a_j = - \sum_{i \neq j} \frac{s_i}{f_k(x_i) - f_k(x_j)}$$

and  $s_1 = \dots = s_n = 2, n = p-1$ , if (4.2) or (4.3) holds and  $s_1 = s_n = 1, s_2 = \dots = s_{n-1} = 2, n = p$ , if (4.4) holds. This follows by a tedious calculation showing that the function on the right hand side of (4.18) satisfies the conditions (4.7) and (4.8) which determine the extremal polynomial of the form (4.5) uniquely.

**Remark 4.6.** Note that by Theorem 4.2 the optimal constant in (1.3) is given by

$$c_{k,m,\infty} = \begin{cases} \left\{ \frac{2}{p} \cot\left(\frac{\pi}{2p}\right) \right\}^{-1} & \text{if } 1 \leq \lfloor \frac{k+1}{2} \rfloor \leq \frac{m}{3} \\ 1 & \text{else} \end{cases}.$$

Therefore if  $m \rightarrow \infty$  it follows for any fixed  $k \in \mathbb{N}$  that

$$\lim_{m \rightarrow \infty} c_{k,m,\infty} = \lim_{p \rightarrow \infty} \left\{ \frac{2}{p} \cot\left(\frac{\pi}{2p}\right) \right\}^{-1} = \frac{\pi}{4}.$$

In other words the optimal constant  $c_{k,m,\infty}$  in the extremal problem (1.5) exhibits the same asymptotic behaviour as the optimal constants  $c_{m,p}$  in the extremal problem (1.4) [see Marshall and Ganzburg (1999)].




Figure 4.2: The solution of the extremal problem (1.2) in the case  $m = 6$ . The left part of the figure shows the extremal polynomials for  $k = 1$  (dotted line) and  $k = 3$  (solid line), while the right part displays the solution for  $k = 2$  (dotted line) and  $k = 4$  (solid line).

**Example 4.7.** Consider the case  $m = 6$ , where Theorem 4.2 can be applied for the cases  $k = 1, 2, 3, 4$ . If  $\ell = 1$  we have  $p = 4$ , and the trigonometric polynomial in (4.5) is given by

$$(4.19) \quad \varphi(t) = \frac{\sqrt{3 + \sqrt{2}}}{2} \left( \sin t + \frac{8 - 5\sqrt{2}}{4} \sin(3t) + \frac{3\sqrt{2} - 4}{4} \sin(5t) \right),$$

$$(4.20) \quad \varphi(t) = \frac{\sqrt{3 + \sqrt{2}}}{2} \left( -\cos t + \frac{8 - 5\sqrt{2}}{4} \cos(3t) - \frac{3\sqrt{2} - 4}{4} \cos(5t) \right)$$

corresponding to the cases  $k = 1$  and  $k = 2$ , respectively. If  $\ell = 2$  the solution of the extremal problem (1.2) is simpler and given by

$$(4.21) \quad \varphi(t) = \frac{\sqrt{3}}{2} \left( \sin(2t) + \frac{1}{6} \sin(6t) \right)$$

$$(4.22) \quad \varphi(t) = \frac{\sqrt{3}}{2} \left( \cos(2t) - \frac{1}{6} \cos(6t) \right)$$

corresponding to the cases  $k = 3$  and  $k = 4$ , respectively. These polynomials are depicted in Figure 4.2. In all other cases the solution of the extremal problem (1.2) is given in Theorem 3.1.

## 5 Uniqueness and examples

In the present section we will demonstrate that the solution of the extremal problem (1.2) [or equivalently (1.3) or (1.5)] is not necessarily unique. Our first result characterizes the set of all extremal polynomials.

**Theorem 5.1.** Let  $\hat{\varphi}(x)$  denote the extremal polynomial described in Theorem 3.1 or 4.2 and define  $\{x_1, \dots, x_N\}$  as the set of all extremal points of  $\hat{\varphi}(x)$ . An arbitrary trigonometric polynomial  $\varphi(x)$  with  $\|\varphi\|_\infty \leq 1$  is a solution of the extremal problem (1.2) if and only if it can be represented as

$$(5.1) \quad \varphi(x) = \hat{\varphi}(x) + \gamma Q(x).$$

Here  $\gamma$  is a real constant and  $Q(x)$  is a trigonometric polynomial with roots  $x_1, \dots, x_N$ , where the multiplicity of a root  $x_i$  is two whenever  $|x_i| \neq \pi$ .

**Proof.** Assume that  $\varphi$  is a solution of the extremal problem (1.2), such that it has the same coefficient  $\vartheta_k$  as  $\hat{\varphi}$ . Then the trigonometric polynomial

$$\tilde{\varphi}(x) = \frac{1}{2}(\varphi(x) + \hat{\varphi}(x))$$

is also a solution of the extremal problem (1.2) because it has the same coefficient of  $\vartheta_k$  and satisfies  $\|\tilde{\varphi}\|_\infty \leq 1$ . If  $\tilde{x}_1, \dots, \tilde{x}_{\tilde{N}}$  are the extremal points of  $\tilde{\varphi}$ , then it follows from the inequality

$$|\tilde{\varphi}(x)| \leq \frac{1}{2}\{|\varphi(x)| + |\hat{\varphi}(x)|\} \leq 1$$

that

$$\{\tilde{x}_1, \dots, \tilde{x}_{\tilde{N}}\} \subset \{x_1, \dots, x_N\}.$$

Moreover,  $\tilde{x}_1, \dots, \tilde{x}_{\tilde{N}}$  are also extremal points of the trigonometric polynomial  $\varphi(x)$ , by the same argument. Note that  $\{f_0, \dots, f_{2m}\}$  is a Chebyshev system on the interval  $[-\pi, \pi)$  and that  $N \leq 2m < 2m + 1$ . As a consequence there exists a unique solution  $A_1, \dots, A_N$  of the system of equations (ii) and (iii) in Theorem 2.1, which implies

$$\{\tilde{x}_1, \dots, \tilde{x}_{\tilde{N}}\} = \{x_1, \dots, x_N\}.$$

Consequently, we obtain  $\tilde{\varphi}(x_j) = \hat{\varphi}(x_j) = \text{sign}(A_j)$ ,  $j = 1, \dots, N$ , and the function

$$\tilde{\varphi}(x) - \hat{\varphi}(x) = \frac{1}{2}(\varphi(x) - \hat{\varphi}(x))$$

has roots at the points  $x_1, \dots, x_N$ , where the multiplicity of a root  $x_i$  is (at least) two, whenever  $|x_i| \neq \pi$ . Therefore the polynomial  $\varphi$  can be written in the form (5.1), which proves the first part of the assertion.

The converse part of the assertion can be proved in a similar way and is omitted for the sake of brevity. □

**Example 5.2.** Assume that  $\frac{m}{3} < \ell = \lfloor \frac{k+1}{2} \rfloor \leq \frac{m}{2}$ , then, if  $k = 2\ell$  is even, by Theorem 3.1 the polynomial  $\hat{\varphi}(x) = \cos(\ell x)$  is a solution of the extremal problem (1.2) and the extremal points are given by  $x_j^* = (j - \ell - 1)\pi/\ell$  ( $j = 1, \dots, 2\ell$ ). These points are roots of multiplicity 2 of the polynomial  $Q(x) = 1 - \cos(2\ell x)$  and a further extremal polynomial is of the form

$$(5.2) \quad \varphi_\gamma(x) = \cos(\ell x) + \gamma(1 - \cos(2\ell x)),$$

where the constant  $\gamma$  is chosen such that  $\|\varphi_\gamma\|_\infty = 1$ . By a substitution of  $u = \cos(\ell x)$  it is easy to see that the possible range for  $\gamma$  is the interval  $[-\frac{1}{4}, \frac{1}{4}]$ . Extremal polynomials of the form (5.2) for various values of  $\ell$  and  $\gamma$  are depicted in Figure 5.1 in the case  $\ell = 2$  and  $\ell = 5$  (note that the solution of the extremal problem (1.2) depends only on  $m$  by the inequality  $\frac{m}{3} < \ell \leq \frac{m}{2}$ ).




Figure 5.1: Various solutions of the form (5.2) for the extremal problem (1.2) with  $k = 2\ell$ . Solid lines  $\gamma = 0$ ; dashed lines:  $\gamma = 1/4$ ; dotted lines:  $\gamma = -1/4$ . The left part of the figure corresponds to the case  $\ell = 2$ , while the right part represents the case  $\ell = 5$ .

It can be shown by similar arguments that in the case  $k = 2\ell - 1$ ;  $m/3 \leq \ell \leq m/2$  trigonometric polynomials of the form

$$(5.3) \quad \varphi_\gamma(x) = \sin(\ell x) + \gamma(1 + \cos(2\ell x)); \quad \gamma \in \left[-\frac{1}{4}, \frac{1}{4}\right]$$

are solutions of the extremal problem (1.2), and some representative examples are depicted in Figure 5.2 (in the case  $\ell = 2$  and  $\ell = 5$ ).

**Example 5.3.** Assume that  $1 \leq \ell = \lfloor \frac{k+1}{2} \rfloor \leq \frac{m}{3}$  and that either  $k = 2\ell - 1$  and  $\frac{m+3\ell}{2\ell}$  is an integer or that  $k = 2\ell$  and  $\frac{m+3\ell}{2\ell}$  is odd. If  $\frac{m+3\ell}{2\ell}$  is an integer we obtain that  $2p - 3 = \frac{m}{\ell}$  and consequently the trigonometric polynomial in (4.5) is of degree  $m$ . In this case it follows from Remark 4.3 that the polynomial  $Q$  in Theorem 5.1 must have  $2\ell(p - 1) = m + \ell$  roots with multiplicity no less than 2 and its degree must be larger than  $m$ . As a consequence, there exists a unique solution of the extremal problem (1.2) in this case.

On the other hand, if  $\frac{m+3\ell}{2\ell}$  is not an integer, we obtain  $\frac{m+3\ell}{2\ell} - 1 < p < \frac{m+3\ell}{2\ell}$  or equivalently  $m - 2\ell < \ell(2p - 3) < m$ . Because the degree of the polynomial in (4.5) is  $\ell(2p - 3) < m$  there exists more than one solution of the extremal problem (1.2). To be more specific consider the case  $m$  even  $k = 2\ell$  and assume that  $p$  is odd, then the extremal polynomial  $\hat{\varphi}$  from Lemma 4.1 is of the form

$$(5.4) \quad \hat{\varphi}(x) = \sum_{j=1}^{p-1} \vartheta_j \cos(\ell(2j - 1)x)$$

Let  $Q$  denote the positive trigonometric polynomial, which has precisely roots of multiplicity two at the extreme points of  $\hat{\varphi}$ . Because  $k = 2\ell$  and  $p$  is odd it follows from Remark 4.3 that the extreme points are not located at the boundary of the interval  $[-\pi, \pi]$  and by the above discussion the degree of  $Q$  is not larger than  $m$ . By Theorem 5.1 the trigonometric polynomial




Figure 5.2: Various solutions of the form (5.3) for the extremal problem (1.2) with  $k = 2\ell - 1$ . Solid lines  $\gamma = 0$ ; dashed lines:  $\gamma = 1/4$ ; dotted lines:  $\gamma = -1/4$ . The left part of the figure corresponds to the case  $\ell = 2$ , while the right part represents the case  $\ell = 5$ .

$\varphi_\gamma(x) = \hat{\varphi}(x) + \gamma Q(x)$  is a solution of the extremal problem (1.2) if and only if  $\|\varphi_\gamma\|_\infty \leq 1$ . In order to derive conditions on the parameter  $\gamma$  such that this inequality is satisfied let  $\bar{x}_1, \dots, \bar{x}_{N_1}$  denote the extreme points of  $\hat{\varphi}(x)$  located in the interval  $[0, \pi)$  with  $\hat{\varphi}(x_j) = 1$ , define the trigonometric polynomial

$$(5.5) \quad U(x) = \prod_{j=1}^{N_1} (\cos x - \cos \bar{x}_j)^2,$$

and consider the functions

$$(5.6) \quad H_1(x) = \frac{\hat{\varphi}(x) - 1}{U(x)}; \quad H_2(x) = \frac{Q(x)}{U(x)}.$$

Note that  $H_1$  and  $H_2$  are trigonometric polynomials and that the function  $H_2$  is nonnegative. The inequality  $\|\varphi_\gamma\|_\infty \leq 1$  is obviously equivalent to the inequalities

$$(5.7) \quad \hat{\varphi}(x) - 1 + \gamma Q(x) = (H_1(x) + \gamma H_2(x))U(x) \leq 0$$

$$(5.8) \quad \hat{\varphi}(x) + 1 + \gamma Q(x) \geq 0$$

for all  $x \in [-\pi, \pi]$ . We consider the case (5.7) in more detail, the second case is treated similarly by an appropriate modification of the definition of the functions  $H_1$  and  $H_2$ . Because  $U(x)$  is a positive function it follows that  $H_1(x)$  is nonpositive and (5.7) is equivalent to the inequality

$$(5.9) \quad H_1(x) + \gamma H_2(x) \leq 0$$

for all  $x \in [-\pi, \pi]$ . Define

$$\bar{\gamma} = \min \left\{ -\frac{H_1(x)}{H_2(x)} \mid x \in [-\pi, \pi] \right\},$$



and note that  $\bar{\gamma} > 0$ , then it is easy to see that (5.9) is satisfied if and only if  $\gamma \leq \bar{\gamma}$ . Similarly, it can be shown that (5.8) holds if and only if  $\gamma \geq -\bar{\gamma}$ . Consequently, the polynomial  $\varphi_\gamma(x) = \hat{\varphi}(x) + \gamma Q(x)$  constructed in this way is a solution of the extremal problem (1.2) if and only if  $|\gamma| \leq \bar{\gamma}$ .

As a concrete example consider the case  $m = 4$ ,  $k = 2\ell = 2$ , where  $p = 3$  and the extremal polynomial defined in (4.5) is given by

$$\hat{\varphi}(x) = \frac{2}{\sqrt{3}} \left\{ \cos x - \frac{\cos(3x)}{6} \right\}.$$

The extremal points of  $\hat{\varphi}$  are  $-\frac{5\pi}{6}$ ,  $-\frac{\pi}{6}$ ,  $\frac{\pi}{6}$ ,  $\frac{5\pi}{6}$  and polynomial  $Q$  is calculated as

$$Q(x) = 16(\cos^2 x - \frac{3}{4})^2 = 2 \cos(4x) - 4 \cos(2x) + 3$$

(note that  $\cos(\frac{\pi}{6}) = -\cos(\frac{5\pi}{6}) = \frac{\sqrt{3}}{2}$ ). The polynomial  $U$  is given by  $U(x) = (\cos x - \frac{\sqrt{3}}{2})^2$ , which yields

$$\begin{aligned} H_2(x) &= 16 \left( \cos x + \frac{\sqrt{3}}{2} \right)^2, \\ H_1(x) &= -\frac{4}{3} \left( 1 + \frac{\cos x}{\sqrt{3}} \right), \end{aligned}$$

and for  $\bar{\gamma}$  the expression

$$\bar{\gamma} = \min \left\{ \frac{1 + (\cos x)/\sqrt{3}}{12(\cos x + \sqrt{3}/2)^2} \mid x \in [-\pi, \pi] \right\} = 1 - \frac{5}{9}\sqrt{3}.$$

Several solutions of the extremal problem (1.2) are depicted in Figure 5.3.

**Example 5.4.** If  $\ell > \frac{m}{2}$ , the solution of the extremal problem (1.2) is given by  $\hat{\varphi}(x) = \cos(\ell x)$  if  $k = 2\ell$  and by  $\hat{\varphi}(x) = \sin(\ell x)$  if  $k = 2\ell - 1$ . In the second case the corresponding polynomial  $Q$  must have (at least)  $2 \cdot 2\ell > 2m$  roots (counted with multiplicities), while in the first case there are  $2\ell - 1$  roots with multiplicity two (corresponding to the extreme points in the interval  $(-\pi, \pi)$  and one root at the point  $\pi$ , which gives (counted with multiplicities)  $2(2\ell - 1) + 1 > 2m$  roots for the polynomial  $Q$ . In both cases the polynomial  $Q$  would not be a trigonometric polynomial of degree  $m$  and therefore the solution of the extremal problem is unique if  $\ell > \frac{m}{2}$ .

We conclude this paper summarizing the discussion presented in the previous examples.

**Theorem 5.5.** *The solution of the extremal problem (1.2) is unique if one of the following conditions is satisfied*

- (a)  $\lfloor \frac{k+1}{2} \rfloor \geq \frac{m}{2}$
- (b)  $1 \leq \lfloor \frac{k+1}{2} \rfloor \leq \frac{m}{3}$ ,  $k$  is even,  $\frac{m+3\ell}{2\ell} \in \mathbb{N}$  and is odd
- (c)  $1 \leq \lfloor \frac{k+1}{2} \rfloor \leq \frac{m}{3}$   $k$  is odd,  $\frac{m+3\ell}{2\ell} \in \mathbb{N}$




Figure 5.3: Various solutions of the form (5.1) of the extremal problem (1.2) for  $m = 4$  and  $k = 2$  ( $\ell = 1$ ). Solid line  $\gamma = 0$ ; dashed line:  $\gamma = -1 + \frac{5}{9}\sqrt{3}$ ; dotted lines:  $\gamma = 1 - \frac{5}{9}\sqrt{3}$ .

In all other cases, the solution of the extremal problem (1.2) is not unique.

**Acknowledgements.** The work of the authors was supported by the Deutsche Forschungsgemeinschaft (SFB 475 Komplexitätsreduktion in multivariaten Datenstrukturen). The authors would also like to thank Isolde Gottschlich, who typed this paper with considerable technical expertise. We are also grateful to T. Erdelyi for some help regarding the related literature and to W.J. Studden, who brought our attention to the book of Singer.

## References

- Achieser, N. I. (1956). *Theory of Approximation*. Ungar Publishing Co., New York.
- Chebyshev, P. L. (1859). *Sur les questions de minima qui se rattachent à la représentation approximative de fonctions*. Mémoires de l'Académie Impériale des Sciences de St-Petersbourg. Sixième Série Sciences Mathématiques et Physiques, **7**, 199–291.
- L.B.W. Jolley (1961). *Summation of series*. New York: Dover Publications, Inc.
- A.J. Marshall, M. Ganzburg (1999). *An extremal problem for trigonometric polynomials*. Proc. Am. Math. Soc. 127, 211-216.
- G.V. Milovanović, D.S. Mitrinović, Th. M. Rassias (1994). *Topics in polynomials: extremal problems, inequalities, zeros*. Singapore: World Scientific.
- Natanson, T. P. (1955). *Konstruktive Funktionentheorie*. Akademie Verlag, Berlin.
- F. Peherstorfer (1982). *On an extremal problem for nonnegative trigonometric polynomials and the characterization of positive quadrature formulas with Chebyshev weight function*. Acta Math. Acad. Sci. Hungar 39, 107-116.
- T.J. Rivlin (1974). *Chebyshev polynomials*. Wiley, New York.
- I.Singer (1970). *Best Approximation in Normed Linear Spaces by Elements of Linear Subspaces*. Springer, N.Y.
- G. Szegő (1959). *Orthogonal polynomials*. Am. Math. Soc. Colloqu. Publ. 23. Providence, RI.