



Multi-attribute decision-making in individual and social choice

Walter Bossert^a, Hans Peters^{b,*}

^a*Department of Economics, University of Nottingham, University Park, Nottingham NG7 2RD, UK*

^b*Department of Quantitative Economics, University of Maastricht, P.O. Box 616, 6200 MD Maastricht, The Netherlands*

Received 1 October 1998; received in revised form 1 August 1999; accepted 1 October 1999

Abstract

This paper provides an analysis of individual and social decision criteria for alternatives that are composed of several attributes. We derive additive and multiplicative criteria for individual decision-making with new axioms and apply these criteria to obtain new justifications of known social choice rules with a bargaining interpretation, namely the generalized utilitarian and Nash social choice functions. Unlike most axiomatizations of bargaining solutions, our approach is, to a large extent, based on the multi-attribute structure of the underlying alternatives and the resulting individual decision criteria instead of axioms that impose restrictions on the choice function directly. © 2000 Elsevier Science B.V. All rights reserved.

JEL classification: D81; D71; C78

1. Introduction

A standard cooperative bargaining solution as introduced by Nash (1950) assigns a utility vector to each pair of a utility possibilities set and a disagreement point (the utility vector resulting if the agents fail to reach an agreement) within a given domain. For example, a typical domain of a bargaining solution consists of all pairs such that the disagreement point is normalized to be the origin and the utility possibilities set is convex and compact. Comprehensiveness of the feasible set frequently is required as well. In this paper, we consider an informationally richer framework by allowing for the possibility that the structure of the set of objects over which the bargaining process takes

*Corresponding author. Tel. +31-43-388-3834; fax: +31-43-388-4874.

E-mail addresses: lezwb@unix.ccc.nottingham.ac.uk (W. Bossert), h.peters@ke.unimaas.nl (H. Peters).

place may matter. In particular, we examine *multi-attribute* bargaining problems. Multiple-issue bargaining — as it is often called — is also discussed in Kalai (1977) in the context of interpreting his step-by-step negotiation axiom; see Kalai (1977, p. 1627) for details. Furthermore, Ponsati and Watson (1997) provide an analysis of multiple-issue bargaining in a cooperative framework. A typical example for a concrete multi-attribute bargaining problem is a wage-employment bargaining model such as the one developed by McDonald and Solow (1981). Multiple-issue bargaining in a non-cooperative setting is discussed, for example, in Busch and Horstmann (1997) and in Fershtman (1990).

To begin with, we provide new axiomatic derivations of additive and multiplicative multi-attribute criteria for an individual decision maker. The preferences of a decision maker over combinations of attributes are additive if they can be represented by a cardinal utility function that can be written as the sum of the utilities of the individual attributes. The multiplicative decision criterion is based on the product of the utilities over attributes. Clearly, these types of preferences possess separability properties with respect to the attributes under consideration. See Fishburn (1970) for a comprehensive treatment of utility theory and decision criteria.

Specifically, we use a weak version of *utility independence* (see, for instance, Keeney and Raiffa, 1976, and the references quoted there) together with an additivity axiom to generate an additive representation of an agent's preferences. Similarly, we show that the multiplicative decision criterion is implied by an analogous utility independence axiom and a condition regarding the existence and properties of worst possible attributes. By employing these new axioms, the paper also makes a contribution to the theory of decision-making in the presence of several attributes. A discussion of separable structures in consumer and producer theory can be found in Blackorby et al. (1978).

Turning from individual to social decision-making, we first characterize efficiency in both the additive and the multiplicative case. This efficiency criterion, which leads to the standard weak Pareto optimality condition, together with some other assumptions, is then used to derive social decision procedures with a bargaining interpretation. In the additive case, weak Pareto optimality and an independence condition regarding the restriction of choices to specific attributes lead to the generalized (not necessarily symmetric) utilitarian social choice functions, provided a mild regularity condition is satisfied. Analogously, in the multiplicative case, generalized Nash social choice functions are obtained. The novel aspect of those axiomatic derivations of well-known social choice functions is that much of the structure is imposed through the individual preferences rather than through axioms that operate directly on the social choice function itself. We therefore provide a decision-theoretic foundation of these commonly used solutions.

2. Multi-attribute preferences

Let \mathbb{N} denote the set of positive integers, and let \mathbb{R} (\mathbb{R}_{++}) be the set of all (positive) real numbers. For $n \in \mathbb{N}$, \mathbb{R}^n is the n -fold Cartesian product of \mathbb{R} . Let $M = \{1, \dots, m\}$ be a set of $m \in \mathbb{N}$ attributes. The sets A_1, \dots, A_m are the corresponding non-empty spaces

of these attributes. Let $A = \times_{j \in M} A_j$. For $a \in A$ and $M' \subseteq M$, let $a_{-M'}$ be the subvector of a that contains the components in $M \setminus M'$ only. If M' is a singleton $\{j\}$, we use the simpler notation a_{-j} instead of $a_{-\{j\}}$. Analogously, for $A' \subseteq A$ and $M' \subseteq M$, $A'_{-M'} = \{a_{-M'} \mid a \in A'\}$, and if M' is a singleton $\{j\}$, we write A'_{-j} instead of $A'_{-\{j\}}$.

We assume there is a decision maker who has a cardinal utility function $u: A \rightarrow \mathbb{R}$. We also assume that there is a fixed m -tuple $a^0 \in A$ with $u(a) \geq u(a^0)$ for all $a \in A$. That is, a^0 is a worst alternative in A .

Let $j \in M$. By fixing the $m - 1$ attributes in $M \setminus \{j\}$ at some given levels $a'_{-j} \in A_{-j}$, the utility function u induces a utility function $u_j: A_j \rightarrow \mathbb{R}$ on the attribute space A_j . The property of utility independence (see Keeney and Raiffa, 1976, in particular Chaps. 5 and 6) requires that this induced utility function is independent of the specific levels of the other attributes — as is already suggested by its notation. To define this property formally, let $\emptyset \neq A' \subseteq A$, and let $j \in M$. A'_j is *utility independent of A'_{-j}* if for every $a'_{-j} \in A'_{-j}$ there exist functions $f_j: A'_j \rightarrow \mathbb{R}$ and $g_j: A'_{-j} \rightarrow \mathbb{R}_{++}$ such that

$$u(a) = f_j(a_{-j}) + g_j(a_{-j})u(a_j, a'_{-j}), \quad \text{for all } a \in A' \tag{1}$$

Clearly, utility independence is a separability property. (1) means that, on the restricted domain A' , values of attribute j can be assessed without knowledge of specific values of the remaining attributes. See Keeney and Raiffa (1976) for a detailed discussion of this and related separability assumptions in this framework.

3. Additive decision criteria

In this section, we show that a specific utility independence requirement and a weak additivity assumption imply that u must be additive.

Restricted a^0 utility independence requires that we can find a chain of $m - 1$ attributes such that, for each attribute j in this chain, A_j is utility independent of all other attributes, where this independence can be restricted to attribute values of a_k^0 for all attributes k that appear before j in the chain. For simplicity of exposition, we formulate this condition for the chain consisting of the first $m - 1$ attributes and leave the obvious generalization to the reader.

Restricted a^0 utility independence:

- A_1 is utility independent of A_{-1} ;
- A_2 is utility independent of $\{a_1^0\} \times A_{-\{1,2\}}$;
- \vdots
- A_{m-1} is utility independent of $\{a_1^0\} \times \{a_2^0\} \times \dots \times \{a_{m-2}^0\} \times A_m$.

This independence axiom is considerably weaker than Keeney and Raiffa's (1976) mutual utility independence because, for each $j \in M \setminus \{1, m\}$, all attributes $k < j$ in the chain can be fixed at a_k^0 . Moreover, only $m - 1$ rather than all m attributes are required to satisfy an independence condition.

The second axiom is a weakening of additive independence (see Keeney and Raiffa, 1976). Again, we state this condition for a chain involving the first $m - 1$ attributes only in order to simplify notation.

Restricted additive independence:

- For all $a_{-1} \in A_{-1}$, there exist $a'_1, a''_1 \in A_1$ such that $u(a'_1, a_{-1}) + u(a''_1, a^0_{-1}) = u(a'_1, a^0_{-1}) + u(a''_1, a_{-1})$ and $u(a'_1, a^0_{-1}) \neq u(a''_1, a^0_{-1})$;
- and, for all $a_{-\{1,2\}} \in A_{-\{1,2\}}$, there exist $a'_2, a''_2 \in A_2$ such that $u(a'_1, a'_2, a_{-\{1,2\}}) + u(a''_2, a^0_{-2}) = u(a'_2, a^0_{-2}) + u(a'_1, a''_2, a_{-\{1,2\}})$ and $u(a'_1, a^0_{-2}) \neq u(a''_2, a^0_{-2})$;
- \vdots
- and, for all $a_m \in A_m$, there exist $a'_{m-1}, a''_{m-1} \in A_{m-1}$ such that $u(a^0_1, \dots, a^0_{m-2}, a'_{m-1}, a_m) + u(a''_{m-1}, a^0_{-(m-1)}) = u(a'_{m-1}, a^0_{-(m-1)}) + u(a^0_1, \dots, a^0_{m-2}, a''_{m-1}, a_m)$ and $u(a'_{m-1}, a^0_{-(m-1)}) \neq u(a''_{m-1}, a^0_{-(m-1)})$.

Analogously to restricted a^0 utility independence, the requirements imposed by restricted additive independence become successively weaker as we progress along the chain of attributes. Restricted additive independence is analogous in spirit to Keeney and Raiffa’s (1976) additive independence (formulated for two attributes only). Additive independence requires that there exist some attribute values $a'_1, a''_1 \in A_1$ and $a'_2, a''_2 \in A_2$ such that $u(a'_1, a'_2) + u(a''_1, a''_2) = u(a'_1, a''_2) + u(a''_1, a'_2)$ and $u(a'_1, a'_2) \neq u(a''_1, a''_2)$ and $u(a'_1, a'_2) \neq u(a''_1, a'_2)$. Our version of the additive independence condition, in contrast, requires the existence of specific attribute values for *several* values of the other attributes, and the worst attribute combination a^0 must be involved. This is the case because we combine the axiom with a rather weak form of utility independence — namely restricted a^0 utility independence — whereas Keeney and Raiffa (1976) employ the much stronger mutual utility independence. Furthermore, note that restricted additive independence only requires two of the utility values involved to be distinct, but additive independence demands two inequalities of that kind.

The following theorem derives the additive decision criterion from the above axioms. That is, we show that, if combined, the two conditions generate a strong separability property — namely additive separability. See, for example, Blackorby et al. (1978) for a thorough discussion of various separability properties. Note that, in order to reformulate the axioms for general chains in this theorem, the same chains have to be used in both axioms. Recall that a^0 is a worst alternative, i.e. $u(a) \geq u(a^0)$ for all $a \in A$.

Theorem 1. *Let u satisfy restricted a^0 utility independence and restricted additive independence. Assume u is normalized so that $u(a_0) = 0$. Then $u(a) = \sum_{j \in M} u(a_j, a^0_{-j})$ for all $a \in A$.*

Proof. Because A_1 is utility independent of A_{-1} , there exist functions f_1 and g_1 such that:

$$u(a) = f_1(a_{-1}) + g_1(a_{-1})u(a_1, a^0_{-1}), \quad \text{for all } a \in A \tag{2}$$

where we chose $a'_{-1} = a^0_{-1}$ in (1). Letting $a_1 = a^0_1$ in (2), we obtain $f_1(a_{-1}) = u(a^0_1, a_{-1})$ for all $a \in A$ because $u(a^0) = 0$. Therefore:

$$u(a) = u(a^0_1, a_{-1}) + g_1(a_{-1})u(a_1, a^0_{-1}), \quad \text{for all } a \in A \tag{3}$$

Because, by restricted additive independence, there exists $a_1 \in A_1$ such that $u(a_1, a^0_{-1}) \neq 0$, we can choose $a_{-1} = a^0_{-1}$ in (3) and use $u(a^0) = 0$ to obtain $g_1(a^0_{-1}) = 1$.

Using restricted additive independence and (3), it follows that, for all $a_{-1} \in A_{-1}$, there exist $a'_1, a''_1 \in A_1$ such that:

$$g_1(a_{-1})[u(a'_1, a^0_{-1}) - u(a''_1, a^0_{-1})] = g_1(a^0_{-1}) [u(a'_1, a^0_{-1}) - u(a''_1, a^0_{-1})]$$

Because, by restricted additive independence, $u(a'_1, a^0_{-1}) \neq u(a''_1, a^0_{-1})$, this implies: $g_1(a_{-1}) = g_1(a^0_{-1}) = 1$ for all $a \in A$. Therefore:

$$u(a) = u(a^0_1, a_{-1}) + u(a_1, a^0_{-1}), \quad \text{for all } a \in A \tag{4}$$

Because A_2 is utility independent of $\{a^0_1\} \times A_{-\{1,2\}}$, there exist functions f_2 and g_2 such that, with $a'_2 = a^0_2$ in (1):

$$u(a^0_1, a_{-1}) = f_2(a^0_1, a_{-\{1,2\}}) + g_2(a^0_1, a_{-\{1,2\}})u(a_2, a^0_{-2}), \quad \text{for all } a \in A \tag{5}$$

Letting $a_2 = a^0_2$ in (5) and using $u(a^0) = 0$, we obtain $f_2(a^0_1, a_{-\{1,2\}}) = u(a^0_1, a^0_2, a_{-\{1,2\}})$ for all $a \in A$. Therefore:

$$u(a^0_1, a_{-1}) = u(a^0_1, a^0_2, a_{-\{1,2\}}) + g_2(a^0_1, a_{-\{1,2\}})u(a_2, a^0_{-2}), \quad \text{for all } a \in A \tag{6}$$

By restricted additive independence, there exists $a_2 \in A_2$ such that $u(a_2, a^0_{-2}) \neq 0$. Therefore, letting $a_{-\{1,2\}} = a^0_{-\{1,2\}}$ in (6) implies $g_2(a^0_{-2}) = 1$.

By restricted additive independence and (6), for all $a_{-\{1,2\}} \in A_{-\{1,2\}}$, there exist $a'_2, a''_2 \in A_2$ such that:

$$g_2(a^0_1, a_{-\{1,2\}})[u(a'_2, a^0_{-2}) - u(a''_2, a^0_{-2})] = g_2(a^0_{-2})[u(a'_2, a^0_{-2}) - u(a''_2, a^0_{-2})]$$

By restricted additive independence, $u(a'_2, a^0_{-2}) \neq u(a''_2, a^0_{-2})$ and, hence, $g_2(a^0_1, a_{-\{1,2\}}) = g_2(a^0_{-2}) = 1$ for all $a \in A$. Therefore:

$$u(a^0_1, a_{-1}) = u(a^0_1, a^0_2, a_{-\{1,2\}}) + u(a_2, a^0_{-2}), \quad \text{for all } a \in A \tag{7}$$

Using (7) in (4), we obtain:

$$u(a) = u(a^0_1, a^0_2, a_{-\{1,2\}}) + u(a_2, a^0_{-2}) + u(a_1, a^0_{-1}), \quad \text{for all } a \in A$$

Repeated application of this argument for the attributes 3, ..., $m - 1$ yields:

$$u(a) = u(a_m, a^0_{-m}) + \dots + u(a_2, a^0_{-2}) + u(a_1, a^0_{-1}) \text{ for all } a \in A. \quad \square$$

An immediate consequence of Theorem 1 is that u can be written as:

$$u(a) = \sum_{j \in M} u_j(a_j), \quad \text{for all } a \in A \tag{8}$$

where for every $j \in M$, $u_j: A_j \rightarrow \mathbb{R}$ is an induced utility function as introduced in Section 2.

4. Multiplicative decision criteria

Keeney and Raiffa (1976) prove that if each attribute space A_j is utility independent of A_{-j} , then the cardinal utility function u can be written as a *multilinear* function — see Keeney and Raiffa (1976, Chaps. 5 and 6) for details. In this section, we derive the special case of a multiplicative decision criterion with a weaker utility independence axiom and an assumption which requires the a_j^0 to lead to a worst alternative even when combined with some values of a_{-j} other than a_{-j}^0 .

To exclude degenerate cases, we assume that there is an $a^* \in A$ with $a_j^* \neq a_j^0$ for all $j \in M$ and $u(a^*) > u(a^0)$. For $j \in M$, define $\bar{A}_{-\{1, \dots, j\}} := \{a \in A_{-\{1, \dots, j\}} \mid a_k \neq a_k^0 \text{ for all } k = j + 1, \dots, m\}$. The utility independence condition is parallel to the one in the previous section, where a^0 is replaced with a^* , and $A_{-\{1, \dots, j\}}$ by $\bar{A}_{-\{1, \dots, j\}}$.

Restricted a^* utility independence:

- A_1 is utility independent of \bar{A}_{-1} ;
- A_2 is utility independent of $\{a_1^*\} \times \bar{A}_{-\{1,2\}}$;
- \vdots
- A_{m-1} is utility independent of $\{a_1^*\} \times \{a_2^*\} \times \dots \times \{a_{m-2}^*\} \times \bar{A}_{-\{1, \dots, m-1\}}$.

The following axiom requires a_j^0 to lead to a worst alternative when combined with certain values of a_{-j} .

Restricted zero independence:

$$u(a') \geq u(a_1^0, a_{-1}), \quad \text{for all } a, a' \in A \tag{9}$$

$$u(a') \geq u(a_1^*, a_2^0, a_{-\{1,2\}}), \quad \text{for all } a, a' \in A \tag{10}$$

\vdots

$$u(a') \geq u(a_1^*, \dots, a_{m-2}^*, a_{m-1}^0, a_m), \quad \text{for all } a, a' \in A \tag{11}$$

Restricted zero independence implies that a_1^0 is a worst-possible value for attribute one in an absolute sense: a_1^0 leads to a worst alternative not only combined with a_{-1}^0 but combined with *any* value of a_{-1} . The remaining restrictions imposed by the axiom are weaker because they apply only to some but not to all values of the remaining attributes.

Requiring restricted zero independence is close to but weaker than imposing the so-called zero condition on attributes $1, \dots, m - 1$, as is done in Miyamoto et al. (1998). The zero condition holds for attribute j if the level a_j^0 for attribute j makes the decision maker indifferent between all combinations of the other attributes.

The above two axioms lead to the multiplicative decision criterion described in the following theorem.

Theorem 2. *Let u satisfy restricted a^* utility independence and restricted zero independence. Assume that u is normalized so that $u(a^0) = 0$ and $u(a^*) = 1$. Then: $u(a) = \prod_{j \in M} u(a_j, a_{-j}^*)$ for all $a \in A$.*

Proof. By (9–11) we must have:

$$u(a_1^0, a_{-1}) = 0, \quad \text{for all } a \in A \tag{12}$$

$$u(a_1^*, a_2^0, a_{-\{1,2\}}) = 0, \quad \text{for all } a \in A \tag{13}$$

⋮

$$u(a_1^*, \dots, a_{m-2}^*, a_{m-1}^0, a_m) = 0, \quad \text{for all } a \in A \tag{14}$$

Because A_1 is utility independent of \bar{A}_{-1} , choosing $a'_{-1} = a_{-1}^*$ in (1) implies that there exist functions f_1 and g_1 such that:

$$u(a) = f_1(a_{-1}) + g_1(a_{-1})u(a_1, a_{-1}^*), \quad \text{for all } a \in A \tag{15}$$

Letting $a_1 = a_1^0$ in (15), (12) implies $f_1(a_{-1}) = 0$ for all $a \in A$.

Now let $a_1 = a_1^*$ in (15). Noting that $f_1(a_{-1}) = 0$ and $u(a^*) = 1$, it follows that $g_1(a_{-1}) = u(a_1^*, a_{-1})$ for all $a \in A$. Therefore:

$$u(a) = u(a_1^*, a_{-1})u(a_1, a_{-1}^*), \quad \text{for all } a \in A \tag{16}$$

Because A_2 is utility independent of $\{a_1^*\} \times \bar{A}_{-\{1,2\}}$, there exist functions f_2 and g_2 such that:

$$u(a_1^*, a_{-1}) = f_2(a_1^*, a_{-\{1,2\}}) + g_2(a_1^*, a_{-\{1,2\}})u(a_2, a_{-2}^*), \quad \text{for all } a \in A \tag{17}$$

[choose $a'_{-2} = a_{-2}^*$ in (1)]. Letting $a_2 = a_2^0$ in (17) and using (13), we obtain $f_2(a_1^*, a_{-\{1,2\}}) = 0$ for all $a \in A$. Now let $a_2 = a_2^*$ in (17) to obtain $g_2(a_1^*, a_{-\{1,2\}}) = u(a_1^*, a_2^*, a_{-\{1,2\}})$ for all $a \in A$. Hence:

$$u(a_1^*, a_{-1}) = u(a_1^*, a_2^*, a_{-\{1,2\}})u(a_2, a_{-2}^*), \quad \text{for all } a \in A \tag{18}$$

Using (18) in (16), we obtain:

$$u(a) = u(a_1^*, a_2^*, a_{-\{1,2\}})u(a_2, a_{-2}^*)u(a_1, a_{-1}^*), \quad \text{for all } a \in A$$

Repeated application of this argument for the attributes 3, ..., $m - 1$ yields:

$$u(a) = u(a_m, a_{-m}^*) \dots u(a_2, a_{-2}^*)u(a_1, a_{-1}^*) \text{ for all } a \in A. \quad \square$$

Again, it follows that there exist induced cardinal utility functions $u_j: A_j \rightarrow \mathbb{R}$ for all $j \in M$ such that:

$$u(a) = \prod_{j \in M} u_j(a_j), \quad \text{for all } a \in A \tag{19}$$

To conclude this section we note that the conditions in Theorem 2 are sufficient but not necessary for a multiplicative representation. For example, take $m = 2$, $A_1 = A_2 = \{1/2, 1\}$, and $u(a_1, a_2) = a_1 a_2$. In this case, (9) is not satisfied.

5. Efficiency

Consider now a situation where we have $n \in \mathbb{N}$ decision makers concerned with the multi-attribute alternatives in A . Let $N = \{1, \dots, n\}$ denote the set of decision makers. We use $u^i: A \rightarrow \mathbb{R}$ to denote the cardinal utility function of individual $i \in N$. The induced utility functions (as in Theorems 1 and 2) on A_j are denoted by $u_j^i: A_j \rightarrow \mathbb{R}$ for all $j \in M$ and all $i \in N$. For each agent $i \in N$, a^{0i} and a^{*i} are the alternatives corresponding to a^0 and a^* in the single-agent case. Note that these alternatives may be agent-specific.

We make the assumption that the set of utility vectors corresponding to outcomes in A , i.e. the set $\{(u^1(a), \dots, u^n(a)) \in \mathbb{R}^n \mid a \in A\}$, is a convex set. In this framework, it is of interest to identify those outcomes that are efficient in the sense that it is impossible to make everyone in N better off. Formally, an outcome $\hat{a} \in A$ is *efficient* if and only if, for all $a \in A$, there exists $i \in N$ such that $u^i(\hat{a}) \geq u^i(a)$. Hence, efficiency here corresponds to the concept of weak Pareto optimality — see Section 6.

The set of efficient outcomes can be characterized in the additive and multiplicative cases discussed in the previous sections. If all agents’ preferences have an additive representation, we obtain the following result. Let Δ^n denote the unit simplex in \mathbb{R}^n .

Theorem 3. *Let each u^i satisfy restricted a^{0i} utility independence and restricted additive independence, and assume $u^i(a^{0i}) = 0$ for every $i \in N$. Then an outcome $\hat{a} \in A$ is efficient if and only if there exists $t = (t^1, \dots, t^n) \in \Delta^n$ such that \hat{a} maximizes $\sum_{i \in N} t^i u^i(a)$ on A and, for all $j \in M$, \hat{a}_j maximizes $\sum_{i \in N} t^i u_j^i(a_j)$ on A_j .*

Proof. Because the set of utility vectors generated by the outcomes in A is convex, $\hat{a} \in A$ is efficient if and only if there exists $t \in \Delta^n$ such that \hat{a} maximizes $\sum_{i \in N} t^i u^i(a)$ on A .

Suppose $\hat{a} \in A$ maximizes $\sum_{i \in N} t^i u^i(a)$ on A for some $t \in \Delta^n$. By (8):

$$u^i(a) = \sum_{j \in M} u_j^i(a_j), \quad \text{for all } i \in N, \quad \text{for all } a \in A$$

Because:

$$\max \left\{ \sum_{i \in N} t^i \sum_{j \in M} u_j^i(a_j) \right\} = \sum_{j \in M} \max \left\{ \sum_{i \in N} t^i u_j^i(a_j) \right\} \tag{20}$$

\hat{a}_j maximizes $\sum_{i \in N} t^i u_j^i(a_j)$ for all $j \in M$

Conversely, suppose \hat{a}_j maximizes $\sum_{i \in N} t^i u_j^i(a_j)$ for all $j \in M$. Using (20), it follows that \hat{a} maximizes $\sum_{i \in N} t^i u^i(a)$ on A . \square

In the multiplicative case, we obtain an analogous result. As in Theorem 2, we now assume the existence of appropriate a^{0i} and a^{*i} for each $i \in N$.

Theorem 4. *Let each u^i satisfy restricted a^{*i} utility independence and restricted zero independence, and assume $u^i(a^{0i}) = 0$ and $u^i(a^{*i}) = 1$ for each $i \in N$. Then an outcome $\hat{a} \in A$ is efficient if and only if there exists $t = (t^1, \dots, t^n) \in \Delta^n$ such that \hat{a} maximizes $\prod_{i \in N} u^i(a)^{t^i}$ on A and, for all $j \in M$, \hat{a}_j maximizes $\prod_{i \in N} u_j^i(a_j)^{t^i}$ on A_j .*

Proof. Again, the convexity of the set of utility vectors generated by the outcomes in A implies that an outcome $\hat{a} \in A$ is efficient if and only if there exists $t \in \Delta^n$ such that \hat{a} maximizes $\prod_{i \in N} u^i(a)^{t^i}$ on A , where we use the convention $0^0 := 1$.

Suppose \hat{a} maximizes $\prod_{i \in N} u^i(a)^{t^i}$ on A for some $t \in \Delta^n$. By (19):

$$u^i(a) = \prod_{j \in M} u_j^i(a_j), \quad \text{for all } i \in N, \quad \text{for all } a \in A$$

and, therefore, \hat{a} maximizes:

$$\prod_{i \in N} \prod_{j \in M} u_j^i(a_j)^{t^i}$$

on A . Suppose first that there exists $k \in N$ such that $u^k(\hat{a})^{t^k} = 0$, in which case $\prod_{i \in N} u^i(a)^{t^i} = 0$. Note that this can occur only if $t^k > 0$ and $u^k(\hat{a}) = 0$. By assumption, $u^i(a^{*i}) = 1 > 0$ for all $i \in N$. By restricted zero independence and the assumption that the set of utility vectors generated is a convex set, there must be an $a^* \in A$ such that $u^i(a^*) > 0$ for all $i \in N$ and thus $\prod_{i \in N} u^i(a^*)^{t^i} > 0$, contradicting the observation that \hat{a} maximizes $\prod_{i \in N} u^i(a)^{t^i}$. Therefore, this case cannot occur, and we must have $u^i(\hat{a})^{t^i} > 0$ for all $i \in N$. Hence:

$$\max \left\{ \prod_{i \in N} \prod_{j \in M} u_j^i(a_j)^{t^i} \right\} = \prod_{j \in M} \max \left\{ \prod_{i \in N} u_j^i(a_j)^{t^i} \right\} \tag{21}$$

and, as in Theorem 3, it follows immediately that \hat{a}_j maximizes $\prod_{i \in N} u_j^i(a_j)^{t^i}$ on A_j for all $j \in M$.

Now suppose \hat{a}_j maximizes $\prod_{i \in N} u_j^i(a_j)^{t^i}$ on A_j for all $j \in M$. Because $u^i(a^{*i}) > 0$ for all $i \in N$, $u_j^i(a_j^{*i}) > 0$ for all $i \in N$ and for all $j \in M$. As before there are $a^{*j} \in A_j$ with $u_j^i(a^{*j}) > 0$ for all $i \in N$ and for all $j \in M$ and, thus, $\prod_{i \in N} u_j^i(a^{*j})^{t^i} > 0$ for all $j \in M$. Therefore, in order for \hat{a}_j to maximize $\prod_{i \in N} u_j^i(a_j)^{t^i}$, it must be the case that $u_j^i(\hat{a}_j)^{t^i} > 0$ for all $i \in N$ and for all $j \in M$. By (21), it follows that \hat{a} maximizes $\prod_{i \in N} u^i(a)^{t^i}$ on A . \square

6. Multi-attribute bargaining problems

The results of the previous sections can be illustrated by applying the additive and multiplicative criteria to specific group decision problems. For example, consider the problem of selecting outcomes from a feasible set of alternatives on the basis of the agents' preferences. Suppose the space of alternatives A is given, and we want to make a

selection from A for all profiles of individual utility functions within a given class. Because A is fixed throughout this section, we simplify notation by suppressing the dependence of the functions considered here on A . A profile of cardinal utility functions is denoted by $U = (u^1, \dots, u^n)$. The utility possibilities set generated by the profile U is given by:

$$\{x \in \mathbb{R}^n \mid \exists a \in A \text{ such that } x^i = u^i(a), \text{ for all } i \in N\}$$

Accordingly, the induced utility possibilities set for attribute $j \in M$ is:

$$\{x \in \mathbb{R}^n \mid \exists a_j \in A_j \text{ such that } x^i = u_j^i(a_j), \text{ for all } i \in N\}$$

Let \mathcal{U} be a set of admissible profiles $U = (u^1, \dots, u^n)$ of cardinal utility functions. Let 2^A be the set of all non-empty subsets of A . A social choice function is a mapping $c: \mathcal{U} \rightarrow 2^A$ such that, for all $U \in \mathcal{U}$ and for all $a, a' \in c(U)$, $u^i(a) = u^i(a')$ for all $i \in N$. That is, we make the assumption commonly used in cooperative models of bargaining that there is a unique utility vector associated with the selected outcomes. Interpreted as a bargaining problem, the disagreement point is given by $(u^1(a^{01}), \dots, u^n(a^{0n})) = (0, \dots, 0)$; recall that a^{0i} denotes the worst alternative for agent $i \in N$. A bargaining solution is a special case of a social choice function, where the only relevant features are the utility possibilities set and the disagreement point of a problem. Since this additional restriction is not necessary for the purposes of this section, we will use the more general choice function c as defined above.

The social choice function c induces a choice function $c_j: \mathcal{U} \rightarrow 2^{A_j}$ for each attribute $j \in M$. Specifically, for all $j \in M$ and all $U \in \mathcal{U}$, $c_j(U) = \{a_j \in A_j \mid a \in c(U)\}$. Note that we do not explicitly require that the same utility vector results for all chosen outcomes in the image of the induced choice functions — this property follows as a consequence of our choice independence axiom defined below.

Using the results of the previous sections, we can derive generalized utilitarian and generalized Nash social choice functions in this framework. In addition to the axioms on individual preferences leading to the additive and multiplicative representations derived above, we impose some restrictions on the choice function c .

First, we require c to be weakly Pareto optimal in the sense that it selects efficient outcomes only. Let $E(U)$ denote the set of efficient outcomes A for the profile $U \in \mathcal{U}$.

Weak Pareto optimality: For all $U \in \mathcal{U}$, $c(U) \subseteq E(U)$.

Furthermore, we impose an independence condition regarding the choice of the individual attributes.

Choice independence: For all $j \in M$, for all $U, \hat{U} \in \mathcal{U}$, if $u_j^i = \hat{u}_j^i$ for all $i \in N$, then $u_j^i(a_j) = u_j^i(\hat{a}_j)$ for all $a_j \in c_j(U)$ and for all $\hat{a}_j \in c_j(\hat{U})$.

Choice independence requires that the selection of each attribute is independent of the values of the remaining attributes and, thus, is another separability condition. Note that choice independence implies that, for a given profile, all attribute values selected by the

corresponding induced social choice function lead to the same utility vector (to see this, set $U = \hat{U}$ in the definition of choice independence).

Finally, we impose a regularity condition.

Regularity: There exist $\bar{U} \in \mathcal{U}$ and $h, k \in M$ such that $h \neq k$ and, for all $j \in \{h, k\}$, there exists a unique tangent hyperplane to the induced utility possibilities set for attribute j at the point $(\bar{u}_j^1(\bar{a}_j), \dots, \bar{u}_j^n(\bar{a}_j))$, where $\bar{a}_j \in c_j(\bar{U})$.

Regularity only requires the existence of *some* profile such that the tangent hyperplane to two induced utility possibilities at the utility vectors induced by the choice function is unique. Note that any bargaining solution defined on a standard domain satisfies a regularity condition of that type.

We call c a generalized utilitarian social choice function if there exists $t \in \Delta^n$ such that, for all $U \in \mathcal{U}$, $c(U)$ is a subset of the set of maximizers of $\sum_{i \in N} t^i u^i(a)$ on A . Analogously, c is a generalized Nash social choice function if there exists $t \in \Delta^n$ such that, for all $U \in \mathcal{U}$, $c(U)$ is a subset of the set of maximizers of $\prod_{i \in N} u^i(a)^{t^i}$ on A .

Theorem 5. *Let \mathcal{U} be the set of all profiles U such that the utility possibilities set associated with U is compact and convex and, for each $i \in N$, there exists $a^{0i} \in A$ such that $u^i(a) \geq u^i(a^{0i}) = 0$ for all $a \in A$, and u^i satisfies restricted a^{0i} utility independence and restricted additive independence. Let c be a social choice function satisfying weak Pareto optimality, choice independence, and regularity. Then c is a generalized utilitarian social choice function.*

Proof. Let $\bar{U} \in \mathcal{U}$ and $h, k \in M$ be as in the definition of the regularity axiom. By weak Pareto optimality and the definition of c , there exists $t \in \Delta^n$ such that \bar{a} is a maximizer of $\sum_{i \in N} t^i \bar{u}^i(a)$ on A for all $\bar{a} \in c(\bar{U})$. By Theorem 3, \bar{a}_j maximizes $\sum_{i \in N} t^i \bar{u}_j^i(a_j)$ on A_j for all $j \in M$. Regularity implies that, for $j \in \{h, k\}$, t is the only vector of coefficients for which \bar{a}_j maximizes this weighted sum on A_j .

Let $U \in \mathcal{U}$ be arbitrary. Let the profile \hat{U} be such that $\hat{u}_h^i = \bar{u}_h^i$ and $\hat{u}_j^i = u_j^i$ for all $i \in N$ and for all $j \in M \setminus \{h\}$. Let $\hat{a} \in c(\hat{U})$. By weak Pareto optimality, \hat{a} maximizes $\sum_{i \in N} \hat{t}^i \hat{u}^i(a)$ on A for some $\hat{t} \in \Delta^n$. Theorem 3 implies that \hat{a}_j maximizes $\sum_{i \in N} \hat{t}^i \hat{u}_j^i(a_j)$ on A_j for all $j \in M$. Choice independence implies that $\hat{u}_h^i(\hat{a}_h) = \hat{u}_h^i(\bar{a}_h)$. By regularity, $\hat{t} = t$.

Now let the profile \tilde{U} be such that $\tilde{u}_k^i = \bar{u}_k^i$ and $\tilde{u}_j^i = u_j^i$ for all $i \in N$ and for all $j \in M \setminus \{k\}$. Let $\tilde{a} \in c(\tilde{U})$. By weak Pareto optimality, \tilde{a} maximizes $\sum_{i \in N} \tilde{t}^i \tilde{u}^i(a)$ on A for some $\tilde{t} \in \Delta^n$. Again, Theorem 3 implies that \tilde{a}_j maximizes $\sum_{i \in N} \tilde{t}^i \tilde{u}_j^i(a_j)$ on A_j for all $j \in M$. Choice independence implies that $\tilde{u}_k^i(\tilde{a}_k) = \tilde{u}_k^i(\bar{a}_k)$. By regularity, $\tilde{t} = t$.

Let $a \in c(U)$. By choice independence, for all $i \in N$, $u_j^i(a_j) = u_j^i(\hat{a}_j)$ for all $j \in M \setminus \{h\}$ and $u_h^i(a_h) = u_h^i(\bar{a}_h)$. This implies that a_j maximizes $\sum_{i \in N} t^i u_j^i(a_j)$ on A_j for all $j \in M$ and, using the argument in the proof of Theorem 3, it follows that a maximizes $\sum_{i \in N} t^i u^i(a)$ on A . \square

Analogously, the generalized Nash social choice functions are implied in the multiplicative case. Because the proof of this result is parallel to the proof of the previous theorem, it is omitted.

Theorem 6. *Let \mathcal{U} be the set of all profiles U such that the utility possibilities set associated with U is compact and convex and, for each $i \in N$, there exist $a^{0i}, a^{*i} \in A$ such that $u^i(a) \geq u^i(a^{0i}) = 0$ for all $a \in A$ and $u^i(a^{*i}) = 1$, and u^i satisfies restricted a^{*i} utility independence and restricted zero independence. Let c be a social choice function satisfying weak Pareto optimality, choice independence, and regularity. Then c is a generalized Nash social choice function.*

An interesting feature of Theorems 5 and 6 is that the resulting social choice functions are such that the attributes can be separated in a bargaining process: it is sufficient to consider each attribute independently of the remaining ones. This considerably facilitates the application of the procedure to actual bargaining situations.

As an illustration, suppose the allocation of m goods in an n -agent economy is to be determined by means of a bargaining process. If the individual preferences satisfy the appropriate independence axioms, the results of this section can be used to justify the use of a generalized utilitarian solution or a generalized Nash solution. By adding a symmetry or anonymity condition, the utilitarian solution and the Nash solution are obtained. See also Kalai (1977) and Ponsati and Watson (1997) for examples and discussions of multiple-issue bargaining situations.

7. Concluding remarks

The results of this paper provide a decision-theoretic foundation for specific social choice procedures. In particular, multi-attribute decision criteria are employed. An assumption underlying our approach is that individual utility functions are of the same structure. One possibility for extending our work would be to examine situations where individual agents may have different types of utility functions over multi-attribute alternatives.

All our results have been formulated and derived for the case of cardinal utility. To a large extent they can almost without modification be stated and derived for the framework of decision making under risk, where lotteries are included and utility is of the von Neumann–Morgenstern type (von Neumann and Morgenstern, 1947). In bargaining this is a quite standard framework and, indeed, it would imply the convexity assumption made in Sections 5 and 6. Theorems 4 and 6 on the multiplicative case, however, would have to be modified, e.g. by requiring that efficient outcomes are always riskless.

Finally, we note that the results for the multiplicative case (the even-numbered theorems) are derived independently of the results for additive structures (the odd-numbered theorems). In contrast, Trockel (1998) derives the Nash bargaining solution by means of an exponential transformation from an additive setup. In order to employ an analogous procedure for the results obtained here, it would be required to formulate the axioms for the multiplicative case by using a structure that exactly parallels that of the additive model.

Acknowledgements

We thank the Nederlandse Organisatie voor Wetenschappelijk Onderzoek (NWO) for financial support under Grant No. B46-363. The comments of two anonymous referees and an associate editor led to substantial improvements.

References

- Blackorby, C., Primont, D., Russell, R.R., 1978. *Duality, Separability, and Functional Structure: Theory and Economic Applications*. North Holland, New York.
- Busch, L.-A., Horstmann, I.J., 1997. Bargaining frictions, bargaining procedures and implied costs in multiple-issue bargaining. *Economica* 64, 669–680.
- Fershtman, C., 1990. The importance of the agenda in bargaining. *Games and Economic Behavior* 2, 224–238.
- Fishburn, P.C., 1970. *Utility Theory for Decision Making*. John Wiley, New York.
- Kalai, E., 1977. Proportional solutions to bargaining situations: Interpersonal utility comparisons. *Econometrica* 45, 1623–1637.
- Keeney, R.L., Raiffa, H., 1976. *Decisions with Multiple Objectives: Preferences and Value Tradeoffs*. John Wiley, New York.
- McDonald, I.M., Solow, R.M., 1981. Wage bargaining and employment. *American Economic Review* 71, 896–908.
- Miyamoto, J., Wakker, P., Bleichrodt, H., Peters, H., 1998. The zero-condition: A simplifying assumption in QALY measurement and multiattribute utility. *Management Science* 44, 839–849.
- Nash, J.F., 1950. The bargaining problem. *Econometrica* 18, 155–162.
- Ponsati, C., Watson, J., 1997. Multiple-issue bargaining and axiomatic solutions. *International Journal of Game Theory* 26, 501–524.
- Trockel, W., 1998. Rationalizability of the Nash Bargaining Solution, Working Paper 291. Institute of Mathematical Economics, University of Bielefeld.
- von Neumann, J., Morgenstern, O., 1947. *Theory of Games and Economic Behavior*, 2nd Edition. Princeton University Press, Princeton.