

School of Economic Sciences

Working Paper Series
WP 2010-3

The Economic Value of Biochar in Crop Production and Carbon Sequestration

By

Suzette P. Galinato, Jonathan K. Yoder, and
David Granatstein

February 2010

The Economic Value of Biochar in Crop Production and Carbon Sequestration

Suzette P. Galinato^{a,*}, Jonathan K. Yoder^b and David Granatstein^c

^a IMPACT Center, School of Economic Sciences, Washington State University, Pullman, WA 99164 USA

^b School of Economic Sciences, Washington State University, Pullman, WA 99164 USA

^c Center for Sustaining Agriculture and Natural Resources, Washington State University, 1100 N. Western Ave., Wenatchee, WA 98801 USA

*Corresponding author. Tel.: +1 509-335-1408; fax: +1 509-335-3958. Email address: sgalinato@wsu.edu.

Abstract

This paper estimates the economic value of biochar application on agricultural cropland for carbon sequestration and its soil amendment properties. In particular, we consider the carbon emissions avoided when biochar is applied to agricultural soil, instead of agricultural lime, the amount of carbon sequestered, and the value of carbon offsets, assuming there is an established carbon trading mechanism for biochar soil application. We use winter wheat production in Eastern Whitman County, Washington as a case study, and consider different carbon offset price scenarios and different prices of biochar to estimate a farm profit. Our findings suggest that it may be profitable to apply biochar as a soil amendment under some conditions if the biochar market price is low enough and/or a carbon offset market exists.

Keywords: Biochar, Carbon sequestration, Crop, Farm profitability, Soil amendment

1. Introduction

Biochar is a charcoal-like material produced by the thermochemical pyrolysis of biomass materials. It is being considered as a potentially significant means of storing carbon for long periods to mitigate greenhouse gases (Laird, 2008). Much of the interest in biochar comes from studies of Amazonian soils that appear to have been amended with biochar, with significant improvements in soil quality and positive effects on crop yields (Lehmann et al., 2004). These changes have persisted for hundreds, if not thousands, of years. It is not yet known how long it takes for biochar to integrate with the soil and express its benefits. However, biochar represents a stable form of carbon and thus provides an intriguing potential carbon storage strategy as a soil amendment.

This study assesses the potential value of the use of biochar as a soil amendment from potential dual benefits of increased crop yields and returns from carbon sequestration, under a set of assumed conditions. We consider the effect of biochar in improving crop productivity by ameliorating the soil acidity. Next, we evaluate and aggregate emissions avoided and carbon sequestered when biochar replaces lime usage in the field. We then calculate the value of carbon offsets by using a low and high price range \$1/MT CO₂ and \$31/MT CO₂, assuming a carbon trading mechanism exists for biochar soil application. As a case study, we focus on wheat production in Washington State and examine farm profitability with and without the application of biochar. Our findings suggest that it may be profitable to apply biochar as a soil amendment under some conditions if the biochar market price is low enough and/or a carbon offset market exists.

The existing literature will be reviewed in detail below. As will be discussed, some of these studies focus on the properties and application rates of biochar and their impacts on agricultural

productivity, and some examine biochar's potential in sequestering carbon. However, this is the first study to our knowledge that links farm profitability with the economic value of biochar as a soil additive and as a source of carbon offset credits.

The following two sections provide a review of existing studies of the impacts of biochar soil application to crop productivity and to carbon (C) sequestration. We rely on the results of these studies in our analysis of the economic value of biochar as a soil amendment. Section 4 describes our methodology, and Section 5 presents estimated costs and returns in crop production for the case of wheat, with and without the application of biochar. Section 6 concludes.

2. Impacts of biochar on crop productivity — related studies

A number of studies have investigated the response of crops to biochar application. Table 1 presents summaries of a limited sample of these studies showing the impacts of biochar on crop response in terms of yield or plant biomass. Observed impacts vary depending on interactions between the types of biochar used, crop studied, soil type, local conditions, among others. Some studies have observed increased crop productivity from using biochar alone (Baum and Weitner, 2006; Chan et al., 2008). Other studies found a more positive crop response when biochar is applied together with fertilizers (Steiner et al., 2007). However, some studies have found negative crop response to biochar soil amendments. For example, Kishimoto and Sugiura (as cited in Chan and Xu, 2009) reported reductions in soybean yields with higher application rates of biochar. Collins (2008) found a decline in the root-shoot ratio of wheat in Quincy sand soil amended with peanut hull biochar and softwood bark biochar compared to unamended soil.¹ An increase in the root-shoot ratio of wheat, however, is found in Hale silt loam soil amended with

¹ The root/shoot ratio is the ratio of below-ground level biomass and above-ground level biomass.

softwood bark biochar. Collins (unpublished data) also found a significant increase in soil water holding capacity on silt loam soils but not on sandy soil. This could potentially increase crop yields in a dryland production region that is often water-limited for yield. Lehmann et al. (2006) discussed a greenhouse study in Columbia where biochar application led to low N availability to crops. Leguminous crops were found to compensate for this due to biological N₂ fixation induced by biochar application. On the other hand, non-legume crops were found to require additional N fertilizers due to low N availability. The above studies are controlled, small-scale experiments. At this point, it is not possible to draw conclusions on the effect of biochar that can be broadly applied, especially in temperate regions with younger soils (compared to highly weathered soils in more tropical environments). Furthermore, biochar itself will not contribute meaningful amounts of nutrients given its high stability, so the potential for biochar to reduce chemical fertilizer requirements remains unclear.² Nonetheless, the evidence from available studies does show that soil application of biochar often affects crop productivity and can be beneficial in some situations.

One consistent effect of biochar amendment has been change in soil pH (most frequently raising it), which implies a liming value of biochar. Collins (2008) found nearly a unit increase in soil pH with biochar derived from herbaceous feedstocks (switchgrass, digested fiber) and 0.5 to 1 unit increase in the soil pH with biochar derived from woody sources (softwood bark, wood pellets) (Table 2). Van Zwieten et al. (2007) also reported an increase in the soil pH and 30 to 40 percent increase in the height of wheat when biochar was applied to an acidic soil. Rondon et al. (2007) credited the improvement of bean productivity due to the elevation of soil pH and other

² In Collins (2008), total nitrogen in the soil is found to increase (although at small amounts) after addition of biochar. This does not, however, imply that a lesser amount of N fertilizer may be needed when biochar is added to the soil. N in biochar is not available to plants. It is fused in the C matrix (H. Collins, personal communication, 2009).

soil nutrients as a consequence of biochar use. Biochar may be considered a potential substitute for agricultural lime, especially in agricultural regions that have acidic soils. However, a unit change in the soil pH would require 0.54 metric ton (MT) to 3.91 metric tons per acre of agricultural lime, depending on the soil type (CPHA, 2002). For example, a sand soil type requires 0.54 MT/acre of lime to increase the soil pH by a unit as compared to about 17 MT/acre biochar³ needed to achieve the same desired change in soil pH (Collins, 2008). Thus, it may not be economically feasible for farmers to use biochar in crop production solely for pH adjustment since it would entail a relatively higher cost compared to agricultural lime. On the other hand, other potential benefits from adding biochar to the soil such as avoided emissions of lime and the capacity of biochar to sequester carbon (to be discussed in the next section) should be considered. It is possible that the economic returns from using biochar may be higher than from using lime after accounting for any other non-pH related plant growth benefits or carbon offset credits, were they to be available to farmers.

3. Biochar carbon sequestration — related studies, policy and program

Biochar is produced through the process of pyrolysis. The three main products of pyrolysis: liquid (bio-oil), biochar, and gas, can influence the global carbon (C) cycle in two ways. First, all three pyrolysis products may be used as an energy source that can displace fossil energy use. Second, if the carbon-rich and stable biochar is produced from a biomass feedstock that removes carbon dioxide (CO₂) from the air via photosynthesis, which would otherwise have decomposed,

³ This refers to Quincy sand soil type. The biochar requirement to raise the pH by a unit depends on the type of char used, e.g., switchgrass, digested fiber, bark, etc. Assuming an average pH increase across the chars of 0.058 pH unit/ton of biochar, it would require about ~17.24 tons of biochar to increase the soil pH by 1 unit.

then char-amended land becomes a carbon sink for more intensive and long-lasting carbon storage.

Lehmann et al. (2006) estimated an annual sequestration of 0.2 Pg C (200 million metric tons) through slash-and-char (instead of slash-and burn) and biochar application to the soil. Furthermore, the study reported that low-temperature pyrolysis of biomass combined with the capture of gas and liquid products for bioenergy production and soil application of biochar, could sequester the equivalent of about 10% of the annual US fossil-fuel emissions.

Laird (2008) proposed a national system of distributed fast pyrolyzers for converting biomass into bio-oil, gas and char. Similar to Lehmann et al. (2006), he assumed that bio-oil and gas are used as energy sources that can displace fossil fuel use, while char was applied to agricultural soils. Assuming the United States can produce 1.1 billion metric tons of biomass per year from harvestable forest and crop lands, the implementation of Laird's proposal could displace 25% of the nation's fossil fuel oil consumption per year. The study also estimated the aggregate carbon credit for fossil fuel displacement and biochar C sequestration to be 10% of the average annual US CO₂-C emissions.

The carbon content of biochar varies depending on the feedstock. Collins (2008) showed biochar carbon content (from slow pyrolysis) ranging from 61% to 80%, the highest being from wood pellets (Table 2). Woody feedstocks (bark, wood pellets) tended to have a higher carbon content compared to herbaceous feedstocks (switchgrass, digested fiber). Based on these figures, approximately 0.61 to 0.80 MT of carbon (or 2.2 to 2.93 MT of CO₂)⁴ is sequestered for every ton of biochar applied to the soil.

⁴ To convert from carbon to carbon dioxide, multiply by 44/12 (~3.67) (Blasing et al., 2004).

Incentives for greenhouse gas mitigation such as carbon market offset credits may tip the scale in favor of biochar as a soil amendment rather than as a renewable energy source. At the international level, the Kyoto Protocol under the United Nations Framework on Climate Change (UNFCCC) only allows C sequestration from afforestation and reforestation in the trading program established under the Clean Development Mechanism (CDM) (UNFCCC-CDM, 2009). Carbon sequestration in agricultural crops and soils is not currently eligible under CDM (Lehmann et al., 2006; FAO, 2009). In the United States, the Chicago Climate Exchange (CCX) has developed standardized rules for soil carbon management offsets in the agricultural sector. Eligible projects are conservation tillage and grass planting, which have to be enrolled with a CCX-registered Offset Aggregator.

In 2008, prices of traded CO₂ offsets on the Chicago Climate Exchange were volatile, ranging from \$1 to \$7.40 per metric ton of CO₂ (CCX, 2008). During the same year, the market prices of CO₂ offsets in the European Climate Exchange varied between \$17 and \$31 per metric ton of CO₂ (ECX, 2008). The differences in price across markets are in part due to the fact that participation in the CCX is currently optional; no entity is legally required to participate in this exchange. However, recent policy discussions at the national level suggest increased momentum toward a binding national carbon market. In the following section, we estimate the value of biochar as an input in crop production and as an instrument in C sequestration. We assume that a carbon market exists for avoided emissions and C sequestration due to use of biochar as soil amendment. Also, for the value of potential CO₂ offset, we use a low and high value of \$1/MT CO₂ and \$31/MT CO₂.

4. Estimated costs and returns for using biochar as soil amendment

In this section, we estimate the value of biochar as a soil amendment and the economic returns to farmers under a set of assumed conditions. The calculation is done in two stages. First, we assess: (a) the avoided emissions from the soil application of biochar instead of agricultural lime, excluding the energy and emissions from transporting and spreading the material; and (b) the amount of carbon sequestered from biochar application. Combined, the emissions avoidance and sequestration effects are counted as CO₂ offsets that can be sold under an assumed set of carbon offset prices. Second, we calculate the profit of crop production given two scenarios —without biochar but with lime application to the soil, and with biochar application as a substitute for lime. We focus on dryland wheat production in the state of Washington as a case study.

4.1. Estimates of avoided emissions

Gaunt and Lehmann (2008) and McCarl et al. (2009) estimated the avoided greenhouse gas emissions of applying biochar to agricultural land in terms of reduced agricultural input requirements due to a crop's improved use of nutrients. This translates to reductions in both fertilizer use and nitrous oxide emissions in fertilized fields. However, the dynamics of the relationship between fertilizers and biochar are not included in our analysis since the effects of biochar on the utilization intensity of fertilizers depend on various factors like the type of crop studied, soil type, soil quality and biochar type. On the other hand, there is strong consensus about biochar's positive effects on soil pH that is similar to liming effects, as exemplified by studies discussed in section 2.

Agricultural lime is commonly applied to soils to ameliorate the soil pH. The recommended rates of lime application in western Washington range from 0.91 metric ton to 4.54 metric tons per acre every 3 to 5 years, roughly (C. Cogger, personal communication, 2009). Less or no lime

is needed in Washington east of the Cascades because the native soil pH is high (R. Koenig, C. Kogger and J. Davenport, personal communications, 2009). However, some soils in eastern Washington with a long history of urea or ammonium-based nitrogen fertilizers have experienced a reduction in pH to a low enough level to justify the need for lime. When there is a soil acidity problem, lime applications range from 0.91 metric ton to 2.72 metric tons per acre (J. Davenport, personal communication, 2009).

West and McBride (2005) estimated the net CO₂ emission from application of agricultural lime at about 0.059 metric ton C (or 0.22 metric ton CO₂) per ton of limestone, based on the chemical reaction of lime in the soil and transportation of lime-derived bicarbonates to the ocean (via leaching and precipitation). This is the amount of emissions that can potentially be avoided by replacing lime with biochar. Using the CO₂ offset price range of \$1 to \$31/MT CO₂, the value of avoided emissions amounts to \$0.22-\$6.82 per metric ton of lime.

4.2 Estimates of biochar carbon sequestration

Biochar from herbaceous and woody feedstock sources are found to have a carbon content of 60.5%-66.7% and 74.5%-80%, respectively. We can assume from these figures that for every ton of biochar applied to the soil, 0.61 to 0.80 ton of carbon (equivalent to 2.2-2.93 tons of CO₂) can be sequestered (Collins, 2008). Using the highest carbon content of the wood-based biochar (i.e., 80%) and the CO₂ offset price range, the approximate value of biochar C sequestration is \$2.93-\$90.83 per metric ton of biochar.

4.4 Costs and returns of crop production

We examine the potential economic returns to farmers if they utilize biochar as a substitute for agricultural lime under three price scenarios: (a) \$114.05 per metric ton based on the energy content of a wood-based biochar; (b) \$87 per metric ton; and (c) \$350.74 per metric ton. The first

value represents the opportunity cost of the foregone use of biochar as energy source. A wood-based biochar has an average energy content of 12,500-12,500 BTU/lb (Dynamotive Energy Systems, 2007). The energy content of the Central Appalachian coal is 12,500 BTU/lb and its price is \$116.38 per metric ton as of 2008 (EIA, 2009). Using the energy content as basis, the combustion value of biochar is 98% that of Central Appalachian coal, or \$114.05/metric ton. The latter two prices are adopted from the estimated break-even prices of biochar in Granatstein et al. (2009).

Wheat, a key economic crop in Washington, belongs to a group of crops that can tolerate slightly acidic (i.e., 6.0-6.5) soil pH (CPHA, 2002). In general, wheat tends to favor soil pH between 6 and 7 (Beegle and Lingenfelter, 2005). We focus on changes in winter wheat yield given changes in the soil pH as a case study. The crop yield is estimated through the following equation adopted from Mahler (1986):

$$\text{Winter Wheat Yield} = -2,960.56 + 1,530 \text{ SOILPH} \quad (1)$$

where *winter wheat yield* is in kilograms per hectare, and *SOILPH* refers to the value of the soil pH of Palouse silt loam. Assuming the base soil pH of 4.5 for this soil type from Collins (2008), increasing the soil pH to 6 would require 2.59 metric tons of limestone per acre (CPHA, 2002) or 30.62 metric tons of biochar per acre (H. Collins, unpublished data).⁵

Using equation (1) with soil pH of 4.5, wheat yield is estimated at about 3,924.44 kg per hectare or 58.36 bushels per acre.⁶ On the other hand, with a soil pH of 6, the estimated wheat

⁵ Note that the impact of char on soil pH depends on the soil type and the type of char. For the Palouse silt loam soil type, an average pH increase across the chars is ~0.049 pH unit/ ton of biochar per acre. Based on this, it would require about 30.62 tons/acre to increase the soil pH from 4.5 to 6 (i.e., increase by 1.5 units).

⁶ Conversion: 1 bushel per acre x 0.06725 = 1 metric ton (or 1,000 kg) per hectare (Prairie Grains Magazine, 2003).

yield is about 6,219.44 kg per hectare or 92.5 bushels per acre. Profits from winter wheat production, with and without the application of biochar, are calculated as follows:

Without biochar or agricultural lime:

$$\text{Profit}_1 = P_W * Q_1 - \text{Total Cost}; \quad (2)$$

Without biochar, with agricultural lime:

$$\text{Profit}_2 = P_W * Q_2 - \text{Total Cost} - P_L * \text{AGLIME}; \quad (3)$$

With biochar, without agricultural lime:

$$\text{Profit}_3 = P_W * Q_2 + \text{COFFSET} - \text{Total Cost} - P_{Bi} * \text{BCHAR}, \quad (4)$$

where P_W refers to the Fall 2008 contract price of winter wheat, which is \$7.50 per bushel (Union Elevator, 2008). Q_1 is the estimated yield of winter wheat given a soil pH of 4.5 in equation (2) and Q_2 in equations (3) and (4) is the estimated wheat yield given a soil pH of 6. The product of P_W and Q gives the *revenue* in Table 3 below. P_L is the price of lime at \$51.53 per metric ton for a 100-lb bag in 2008 (Steve Eckhart, J.A. Jack and Sons, personal communication, 2009). *AGLIME* refers to the application rate of agricultural lime (2.59 metric tons per acre); P_B means the biochar price; *BCHAR* represents the application rate of biochar (30.62 metric tons per acre); and *COFFSET* is the value of carbon offset from avoided emissions and biochar C sequestration by replacing lime with biochar. *Total Cost* denotes the sum of fixed cost and variable cost of winter wheat crop production based on the Eastern Whitman County 2008 Enterprise Budget (Painter, unpublished), exclusive of lime or biochar cost.

Table 3 shows the estimated profits given the addition of agricultural lime or biochar to the soil and different price scenarios. As discussed above, the yield of wheat is higher when the soil pH improves; hence, the revenue is higher with the application of lime or biochar than without.

A farmer will gain a profit even when there is an additional cost of agricultural lime. When biochar replaces agricultural lime, on the other hand, getting a profit or a loss will depend on the price of biochar and value of sequestered carbon. Without a carbon offset market (COFFSET is zero in equation 4), the price of biochar has to be about \$9.10/MT in order for a farmer to break even (profit = 0)⁷ and about \$4.34/MT for profit to be equal to that of agricultural lime scenario, excluding the transportation and application costs of biochar.

Suppose now that a carbon offset market exists. At \$1/MT CO₂, the farmer loses income given any of the biochar price scenarios; and at \$31/MT CO₂, losses are also incurred if the price of biochar is high, i.e., at \$351/MT or \$114/MT. This means that the income from offsets is not enough to support the adoption of biochar in agricultural production. If, however, the price of biochar goes down to \$87/MT and the carbon offset is priced at \$31/MT CO₂, a profit is gained and it is higher than the case where agricultural lime is used instead. This implies that when the price of biochar is low enough, the income derived from carbon offsets can outweigh the cost of biochar. It should be noted that the production cost does not include the cost of transporting and applying lime or biochar to agricultural land (machinery and labor cost). Including these would likely further drive up the estimated losses or decrease any profit earned.

The differences in potential profit given varying prices of biochar and carbon offset are further illustrated in Figure 1. Profit declines as the price of biochar increases, holding other things constant. When the carbon offset price is \$31/MT CO₂, the farmer will break even if the price of biochar is approximately \$100.52/MT. The farmer's estimated profit with biochar application will be equal to the profit with agricultural lime application (\$146/acre) if the price of

⁷ From equation (4), the price of biochar (P_B^*) at which profit is zero is derived by calculating:
 $P_B^* = (P_W * Q + COFFSET - \text{Total Cost}) / BCHAR.$

biochar is about \$95.75/MT. On the other hand, at a carbon offset price of \$1/MT CO₂, a profit of about \$62.84 per acre is gained when the price of biochar is \$10/MT. At this lower offset price, the farmer will break even if the price of biochar is approximately \$12.05/MT.

5. Conclusions

Our quantitative analyses focus on using biochar as a soil additive and its potential carbon sequestration benefits for agricultural uses. We find that biochar soil application can be economically feasible given the following scenarios:

- If there exists a carbon market that recognizes the avoided emissions and carbon sequestration due to the application of biochar to agricultural soils. This is a necessary condition if biochar will be promoted as a technology for carbon sequestration; and
- If the market price of biochar is low enough so that a farmer will earn a profit after applying biochar to the crop field (i.e., in our case study, lower than \$12.05/MT and \$100.52/MT when the price of carbon offset is \$1/MT CO₂ and \$31/MT CO₂, respectively).

It is clear that biochar has potential as a soil amendment and its value as such would likely increase as social and regulatory interest in carbon sequestration increases because of the longevity of carbon in the soil. However, more substantial increases in crop production need to be documented across a range of crops and soils that can add value to the farm at a level beyond the estimated value of biochar for carbon sequestration. Under the current economic situation, growers are unlikely to adopt biochar use without greater payback. Also at this time, even if growers found biochar beneficial, they could face difficulty in sourcing quantities large enough for farm application.

Many niche opportunities for biochar use are also possible, including soil amendment and compost use outside of agriculture such as urban gardens, lawns, parks, and ball fields. Biochar can be suitable as a precursor to activated carbon commonly utilized in industrial filtration process (Azargohar and Dalai, 2006) like municipal wastewater treatment (e.g. Bansode et al., 2003, Ng et al., 2003) and other water and air filtering systems (Kearns, 2008; Lima et al., 2008). Biochar can also be used as an energy source; as combustion fuel to power the pyrolysis process; as a gasifier feedstock (Boateng, 2007; Polagye et al., 2007); or for water heating and cooking (IBI, 2009; Johannes, 2008). The extent of developing these markets, of course, depends on many factors associated with the cost of biochar production relative to existing alternatives, as well as the relative effectiveness of biochar from pyrolysis for intended uses. The question is whether or not it would be more economically valuable to use biochar as a soil amendment rather than for energy production or other alternative uses. For instance, it may be more economically attractive to burn it to generate energy (i.e., as a substitute for coal) if the energy content of biochar is high. However, the process loses the added benefits of applying biochar to soils, such as gains in agricultural productivity due to soil quality improvement and payments for carbon sequestration.

Acknowledgments

We thank the Washington State Department of Ecology for funding the project, “Use of Biochar from the Pyrolysis of Waste Organic Material as a Soil Amendment”. We also thank Dr. Hal Collins for providing us the data on the carbon content of biochar, the soil scientists at Washington State University for additional references about soil fertility in eastern Washington and Mark Fuchs for his valuable comments and suggestions.

References

- Azargohar, R., Dalai, A. K., 2006. Biochar as a precursor of activated carbon. *Applied Biochemistry and Biotechnology* 129–132, 762-773.
- Bansode, R., Losso, J., Marshall, W., Rao, R., Portier, R., 2003. Pecan shell-based granular activated carbon for treatment of chemical oxygen demand (COD) in municipal wastewater. *Bioresource Technology* 94, 129-135.
- Baum, E., Weitner, S., 2006. Biochar application on soils and cellulosic ethanol production. Clean Air Task Force, Boston, MA, U.S.A.
- Beegle, D.B., Lingenfelter, D.D., 2005. Soil acidity and aglime. *Agrifacts 3*. College of Agricultural Sciences, Cooperative Extension, Pennsylvania State University.
<http://cropsoil.psu.edu/Extension/Facts/AgFact3.pdf>.
- Blackwell, P., Riethmuller, G., Collins, M., 2009. Biochar application to the soil. In: J. Lehmann and S. Joseph (Eds.), *Biochar for Environmental Management: Science and Technology*. Earthscan Publications Ltd., United Kingdom, pp. 207-222.
- Blasing, T.J., Broniak, C.T., Marland, G., 2004. Estimates of monthly CO₂ emissions and associated ¹³C/¹²C values from fossil-fuel consumption in the U.S.A. In: *Trends: A Compendium of Data on Global Change*, Carbon Dioxide Information Analysis Center. Oak Ridge National Laboratory, U.S. Department of Energy, Oak Ridge, TN, U.S.A.
- Boateng, A. A., 2007. Characterization and thermal conversion of charcoal derived from fluidized-bed fast pyrolysis oil production of switchgrass. *Industrial & Engineering Chemistry Research* 46(26), 8857-8862.
- California Plant Health Association (CPHA). 2002. *Western Fertilizer Handbook*, ninth ed. Interstate Publishers, Inc., Danville, IL., U.S.A.

- Chan, K.Y., Van Zwieten, L., Meszaros, I., Downie, A., Joseph, S., 2008. Using poultry litter biochars as soil amendments. *Australian Journal of Soil Research* 46(5), 437–444.
- Chan, K.Y., Xu, Z., 2009. Biochar: nutrient properties and their enhancement. In: J. Lehmann and S. Joseph (Eds.). *Biochar for Environmental Management: Science and Technology*. Earthscan Publications Ltd., United Kingdom, pp. 67-81.
- Chicago Climate Exchange (CCX), 2008. December 2008 market summary. *CCS Market Report* 5(12), 1-4.
http://www.chicagoclimatex.com/docs/publications/CCX_carbonmkt_V5_i12_dec2008.pdf.
- Collins, H., 2008. Use of biochar from the pyrolysis of waste organic material as a soil amendment: laboratory and greenhouse analyses.” A quarterly progress report prepared for the Biochar Project (December 2008).
- Dynamotive Energy Systems, 2007. Table 1: Comparison of fuel properties-wood based pyrolysis fuel.”
- Energy Information Administration (EIA), 2009. Coal news and market reports, average weekly coal commodity spot prices, January to December 2008.”
<http://www.eia.doe.gov/cneaf/coal/page/coalnews/cnmarchive.html>.
- European Climate Exchange (ECX), 2008. Certified emission reduction futures contracts — 2008 historic data. <http://www.ecx.eu/CER-Futures>.
- Food and Agriculture Organization (FAO), Forest Resources Division, 2009. Afforestation and reforestation projects under the Clean Development Mechanism of the Kyoto Protocol. Fact sheet. <http://www.fao.org/forestry/media/8953/1/0/>.

- Gaunt, J.L., Lehmann, J., 2008. Energy balance and emissions associated with biochar sequestration and pyrolysis bioenergy production. *Environmental Science and Technology* 42(11), 4152–4158.
- Glaser, B., Lehmann, J., Zech, W., 2002. Ameliorating physical and chemical properties of highly weathered soils in the tropics with charcoal – a review. *Biology and Fertility of Soil* 35, 219–230.
- Granatstein, D., Kruger, C.E., Collins, H., Galinato, S., Garcia-Perez, M., Yoder, J., 2009. Use of biochar from the pyrolysis of waste organic material as a soil amendment. Final Project Report. Center for Sustaining Agriculture and Natural Resources, Washington State University, Wenatchee, WA. 181 pp. <http://www.ecy.wa.gov/biblio/0907062.html>.
- International Biochar Initiative (IBI), 2009. Biochar kiln designs for small farms. <http://www.biochar-international.org/projectsandprograms/memberprojects.html>.
- Johannes, H., 2008. Energy efficient stoves that burn biochar and biosmoke only and can save the world's forests. <http://www.hedon.info/BP20:BiocharBriquettingAndBurning>.
- Kearns, J., 2008. NGOs team up to offer climate solutions, enhanced sustainable agriculture, and clean drinking water.” <http://globalclimatesolutions.org/2008/10/25/ngos-team-up-to-offer-climate-solutions-enhanced-sustainable-agriculture-and-clean-drinking-water/>.
- Laird, D.A., 2008. The charcoal vision: A win-win-win scenario for simultaneously producing bioenergy, permanently sequestering carbon, while improving soil and water quality. *Agronomy Journal* 100(1), 178-181.
- Lehmann, J., Kern, D.C., Glaser, B., Woods, W.I. (eds.), 2004. *Amazonian Dark Earths: Origin, Properties, Management*. Kluwer Academic Publishers, New York, 523 pp.

- Lehmann, J., Gaunt, J., Rondon, M., 2006. Bio-char sequestration in terrestrial ecosystems — a review. *Mitigation and Adaptation Strategies for Global Change* 11, 403-427.
- Lima, I.M., McAloon, A., Boateng, A.A., 2008. Activated carbon from broiler litter: process description and cost of production. *Biomass and Bioenergy* 32, 568-572.
- Mahler, R.L., 1986. Evaluation of soil pH manipulation on crop production in northern Idaho. *Communications in Soil Science and Plant Analysis* 17(9), 905-919.
- McCarl, B.A., C. Peacocke, R. Chrisman, C.-C. Kung, and R.D. Sands. 2009. Economics of biochar production, utilisation and GHG offsets. In: J. Lehmann and S. Joseph (Eds.). *Biochar for Environmental Management: Science and Technology*. Earthscan Publications Ltd., United Kingdom, pp. 341-356.
- Ng, C., Marshall, W.E., Rao, R.M., Bansode, R.R., Losso, J.N., 2003. Activated carbon from pecan shells: process description and economic analysis. *Industrial Crops and Products* 17, 209–217.
- Painter, K., Unpublished. 2008 Crop Rotation Budgets, Over 18" Precipitation Zone Under Conventional Tillage, Whitman County, Washington.
- Polagye, B.L., Hodgson, K.T., Maltea, P.C., 2007. An economic analysis of bio-energy options using thinnings from overstocked forests. *Biomass and Bioenergy*, 31, 105-125.
- Prairie Grains Magazine, 2003. Grain math weights, measures and conversion factors.
<http://www.smallgrains.org/springwh/June03/weights/weights.htm>.
- Rondon, M. A., Lehmann, J., Ramírez, J., Hurtado, M., 2007. Biological nitrogen fixation by common beans (*Phaseolus vulgaris* L.) increases with bio-char additions. *Biology and Fertility of Soils* 43(6), 699-708.

- Steiner, C., Teixeira, W.G., Lehmann, J., Nehls, T., Vasconcelos de Macêdo, J.L., Blum, W.E.H., Zech, W., 2007. Long term effects of manure, charcoal and mineral fertilization on crop production and fertility on a highly weathered Central Amazonian upland soil. *Plant Soil* 291, 275-290.
- Union Elevator and Warehouse Co., 2008. Cash Prices: FOB Lind — August 2008 price posted on July 15, 2008. <http://www.unionelevator.com/>.
- United Nations Framework Convention on Climate Change, Clean Development Mechanism (UNFCCC-CDM), 2009. Methodologies for afforestation and reforestation CDM project activities. <http://cdm.unfccc.int/methodologies/ARmethodologies/index.html>.
- Van Zwieten, L., 2007. Research confirms biochar in soils boosts crop yields. <http://biopact.com/2007/06/research-confirms-biochar-in-soils.html>.
- West, T.O., McBride, A.C., 2005. The contribution of agricultural lime to carbon dioxide emissions in the United States: dissolution, transport and net emissions. *Agriculture Ecosystems and Environment* 108, 145-154.
- Young, M.J., Johnson, J.E., Abrams, M.D., 1996. Vegetative and Edaphic Characteristics on Relic Charcoal Hearths in the Appalachian Mountains. *Plant Ecology* 125(1), 43-50.

Table 1

Summary of studies on biochar used as a soil amendment.

Author(s)	Location	Soil type	Type of biochar	Biochar application rate	Crop
Kishimoto and Sugiura (1985) ^{a,b}	Japan	Volcanic ash soil, loam	Unknown wood	0, 0.2, 2.0 and 6.1 t/acre	Soybean
Crop response: At 0.2 t/acre, increased yield by 51%. At 2 t/acre and 6.1/acre, reduced yield by 37% and 71%, respectively. Reductions were attributed to micronutrient deficiency induced by an increase in pH.					
Mikan and Abrams (1995) ^b	United States (Pennsylvania)	Forest area on relic charcoal hearths	Wood for charcoal production	Unknown	Vegetation in hearth and non-hearth areas compared after 110 years
Crop response: Tree density and basal area were reduced by 40%.					
Young et al. (1996)	United States (Appalachian mountains)	Forest area on relic charcoal hearths	Wood for charcoal production	Unknown	Trees
Crop response: Lower overstory tree cover and density on relic charcoal hearths than on adjacent, non-hearth areas. The richness and diversity of overstory and understory tree cover as well as ground vegetation were consistently lower on hearths.					
Glaser et al. (2002)	Brazil	Xanthic Ferralsol	Secondary forest wood	0, 27.2 and 54.7 t/acre	Rice, Cowpea
Crop response: At application rate of 27.2 t/acre, biomass increased by: 20%, rice; 50%, cowpea compared to control treatment where no biochar was applied. At application rate of 54.7 t/acre, biomass of cowpea increased by 100%.					
Steiner (2006) ^c	Brazil	Xanthic Ferralsol	Wood	4.5 t/acre	Banana
Crop response: Reduced soil acidity and increased K uptake					
Yamamoto et al. (2006) ^c	Indonesia	Acid soil	Bark	6.1 t/acre	Maize
Crop response: Higher yields with biochar and fertilizer, than fertilizer alone					
Steiner et al. (2007)	Brazil	Xanthic Ferralsol	Secondary forest wood	4.5 t/acre	Rice, Sorghum
Crop response: Charcoal plus mineral fertilizer improved yield by a factor of 1.5-2 and improved stover by a factor of 1.3-1.4. Using charcoal plus compost and/or fertilizer, yields are consistently greater (i.e., 4 to 12 times greater) compared to using fertilizer alone.					

Sources: ^a Adopted from Glaser et al. (2002); ^b Adopted from Chan and Xu (2009); ^c Adopted from Blackwell et al. (2009).

Table 1 (continued)

Summary of studies on biochar used as a soil amendment.

Author(s)	Location	Soil type	Type of biochar	Biochar application rate	Crop
Van Zwieten (2007)	Australia	Semi-tropical soil		4 t/acre	Wheat, soybeans
Crop response: Wheat: biomass tripled. Soybeans: biomass more than doubled. Percentage increase in biomass is the same when nitrogen fertilizer is applied together with biochar. Biochar raised soil pH at about 1/3 the rate of lime.					
Van Zwieten et al. (2007) ^b	Australia	Ferrosol	Paper mill sludge	4 t/acre	Wheat
Crop response: 30-40% increase in wheat height in acidic soil but not in alkaline soil. Response was attributed mainly to the liming value of biochar.					
Collins (2008)	Washington	Quincy sand, Hale silt loam	Peanut hull (PH), fir bark (SB)	0, 5, 10 and 20 t/acre	Wheat
Crop response: Quincy: Root-shoot ratio of wheat decreased in all application rates of biochar. Hale: Using PH, decline in root-shoot ratio of wheat at 10 t/acre of biochar compared to nil; no change at 5 t/acre and 20 t/acre. Hale: Using SB, root-shoot ratio of wheat increased in all treatments. 0.5 to 1 unit increase in soil pH due to biochar addition					
Chan et al. (2008)	Australia	Alfisol	Poultry litter	0, 4, 10.1 and 20.2 t/acre	Radish
Crop response: With biochar, without N fertilizer: yield increased from 42% at 4 t/acre of biochar to 96% at 20.2 t/acre of biochar, relative to the yield from unamended control.					
Van Zwieten et al. (2008) ^c	Australia	Ferrosol	Poultry litter (PL), Paper mill waste (PM)	Maize: 0.2-20.2 t/acre PL. Beans: 4 t/acre PL and PM versus 1.2 t/acre lime	Maize, Faba beans
Crop response: Maize: 51% yield increase at 4 t/acre; and 109% yield increase at 20.2 t/acre compared to nil. Beans: Yields are highest with biochar plus fertilizer, compared to biochar alone. PL biochar outperformed lime amendment.					

Sources: ^a Adopted from Glaser et al. (2002); ^b Adopted from Chan and Xu (2009); ^c Adopted from Blackwell et al. (2009)

Table 2

Selected characteristics of six biochars (slow pyrolysis at 500oC) used in the laboratory analyses.

Source of biochar	Biochar Characteristics					
	C	N	S	C:N	C:S	pH
	----- % -----					
Switchgrass	60.5	2.06	0.20	30	350	9.4
Digested fiber	66.7	2.23	0.30	30	228	9.3
Peanut hull	70.6	1.74	0.04	41	1203	9.6
Bark (UGA)	74.5	0.34	0.03	220	2833	7.6
Softwood bark	77.8	0.44	0.06	176	1482	8.4
Wood Pellets	80.0	0.14	0.04	588	1855	7.4
Activated Charcoal	87.3	0.47	0.80	186	114	9.1

Source: Collins (2008).

Note: Activated charcoal is included as a standard analysis and comparison to biochars.

Table 3

Comparison of profits from winter wheat production^a (US\$ per acre), with and without biochar application.

Scenario	Revenue	CO ₂ Offset Value ^b	Total Cost ^c	Cost of Ag Lime ^d	Cost of Biochar ^d	Profit ^e
Without biochar or agricultural lime application	\$438	—	\$415	—	—	\$23
With ag lime application	\$694	—	\$415	\$133	—	\$146
With biochar application, when offset price is \$1/MT CO ₂ and the price of biochar (P _B) is...						
P _{B1} = \$350.74/metric ton ^f	\$694	\$90	\$415	—	\$10,740	-\$10,371
P _{B2} = \$114.05/metric ton ^g	\$694	\$90	\$415	—	\$3,492	-\$3,123
P _{B3} = \$87/metric ton ^f	\$694	\$90	\$415	—	\$2,664	-\$2,295
With biochar application, when offset price is \$31/MT CO ₂ and the price of biochar (P _B) is...						
P _{B1} = \$350.74/metric ton ^f	\$694	\$2,799	\$415	—	\$10,740	-\$7,662
P _{B2} = \$114.05/metric ton ^g	\$694	\$2,799	\$415	—	\$3,492	-\$414
P _{B3} = \$87/metric ton ^f	\$694	\$2,799	\$415	—	\$2,664	\$414

Figures for the revenue, CO₂ offset value, cost and profit are rounded to the nearest whole number.

^a The assumed base soil pH is 4.5. Biochar or agricultural lime application is intended to raise the assumed soil pH to 6.

^b CO₂ Offset Value = 90.29 metric tons of CO₂ offset per acre from avoided emissions of lime and biochar C sequestration (see Appendix) *times* the price of CO₂ offset (\$1 or \$31 per MT CO₂).

^c From 2008 Enterprise Budget for Eastern Whitman County, Conventional Tillage (Painter, unpub.). To illustrate the estimation of a farmer's profits with and without ag lime or biochar application, we chose Eastern Whitman County as example based on Mahler et al. (1985). The study found that the pH of soils in eastern Washington had significantly declined. By 1980, more than 65 percent in Whitman County had a soil pH less than 6.

^d Excludes the cost of applying lime or biochar to agricultural land (machinery and labor cost).

^e Profit = Revenue + CO₂ offset value – Total Cost – Ag Lime Cost – Biochar Cost. All are in US\$ per acre.

^f Obtained from Granatstein et al. (2009)

^g Based on the energy content of a wood-based biochar.




Fig. 1. Profit given the low/high price of CO₂ offset and different prices of biochar.

Appendix

Assumptions used to calculate the value of CO₂ offset

1. Case study: Eastern Whitman County, a high precipitation region (more than 18" per year)
2. Type of soil: Silt loam. We use the base pH = 4.5 in Collins (2008) for Palouse silt loam. To increase the soil pH of silt loam by 1.5 units (i.e., from 4.5 to 6), the requirements are:
 - 2.85 short tons of lime per acre (or 2.59 metric tons/acre)^{*}; or
 - 33.75 short tons of biochar per acre (or 30.62 metric tons/acre)[†].

Note the conversion 1 metric ton (MT) = 1.10231131 short tons.

3. Emissions avoidance and carbon sequestration from using biochar as soil amendment:
 - Avoided emissions for not using lime = 0.22 MT CO₂ per metric ton of limestone[‡]
 - Biochar C sequestration = 0.8 ton per metric ton of carbon or 2.93 MT CO₂[§] per metric ton of biochar applied to the soil^{**}
4. Estimated carbon sequestration per acre (CO₂ offset per acre) given application of biochar to cropland:
 - Avoided emissions for not using lime =
 $2.59 \text{ MT lime} \times (0.22 \text{ MT CO}_2/\text{metric ton of lime}) \approx 0.57 \text{ MT CO}_2$
 - Biochar C sequestration =
 $30.62 \text{ metric ton of biochar} \times (2.93 \text{ MT CO}_2/\text{metric ton of biochar}) \approx 89.72 \text{ MT CO}_2$
 - Total value of CO₂ offset per acre $\approx 90.29 \text{ MT CO}_2$

^{*} Source: California Plant Health Association (CPHA). 2002. Western Fertilizer Handbook, 9th edition. Danville, Illinois: Interstate Publishers, Inc.

[†] Source: H. Collins, unpublished data. Palouse silt loam soil analysis, biochar analyses.

[‡] Source: West, Tristram O. and Allen C. McBride. 2005. "The Contribution of Agricultural Lime to Carbon Dioxide Emissions in the United States: Dissolution, Transport and Net Emissions." *Agriculture Ecosystems and Environment*, 108:145-154.

[§] To convert from carbon to carbon dioxide, multiply by 44/12 (~3.67) (Blasing, T.J. *et al.*, 2004. Estimates of Monthly CO₂ emissions and Associated 13C/12C Values from Fossil Fuel Consumption in the U.S.A. Available at: http://cdiac.ornl.gov/trends/emis_mon/emis_mon_co2.html.

^{**} Based on biochar content of pine pellets. Source: Collins, Hal. 2008. "Use of Biochar from the Pyrolysis of Waste Organic Material as a Soil Amendment: Laboratory and Greenhouse Analyses." Quarterly report prepared for the Biochar Project (December 2008).