

Charging NO<sub>x</sub> Emitters for Health Damages:  
An Exploratory Analysis

by

Denise L. Mauzeral, Princeton University  
Babar Sultan, Harvard Medical School  
Namsoug Kim, Princeton University  
David F. Bradford, Princeton University, NYU, and CESifo

CEPS Working Paper No. 103  
September 2004

Keywords: surface ozone, NO<sub>x</sub> emissions, point sources, health impacts, mortality, morbidity, cap-and-trade.

JEL classification codes: Q5, H1.

# Charging NO<sub>x</sub> Emitters for Health Damages: An Exploratory Analysis

Denise L. Mauzerall<sup>1,2,\*</sup>, Babar Sultan<sup>3,4</sup>, Namsoug Kim<sup>1</sup> and David F. Bradford<sup>1,3, 5,6,7</sup>

\* Corresponding author, mauzeral@princeton.edu

<sup>1</sup>Science, Technology and Environmental Policy program  
Woodrow Wilson School of Public and International Affairs  
Princeton University, Princeton, NJ 08544

<sup>2</sup>Department of Geosciences, Princeton University

<sup>3</sup>Economics Department, Princeton University

<sup>4</sup>Presently at: Harvard Medical School, Boston, MA

<sup>5</sup>National Bureau of Economic Research, Cambridge, MA

<sup>6</sup>New York University School of Law

<sup>7</sup>CESifo, Munich, Germany

December 14, 2004

An earlier version of this paper has appeared as NBER working paper 10824; a revised version, with different title, has been submitted to *Atmospheric Environment*. We thank Winston Hao and Gopal Sistla of the New York State Department of Environmental Conservation for providing us with the version of the CAMx model used by the Ozone Transport Commission. We thank Arlene Fiore, Sarah Fox, Don Fullerton, Michael Greenstone, Michael Herz, Larry Horowitz, Jesse Rothstein, Wolfram Schlenker, Quansong Tong, and Jason West for helpful comments. We are pleased to acknowledge the Woodrow Wilson School of Public and International Affairs and the Center for Economic Policy Studies at Princeton University for financial support.

## ABSTRACT

We present a proof-of-concept analysis of the measurement of the health damage of ozone ( $O_3$ ) produced from nitrogen oxides ( $NO_x = NO + NO_2$ ) emitted by individual large point sources in the eastern United States. We use a regional atmospheric model of the eastern United States, the Comprehensive Air Quality Model with Extensions (CAMx), to quantify the variable impact that a fixed quantity of  $NO_x$  emitted from individual sources can have on the downwind concentration of surface  $O_3$ , depending on temperature and local biogenic hydrocarbon emissions. We also examine the dependence of resulting ozone-related health damages on the size of the exposed population. The investigation is relevant to the increasingly widely used “cap and trade” approach to  $NO_x$  regulation, which presumes that shifts of emissions over time and space, holding the total fixed over the course of the summer  $O_3$  season, will have minimal effect on the environmental outcome. By contrast, we show that a shift of a unit of  $NO_x$  emissions from one place or time to another could result in large changes in the health effects due to ozone formation and exposure. We indicate how the type of modeling carried out here might be used to attach externality-correcting prices to emissions. Charging emitters fees that are commensurate with the damage caused by their  $NO_x$  emissions would create an incentive for emitters to reduce emissions at times and in locations where they cause the largest damage.

Key words: surface ozone,  $NO_x$  emissions, point sources, health impacts, mortality, morbidity, cap-and-trade.

## 1. Introduction

Pollution in economic theory is essentially defined by externalities, negative impacts of one economic agent on others that are not reflected in market prices. An important input to corrective policy is the size of the externality, an estimation of the amount of damage done by the externality-causing action. In this paper we lay out a method of estimating the damage to human health due to exposure to ozone ( $O_3$ ) formed as a result of emission of nitrogen oxides ( $NO_x = NO + NO_2$ ) from individual large stationary sources in the eastern United States. We also present quantitative estimates for cases selected to draw attention to sources of variation in health damage that the present thrust of  $NO_x$  emissions regulation neglects. We apply the modifier “proof of concept” to the estimates to reflect the need for further testing and refinement of the modeling components and for incorporating effects other than  $O_3$  -exposure. The longer term objective is to associate an estimate of external costs due to emissions from any point source as a function of the prevailing weather conditions and populations exposed. The initial calculations suggest the possibility of substantial gains (ie. lower total damages for a given quantity of emissions) from regulatory redesign.

$NO_x$  contributes to a wide range of environmental effects including the formation of acid rain (via formation of nitric acid),  $PM_{2.5}$  (via formation of secondary particulates such as ammonium nitrate) with resulting health impacts and contributions to regional haze, eutrophication of aquatic ecosystems (via addition of excess nitrogen), and elevated  $O_3$  concentrations (via reaction with hydrocarbons and carbon monoxide) with resulting impacts on health and agriculture. Due to recent regulatory emphasis on controlling  $O_3$  through  $NO_x$  “cap-and-trade” programs, we focus here on the contribution of  $NO_x$  emissions from point sources to ambient  $O_3$  concentrations and the resultant damage to human health.

“Cap and trade” is a mechanism for regulating  $NO_x$  emissions, whereby a limit, or cap, is placed on total emissions from large point sources (typically during the “ozone season” from May to October each year). Regulated sources (mostly electric power generating plants) are obliged to have an allowance for each unit (ton) of  $NO_x$  emitted. In practice, allowances are initially allocated to sources based on their historical operating levels. A source that does not

own sufficient allowances for its needs can purchase them from others at a price agreed upon by the trading parties. The trading of allowances is expected to lower the total cost of whatever level of emissions control is embodied in the cap relative to a prescriptive type of regulatory control. A premise of such a system is that, because the total emissions are unaffected, such an exchange has no environmental significance. However, this premise may not be realized in any given situation and when it fails an exchange of allowances may have either positive or negative environmental effects.. Our damage estimates allow us to isolate instances where a shift in  $\text{NO}_x$  emissions can have large effects on environmental damage.

To estimate the damages to human health resulting from  $\text{NO}_x$  emissions we use a model of the chemical and transport processes in the troposphere to calculate the increase in  $\text{O}_3$  concentration, at each time and location, that results from a given incremental quantity of  $\text{NO}_x$  emitted from a power plant under different conditions. These  $\text{O}_3$  concentration changes are combined with geographically specific demographic data to estimate the increase in mortality and respiratory illness that results from that increase in  $\text{O}_3$ . Specifically, we examine how temperature variations within a single summer month can have a significant effect on the quantity of  $\text{O}_3$  produced from a unit of  $\text{NO}_x$ . We also examine how the same quantity of  $\text{NO}_x$  emissions from a power plant can result in very different  $\text{O}_3$  production depending on the quantity of biogenic hydrocarbons emitted in the surrounding region. Finally, we show how the health damage that elevated  $\text{O}_3$  concentrations cause increases as the size of the exposed population increases.

In the section relating these exercises to the design of policy, we draw attention to the standard economic argument for presenting firms with the “right” prices of the resources they use. In this case the resource used is  $\text{NO}_x$  emissions, for which, in the locations we study, the price charged is a combination of the price of emission allowances and the implicit price effect of applicable environmental regulations. The current regulatory regime diverges from one that would connect  $\text{NO}_x$  emission choices with their economic consequences in at least two respects. First, the present regulatory regime gives little systematic incentive to shift  $\text{NO}_x$  emission from times and places where it has a large negative externality in the form of health impacts to times and places where the externality is smaller. Our analysis describes the benefits of considering

this variability. Second, the regulatory system is focused on avoiding exceedences of limits on ambient concentrations in order to comply with the National Ambient Air Quality Standards (NAAQS). The epidemiological studies on which we base our estimates of damages indicate that health impacts increase linearly with increasing concentration [EPA, 2003; Steib *et al.*, 2002]. A threshold limit on ambient O<sub>3</sub> concentration is a related but distinct objective to the avoidance of mortality and morbidity that a standard economic approach would imply because it is not concerned with damages which result from exposures to concentrations below the NAAQS.

In section 2 we provide background on the science of O<sub>3</sub> formation from NO<sub>x</sub> emissions and on the regulatory history of NO<sub>x</sub> and O<sub>3</sub>. In section 3 we describe CAMx (Comprehensive Air Quality Model with Extensions), the regional air quality model we use to quantify the O<sub>3</sub> produced from NO<sub>x</sub> emitted from point sources, evaluate its ability to reproduce observed O<sub>3</sub> concentrations, and describe the method we use to quantify morbidities and mortalities resulting from O<sub>3</sub> exposure. Section 4 uses CAMx to demonstrate the differences in O<sub>3</sub> production resulting from variability in local temperature and biogenic hydrocarbon emissions, as well as the importance of the location of high population densities relative to emission sites in determining resulting mortality and respiratory morbidity. These exercises demonstrate the potential environmental impact of an exchange of emissions from one time, place, or weather condition to another, expressed in terms of lives lost or sickness experienced. In the fifth section we estimate the monetary value of these environmental impacts, using valuation factors taken from the literature. Section 6 discusses the policy advantages of a damage based incentive system to control NO<sub>x</sub> emissions from point sources. Section 7 summarizes our conclusions and highlights needed future research.

## **2. Scientific and Regulatory Background**

### **2.1. Ozone Chemistry**

O<sub>3</sub> is a pollutant that is formed in the lower atmosphere (troposphere) from a complex series of sunlight-driven reactions between NO<sub>x</sub>, carbon monoxide (CO), volatile organic

compounds (VOCs) which are largely hydrocarbons, and methane (CH<sub>4</sub>). The primary source of NO<sub>x</sub> to the troposphere is fossil-fuel combustion. In 1998 in the United States, electric utilities accounted for 25% of total NO<sub>x</sub> emissions, transportation for 53% and industrial sources for 12% [EPA, 2000]. VOCs are emitted from a range of human activities, including fossil-fuel combustion (excepting high-temperature combustion), direct evaporation of fuel and solvents, and chemical manufacturing. Terrestrial vegetation also provides a large natural source of hydrocarbons in summer with isoprene being the biogenic hydrocarbon emitted in the largest quantity. The emission of biogenic hydrocarbons varies greatly by location as indicated in Figure 4. Biogenic hydrocarbon emissions reach a maximum at approximately 40°C and decrease following a bell shaped curve on either side of 40°C [Pierce *et al.*, 1998]. Difficulty regulating O<sub>3</sub> occurs because in regions of high NO<sub>x</sub> (primarily urban centers and power plant plumes), O<sub>3</sub> formation is limited by the availability of hydrocarbons. In regions of low NO<sub>x</sub> (primarily rural areas with abundant emission of natural hydrocarbons), O<sub>3</sub> formation is limited by the availability of NO<sub>x</sub> [Sillman *et al.*, 1990].

Ozone production from NO<sub>x</sub> emitted in power plant plumes varies depending on the availability of hydrocarbons, the magnitude of the NO<sub>x</sub> emission rate, and meteorological conditions. A power plant plume contains large quantities of NO<sub>x</sub>, little CO and virtually no VOCs. Hence O<sub>3</sub> production from NO<sub>x</sub> in a power plant plume is limited by the availability of VOCs in the surrounding environment as it dilutes. Data from aircraft transects of power plant plumes indicate that higher availability of biogenic hydrocarbons, principally isoprene, result in higher quantities of O<sub>3</sub> produced from the NO<sub>x</sub> in the plume [Ryerson *et al.*, 2001]. In addition, as the size of the plume increases, the ozone production efficiency (the number of molecules of O<sub>3</sub> produced per molecule of NO<sub>x</sub> consumed [Liu *et al.*, 1987]) decreases. Therefore, large concentrated plumes can result in less net O<sub>3</sub> formation than small plumes per NO<sub>x</sub> molecule emitted because of the more rapid conversion of NO<sub>x</sub> in concentrated plumes to HNO<sub>3</sub> [Ryerson *et al.*, 2001]. Hence, in our later analysis all of our cases use large power plants of approximately the same size.

In addition, O<sub>3</sub> production occurs more rapidly under conditions of stagnant high pressure which are frequently accompanied by high temperatures and sunny conditions [Logan, 1989;

Vukovich, 1995]. As a result, the probability of exceeding the NAAQS for O<sub>3</sub> increases with daily maximum temperature [Lin *et al.*, 2001]. In our analysis temperature is used as a proxy for stagnant meteorological conditions.

## **2.2. NO<sub>x</sub> and O<sub>3</sub> regulatory history**

Legislative and regulatory efforts to control ambient O<sub>3</sub> concentrations started with the passage of the 1970 Clean Air Act (CAA). In 1971 EPA established a NAAQS for O<sub>3</sub> of a 1-hour maximum daily average concentration not to exceed 0.08ppm which in 1979 was revised to 0.12ppm [EPA, 2004a]. Based on scientific evidence of the adverse effect of O<sub>3</sub> on health, in 1997 EPA again revised the O<sub>3</sub> NAAQS and required that a 3-year average of the annual fourth highest daily 8-hour average concentration not exceed 0.08 ppm [EPA, 2004a; NRC, 2004]. Under the CAA, a state with monitoring stations that fail to meet the NAAQS is regarded as “out of attainment” and is obliged to specify in its State Implementation Plan (SIP) how it will attain the NAAQS in the future. No incentive presently exists to reduce O<sub>3</sub> further if a state is in compliance. O<sub>3</sub> has a lifetime of approximately 2 days in the boundary layer of the eastern United States in summer [Fiore *et al.*, 2002] and hence may be transported across state boundaries. Inter-state transport can make it difficult for states that fail to meet the NAAQS to become compliant by simply reducing in-state emissions.

Economists have been encouraging the use of market-based policies for controlling pollution as an alternative to more traditional command-and-control approaches (such as uniform emission standards or specific technology requirements) for some time. In the late 1960s, discussion of the concept of markets in pollution rights began in academic circles and entered the U.S. policy world in 1975 [Atkinson and Tietenberg, 1987]. Economic theory predicts that tradeable emission permits will induce emitters to find a cost-minimizing emission allocation when a cap on total emissions is specified [NRC, 2004]. Emitters that can reduce emissions easily do so and sell permits to those for whom reducing emissions is more costly.

In an attempt to control regional O<sub>3</sub> concentrations and to address inter-state transport of O<sub>3</sub> and its precursors to facilitate the compliance of down-wind states with the O<sub>3</sub> NAAQS, a coalition of 13 northeastern states formed the Ozone Transport Commission (OTC) and created a



regional NO<sub>x</sub> cap-and-trade program. The OTC “NO<sub>x</sub> Budget program” ran from 1999-2002 during which time the total NO<sub>x</sub> emissions from large point sources within the 13-state coalition were reduced approximately 50% from 1990 baseline levels [EPA, 2004b]. The period of NO<sub>x</sub> emission reduction from power plants coincided with a period of economic growth during which emissions from other sources (ex. vehicles) may have increased. A comparison of surface NO concentrations between 10am and 2pm over the summer season pre- and post- the 1999 cap, made as part of the EPA-AIRS measurement network, does not indicate the expected decrease in NO concentrations. Nor does a similar comparison of O<sub>3</sub> concentrations over the summer season pre- and post- the 1999 cap show a decrease in O<sub>3</sub> concentrations resulting from a decrease in NO<sub>x</sub> emissions from the power plants as a result of the cap (analysis not shown).

In 2004 the OTC program was replaced by the NO<sub>x</sub> State Implementation Plan (SIP) Call budget trading program, developed in response to the EPA’s call for SIPs to reduce the transport of O<sub>3</sub> over large regions of the country. Under the SIP Call program the trading region has expanded to 19 eastern states and total summer emissions from electric power generators are planned to be reduced by approximately 80% from 1990 levels [*EvoMarkets*, 2003].

The theoretical advantages of an emissions cap and trade system depend on the assumption that only the total amount emitted matters. Under this condition, a shift of emissions from one time or place to another will have no environmental consequences and minimizing the cost of abatement is the object of the plan. Recognizing the general problem, the “NO<sub>x</sub> Budget program” incorporated “flow control” to require that allowances banked from one summer season for use in subsequent summers be surrendered on a 2:1 basis if the bank became larger than 10 percent of the regional budget. The flow control rule inhibits excessively large emissions in any single summer but it is not clear whether the net environmental result (measured by health and mortality impacts) is likely to be positive. Our analysis suggests the importance of designing rules to take into account the environmental effects of variations in the time or location of emissions in a single summer season, let alone between seasons.

### 3. Methods

#### 3.1. CAMx model description


We utilize the Comprehensive Air Quality Model with extensions (CAMx) version 3.0, a regional Eulerian photochemical grid model [ENVIRON, 2000]. CAMx simulates the emission, dispersion, chemical and photochemical reactions and removal of tropospheric ozone and its precursors in the lower troposphere by solving the continuity equation for each chemical species in a three-dimensional grid. The Eulerian continuity equation relates the time dependency of the average species concentration in each grid cell volume to the sum of all physical and chemical processes operating on that volume.

We use the CAMx model in a configuration previously used by the Ozone Transport Assessment Group (OTAG) to solve for the concentration distribution of ozone and its primary precursors. Model resolution is 0.33° latitude by 0.5° longitude extending in 5 vertical layers (100, 500, 1500, 2500, and 4000 meters above ground level) from the surface to 4 km. CAMx includes a plume-in-grid sub-model to simulate the chemistry of NO<sub>x</sub> plumes emitted from point sources such as power plants. The plume-in-grid sub-model tracks individual plume segments, accounting for dispersion and inorganic chemical evolution, until the puff mass can be adequately resolved by the grid box in which it resides at which time the remaining NO<sub>x</sub> is released to the grid. This CAMx configuration uses the Carbon Bond IV (CB-IV) chemical mechanism with updated radical termination and isoprene oxidation mechanisms based on [Carter, 1996] including 25 gas species and 96 reactions. Winds are generated by the Regional Atmospheric Modeling System (RAMS) with horizontal advection provided by the diffusive-corrected algorithm of [Smolarkiewicz, 1983]. Boundary conditions are set to clean background concentrations for all constituents on all four sides and at the top of the domain (O<sub>3</sub> = 35ppbv) with a day/night diurnal cycle imposed on short-lived species. Hourly surface emissions are included from elevated point sources, gridded low-level point, mobile, biogenic (from the BEIS2 model) and anthropogenic area emissions in the eastern United States. The BEIS2 model includes the dependence of isoprene emissions on temperature, solar radiation, ecosystem extent and leaf area [Geron *et al.*, 1994; Guenther *et al.*, 1994; Pierce *et al.*, 1998]. However, [Palmer


*et al.*, 2003] inferred from GOME satellite measurements of formaldehyde that BEIS2 underpredicts isoprene emissions by a factor of two. CAMx also includes wet and dry deposition [Wesley, 1989] of gases. Emissions and meteorological fields for the model were obtained from the New York State Department of Environmental Conservation (NYSDEC) [Sistla and Hao, personal communication, 2001]. Model simulations were conducted for July 7-17, 1995, with the first two days of the simulation used for model spin-up.

### 3.2. CAMx model / EPA-AIRS O<sub>3</sub> data comparison

To evaluate the ability of this version of CAMx to simulate accurately ambient O<sub>3</sub> concentrations within the region, we compared concentrations of O<sub>3</sub> measured as part of the EPA-AIRS surface measurement network with O<sub>3</sub> concentrations obtained from the CAMx simulation. Locations of the EPA-AIRS O<sub>3</sub> measurement sites for July 1995 are shown in Figure 1. Figure 2 shows a comparison of 1-hour daily maximum O<sub>3</sub> concentrations calculated by the CAMx model for July 9-17, 1995 and measured on the same dates by the EPA-AIRS network. A comparison of 1-hour maximum concentrations is shown because the epidemiological dose-response functions used later in this paper to estimate both mortality and morbidity use this value.



**Figure 1.** EPA-AIRS O<sub>3</sub> measurement site locations within the CAMx domain, July 1995.



**Figure 2.** Comparison of 1-hr. daily maximum O<sub>3</sub> surface concentrations for July 9-17, 1995 from the EPA-AIRS measurement network and CAMx simulation. Rural, suburban and rural labels refer to the EPA-AIRS designation for individual measurement sites. O<sub>3</sub> concentrations simulated in a surface CAMx grid box are compared with the measured O<sub>3</sub> concentrations at the EPA-AIRS network stations that fall inside the grid box. Multiple stations within one category (urban, suburban or rural) that fall within one grid box are averaged together for the comparison.

Statistical analysis of the results in Figure 2, with  $P_i$  indicating the predicted CAMx O<sub>3</sub> values and  $O_i$  indicating the observed EPA-AIRS O<sub>3</sub> concentrations, result in the following summary statistics:

$$\text{Root mean square error (RMSE)} = \left[ \frac{1}{N} \sum_{i=1}^N (P_i - O_i)^2 \right]^{\frac{1}{2}} = 19.6 \text{ ppb};$$

$$\text{Mean Bias (MB)} = \frac{1}{N} \sum_{i=1}^N (P_i - O_i) = 0.62 \text{ ppb};$$

$$\text{Mean Normalized Bias (MNB)} = \frac{1}{N} \sum_{i=1}^N \frac{(P_i - O_i)}{O_i} = 6.0 \%$$

This evaluation indicates that although there is random error in the maximum daily O<sub>3</sub> concentrations calculated by CAMx the model is without significant bias.

We use the model to predict the change in ozone concentration (measured by the 1-hour daily maximum) over time over the whole model domain that would result from a specified quantum change in NO<sub>x</sub> emissions at a specified time at a given power plant location. A simple regression of  $O_i$  on  $P_i$  in the data in Figure 1, yields the slope and intercept estimates of 0.646 (56.43) and 26.0 (28.54), where the figures in parentheses are the t statistics for the difference from zero (adjusted R-square .42). The implication of the regression is that a unit difference in the O<sub>3</sub> prediction of the model (from one place or time to another within the data set) corresponds on average to a .646 difference in measured ozone concentration between the two time or location points.

Even with the limited data set, one could explore other formulations of the problem of predicting measured ozone. An obvious “experiment” is to allow for a different intercept coefficient at every location, on the thought that there are likely to be persistent differences in receptor conditions that are not captured by CAMx. A location fixed effects multivariate regression of  $O_i$  on  $P_i$  (a regression of  $O_i$  on  $P_i$  plus a dummy variable for each receptor location), using the data in Figure 1, yields a slope coefficient of 0.613 (47.29) (adjusted R-square .54), interestingly, very close to the coefficient in the simple regression. These exercises (calculations thanks to Wolfram Schlenker) just scratch the surface of a thorough investigation of the uses of CAMx for the prediction we make in this analysis. (For example, with a longer observation period, one could allow for receptor location-specific slope coefficients.)

The slope coefficient (change in measured per unit change in model-generated ozone concentration) in these regression analyses are of central interest to us. As we explain, we focus on the difference between the O<sub>3</sub> concentrations generated by CAMx for all receptor locations in a standard run (which uses the input data underlying Figure 1) and in a set of alternative runs

(which perturb the input emissions at strategically chosen times and places). The underlying assumption is that the differences in model-generated ozone concentrations are good predictors of the differences in measured concentrations that would have been experienced, had the real world input data been different in the prescribed manner. The regression analysis of the use of model results in predicting measured outcomes would suggest, instead, taking as the basis for predicting the impacts the model differences multiplied by the slope coefficient (.646 or .613 in the case of the mentioned regressions).

Because we have found in discussions with colleagues that there is no consensus on the validity of such an adjustment, and because, in any case, it is just one of many refinements to what we emphasize is a proof-of-concept study, for the remainder of this paper we do not make any allowance for these issues.

### **3.3. Model Simulations**

The two goals of the CAMx simulations presented in this paper are first to demonstrate the spatial and temporal heterogeneity of tropospheric ozone production and the variability in resulting health damages (morbidity and mortality) that may result from a given amount of NO<sub>x</sub> emitted from the same location under different meteorological conditions or from different locations with varying biogenic hydrocarbon emissions or downwind populations. The second objective is to provide a proof-of-concept prototype of the feasibility of conducting a full integrated assessment that links NO<sub>x</sub> emissions to O<sub>3</sub> concentrations to human exposure, morbidity, mortality and damage valuations.

To demonstrate the spatial and temporal heterogeneity of tropospheric ozone production, we conduct a set of CAMx regional simulations for seven cases. First, we conduct a standard simulation that includes all regional emissions. This is followed by six perturbation simulations. In order to quantify the impact of NO<sub>x</sub> emissions from individual power plants on O<sub>3</sub> concentrations, we reduce the NO<sub>x</sub> emissions from each plant by 62.5 tons (125,000 lbs) in a 24-hour period ( $1.77 \times 10^6$  moles where NO<sub>x</sub> emissions from power plants were assumed to be 90% NO and 10% NO<sub>2</sub>) and calculate the difference in maximum O<sub>3</sub> concentration between the

standard simulation and each perturbation simulation. The power plants used for this experiment are among the largest 5% of power plants in the United States each of which emits approximately 100 tons of NO<sub>x</sub> per day. 62.5 tons of NO<sub>x</sub> was chosen for the analyses because it is sufficiently large to generate a discernable effect and sufficiently small that a single large power plant may emit this quantity or more in a 24-hour period. The perturbation simulations were chosen to demonstrate the variability in O<sub>3</sub> production resulting from identical NO<sub>x</sub> emissions from power plants produced during conditions of high and low temperature at the same location and in locations of high and low biogenic hydrocarbon emissions. They were also chosen to demonstrate the dependence of total health effects on the population density in the regions downwind of the emissions. Figures 3, 5 and 7 show the increase in maximum surface O<sub>3</sub> concentrations over the model domain between the standard simulation and perturbation simulations summed over the day of emission and the subsequent 3 days. After 4 days the power plant emissions from the 24-hour period have a non-discernable effect on regional O<sub>3</sub> concentrations. The total increase in 1-hour maximum O<sub>3</sub> is shown because it is incorporated in the dose-response functions used to estimate the mortality and morbidity that result from exposure to O<sub>3</sub>.

### **3.4. Estimating Mortality and Morbidity**

The relationship between O<sub>3</sub> exposure and resulting health effects can be estimated from epidemiological studies. Most epidemiological research has made use of time-series analyses that have focused on the effects of short-term exposures to a pollutant by using records of hospital admissions and deaths that occur following periods of elevated concentrations. These studies are unable to include the negative effects of chronic or long-term exposure and hence presumably underestimate the total health damage due to exposure. Cohort studies, which follow a population over time and do include both acute and chronic impacts, have been conducted for PM<sub>2.5</sub> [Dockery et al., 1993; Pope et al., 2002; Pope et al., 1995] but have not focused on O<sub>3</sub>. Hence, in relying in this analysis on time-series studies only (specifically, on meta-analyses that pool the results of many regional time-series studies [Steib et al., 2002]) we

use an arguably conservative estimate of the mortality and respiratory morbidity that results from elevated O<sub>3</sub> exposure.

The change in the incidence of mortality or morbidity due to exposure to air pollution (here O<sub>3</sub>) is usually estimated with the following function [EPA, 1999]:

$$\Delta M = Y_o \left(1 - e^{-\beta \Delta O_3}\right) * population \quad (1)$$

Here  $\Delta M$  is the change in the number of mortalities during a 24-hour period, or change in incidence of hospital admissions for respiratory disease, relative to a baseline number of deaths or incidence of respiratory disease. For mortality,  $Y_o$  is the baseline incidence of daily non-accidental deaths per person of any age. For morbidity,  $Y_o$  is the daily average hospital admission rate for diseases of the respiratory system. Epidemiological studies often report a relative risk (RR) (the percentage increase in mortality or morbidity) due to a specific increase in pollutant concentration. The concentration-response coefficient,  $\beta$ , can be derived from the reported relative risk and the change in pollutant concentration relative to a baseline (here  $\Delta O_3$ ), by solving:

$$\beta = \frac{\ln(RR)}{\Delta O_3} \quad (2)$$

We discuss the  $\beta$  values we use for mortality and morbidity in the following sub-sections. We obtain the population in equation (1) from 1995 world population data gridded at 0.04167 latitude by longitude resolution [CIESIN *et al.*, 2000]. Population data for the eastern United States is regridded at the CAMx resolution of 0.3 degrees latitude by 0.5 degrees longitude (approximately (36 km)<sup>2</sup> in the United States). The population within each CAMx grid box is used to calculate the mortality and morbidity that result from the change in O<sub>3</sub> concentrations in each grid box. Mortality and morbidity incidence is summed over all affected grid boxes over all days in which the O<sub>3</sub> concentration was enhanced due to (or, in some locations, reduced by) the NO<sub>x</sub> emitted from the individual power plants.

### 3.4.1. O<sub>3</sub> Mortality

To obtain the relative risk of mortality for an increase in O<sub>3</sub> concentration resulting from individual power plant emissions, we use a pooled estimate of percent excess mortality derived



from a compilation of 25 epidemiological time-series analyses of daily 1-hour maximum O<sub>3</sub> concentrations during the warm season [Steib *et al.*, 2002]. The pooled estimate of RR is 1.016 for a 31.2 ppb increase in 1 hour maximum daily O<sub>3</sub> concentration, which translates into a  $\beta$  value of 0.000509 [Steib *et al.*, 2002]. We estimate  $Y_0$  from the total U.S. population in 1995 and calculate an average daily mortality rate of  $2.4 \times 10^{-5}$  (2,312,132 deaths/ 261,638,000 persons / 365 days) [CDC-NCHS, 2003; USCB, 2000].

### 3.4.2. O<sub>3</sub> Morbidity


Hospital admissions for respiratory ailments have been found to be substantially and consistently associated with O<sub>3</sub> exposure. A synthesis of O<sub>3</sub> hospital admission time-series studies, compiled by Thurston and Ito [1999] is used here to estimate the effect of O<sub>3</sub> on morbidity. Data on hospital admissions for respiratory disease in the United States are not available. Therefore, to calculate  $Y_0$  we assume that the average daily rate of hospital admissions for respiratory disease is approximately equal to the rate of hospital discharge following a diagnosis of a respiratory ailment which was  $3.41 \times 10^{-5}$  discharges per day in 1998 (124.6 discharges / 10,000 population / 365 days), the year for which such data are available [CDC-NCHS, 1998]. This value is of similar magnitude to values obtained in regional studies of daily hospital admission for respiratory diseases [EPA, 1999]. The estimate for hospital admissions for major respiratory ailments due to O<sub>3</sub> exposure for all ages was a RR = 1.18 per 100 ppb increase in daily 1-hour maximum O<sub>3</sub> (95% CI, 1.10-1.26) [Thurston and Ito, 1999] which translates into a  $\beta$  value of 0.00166. In cases where studies in the meta-analysis used other measures of O<sub>3</sub> exposure besides a daily 1-hour maximum concentration, (such as an 8-hour average O<sub>3</sub> concentration) the ratio of the daily 1-hour maximum to the other measures used in the original analyses were used to compute the increment comparable to a 100 ppb change in daily 1-hour maximum O<sub>3</sub> concentrations [Thurston and Ito, 1999].

## 4. Results

Our results demonstrating the impacts of variability in temperature and biogenic hydrocarbon emissions on O<sub>3</sub> production, and the importance of population density in determining resulting mortalities and respiratory morbidity are summarized below. These factors were chosen for examination because of the expected size of their influence and because of the ability of our model to simulate them. Additional factors may also influence O<sub>3</sub> production from large point sources such as the concentration of NO<sub>x</sub> in the plume, cloudiness, time of day of emission, proximity of the NO<sub>x</sub> emission to other anthropogenic emissions, and other factors. Analysis of these additional factors is left for the future.

### 4.1. Temperature variability -- Effect on O<sub>3</sub> production and resulting mortality and morbidity

First, to examine the temporal heterogeneity of tropospheric ozone production due to temperature variability within a short period of time, a point source was selected which experienced a range of temperatures between July 9 and 17, 1995. The effects on O<sub>3</sub> production of identical emissions of NO<sub>v</sub> from a power plant in eastern Pennsylvania on July 9, 1995 and July 14, 1995 when the 24-hour period temperature ranged over  $72 \pm 7.7^\circ\text{F}$  and  $83.5 \pm 6.3^\circ\text{F}$ , respectively are shown in Figures 3a and 3b. The average increase in 1-hour maximum surface O<sub>3</sub> over the model domain between the standard simulation and perturbation simulations on the day of emission and the subsequent 3 days, was 0.021 ppbv during the cool period and 0.036 ppbv during the warm period. The increase in 1-hour maximum O<sub>3</sub> is shown in Figure 3 because it is used in the dose-response functions to estimate O<sub>3</sub> induced mortality and morbidity. The model predicts that the same amount of NO<sub>x</sub> emitted in the warmer period results in more O<sub>3</sub> which leads to twice the mortalities as in the cooler period -- 0.185 versus 0.354 persons as shown in Figures 3c and 3d. Likewise, the number of cases of respiratory morbidity due to O<sub>3</sub> exposure was also nearly twice as high during the warm period -- 0.866 versus 1.654 cases as is shown in Figures 3e and 3f.




**Figure 3.** O<sub>3</sub> produced by an identical amount of NO<sub>x</sub> emitted from the same power plant in eastern Pennsylvania in a 24-hour period on a) 9 July 1995, a cool day (296°K ± 4.3°K;

72° ± 7.7°F) and b) 14 July 1995, a warm day (302°K ± 3.5°K; 83.5 ° ± 6.3 ° F) and the subsequent 3 days until O<sub>3</sub> production from these NO<sub>x</sub> emissions cease. The “X” in each figure marks the power plant location. The ‘average increase in 1-hour maximum O<sub>3</sub>’ is the total increase in maximum surface O<sub>3</sub> concentrations over the model domain between the standard simulation and perturbation simulations on the day of emission and the subsequent 3 days. Beyond 3 days the power plant emissions cease to have a significant influence. Here figures c) and d) show the mortalities and e) and f) show the morbidities that are estimated to result from the increase in daily maximum O<sub>3</sub> occurring in a) and b), respectively. These simulations show that, in violation of the preconditions for the use of a cap and trade system, a shift of a fixed amount of NO<sub>x</sub> emissions from a single source between the two closely separated time points could have a substantial effect on the quantities of O<sub>3</sub> that are produced, and the consequent health effects. In fact, since energy demand is usually larger when temperatures are high, emissions are also likely to be larger during periods of high temperature when the model indicates that more O<sub>3</sub> is produced per unit NO<sub>x</sub> emitted. An emitter using emission allowances at two different times presently has no reason to favor the time when the resulting damage is lower. Trading allowances (or variability in the timing of the use of allowances from a single source) to emit at a price that does not reflect the difference in damage will lead to too much emission in the high-damage conditions relative to the low-damage conditions. In the example, the temperature at the emission source and at downwind locations makes an important difference. The question is naturally raised, whether an alternative regulatory system might be developed that provides an appropriate incentive for emitters to reduce their emissions during times of elevated temperature within the ozone season.


#### **4.2. Biogenic hydrocarbon emission variability -- Effect on O<sub>3</sub> production and resulting mortality and morbidity**

Isoprene is a hydrocarbon that is naturally emitted from deciduous tree species, particularly oak, poplar and gum trees. In the United States these species are most abundant in the south-eastern part of the country and are absent in most of the west. This results in large spatial variability in biogenic VOC emissions in the United States, as is shown in Figure 4. In addition, isoprene emissions increase with increasing temperature and sunlight. A unit of NO<sub>x</sub> emitted from a power plant in a region of high isoprene emissions can result in larger O<sub>3</sub> production than the same amount emitted in a region of low isoprene emissions [Ryerson *et al.*, 2001]. To examine the impact that biogenic hydrocarbon emissions, particularly isoprene, can have on O<sub>3</sub> production, we examined the quantity of O<sub>3</sub> produced from identical emissions of NO<sub>x</sub> by two different power plants, one in the south (Mississippi) and the other in the northern

mid-west (Indiana) of the U.S. To control for the effect of temperature on isoprene emissions and on O<sub>3</sub> production, we chose 24-hour periods during which the mean temperature at the two sites was nearly identical: 87.2° F ± 5.9° F in the high isoprene region and 87.9° F ± 6.3° F in the low isoprene region. The regional average increase in 1-hour maximum O<sub>3</sub>, as shown in Figure 5a and 5b, was 0.033 ppbv in the region with low emissions and over three times higher at 0.102 ppbv in the region of high emissions. Due to the apparent factor of two under-prediction of isoprene emissions by BEIS2 [Palmer *et al.*, 2003], the quantity of O<sub>3</sub> produced in our simulation in the high-isoprene region may be under-predicted.



**Figure 4.** July emissions of isoprene in the United States (monthly mean of 24-hour daily averages). This figure was created from a simulation using the Community Multi-scale Air Quality Modeling System (CMAQ) [Byun and Ching, 1999].



**Figure 5.** O<sub>3</sub> produced by an identical amount of NO<sub>x</sub> emitted on 10 July 1995 from a power plant in a) a region of relatively low isoprene emissions and b) a region of relatively high isoprene emissions. The temperatures at both locations were nearly identical (87.9° F ±

6.3° F and 87.2° F ± 5.9° F in the low and high isoprene regions respectively). The rest of the notation is as described in Figure 3.


In this example, the region of higher isoprene emissions was also a region of lower population density (see Figure 6). Despite this fact, the 3.5 times larger quantity of O<sub>3</sub> produced in the region of higher isoprene emissions resulted in both 30% more mortalities (0.166 versus 0.121 deaths) and morbidities (0.774 versus 0.568 cases) than resulted from the same increase in NO<sub>x</sub> emissions in the region of lower isoprene emissions (see Figures 5c-f). By choosing locations where the region of high isoprene corresponds to a region of lower population, this scenario is intentionally conservative and is hence understating the potential importance of variability of isoprene concentration on resulting O<sub>3</sub> mortalities.

A regulatory scheme that ignores the spatial variability of biogenic hydrocarbon emissions when regulating NO<sub>x</sub> may miss an opportunity to decrease O<sub>3</sub> concentrations and could inadvertently increase mortalities and morbidities by trading from a region of relatively low isoprene emissions (north-east) to high emissions (south-east). In particular, trades between regions under a cap-and-trade program in which all NO<sub>x</sub> emissions from large point sources in a region are treated identically, could have a substantial effect (up or down, depending on the direction of the exchange) on the quantities of O<sub>3</sub> that are produced, and the consequent health effects. A regulatory system that provides incentives for emitters to reduce their emissions in regions of high isoprene emissions would be more effective at reducing ambient O<sub>3</sub> concentrations [Ryerson *et al.*, 2001]. If these reductions coincided with high population densities the reductions would also be more effective at reducing adverse effects on exposed populations than would a regulatory program that assumes emissions in all locations have identical effects. We discuss the effect of downwind population density next.

#### **4.3. Downwind Population Density -- Effect on resulting mortality and morbidity**


Population density in the eastern United States is shown in Figure 6. The north-eastern Washington D.C. to Boston corridor has among the highest population densities in the United

States. To examine the impact that equivalent reductions of  $\text{NO}_x$  would have from point sources upwind of regions of high and low population density, point sources in the high-population density northeast (Maryland) and lower-population density mid-Atlantic region (North Carolina) were selected. However, average isoprene emissions are substantially larger in North Carolina than in Maryland. Thus, despite controlling for temperature, the average increase in 1-hour maximum  $\text{O}_3$ , as shown in Figures 7a and 7b, was more than twice as great (0.081 ppbv versus 0.035 ppbv) when emissions originated from North Carolina in the low population region (Figure 7a) than from Maryland in the high population region (Figure 7b). Despite higher maximum  $\text{O}_3$  concentrations in the low population region, the resulting incidence of mortalities shown in Figures 7b and 7d (0.592 versus 0.268 deaths) and respiratory morbidity shown in Figure 7c and 7e (2.771 versus 1.252 cases), is much lower. Thus, for the same  $\text{NO}_x$  emission and a smaller  $\text{O}_3$  increase, the damage (i.e., higher mortality and morbidity) was substantially greater in the high population region.



**Figure 6.** Population density in the eastern United States [*CIESIN et al.*, 2000].





**Figure 7.** O<sub>3</sub> produced by an identical amount of NO<sub>x</sub> emitted on 10 July 1995 from a power plant a) upwind from a relatively low population region and b) upwind of a high population region. The mean temperature at the two sites was nearly identical: 83.9°F ±

6.3°F in the high population region and 83.4°F ± 5.8°F in the low population region. The rest of the notation is as described in Figure 3.

Regulation of NO<sub>x</sub> emissions that ignores the variability of downwind population density misses an opportunity to decrease the health effects of emissions. As in the case of variations in biogenic hydrocarbon emissions from place to place, trades between regions under a cap-and-trade program in which all NO<sub>x</sub> emissions from large point sources in a region are treated identically could have a substantial effect (up or down, depending on the direction of the exchange) on health outcomes. A regulatory system that provides incentives for emitters to reduce their emissions in regions of high downwind population density would be more effective at reducing the adverse effects on exposed populations than would a regulatory program that assumes emissions in all locations will have identical effects. Variations in daily wind direction can determine whether a particular NO<sub>x</sub> source is upwind or downwind from a large population center. Ideally, a chemical weather forecasting system could be used to predict a few days in advance what the likely damage that NO<sub>x</sub> emitted today would cause. Emitters could then take that information into account when making decisions as to whether to control their emissions, import power from elsewhere, or purchase or sell permits to cover the damage that their emissions were projected to cause.

## **5. Monetizing health costs resulting from O<sub>3</sub> produced from point source NO<sub>x</sub> emissions**

In the standard economic analysis of externalities, the problem that externalities cause is an inefficient allocation of resources. Much as a producer of electricity would be expected to use “too much” coal if it were provided free of charge (and as a result, in a competitive system, the price of electricity will be “too low”), a producer of electricity who is charged nothing for NO<sub>x</sub> emissions may be expected to emit “too much” NO<sub>x</sub>. A standard corrective in the economic theory of pollution is to determine the “right” price for the unpriced input (here NO<sub>x</sub> emissions), and to charge the firms that use NO<sub>x</sub> emissions in production via a fee or tax. The argument is that the producing firm “should” pay the price for these inputs, just as the firm pays for other

inputs, not as a moral matter but as a condition for avoiding a use of resources that result in a loss of potential value to the ultimate consumers of goods and services and to the general public.

Our analysis indicates potentially large differences in the damage done by incremental  $\text{NO}_x$  emissions from sources at different locations and under different regional VOC and meteorological conditions, where the impacts have been expressed in terms of change in mortality and morbidity risks. The logic of correcting a negative externality calls for expressing these effects in monetary terms, so that they can be compared with the cost of corrective actions. In the case of health impacts, the needed valuation is the willingness of people to pay for reduced risks of suffering them.

A large literature addresses the estimation of the monetary value of health effects. A deeper analysis than the one we undertake here would consider issues such as whether the willingness to pay for reduced risk of health effects depends on the age of the affected people, their incomes, number of dependents, etc. For purposes of the present analysis, we take off-the-shelf figures for the value of a statistical life or an illness. Mortality valuation estimates, obtained from 26 different studies that use both contingent valuation and labor market valuation methods to approximate the willingness to pay to avoid the occurrence of a premature death, give values of a statistical life that range from \$0.6 million - \$13.5 million with a mean of \$4.8 million in 1990 U.S. dollars [EPA, 1999; Viscusi, 1992]. The model simulations represent  $\text{O}_3$  production during July 1995, so in order to calculate the value of the damages caused by the  $\text{NO}_x$  emissions in our 1995 simulations in contemporary terms, we use the 17% increase in the consumer price index (CPI) between 1990 and 1995 to adjust the mean value of a death to 1995\$5.6 million [DoL, 2004]. Due to the increase in CPI between 1995 and 2004 these same deaths, had they occurred in 2004, would have a valuation 24 % higher than the 1995 value [DoL, 2004].

For the specified identical quantities of  $\text{NO}_x$  emitted from individual power plants over 24-hour periods, mortalities resulting from the  $\text{O}_3$  produced over the subsequent several days ranged from a minimum of approximately 0.1 deaths in the low isoprene region to 0.6 deaths when emitted upwind from a region of high population density. Using the mean inflation-adjusted value of 1995\$5.6 million/life, the monetary cost of the deaths in our six cases varies

between a low of \$670,000 to a high of \$3.3 million. The NO<sub>x</sub> emission change (62.5 tons) that gives rise to these impacts is comparable to shutting down a large power plant for a day. For an emission of this magnitude, the estimated value of the impact per incremental ton emitted ranges in the six cases from \$10,700 to \$52,200.

The literature on the cost of morbidity due to an illness indicates substantial variation depending on the illness. Chronic bronchitis is usually estimated to have the highest cost and a restricted activity day among the lowest. We utilize a morbidity dose-response function to estimate the relative risk of a hospital admission due to an increase in daily 1-hour maximum O<sub>3</sub> concentrations. We therefore use a health valuation for all respiratory hospital admissions of 1990\$6100/case [EPA, 1997] which adjusted for inflation is 1995\$7020/case. Thus, for the identical quantities of NO<sub>x</sub> used in our exercises, emitted from individual power plants over 24-hour periods, the calculated morbidities range from 0.57 cases for the low isoprene region to 2.8 cases for the high population region. The implied estimated costs of total respiratory effects range from 1995\$4000 to 1995\$19,700 or from \$64. to \$315. per ton NO<sub>x</sub> emitted. Given valuations found in the literature, the morbidity impacts of NO<sub>x</sub> emissions are tiny when compared with the mortality risk effects.

## **6. Policy Implications**

The implications of our investigation for policy design are potentially wide-ranging. Focusing only on the mortality effects, which our calculations suggest are of greater policy significance, the estimated impact of an incremental ton of NO<sub>x</sub> emitted ranges from 0.00195 to 0.00949 fatalities (a factor of almost 5) depending on the ambient temperature and location of emission. By shifting 13.2 tons of NO<sub>x</sub> emissions per day from the (relatively) high-damage to lower-damage location over a ten-day period one could avoid the loss of approximately one life (on average).

The figures also suggest a significant influence of meteorological conditions on the health impact of incremental NO<sub>x</sub> emissions. The mortality impact of emissions from a single location may vary by a factor of nearly two as the temperature varies within a short span of time. In our simulations we calculated 0.00566 and 0.00296 deaths per incremental ton of NO<sub>x</sub> emitted for

the high and low temperature scenarios, respectively. Reducing O<sub>3</sub> generation and thus saving human lives might be brought about, for example, by reallocating electricity production in space to reduce NO<sub>x</sub> emissions under high-temperature stagnant weather conditions and by utilizing pollution control devices more intensely in locations of high population exposure. The use of a chemical weather forecast system would assist in predicting what damages on a given day are likely to be given that days weather conditions.

Placing monetary value on the damages that result from incremental NO<sub>x</sub> emissions (taking into account, as we do here, only the contribution to downwind O<sub>3</sub> formation and neglecting the health damages from secondary PM formation) suggests that the magnitudes are significant in relation to the costs of abating the emissions. Still considering only fatality effects, and using mean off-the-shelf estimates of the value of a statistical life saved, the estimated damage per incremental ton of emissions in our scenarios ranges from 1995\$10,700 to 1995\$52,200. To get a sense of what these figures mean, we compare them to the price of NO<sub>x</sub> emission allowances under the SIP Call cap and trade system. One allowance permits an affected source to emit one ton of NO<sub>x</sub> between May 1 and September 30 for a given year. As reported at <http://www.evomarkets.com/> on 22 July 2004, the last price for NO<sub>x</sub> emission allowances for 2004 was \$2,550, for 2005, \$3,750, and for 2006, \$3,200. Using a CPI adjustment of 0.8 [DoL, 2004], these prices are equivalent to 1995\$2040., 1995\$3000., 1995\$2560., respectively. These figures can be read as measures of the cost of incremental NO<sub>x</sub> abatement during the ozone season (*given* the existing regulations governing power plants, which have complex implications for the marginal cost of abatement). Economic theory predicts that the marginal cost of abatement of NO<sub>x</sub> will be driven to the market price of allowances, with the important proviso that the marginal cost may exceed the allowance price for all levels of abatement. Although the allowance prices cannot be directly compared with our estimate of the marginal damage due to NO<sub>x</sub> emissions (since both the year and the underlying emissions picture is different), the figures certainly suggest the possibility that net gains are possible from more stringent, and as significantly, more effective, NO<sub>x</sub> control.

The textbook remedy for a defect in the allocation of resources that results in an environmental externality is a corrective tax or fee (usually called a Pigovian tax after the British

economist A.C. Pigou, who first elaborated the idea) on the action that gives rise to the externality. The damage done by a marginal unit of NO<sub>x</sub> emissions under a variety of conditions would, under ideal competitive conditions, be exactly the emission fees called for in theory to correct for the polluting effect. Taking those fees into account would induce NO<sub>x</sub> emitters to make decisions about how much, when, and where to emit that are “correct” in the usual sense used in economics, that is, in the same sense that the wages of workers and prices of fuel induce correct decisions about the quantity of these goods and services to use. The marginal damages from our calculations can be thought of as preliminary estimates of the needed Pigovian fees.

An important question, which we do not address here, is how much difference it would make if NO<sub>x</sub> emitters had to pay for the damage they cause. In economists’ jargon, a lot depends on elasticities. If, for example, raising the price of electricity to reflect the health costs implied by our analysis would not affect the amount of electricity demanded (the price elasticity of demand is low); if raising the cost to producers of emitting NO<sub>x</sub>, perhaps varying it substantially over short periods of time, would have little effect on their decisions about how much or where to produce (the price elasticity of producer demand for NO<sub>x</sub> emissions as an input to producing electricity is low), then the substitutions of the sort described above as possible tradeoffs would not occur, or would not occur to significant degree. If, however, as experience suggests is true in most areas of economic activity, there is substantial flexibility in the time and place of production, and substantial flexibility in the commercial and household use of electricity, especially given time to adjust to a different price regime, then large gains are to be had from a regulatory regime that provides the “right” incentives to respond to differences at the margins we have identified. These differences include: meteorological conditions from the perspective of a given source, differences in the biogenic VOC emissions in source areas, and differences in the population density downwind of emission sources.

The tendency in the recent evolution of NO<sub>x</sub> regulation has been to rely on cap and trade systems. In particular, under the recent OTC and NO<sub>x</sub> SIP Call trading regimes implemented in the northeastern United States, fixed totals of emissions are set for the entire ozone season. Economic theory predicts that trading among sources will tend to equalize the marginal cost of abatement across emitters, where abatement is measured by reductions in emissions at any point

in the ozone season. This system misses the distinctions identified in this paper that determine the downstream health impacts of marginal emissions: differences in meteorological conditions within the ozone season, variations in biogenic hydrocarbon emissions and downwind population densities. Variations on cap and trade systems that would address these shortcomings are possible. For example, a modified cap-and-trade program with “zones” could be adopted in which trading in only one direction is permitted (eg. from regions upwind to downwind of high population centers, from regions of high to low emissions of isoprene, etc.). The same sort of modeling work as is carried out in this paper could be used in the design of such alternatives to guide producers to reduce their emissions at times and in places where the emissions cause the most damage.

## **7. Conclusions and Future Work**

In this paper we present a proof-of-concept use of an integrated assessment that combines an atmospheric model with economic and demographic information to estimate the health damage externality due to ozone ( $O_3$ ) formation from nitrogen oxide ( $NO_x = NO + NO_2$ ) emissions from large point sources. We use a regional atmospheric model of the eastern United States (CAMx) to examine the variation in the amount of  $O_3$  produced as a result of locating, at times and places with strategically chosen differences in temperature and local biogenic hydrocarbon emissions, a fixed reduction in the nitrogen oxides emitted from power plants within a 9-day period in July 1995.. Results of the CAMx model indicate that for the same  $NO_x$  emission the  $O_3$  produced can vary by more than a factor of three. In addition, we showed that the variation in health damages that the resulting  $O_3$  causes depends strongly on the size of the exposed population. Within our span of days and conditions examined, variation in  $O_3$  production and downwind population can together result in a factor of six difference in resulting mortalities for an identical change in the quantity of  $NO_x$  emitted.

The increasingly widely adopted emission cap-and-trade approach has been very successful at reducing total  $NO_x$  emissions from large point sources [EPA, 2004b]. However, because it does not control for the location or time during the summer that emissions take place

nor for the resulting damages, it is less successful at minimizing the damages that result from the emissions permitted under the cap. To reduce total damage, consideration could be given to a system of fees for emitters to provide incentives for them to reduce NO<sub>x</sub> emissions at times and in locations where health damages are greatest. This could be achieved by coupling a regional atmospheric chemistry model that calculates O<sub>3</sub> concentrations obtained from NO<sub>x</sub> emitted from individual point sources with estimates of resulting health damages and their economic cost. Such a system would involve a transformation of air quality regulation as it would create an incentive system to reduce total damages in an economically efficient manner rather than simply requiring a reduction in total emissions or the achievement of a particular uniform air quality goal regardless of location or resulting damage.

Although we have suggested the possibility of using model-derived estimates for purposes of setting emission fees, the details of such a system merit examination. Since the fees are to depend on weather and other conditions, would they be set after the fact (making the emitters, in effect, use the model in forecast mode to make their decisions), or would forecasts be used by the regulator to set fees in advance (based on expected damages)? Regardless of which approach were used, refinements in chemical weather forecasting systems would clearly be beneficial to enable the regulatory system to estimate more accurately appropriate corrective fees, providing emitters with the information they need to adjust their power production and NO<sub>x</sub> emission plans to minimize total costs which include mortality and morbidity risks as well as abatement measures.

In addition, to account more fully for the health effects of power plant NO<sub>x</sub> emissions it is necessary to include the formation of secondary particulates (PM) and their impact on mortality and morbidity. Secondary PM, in addition to O<sub>3</sub>, are formed from NO<sub>x</sub> emissions in winter as well as summer. Using an integrated assessment model without chemistry, analyses of the benefits of an annual (rather than just summer) NO<sub>x</sub> emissions cap were found to yield substantial net benefits due to reduction of PM<sub>2.5</sub> concentrations and its detrimental health effects [Burtraw *et al.*, 2003; Burtraw *et al.*, 2001]. Additional analyses that include both gas and aerosol chemistry and an examination of the effects that spatial and temporal variability in emissions have on PM<sub>2.5</sub> formation and resulting health damages and costs would be useful.



Such an analysis would permit a determination of the total health damage NO<sub>x</sub> emissions cause; the fees that an emitter might be charged for the total damage could thus be determined.

## References

- Atkinson, S.E., and T.H. Tietenberg, Economic implications of emissions trading rules for local and regional pollutants, *Canadian Journal of Economics*, 87, 370-386, 1987.
- Burtraw, D., R. Bharvirkar, and M. McGuinness, Uncertainty and the net benefits of emissions reductions of nitrogen oxides from electricity generation, *Land Economics*, 79 (3), 382-401, 2003.
- Burtraw, D., K.L. Palmer, R. Bharvirkar, and A. Paul, Cost Effective Reduction of NOx Emissions from Electricity Generation, *Journal of the Air and Waste Management Association*, 51, 1476-1489, 2001.
- Byun, D.W., and J.K.S. Ching, Science Algorithms of the EPA Models-3 Community Multiscale Air Quality (CMAQ) Modeling System, U.S. Environmental Protection Agency, 1999.
- Carter, W.P.L., Condensed atmospheric photooxidation mechanisms for isoprene, *Atmos. Environ.*, 30, 4275-4290, 1996.
- CDC-NCHS, 1998, Rate of discharges from short-stay hospitals, by age and first-listed diagnosis: United States, 1998, Center for Disease Control (CDC), National Center for Health Statistics, accessed: February 3, 2003, <http://www.cdc.gov/nchs/about/major/hdasd/nhdstab.htm>.
- CDC-NCHS, Deaths from 282 Selected Causes by 5-Year Age Groups, Race, and Sex: Each State and the District of Columbia, 1995, 1996, 1997, and 1998, Center for Disease Control, National Center for Health Statistics, [http://www.cdc.gov/nchs/data/mortab/98gm3\\_01.pdf](http://www.cdc.gov/nchs/data/mortab/98gm3_01.pdf), 2003.
- CIESIN, IFPRI, and WRI, Gridded Population of the World (GPW), Version 2, Center for International Earth Science Information Network (CIESIN); International Food Policy Research Institute (IFPRI); World Resources Institute (WRI), <http://sedac.ciesin.columbia.edu/plue/gpw>, 2000.
- Dockery, D.W., C.A. Pope, III, X. Xu, J.D. Spengler, J.H. Ware, M.E. Fay, B.G. Ferris, and F.E. Speizer, An Association between Air Pollution and Mortality in Six U.S. Cities, *N. Eng. J. Med.*, 329, 1753-1759, 1993.
- DoL, 2004, Consumer Price Indexes, U.S. Department of Labor, Bureau of Labor Statistics, accessed: June 3, 2004, <http://www.bls.gov/cpi/home.htm#data>.
- ENVIRON, User's Guide: Comprehensive Air Quality Model with Extensions (CAMx) version 3.00, 2000.
- EPA, The benefits and costs of the Clean Air Act, 1970-1990, United States Environmental Protection Agency Office of Air and Radiation Policy, <http://www.epa.gov/airprog/oar/sect812/>, 1997.
- EPA, U.S., Benefits and Costs of the Clean Air Act: 1990-2010, U.S. Environmental Protection Agency, <http://www.epa.gov/air/sect812/copy99.html>, 1999.
- EPA, U.S., 2000, National Air Quality and Emission Trends Report:1998, U.S. Environmental Protection Agency, accessed: May 19, 2004, <http://www.epa.gov/oar/aqtrends.html/>.
- EPA, U.S., Benefits and Costs of the Clean Air Act 1990 - 2020: Revised Analytical Plan For EPA's Second Prospective Analysis, U.S. Environmental Protection Agency, <http://www.epa.gov/air/sect812/blueprint.html>, 2003.

- EPA, U.S., 2004a, Ozone and Health - A Timeline, U.S. Environmental Protection Agency, accessed: May 19, 2004, <http://www.epa.gov/ozonedesignations/timeline.htm>.
- EPA, U.S., 2004b, Ozone Transport Commission (OTC) NOx Budget Program, U.S. EPA Clean Air Markets - Programs and Regulations, accessed: July 20, 2004, <http://www.epa.gov/airmarkets/otc/>.
- EvoMarkets, 2003, Environmental Markets - EPA SIP NOx trading program, accessed: July 20, 2004, <http://www.evomarkets.com/>.
- Fiore, A.M., D.J. Jacob, I. Bey, R.M. Yantosca, B.D. Field, and A.C. Fusco, Background ozone over the United States in summer: Origin, trend, and contribution to pollution episodes, *J. Geophys. Res.*, 107 (D15), 10.1029, 2002.
- Geron, C.D., A.B. Guenther, and T.E. Pierce, An improved model for estimating emissions of volatile organic compounds from forests in the eastern United States, *J. Geophys. Res.*, 94, 12,925-12,956, 1994.
- Guenther, A., P. Zimmerman, P. Harley, R. Monson, and R. Fall, Isoprene and monoterpene emission rate variability: Model evaluations and sensitivity analysis., *J. Geophys. Res.*, 94, 12,925-12,956, 1994.
- Lin, C.-Y.C., D.J. Jacob, and A. Fiore, M., Trends in exceedances of the ozone air quality standard in the continental United States, 1980-1998, *Atmos. Environ.*, 35, 3217-3228, 2001.
- Liu, S.C., M. Trainer, F.C. Fehsenfeld, D.D. Parrish, E.J. Williams, D.W. Fahey, G. Hubler, and P.C. Murphy, Ozone production in the rural troposphere and the implications for regional and global ozone distributions, *J. Geophys. Res.*, 92, 4191 - 4207, 1987.
- Logan, J.A., Ozone in rural areas of the United States, *J. Geophys. Res.*, 94 (D6), 8511-8532, 1989.
- NRC, Air Quality Management in the United States, National Research Council of the National Academies, Washington D.C., 2004.
- Palmer, P.I., D.J. Jacob, A.M. Fiore, R.V. Martin, K. Chance, and T.P. Kurosu, Mapping of isoprene emissions over North America using formaldehyde column observations from space, *J. Geophys. Res.*, 108 (D6), doi:10.1029/2002JD002153, 2003.
- Pierce, D., C. Geron, L. Bender, R. Dennis, G. Tonnesen, and A. Guenther, Influence of increased isoprene emissions on regional ozone modeling, *J. Geophys. Res.*, 103 (D19), 25,611-25,629, 1998.
- Pope, C.A., III, R.T. Burnett, M.J. Thun, E.E. Calle, D. Krewski, K. Ito, and G.D. Thurston, Lung Cancer, Cardiopulmonary Mortality and Long-Term Exposure to Fine Particulate Air Pollution, *J. Amer. Med. Assoc.*, 287 (9), 1132-1141, 2002.
- Pope, C.A., III, M.J. Thun, M.M. Namboodri, D.W. Dockery, J.S. Evans, F.E. Speizer, and C.W. Heath, Particulate Air Pollution as a Predictor of Mortality in a Prospective Study of U.S. Adults, *Amer. J. Resp. Crit. Care Med.*, 151, 669-674, 1995.
- Ryerson, T.B., M. Trainer, J.S. Holloway, D. Parrish, L.G. Huey, D.T. Sueper, G.J. Frost, S.G. Donnelly, S. Schauffler, E.L. Atlas, W.C. Kuster, P.D. Goldan, G. Hubler, J.F. Meagher, and F.C. Fehsenfeld, Observations of ozone formation in power plant plumes and implications for ozone control strategies, *Science*, 292, 719-723, 2001.
- Sillman, S., J.A. Logan, and S.C. Wofsy, The sensitivity of ozone to nitrogen oxides and hydrocarbons in regional ozone episodes, *J. Geophys. Res.*, 95, 1837 - 1851, 1990.

- Smolarkiewicz, P.K., A Simple positive definite advection scheme with small implicit diffusion, *Mon. Wea. Rev.*, *111*, 479-486, 1983.
- Steib, D.M., S. Judek, and R.T. Burnett, Meta-Analysis of Time-Series Studies of Air Pollution and Mortality: Effects of Gases and Particles and the Influence of Cause of Death, Age, and Season, *J. Air & Waste Manage. Assoc.*, *52*, 470-484, 2002.
- Thurston, G.D., and K. Ito, Epidemiological Studies of Ozone Exposure Effects, in *Air Pollution and Health*, edited by S. Holgate, J. Samet, H. Koren, and R. Maynard, Academic Press, 1999.
- USCB, Annual Population Estimates by Age Group and Sex, Selected Years from 1990 to 2000, U.S. Census Bureau, Population Division, <http://eire.census.gov/popest/archives/national/nation2/intfile2-1.txt>, 2000.
- Viscusi, W.K., Fatal tradeoffs: public and private responsibilities for risks, 306 pp., Oxford University Press, New York, 1992.
- Vukovich, F.M., Regional-scale boundary layer ozone variations in the eastern United States and their association with meteorological variations, *Atmos. Environ.*, *29* (17), 2259-2273, 1995.
- Wesley, M.L., Parameterization of surface resistances to gaseous dry deposition in regional-scale numerical models, *Atmos. Environ.*, *23*, 1293-1304, 1989.