

Center


Discussion paper

CBM
R
8414
8414
1993
12

for
omic Research



* C I N 0 0 6 7 2 *



No. 9312

**MARGINALIZATION AND CONTEMPORANEOUS
AGGREGATION IN MULTIVARIATE
GARCH PROCESSES**

by Theo Nijman
and Enrique Sentana

February 1993

ISSN 0924-7815

MARGINALIZATION AND CONTEMPORANEOUS AGGREGATION
IN MULTIVARIATE GARCH PROCESSES

Theo Nijman
Enrique Sentana
February 1993*

Abstract

In this paper we first of all show that contemporaneous aggregation of independent univariate GARCH processes yields a process which satisfies the weak GARCH conditions introduced by Drost and Nijman (1992). Subsequently we analyze the dependence of the parameters in the aggregate on the parameters in the underlying models and present numerical results for the aggregation of two GARCH(1,1) processes with identical "persistence" parameters, and a GARCH(1,1) process with conditionally homoskedastic white noise. We show that the variance parameters after aggregation depend on the underlying variance and kurtosis parameters. Subsequently, we generalize the results by showing that a linear combination of variables generated by a multivariate GARCH process will also be weak GARCH and analyzing the parameters in this weak GARCH process. We also derive the marginal (weak GARCH) processes implied by several multivariate GARCH processes. The results explain why GARCH is found in univariate as well as multivariate series and can be used to facilitate the specification of multivariate GARCH processes.

* Dept. of Econometrics, Tilburg University, P.O.Box 90153, 5000 LE Tilburg, The Netherlands and CEMFI, Casado del Alisal 5, 28014 Madrid, Spain, respectively. This research was initiated when the second author was affiliated at the L.S.E. and the first author was visiting there. The authors benefitted from financial support of the Royal Netherlands Academy of Arts and Sciences (K.N.A.W.) and the E.S.R.C. respectively.

1. Introduction

It is a well known stylized fact that many financial time series such as stock returns, exchange rates or interest rates exhibit conditional heteroskedasticity. For univariate time series, the ARCH model proposed by Engle (1982) or the GARCH model proposed by Bollerslev (1986) are usually used to parametrize the conditional heteroskedasticity. Multivariate extensions of these models have been proposed e.g. by Engle (1987), Attanasio and Edey (1987), Bollerslev, Engle and Wooldridge (1988), Diebold and Nerlove (1989), Baba, Engle, Kraft and Kroner (1989), Engle, Ng and Rothschild (1990) and King, Sentana and Wadhvani (1991). An excellent survey of the many applications of GARCH processes is provided by Bollerslev, Chou and Kroner (1992).

In the existing literature univariate or multivariate GARCH models are typically assumed for the time series under consideration, without wondering whether these assumptions are consistent with GARCH assumptions for other time series and/or at other frequencies. GARCH models have been estimated e.g. for the (log) returns in the Deutsche Mark / US Dollar exchange rate, the US Dollar / Japanese Yen rate and the Deutsche Mark / Japanese Yen rate. As the returns on the third exchange rate are simply the sum of the returns on the first two exchange rates, the GARCH models for these exchange rates implicitly specify a model for the third exchange rate as well. Similarly GARCH models have been fitted to returns on individual stocks as well as to returns on portfolios, but once more the relation between the models for the individual stocks and the one for the portfolio has not been considered explicitly. Also multivariate GARCH models are used nowadays to describe e.g. the joint behaviour of interest rates, exchange rates and stock returns. Little attention has been paid however to the implication of such a multivariate model for the univariate processes of the components. In this paper we derive the properties of linear combinations of variables generated by a multivariate GARCH process as well as the properties of marginal processes implied by a multivariate GARCH assumption. We show that the parametric structure of the commonly used GARCH models is lost by taking linear combinations or by marginalizing. Nevertheless we show that the linear combinations and marginal processes will still be weak GARCH processes as

defined by Drost and Nijman (1992). These results explain why GARCH is found in univariate as well as multivariate series and can be used to facilitate the specification of multivariate GARCH processes. Throughout this paper we restrict ourselves to bivariate models with GARCH(1,1) variances in order to keep the algebra simple. Our general framework however applies equally well to more general cases.

Consider a univariate time series $\{y_t\}$ which is stationary and symmetric with finite fourth moments. Drost and Nijman (1992) have defined $\{y_t\}$ to be weak GARCH(1,1) if in the recursion

$$\sigma_t^2 = \psi + \beta \sigma_{t-1}^2 + \alpha y_{t-1}^2 \quad (1)$$

the parameters ψ , α and β can be chosen such that

$$P[y_t \mid y_{t-1}, y_{t-2}, \dots] = 0 \quad (2)$$

and

$$P[y_t^2 \mid y_{t-1}, y_{t-2}, \dots] = \sigma_t^2 \quad (3)$$

where $P[x_t \mid z_{t-1}, z_{t-2}, \dots]$ denotes the best linear predictor of x_t in terms of a constant and values of z_{t-i} and z_{t-i}^2 ($i = 1, 2, \dots$). Drost and Nijman (1992) have shown that GARCH models which assume that y_t/σ_t is i.i.d. as proposed by Bollerslev (1986), which they refer to as strong GARCH models, are not closed under temporal aggregation. A strong GARCH assumption e.g. at the daily frequency is inconsistent with a strong GARCH assumption at the weekly frequency. In addition they have shown that the class of weak GARCH processes is closed under temporal aggregation. In this paper we complement their results by showing that contemporaneous aggregation of independent univariate GARCH processes yields a weak GARCH process. We analyze the dependence of the parameters in the aggregate process on the parameters in the underlying models. Subsequently, we generalize this result by showing that a linear combination of variables generated by a multivariate GARCH process will also be weak GARCH and analyzing the parameters in this weak

GARCH process. Finally we derive the marginal weak processes implied by multivariate GARCH processes. The parameters in weak GARCH processes can easily be estimated consistently. Moreover simulation experiments suggest that in many cases the commonly used estimators, which are ML under the assumption that the model is strong GARCH and the conditional distribution normal, converge to values close to the weak GARCH parameters in large samples.

The plan of this paper is as follows. In section 2 we will consider aggregation of independent GARCH processes. Numerical results are presented on the GARCH parameters that arise through aggregation of a GARCH process with conditional homoskedastic noise, as well as through the aggregation of two GARCH processes with identical "persistence" parameters. In section 3 we introduce an unrestricted bivariate GARCH(1,1) model and check that this structure is preserved under linear transformation. Moreover we show how marginal processes can be derived from multivariate representations. In section 4, diagonal models, conditionally orthogonal models and factor GARCH models, which all are special cases of the general set-up in section 3, are analyzed in more detail. Numerical results are presented on the impact of marginalization on the properties of linear combinations of variables generated by these models. Section 5 contains the results of a simulation experiment on the properties of the commonly used quasi maximum likelihood estimator in cases where data are generated by aggregating independent univariate GARCH processes or by marginalizing from multivariate GARCH processes. Finally section 6 concludes. Some technicalities are outlined in appendices.

2. Contemporaneous aggregation of independent univariate GARCH processes

In this section we consider the simple case of aggregation of two independent GARCH(1,1) processes. The results derived in this section are special cases of those in sections 3 and 4, but are derived here in a more straightforward and intuitive manner. Moreover the assumption of conditional normality which will be made in sections 3 and 4 to simplify the algebra can be avoided here.

Consider variables y_{1t} and y_{2t} which are both generated by (strong) GARCH(1,1) models i.e.,

$$y_{it} = \sigma_{it} \xi_{it} \quad (i = 1,2), \quad (4)$$

$$\xi_{it} \text{ i.i.d.}, \quad E \xi_{it} = 0, \quad E \xi_{it}^2 = 1, \quad E \xi_{it}^4 = \kappa_i.$$

$$\sigma_{it}^2 = \nu_i + \beta_i \sigma_{i,t-1}^2 + \alpha_i y_{it-1}^2. \quad (5)$$

It is well known that substitution of $\sigma_{it}^2 = y_{it}^2 - \eta_{it}$ yields an ARMA(1,1) model for y_{it}^2 ,

$$y_{it}^2 = \sigma_i^2 + [1 - (\alpha_i + \beta_i)L]^{-1} [1 - \beta_i L] \eta_{it}, \quad (6)$$

where L is the lag-operator defined by $L y_{it} = y_{it-1}$ and where $\sigma_i^2 = E y_{it}^2 = \nu_i (1 - \alpha_i - \beta_i)^{-1}$ and

$$\eta_{it} = y_{it}^2 - E[y_{it}^2 | I_{t-1}] = (\xi_{it}^2 - 1) \sigma_{it}^2. \quad (7)$$

From (6) one easily obtains

$$\begin{aligned} (y_{1t} + y_{2t})^2 &= \sigma_1^2 + \sigma_2^2 + [1 - (\alpha_1 + \beta_1)L]^{-1} [1 - \beta_1 L] \eta_{1t} + \\ &\quad + [1 - (\alpha_2 + \beta_2)L]^{-1} [1 - \beta_2 L] \eta_{2t} + 2 y_{1t} y_{2t}. \end{aligned} \quad (8)$$

As η_{1t} , η_{2t} and $y_{1t} y_{2t}$ are mutually uncorrelated and none of these three variables is autocorrelated, (8) implies that the sum of two independent strong GARCH(1,1) processes is weak CARCH(2,2). This result is obviously related to the well known result (see e.g. Lütkepohl (1984)) that the sum of two independent ARMA(1,1) processes is ARMA(2,2). One important difference with the ARMA case is the presence of the cross product term in the right hand side of (8), which complicates the derivation of the GARCH parameters for the aggregate series as we will see below.

Obviously equation (8) only yields an upper bound on the orders of the weak GARCH process. In two important special cases, though, aggregation of

GARCH(1,1) processes for which $\alpha_1 + \beta_1 = \alpha_2 + \beta_2$ and aggregation of a GARCH(1,1) process with conditionally homoskedastic white noise, the aggregate will in fact be weak GARCH(1,1). For these special cases we will derive the weak GARCH parameters in this section. The general case is discussed in section 4. The value of $\alpha_1 + \beta_1$ has often been referred to as the persistence of shocks to the volatility in the GARCH process. As this terminology can easily be criticized we will put this terminology between quotation marks whenever it is used. In fact the mean lag of the ARMA model in (6) is probably a much better measure of persistence.

If the "persistence" parameters in the two independent GARCH processes which are added coincide, i.e. $\alpha_1 + \beta_1 = \alpha_2 + \beta_2 = \gamma$, equation (8) reduces to

$$[1-\gamma L] (y_{1t} + y_{2t})^2 = \tilde{\psi} + [1-\beta_1 L] \eta_{1t} + [1-\beta_2 L] \eta_{2t} + 2 [1-\gamma L] y_{1t} y_{2t}, \quad (9)$$

with $\tilde{\psi} = (\psi_1 + \psi_2)$. Because of the properties of η_{1t} , η_{2t} and $y_{1t} y_{2t}$ the right hand side will be a first order moving average process, say $(1-\lambda L)\omega_t$. Rewriting (9) yields

$$(1-\gamma L) (y_{1t} + y_{2t})^2 = \tilde{\psi} + (1 - \lambda L) \omega_t, \quad (10)$$

where

$$\lambda (1+\lambda^2)^{-1} = - E u_t u_{t-1} / E u_t^2,$$

$$u_t = [1-\beta_1 L] \eta_{1t} + [1-\beta_2 L] \eta_{2t} + 2 [1-\gamma L] y_{1t} y_{2t},$$

$$\omega_t = (1 - \lambda L)^{-1} u_t.$$

Once λ is known, it is clear from the comparison with (6) that the weak GARCH parameters for $y_{1t} + y_{2t}$ are simply $(\tilde{\beta}, \tilde{\alpha}) = (\lambda, \gamma - \lambda)$. Note that the value of the parameter $\alpha + \beta$ is the same in the aggregate process as in the two underlying processes. The remaining problem is the determination of λ . From (10) one has

$$E u_t^2 = (1+\beta_1^2) E \eta_{1t}^2 + (1+\beta_2^2) E \eta_{2t}^2 + 4(1+\gamma^2) E y_{1t}^2 y_{2t}^2 \quad (11)$$

and

$$- E u_t u_{t-1} = \beta_1 E \eta_{1t}^2 + \beta_2 E \eta_{2t}^2 + 4\gamma E y_{1t}^2 y_{2t}^2. \quad (12)$$

Obviously $E y_{1t}^2 y_{2t}^2 = \sigma_1^2 \sigma_2^2$ by independence. Moreover $E \eta_{it}^2$ can be expressed as a function of σ_i^2 , α_i , β_i and the kurtosis of the error term $\kappa_i = E \xi_{it}^4$,

$$\begin{aligned} E \eta_{it}^2 &= E (\xi_{it}^2 - 1)^2 E \sigma_{it}^4 \\ &= (\kappa_i - 1) \sigma_i^4 \{1 - (\beta_i + \alpha_i)^2\} / (1 - \beta_i^2 - 2\alpha_i \beta_i - \alpha_i^2 \kappa_i), \end{aligned} \quad (13)$$

where the second equality follows from a straightforward generalization of the way in which Bollerslev (1986) computed the fourth moment of y_{it} . As $\lambda(1+\lambda^2)^{-1} = \lambda^{-1}(1+\lambda^{-2})^{-1}$, the quadratic equation $\lambda(1+\lambda^2)^{-1} = -Eu_t u_{t-1}/Eu_t^2$ yields a unique solution for λ within the unit circle.

Some illustrative numerical results on the aggregation of independent GARCH(1,1) processes with coinciding "persistence" parameters are reported in Table 1. In the first five lines of the table the values α_i and β_i do not differ between the two processes. The results in (9)-(13) indicate that for $y_{1t} + y_{2t}$ the coefficient on the lagged variance term, β , will exceed $\beta_1 = \beta_2$ while $\alpha < \alpha_i$. The numerical values for α and β depend on the variance ratio and the kurtosis parameters. If the variances of the two processes differ a lot or if at least one of the processes has fat tails, the weak GARCH parameters in the aggregate will be close to those in the underlying processes. The last three lines in the table show that the variance ratio can in fact have a very large impact on the weak GARCH parameters. Note that in the case considered in the last three lines the fourth moments condition is violated if $\kappa_i = 9$.

Table 1 Weak GARCH parameters α and β for $y_t = y_{1t} + y_{2t}$ in (1)-(3) if y_{it} ($i=1,2$) is generated by (4)-(5), assuming $\alpha_1 + \beta_1 = \alpha_2 + \beta_2 = \bar{\alpha} + \bar{\beta}$.

β_1	α_1	κ_1	β_2	α_2	κ_2	σ_1^2/σ_2^2	β	α
0.90	0.05	3.00	0.90	0.05	3.00	1.00	0.920	0.030
0.90	0.05	3.00	0.90	0.05	3.00	4.00	0.912	0.038
0.90	0.05	9.00	0.90	0.05	3.00	1.00	0.910	0.040
0.90	0.05	9.00	0.90	0.05	3.00	4.00	0.903	0.047
0.90	0.05	9.00	0.90	0.05	9.00	1.00	0.907	0.043
0.50	0.35	3.00	0.80	0.05	3.00	1.00	0.569	0.281
0.50	0.35	3.00	0.80	0.05	3.00	4.00	0.516	0.334
0.50	0.35	3.00	0.80	0.05	3.00	0.25	0.705	0.145

A second special case of (8) that is worth considering explicitly is aggregation of a GARCH(1,1) variable and conditionally homoskedastic white noise. In this case (8) reduces to

$$\begin{aligned}
 [1 - (\alpha_1 + \beta_1)L] (y_{1t} + y_{2t})^2 = & \tilde{\psi} + [1 - \beta_1 L] \eta_{1t} + [1 - (\alpha_1 + \beta_1)L] \eta_{2t} \\
 & + 2 [1 - (\alpha_1 + \beta_1)L] y_{1t} y_{2t}. \quad (14)
 \end{aligned}$$

Equation (14) shows that the aggregate will in this case be weak GARCH(1,1) and that the "persistence" parameter of the GARCH process is not affected by adding conditionally homoskedastic noise. Along the lines sketched above, the weak GARCH parameters can easily be determined as functions of $(\sigma_1^2, \alpha_1, \beta_1, \sigma_2^2, \kappa_1, \kappa_2)$. Table 2 shows, as could be expected on intuitive grounds, that the GARCH parameters are affected much more by adding noise if the signal to noise ratio is low than when it is high. Also fat tails in the GARCH process lead to a smaller impact of the noise, while fat tailed noise yields larger deviations from the underlying variance parameters.

Table 2 Weak GARCH parameters α and β for $y_t = y_{1t} + y_{2t}$ in (1)-(3) if y_{it} ($i=1,2$) is generated by (4)-(5), assuming $\alpha_2 = \beta_2 = 0$.

β_1	α_1	κ_1	κ_2	σ_1^2/σ_2^2	β	α
0.90	0.05	3.00	3.00	1.00	0.933	0.017
0.90	0.05	3.00	3.00	4.00	0.914	0.036
0.90	0.05	3.00	3.00	0.25	0.947	0.003
0.90	0.05	9.00	3.00	1.00	0.915	0.035
0.90	0.05	3.00	9.00	1.00	0.940	0.010
0.50	0.35	3.00	3.00	1.00	0.571	0.279
0.50	0.35	3.00	3.00	4.00	0.517	0.333
0.50	0.35	3.00	3.00	0.25	0.721	0.129

3. Marginalization and aggregation in multivariate GARCH models: the general case.

In the previous section we restricted ourselves to aggregation of independent GARCH processes. In many applications, e.g. to exchange rates or stock returns, the independence assumption is very strong. For that reason we therefore consider in this section the aggregation of two processes whose joint distribution belongs to a class of bivariate GARCH(1,1) models. We first check that the class is closed under linear transformation and discuss the properties of the weak GARCH processes that arise after marginalization.

The class of bivariate GARCH models that we consider is

$$\begin{bmatrix} y_{1t} \\ y_{2t} \end{bmatrix} = \Sigma_t^{1/2} \begin{bmatrix} \xi_{1t} \\ \xi_{2t} \end{bmatrix}, \quad \begin{bmatrix} \xi_{1t} \\ \xi_{2t} \end{bmatrix} \sim \text{i.i.d. } N(0, I_2), \quad (15)$$

where

$$\Sigma_t = \begin{bmatrix} \sigma_{11t} & \sigma_{12t} \\ \sigma_{12t} & \sigma_{22t} \end{bmatrix},$$

and

$$\begin{aligned}
 \begin{bmatrix} \sigma_{11t} \\ \sigma_{12t} \\ \sigma_{22t} \end{bmatrix} &= \begin{bmatrix} \psi_{11} \\ \psi_{12} \\ \psi_{22} \end{bmatrix} + \begin{bmatrix} b_{11} & b_{12} & b_{13} \\ b_{21} & b_{22} & b_{23} \\ b_{31} & b_{32} & b_{33} \end{bmatrix} \begin{bmatrix} \sigma_{11t-1} \\ \sigma_{12t-1} \\ \sigma_{22t-1} \end{bmatrix} + \\
 &+ \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} \begin{bmatrix} y_{1t-1}^2 \\ y_{1t-1}y_{2t-1} \\ y_{2t-1}^2 \end{bmatrix}. \quad (16)
 \end{aligned}$$

As we assume throughout that Σ_t is a positive-semidefinite matrix (see Baba et al. (1989) for sufficient conditions), it can be interpreted as $\text{Var}_{t-1}(y_t)$ where $y_t' = (y_{1t}, y_{2t})$. In the sequel we also assume that the parameters in (16) are such that fourth moments of y_t exist. In a more compact matrix notation (16) can be written as

$$s_t = \psi + B s_{t-1} + A z_{t-1}, \quad (17)$$

where $s_t' = (\sigma_{11t}, \sigma_{12t}, \sigma_{22t}) = \{\text{vech}(\Sigma_t)\}'$, $\psi' = (\psi_{11}, \psi_{12}, \psi_{22})$ and $z_t' = (y_{1t}^2, y_{1t}y_{2t}, y_{2t}^2) = \{\text{vech}(y_t y_t')\}'$. The model in (15) and (17) is a particular case of the multivariate GARCH(p,q) model considered in Bollerslev, Engle and Wooldridge (1988). If $b_{ij} = 0$ ($\forall i, j$) one obtains the original bivariate ARCH(1) specification of Kraft and Engle (1982). Other interesting special cases are the diagonal and factor GARCH model, and models which yield conditional orthogonality. These special cases will be considered in the next section.

An important property of the bivariate GARCH model in (16) is that its structure is preserved under linear transformation (see Baba et al. (1989)). In appendix A we show that if P is a non-singular matrix, the vector $x_t = P y_t$ will also satisfy the bivariate GARCH model in (15) and (17) with the parameters ψ , B and A replaced by $\tilde{\psi}$, \tilde{B} and \tilde{A} defined as

$$\begin{aligned}
 \tilde{\psi} &= D^+ (P\Theta P) D \psi, \\
 \tilde{B} &= D^+ (P\Theta P) D B D^+ (P^{-1}\Theta P^{-1}) D, \\
 \tilde{A} &= D^+ (P\Theta P) D A D^+ (P^{-1}\Theta P^{-1}) D,
 \end{aligned} \quad (18)$$

where D is the duplication matrix defined by $\text{vec}(\Sigma_t) = D \text{vech}(\Sigma_t)$ and D^+ is its Moore-Penrose inverse.

A second important aspect in the analysis of the bivariate GARCH model is the derivation of the implied marginal processes. Marginalization is required as the second step in assessing the impact of contemporaneous aggregation on variables that are generated by the multivariate GARCH processes, but it is moreover of substantial interest in its own right and can be used to facilitate the specification of multivariate GARCH processes. In order to discuss marginalization define $\eta_t = z_t - E_{t-1}[z_t]$. Substitution in (17) yields the trivariate ARMA(1,1) process

$$[I - (A+B)L] z_t = \psi + [I - BL] \eta_t. \quad (19)$$

The marginal process for y_{1t} can be obtained from the first row of the final form of the model (see e.g. Zellner and Palm (1974)),

$$\det \{I - (A+B)L\} z_t = \psi^* + \text{adj} \{I - (A+B)L\} (I - BL) \eta_t \quad (20)$$

where $\psi^* = \text{adj} \{I - (A+B)L\} \psi$ and where \det and adj refer to the determinant and adjoint matrix respectively. As the determinant is a third order polynomial in the lag operator and the elements of the adjoint matrix are second order polynomials in the lag operator, (20) shows that the marginal process for y_{1t} will be at most weak GARCH(3,3). As we shall see in the next section, though, in many cases of interest a reduction of these orders can be obtained. Of course a reduction through coincidental parameter cancellation is always a possibility.

To find the weak GARCH parameters, we proceed as in (10) and write the ARMA(3,3) model for y_{1t}^2 as

$$(1 - \pi_1 L - \pi_2 L^2 - \pi_3 L^3) y_{1t}^2 = c + (1 - \lambda_1 L - \lambda_2 L^2 - \lambda_3 L^3) \omega_t; \quad E \omega_t^2 = \sigma_\omega^2. \quad (21)$$

The autoregressive coefficients in the ARMA model in squares, which correspond to the persistence parameters in the weak GARCH model, can easily be found using a well known result for characteristic equations,

$$\det \{I-(A+B)L\} = -\det(A+B) L^3 + p(A+B) L^2 - \text{tr}(A+B) L + 1 \quad (22)$$

with

$$p(A+B) = \det(A_{11}^- + B_{11}^-) + \det(A_{22}^- + B_{22}^-) + \det(A_{33}^- + B_{33}^-) \quad (23)$$

where A_{ij}^- is the 2×2 matrix obtained from A by eliminating row i and column j . The determination of σ_ω^2 and of the MA coefficients λ_1 is less straightforward. These coefficients can be obtained by equating the variance and first three autocovariances of $(1-\lambda_1 L - \lambda_2 L^2 - \lambda_3 L^3)\omega_t$ with those of the first element of the right hand side of (20). Given that η_t is serially uncorrelated with zero mean, only its contemporaneous variance-covariance matrix is required. This is derived in appendix B.

4. Marginalization and aggregation in multivariate GARCH models: special cases.

Given the large number of parameters involved, restrictions have usually been imposed in order to estimate the bivariate GARCH model. Three special cases will be analyzed in three subsections: the diagonal model, models in which y_{1t} and y_{2t} are conditionally orthogonal and the factor GARCH model.

4.1 The diagonal GARCH model

A first special case that has been popular is the diagonal GARCH model estimated e.g. by Attanasio and Edey (1987) and Bollerslev, Engle and Wooldridge (1988) where A and B are both assumed to be diagonal. If moreover $a_{22} = b_{22} = 0$ the independent case that has been analyzed in section 2 is obtained. Equation (20) shows that the marginal process for y_{1t} arising from a diagonal GARCH model will be weak GARCH with parameters $(\alpha, \beta) = (a_{11}, b_{11})$, irrespective of the fact whether the two processes are independent, i.e. whether $a_{22} = 0$ and $b_{22} = 0$, or not. The stronger result that the marginal process arising from a diagonal model will be strong GARCH can easily be verified directly.

The general results in section 3 show that linear combinations $\delta_1 y_{1t} + \delta_2 y_{2t}$ will be weak GARCH(2,2) if y_{1t} and y_{2t} are generated by independent GARCH(1,1) processes, which is the result that has been obtained in section 2 along different lines. Linear combinations of variables generated by the diagonal GARCH model on the other hand will be weak GARCH(3,3) in general as there is no reason to expect cancelling roots in (21) in this case. This is caused by the fact that the class of diagonal GARCH processes is not closed under linear transformation.

4.2 The conditionally orthogonal case

A second special case arises when y_{1t} and y_{2t} are conditionally orthogonal, so that $\sigma_{12t} = 0$ ($\forall t$) i.e. $a_{21} = a_{22} = a_{23} = b_{21} = b_{22} = b_{23} = b_{12} = b_{32} = \psi_{12} = 0$. The model that arises is (15) with $\sigma_{12t} = 0$.

$$\begin{aligned} \sigma_{11t} = & \psi_{11} + b_{11}\sigma_{11t-1} + b_{13}\sigma_{22t-1} + \\ & + a_{11}y_{1t-1}^2 + a_{12}y_{1t-1}y_{2t-1} + a_{13}y_{2t-1}^2 \end{aligned} \quad (24)$$

and an analogous expression for σ_{22t} . It is not hard to check that in general the marginal process for y_{1t} will be weak GARCH(2,2). Obviously the model reduces to GARCH(1,1) if $b_{13} = a_{12} = a_{13} = 0$. Interestingly this arises in other cases as well, e.g. if y_{2t} is homoskedastic, i.e. $b_{31} = b_{33} = a_{31} = a_{32} = a_{33} = 0$. Also, if y_{2t} only depends on its own squared values ($b_{31} = a_{31} = a_{32} = 0$) and moreover $b_{13} = a_{13} = 0$, weak GARCH(1,1) is obtained as the marginal process for y_{1t} whether or not its conditional variance depends on lagged values of $y_{1t}y_{2t}$. The results in section 3 can be used to show that linear combinations $\delta_1 y_{1t} + \delta_2 y_{2t}$ of conditionally orthogonal processes y_{1t} and y_{2t} will be weak GARCH(2,2), as in the independent case. Remember that in the general case the linear combination is GARCH(3,3). In the independent case the marginal process will be GARCH(1,1), while it is GARCH(2,2) in the conditionally orthogonal case. Similar the order of the GARCH process of linear combinations of the two variables distinguishes conditionally orthogonal processes from the general case. The above results

can be used in principle to distinguish between independent and conditionally orthogonal processes using univariate methods only, although this may be difficult to do in practice.

4.3 The factor GARCH model

The restrictions in the third important special case, the factor GARCH model, are more subtle than in the previous two subsections. In the one factor case, the factor GARCH model as defined by Engle (1987) and Engle, Ng and Rothschild (1990) imposes

$$\Sigma_t = \Psi - \varphi \lambda' \Psi \varphi' \lambda + \varphi \text{Var}_{t-1}(\lambda' y_t) \varphi', \quad (25)$$

and

$$\text{Var}_{t-1}(\lambda' y_t) = \lambda' \Psi \lambda + \tilde{\beta} \text{Var}_{t-2}(\lambda' y_{t-1}) + \tilde{\alpha} (\lambda' y_{t-1})^2, \quad (26)$$

where $\varphi' = (\varphi_1, \varphi_2)$ and $\lambda' = (\lambda_1, \lambda_2)$ are two dimensional vectors which are normalized such that $\varphi' \lambda = 1$. It is not difficult to see that this model is a special case of the bivariate GARCH model in (16) with

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} = \tilde{\alpha} \begin{pmatrix} \varphi_1^2 \lambda_1^2 & 2\varphi_1^2 \lambda_1 \lambda_2 & \varphi_1^2 \lambda_2^2 \\ \varphi_1 \varphi_2 \lambda_1^2 & 2\varphi_1 \varphi_2 \lambda_1 \lambda_2 & \varphi_1 \varphi_2 \lambda_2^2 \\ \varphi_2^2 \lambda_1^2 & 2\varphi_2^2 \lambda_1 \lambda_2 & \varphi_2^2 \lambda_2^2 \end{pmatrix}. \quad (27)$$

and $B = \tilde{\beta} \tilde{\alpha}^{-1} A$. According to (26), the "factor representing portfolio" $\lambda' y_t$ is generated by a strong GARCH(1,1) model, with parameters $\lambda' \Psi \lambda$, $\tilde{\beta}$ and $\tilde{\alpha}$. The marginal process for any linear combination $\delta' y_t$ where y_t is generated by a factor GARCH model can be derived using the results in appendices A and B.

An alternative extension of standard (i.e. conditionally homoskedastic) factor analysis models to time-varying variances was introduced by Diebold and Nerlove (1989) and extended by King, Sentana and Wadhvani (1991). Their model is a standard latent factor model with unobservable common factors which are generated by strong GARCH processes,

$$y_t = \varphi f_t + w_t \quad (28)$$

$$\begin{bmatrix} f_t \\ w_t \end{bmatrix} = \begin{bmatrix} \omega_t & 0 \\ 0 & \Phi \end{bmatrix} \begin{bmatrix} f_t^* \\ w_t^* \end{bmatrix} \quad \text{with} \quad \begin{bmatrix} f_t^* \\ w_t^* \end{bmatrix} \sim \text{i.i.d. } N(0, I_3) \quad (29)$$

and

$$\omega_t = \theta + \tilde{\beta} \omega_{t-1} + \tilde{\alpha} f_{t-1}^2 \quad (30)$$

The conditionally heteroskedastic latent factor model in (28)-(30) is not quite the same as the factor GARCH in (15), (25) and (26). However, both models are very closely related. In Sentana (1993) it is shown that factor GARCH models can be written as conditionally heteroskedastic latent factor models in which the latent factors are weak GARCH processes. Moreover it is shown there that the conditionally heteroskedastic factor model can be written in the form of the factor GARCH model, provided the assumption in the factor GARCH model that the factor is generated by a strong GARCH model is replaced by the assumption that it is generated by a weak one.

In order to illustrate the results on marginalization and linear transformation in factor GARCH models, we computed the weak GARCH parameters for several linear combinations $\delta_1 y_{1t} + \delta_2 y_{2t}$ for a number of values of the parameters in the factor GARCH model. First of all we normalized in such a way that $E f_t^2 = 1$ and $E y_{it}^2 = 1$ ($i=1,2$). It can be checked that given these normalizations all parameters in (25) and (26), and thereby the weak GARCH parameters, are fully determined by $\tilde{\alpha}$ and $\tilde{\beta}$ in (26) and the signal to noise ratios $R_i = \text{Var}(\varphi_i f_t) / \text{Var}(y_{it}) = \varphi_i^2$ in (28). The results are presented in table 3. For two specifications of the conditional heteroskedasticity in the factor and two pairs of signal to noise ratios, the weak GARCH parameters for the marginal processes for y_{1t} and y_{2t} as well as for linear combinations are presented. We have already noted that $\lambda' y_t$ will be strong GARCH with parameters $\tilde{\beta}$ and $\tilde{\alpha}$. The value of λ_2/λ_1 which corresponds to $R_1 = .75$ and $R_2 = .70$ is .913, while a value of .236 is implied by $R_1 = .95$ and $R_2 = .10$. The results in table 3 show that when the weight on y_{2t} is increased keeping the weight on y_{1t} fixed to unity, the weak GARCH parameter

Table 3. Weak GARCH parameters α and β for $\delta_1 y_{1t} + \delta_2 y_{2t}$ in (1)-(3) if the data are generated by the factor GARCH model in (15), (25) and (26).

GARCH param. of factor		signal to noise ratios		weight in linear comb.		weak GARCH parameters	
$\tilde{\beta}$	$\tilde{\alpha}$	R_1	R_2	δ_1	δ_2	β	α
.80	.10	.75	.70	1	0	.834	.067
				1	.5	.804	.096
				1	.913	.800	.100
				1	1	.800	.100
				.5	1	.807	.093
				0	1	.841	.059
.10	.40	.75	.70	1	0	.216	.284
				1	.5	.113	.387
				1	.913	.100	.400
				1	1	.100	.400
				.5	1	.123	.377
				0	1	.240	.260
.80	.10	.95	.10	1	0	.807	.093
				1	.236	.800	.100
				1	.5	.806	.094
				1	1	.833	.067
				.5	1	.868	.032
				0	1	.898	.002
.10	.40	.95	.10	1	0	.122	.378
				1	.236	.100	.400
				1	.5	.121	.379
				1	1	.211	.289
				.5	1	.349	.152
				0	1	.491	.009

α will increase upto the value $\bar{\alpha}$, which is reached when the weight on y_{2t} equals λ_2/λ_1 . Further increases in the weight on y_{2t} appear to yield smaller values for α . This property can in principle be used in finding factor representing portfolios using univariate analysis only. Note that e.g. in Engle, Ng and Rothschild (1990) the relative weights in the factor representing portfolios are assumed to be known a priori.

5. Probability limit of the standard QMLE estimator

In the previous sections we have described how weak GARCH parameters can be obtained for sums of independent strong GARCH processes as well as for marginal processes implied by a number of multivariate GARCH processes. The weak GARCH parameters can of course easily be estimated consistently from the autocorrelations of the squared observations. The estimators obtained in this way are probably not very efficient. Moreover many empirical studies in the literature are based on the use of a quasi maximum likelihood estimator (QMLE) under the assumption of a strong GARCH process and conditional normality. For both reasons we consider the properties of this QMLE estimator in this section. In Drost and Nijman (1992) it was noted that if data were generated by strong GARCH processes at one frequency and QMLE estimates were computed at lower frequencies, the parameter estimates tended to be very close to the weak GARCH parameters. In tables 4 and 5 we report results on a similar simulation experiment where we concentrate on aggregation and marginalization. Note that Weiss (1986) has shown that the QMLE estimator is consistent under weaker conditions than the validity of conditionally normal strong GARCH assumption, which are referred to as the semi-strong GARCH conditions in Drost and Nijman (1992). The properties of the QMLE estimator under weak GARCH have not yet been established, so that we have to rely on Monte Carlo results.

In the first row of table 4 e.g. we consider QMLE estimation of the sum of independent conditionally normal GARCH processes with variance parameters $\beta_1 = 0.9$ and $\alpha_1 = 0.05$ and equal (unconditional) variances. In the last two columns of the table we present the averages over 20 samples of the QMLE estimates obtained from 80,000 generated data. Given the very low

Table 4: Weak GARCH parameters and estimated plim of standard QMLE estimator for $y_t = y_{1t} + y_{2t}$ if y_{it} ($i=1,2$) is generated by (4)-(5)

β_1	α_1	κ_1	β_2	α_2	κ_2	σ_1^2/σ_2^2	β	α	plim	plim
									$\hat{\beta}$	$\hat{\alpha}$
0.90	0.05	3.00	0.90	0.05	3.00	1.00	0.920	0.030	0.920	0.029
0.90	0.05	3.00	0.90	0.05	3.00	4.00	0.912	0.038	0.910	0.037
0.90	0.05	9.00	0.90	0.05	3.00	1.00	0.910	0.040	0.916	0.036
0.90	0.05	9.00	0.90	0.05	3.00	4.00	0.903	0.047	0.909	0.041
0.90	0.05	9.00	0.90	0.05	9.00	1.00	0.907	0.043	0.909	0.040
0.50	0.35	3.00	0.80	0.05	3.00	1.00	0.569	0.281	0.690	0.125
0.50	0.35	3.00	0.80	0.05	3.00	4.00	0.516	0.334	0.581	0.234
0.50	0.35	3.00	0.80	0.05	3.00	0.25	0.705	0.145	0.775	0.062
0.50	0.35	3.00	0	0	3.00	1.00	0.571	0.274	0.685	0.112
0.50	0.35	3.00	0	0	3.00	4.00	0.517	0.333	0.582	0.232
0.50	0.35	3.00	0	0	3.00	0.25	0.721	0.129	0.783	0.038

Table 5: Weak GARCH parameters and estimated plim of standard QMLE estimator for $\delta_1 y_{1t} + \delta_2 y_{2t}$ if data are generated by the factor GARCH model in (17), (27) and (28)

GARCH param. of factor		signal to noise ratios		weight in linear comb.		weak GARCH parameters		plim	plim
$\tilde{\beta}$	$\tilde{\alpha}$	R_1	R_2	δ_1	δ_2	β	α	$\hat{\beta}$	$\hat{\alpha}$
.80	.10	.75	.70	1	0	0.834	0.067	0.820	0.075
				1	.5	0.804	0.096	0.803	0.096
				1	.913	0.800	0.100	0.800	0.100
				1	1	0.800	0.100	0.799	0.100
				.5	1	0.807	0.093	0.805	0.094
				0	1	0.841	0.059	0.828	0.070
.80	.10	.95	.10	1	0	0.807	0.093	0.808	0.092
				1	.236	0.800	0.100	0.804	0.098
				1	.5	0.806	0.094	0.807	0.092
				1	1	0.833	0.067	0.829	0.067
				.5	1	0.868	0.032	0.869	0.031
				0	1	0.898	0.002	0.807	0.003

variability of the estimates over the 20 samples these must be very precise estimates of the true probability limit of the QMLE estimators. The estimated probability limit is 0.920 for $\hat{\beta}$ and 0.029 for $\hat{\alpha}$, which is very close to the weak GARCH parameters that are presented in table 1 and in columns eight and nine of table 5. The same holds true for the cases presented in lines 2 to 5 in the table. These results are all in line with the evidence on temporal aggregation in Drost and Nijman (1992). Table 4 also contains a number of cases however where the estimated probability limit of the QMLE estimators differs substantially from the weak GARCH parameters. The probability limit of the QMLE estimate of the "persistence" parameter $\hat{\alpha} + \hat{\beta}$ is roughly in line with the value suggested by the weak GARCH parameters, although the underestimation is clearly significant. The plim of the estimated individual coefficients differs considerably from the weak GARCH parameters. Of course the question why the QMLE estimators is approximately consistent in some cases and clearly inconsistent in others is an important topic for future research, that is however outside the scope of this paper.

In table 5 results similar to those in table 4 are reported for the factor GARCH model. The estimated probability limits are, for the cases considered, usually close to the weak GARCH parameters. In particular the fact that α is largest at the linear combination which yields a strong GARCH process is reflected in the simulated results. The results in the last line of the table are probably caused by the fact that β is hardly identified if α is very close to zero.

6. Conclusions

In this paper we have discussed contemporaneous aggregation of independent univariate GARCH processes as well as marginalization and contemporaneous aggregation in more general multivariate GARCH processes. We have shown that the class of strong GARCH processes, which is typically assumed in applied work, is not closed under these transformations. We have also shown that the weak GARCH conditions are satisfied in all these cases and have discussed the relation between the various models and their parameter values. The results explain why GARCH will be found e.g. when analyzing exchange rate returns which are the sum of two underlying conditionally heteroskedastic exchange rate returns or when analyzing the return on a portfolio of stocks which are conditionally heteroskedastic.

Appendix A The impact of linear transformations on the parameters in the bivariate GARCH model

In order to derive the impact of linear transformation on the parameters in the bivariate GARCH model we use the duplication matrix D defined by $\text{vec}(\Sigma_t) = D \text{vech}(\Sigma_t)$. It is straightforward to check that $\text{vech}(\Sigma_t) = D^+ \text{vec}(\Sigma_t)$. Equation (17) can now be rewritten as

$$\text{vec}(\Sigma_t) = D\psi + D B D^+ \text{vec}(\Sigma_{t-1}) + D A D^+ \text{vec}(y_{t-1}y'_{t-1}) \quad (\text{A.1})$$

If $x_t = P y_t$ where P is a non-singular matrix, and $\Sigma_t^* = \text{Var}_{t-1}(x_t)$ the properties of the vec operator imply

$$\text{vec}(\Sigma_t^*) = \text{vec}(P\Sigma_t P') = (P\theta P) \text{vec}(\Sigma_t)$$

and

$$\begin{aligned} \text{vec}(\Sigma_t^*) &= (P\theta P)D\psi + (P\theta P)DBD^+ \text{vec}(\Sigma_{t-1}) + (P\theta P) DAD^+ \text{vec}(y_{t-1}y'_{t-1}) \\ &= (P\theta P)D\psi + (P\theta P)DBD^+(P^{-1}\theta P^{-1})\text{vec}(\Sigma_{t-1}^*) \\ &\quad + (P\theta P) DAD^+ (P^{-1}\theta P^{-1}) \text{vec}(x_{t-1}x'_{t-1}). \end{aligned} \quad (\text{A.2})$$

Premultiplying (A.2) with the Moore-Penrose inverse of D yields the required result.

Appendix B Determination of the variance of η_t in (19).

In this appendix we will derive the variance-covariance matrix of η_t defined in (19) which is required to determine the weak GARCH parameters in the marginal process for y_{1t} in (21). Note first of all that $\text{Var}(\eta_t) = E \eta_t \eta_t'$ as $E \eta_t = 0$. Moreover as $E_{t-1} z_t = s_t$ the law of iterated expectations yields

$$\text{Var}(\eta_t) = E \eta_t \eta_t' = E z_t z_t' - E s_t s_t'. \quad (\text{A.3})$$

The typical element of $\text{Var}(\eta_t)$ therefore is $E(E_{t-1}(y_{it}y_{jt}y_{kt}y_{lt}) - \sigma_{ijkt}) - \sigma_{ijkt}$. But because the conditional distribution is assumed to be normal in sections 3 and 4 (see (15)) we can use a well known result on fourth moments of a bivariate normal variable $E_{t-1}(y_{it}y_{jt}y_{kt}y_{lt}) = \sigma_{ijkt} + \sigma_{ikt}\sigma_{jlt} + \sigma_{ilt}\sigma_{jkt}$ to show that

$$E \eta_{ijkt} \eta_{ijkl} = E (\sigma_{ikt}\sigma_{jlt} + \sigma_{ilt}\sigma_{jkt}). \quad (\text{A.4})$$

The remaining problem is to determine $h_{ijkl} = E \sigma_{ijkt}$. We solve this problem by constructing a six dimensional linear system with six unknowns, $h = E \text{vech}(s_t s_t')$. From (17) we know that

$$\begin{aligned} E s_t s_t' = & \psi \psi' + \psi E z_{t-1}' A' + A E z_{t-1} \psi' + \psi E s_{t-1}' B' + B E s_{t-1} \psi' + (\text{A.6}) \\ & + A E z_{t-1}' z_{t-1}' A' + B E s_{t-1}' s_{t-1}' B' + A E z_{t-1}' s_{t-1}' B + B E s_{t-1}' z_{t-1}' A'. \end{aligned}$$

Now note first of all that $E z_{t-1} = E s_{t-1} = [I - (A+B)]^{-1} \psi$. Moreover, by the law of iterated expectations, $E s_{t-1}' z_{t-1}' = E s_{t-1}' s_{t-1}'$. As finally $E z_{t-1}' z_{t-1}'$ is also a linear function of the six unknowns in h , as we have seen above, (A.9) yields a six dimensional linear system for $\text{vech}(E s_t s_t')$ in the six unknown elements of h .

References

- Attanasio, O. and Edey, M. (1987), "Non-constant variances and foreign exchange risk: an empirical study", London School of Economics, CLE DP 285.
- Baba, Y., Engle, R.F., Kraft, D.F., and K.F. Kroner (1989), "Multivariate simultaneous generalized ARCH", Department of Economics, UCSD, DP 89-57.
- Bollerslev, T. (1986), "Generalized autoregressive conditional heteroskedasticity", Journal of Econometrics, 31, 307-327.
- Bollerslev, T., Engle, R.F., and J.M. Wooldridge (1988), "A capital asset pricing model with time varying covariances", Journal of Political Economy, 96, 116-131.
- Bollerslev, T., Chou, R.Y., and K.F. Kroner (1992), "ARCH modeling in finance: a review of the theory and empirical evidence", Journal of Econometrics, 52, 5-59.
- Diebold, F.X. and M. Nerlove (1989), "The dynamics of exchange rate volatility: a multivariate latent factor ARCH model", Journal of Applied Econometrics, 4, 1-22.
- Drost, F.C. and Th. E. Nijman (1992), "Temporal aggregation of GARCH processes", Center discussion paper 9240 (revised version of DP 9066), Tilburg University, forthcoming in Econometrica.
- Engle, R.F. (1982), "Autoregressive conditional heteroscedasticity with estimates of the variance of United Kingdom inflation", Econometrica, 50, 987-1007.
- Engle, R.F. (1987), "Multivariate ARCH with factor structures: cointegration in variance", Discussion paper University of California, San Diego.

- Engle, R.F., V. Ng and M. Rothschild (1990), "Asset pricing with a factor ARCH covariance structure: empirical estimates of treasury bills", Journal of Econometrics, 45, 213-238.
- King M., E. Sentana and S. Wadhvani (1991), "Volatility and links between national stock markets", mimeographed, Financial Markets Group, London School of Economics.
- Kroner, K. (1987), "Estimating and testing factor GARCH models", mimeographed, UCSD.
- Lütkepohl, H. (1984), "Linear transformations of vector ARMA processes", Journal of Econometrics, 26, 283-293.
- Sentana, E. (1992), "Factor representing portfolios in large asset markets", Financial Markets Group, London School of Economics, DP 135.
- Sentana, E. (1993), "The relation between conditionally heteroskedastic latent factor models and factor GARCH models", in progress.
- Weiss, A.A. (1986), "Asymptotic theory for ARCH models: estimation and testing", Econometric Theory, 2, 107-131.
- Zellner, A. and F.C. Palm (1974), "Time series analysis and simultaneous equations econometric models", Journal of Econometrics, 2, 17-54.

Discussion Paper Series, CentER, Tilburg University, The Netherlands:

(For previous papers please consult previous discussion papers.)

No.	Author(s)	Title
9146	Th. van de Klundert and S. Smulders	Reconstructing Growth Theory: A Survey
9147	J. Greenberg	On the Sensitivity of Von Neuman and Morgenstern Abstract Stable Sets: The Stable and the Individual Stable Bargaining Set
9148	S. van Wijnbergen	Trade Reform, Policy Uncertainty and the Current Account: A Non-Expected Utility Approach
9149	S. van Wijnbergen	Intertemporal Speculation, Shortages and the Political Economy of Price Reform
9150	G. Koop and M.F.J. Steel	A Decision Theoretic Analysis of the Unit Root Hypothesis Using Mixtures of Elliptical Models
9151	A.P. Barten	Consumer Allocation Models: Choice of Functional Form
9152	R.T. Baillie, T. Bollerslev and M.R. Redfearn	Bear Squeezes, Volatility Spillovers and Speculative Attacks in the Hyperinflation 1920s Foreign Exchange
9153	M.F.J. Steel	Bayesian Inference in Time Series
9154	A.K. Bera and S. Lee	Information Matrix Test, Parameter Heterogeneity and ARCH: A Synthesis
9155	F. de Jong	A Univariate Analysis of EMS Exchange Rates Using a Target
9156	B. le Blanc	Economies in Transition
9157	A.J.J. Talman	Intersection Theorems on the Unit Simplex and the Simplotope
9158	H. Bester	A Model of Price Advertising and Sales
9159	A. Özcam, G. Judge, A. Bera and T. Yancey	The Risk Properties of a Pre-Test Estimator for Zellner's Seemingly Unrelated Regression Model
9160	R.M.W.J. Beetsma	Bands and Statistical Properties of EMS Exchange Rates: A Monte Carlo Investigation of Three Target Zone Models Zone Model
9161	A.M. Lejour and H.A.A. Verbon	Centralized and Decentralized Decision Making on Social Insurance in an Integrated Market Multilateral Institutions
9162	S. Bhattacharya	Sovereign Debt, Creditor-Country Governments, and

No.	Author(s)	Title
9163	H. Bester, A. de Palma, W. Leininger, E.-L. von Thadden and J. Thomas	The Missing Equilibria in Hotelling's Location Game
9164	J. Greenberg	The Stable Value
9165	Q.H. Vuong and W. Wang	Selecting Estimated Models Using Chi-Square Statistics
9166	D.O. Stahl II	Evolution of Smart _n Players
9167	D.O. Stahl II	Strategic Advertising and Pricing with Sequential Buyer Search
9168	T.E. Nijman and F.C. Palm	Recent Developments in Modeling Volatility in Financial Data
9169	G. Asheim	Individual and Collective Time Consistency
9170	H. Carlsson and E. van Damme	Equilibrium Selection in Stag Hunt Games
9201	M. Verbeek and Th. Nijman	Minimum MSE Estimation of a Regression Model with Fixed Effects from a Series of Cross Sections
9202	E. Bomhoff	Monetary Policy and Inflation
9203	J. Quiggin and P. Wakker	The Axiomatic Basis of Anticipated Utility; A Clarification
9204	Th. van de Klundert and S. Smulders	Strategies for Growth in a Macroeconomic Setting
9205	E. Siandra	Money and Specialization in Production
9206	W. Härdle	Applied Nonparametric Models
9207	M. Verbeek and Th. Nijman	Incomplete Panels and Selection Bias: A Survey
9208	W. Härdle and A.B. Tsybakov	How Sensitive Are Average Derivatives?
9209	S. Albæk and P.B. Overgaard	Upstream Pricing and Advertising Signal Downstream Demand
9210	M. Cripps and J. Thomas	Reputation and Commitment in Two-Person Repeated Games
9211	S. Albæk	Endogenous Timing in a Game with Incomplete Information
9212	T.J.A. Storcken and P.H.M. Ruys	Extensions of Choice Behaviour
9213	R.M.W.J. Beetsma and F. van der Ploeg	Exchange Rate Bands and Optimal Monetary Accommodation under a Dirty Float

No.	Author(s)	Title
9214	A. van Soest	Discrete Choice Models of Family Labour Supply
9215	W. Güth and K. Ritzberger	On Durable Goods Monopolies and the (Anti-) Coase-Conjecture
9216	A. Simonovits	Indexation of Pensions in Hungary: A Simple Cohort Model
9217	J.-L. Ferreira, I. Gilboa and M. Maschler	Credible Equilibria in Games with Utilities Changing during the Play
9218	P. Borm, H. Keiding, R. Mclean, S. Oortwijn and S. Tijs	The Compromise Value for NTU-Games
9219	J.L. Horowitz and W. Härdle	Testing a Parametric Model against a Semiparametric Alternative
9220	A.L. Bovenberg	Investment-Promoting Policies in Open Economies: The Importance of Intergenerational and International Distributional Effects
9221	S. Smulders and Th. van de Klundert	Monopolistic Competition, Product Variety and Growth: Chamberlin vs. Schumpeter
9222	H. Bester and E. Petrakis	Price Competition and Advertising in Oligopoly
9223	A. van den Nouweland, M. Maschler and S. Tijs	Monotonic Games are Spanning Network Games
9224	H. Suehiro	A "Mistaken Theories" Refinement
9225	H. Suehiro	Robust Selection of Equilibria
9226	D. Friedman	Economically Applicable Evolutionary Games
9227	E. Bomhoff	Four Econometric Fashions and the Kalman Filter Alternative - A Simulation Study
9228	P. Borm, G.-J. Otten and H. Peters	Core Implementation in Modified Strong and Coalition Proof Nash Equilibria
9229	H.G. Bloemen and A. Kapteyn	The Joint Estimation of a Non-Linear Labour Supply Function and a Wage Equation Using Simulated Response Probabilities
9230	R. Beetsma and F. van der Ploeg	Does Inequality Cause Inflation? - The Political Economy of Inflation, Taxation and Government Debt
9231	G. Almekinders and S. Eijffinger	Daily Bundesbank and Federal Reserve Interventions - Do they Affect the Level and Unexpected Volatility of the DM/\$-Rate?
9232	F. Vella and M. Verbeek	Estimating the Impact of Endogenous Union Choice on Wages Using Panel Data

No.	Author(s)	Title
9233	P. de Bijl and S. Goyal	Technological Change in Markets with Network Externalities
9234	J. Angrist and G. Imbens	Average Causal Response with Variable Treatment Intensity
9235	L. Meijdam, M. van de Ven and H. Verbon	Strategic Decision Making and the Dynamics of Government Debt
9236	H. Houba and A. de Zeeuw	Strategic Bargaining for the Control of a Dynamic System in State-Space Form
9237	A. Cameron and P. Trivedi	Tests of Independence in Parametric Models: With Applications and Illustrations
9238	J.-S. Pischke	Individual Income, Incomplete Information, and Aggregate Consumption
9239	H. Bloemen	A Model of Labour Supply with Job Offer Restrictions
9240	F. Drost and Th. Nijman	Temporal Aggregation of GARCH Processes
9241	R. Gilles, P. Ruys and J. Shou	Coalition Formation in Large Network Economies
9242	P. Kort	The Effects of Marketable Pollution Permits on the Firm's Optimal Investment Policies
9243	A.L. Bovenberg and F. van der Ploeg	Environmental Policy, Public Finance and the Labour Market in a Second-Best World
9244	W.G. Gale and J.K. Scholz	IRAs and Household Saving
9245	A. Bera and P. Ng	Robust Tests for Heteroskedasticity and Autocorrelation Using Score Function
9246	R.T. Baillie, C.F. Chung and M.A. Tieslau	The Long Memory and Variability of Inflation: A Reappraisal of the Friedman Hypothesis
9247	M.A. Tieslau, P. Schmidt and R.T. Baillie	A Generalized Method of Moments Estimator for Long-Memory Processes
9248	K. Wämeryd	Partisanship as Information
9249	H. Huizinga	The Welfare Effects of Individual Retirement Accounts
9250	H.G. Bloemen	Job Search Theory, Labour Supply and Unemployment Duration
9251	S. Eijffinger and E. Schaling	Central Bank Independence: Searching for the Philosophers' Stone
9252	A.L. Bovenberg and R.A. de Mooij	Environmental Taxation and Labor-Market Distortions

No.	Author(s)	Title
9253	A. Lusardi	Permanent Income, Current Income and Consumption: Evidence from Panel Data
9254	R. Beetsma	Imperfect Credibility of the Band and Risk Premia in the European Monetary System
9301	N. Kahana and S. Nitzan	Credibility and Duration of Political Contests and the Extent of Rent Dissipation
9302	W. Güth and S. Nitzan	Are Moral Objections to Free Riding Evolutionarily Stable?
9303	D. Karotkin and S. Nitzan	Some Peculiarities of Group Decision Making in Teams
9304	A. Lusardi	Euler Equations in Micro Data: Merging Data from Two Samples
9305	W. Güth	A Simple Justification of Quantity Competition and the Cournot-Oligopoly Solution
9306	B. Peleg and S. Tijs	The Consistency Principle For Games in Strategic Form
9307	G. Imbens and T. Lancaster	Case Control Studies with Contaminated Controls
9308	T. Ellingsen and K. Wärneryd	Foreign Direct Investment and the Political Economy of Protection
9309	H. Bester	Price Commitment in Search Markets
9310	T. Callan and A. van Soest	Female Labour Supply in Farm Households: Farm and Off-Farm Participation
9311	M. Pradhan and A. van Soest	Formal and Informal Sector Employment in Urban Areas of Bolivia
9312	Th. Nijman and E. Sentana	Marginalization and Contemporaneous Aggregation in Multivariate GARCH Processes

P.O. BOX 90153, 5000 LE TILBURG, THE NETHERLANDS

Bibliotheek K. U. Brabant



17 000 01117401 9