

Endogeneity and Heterogeneity in LDV Panel Data Models*

Jacob N. Arendt

Institute of Local Government Studies, Nyropsgade 37, 1602 Copenhagen V, Denmark

November 2001

Abstract

We extend three existing cross-sectional limited dependent variable (LDV) estimators, that allow for endogenous regressors, to a panel data model. We focus on estimation of effects of time-invariant endogenous regressors, since to our knowledge, besides joint maximum likelihood, no other alternative consistent parametric estimators than the ones suggested here exist. We compare their small sample performance of estimates of marginal effects to i.i.d. LDV estimators as well as to linear estimators by means of Monte Carlo Studies. Some notable differences in the performance of the LDV estimators appear. One estimator, the 2SIV, performs reasonably well in terms of bias, even with weak instruments. Another type, the AGLS estimators, have a large small sample bias when no endogeneity is present. The 2SCML estimators seem to perform reasonable in most scenarios even under some types of misspecification. In addition, 2SLS performed relatively well, but had a substantial MSE with weak instruments and substantial bias in misspecified scenarios. Although potentially important because of heterogeneity bias, our extension of LDV models to the panel case did not give improvements in small sample performance over the cross-sectional estimators.

JEL: C33; C34; C35

Keywords: Two-Step Estimation; Panel Data; Endogenous Regressor; Time-Invariant Regressor; Linear Approximation;

*e-mail: jna@akf.dk. The first draft was written while the author was at Institute of Economics, University of Copenhagen. I appreciate comments from participants at the Ph.D.-workshop in Rungstedgaard, 2001. I also thank Martin Browning for suggesting me to do this study, and Bo Honoré and Dan-Olof Rooth for useful suggestions.

1. INTRODUCTION

The focus of this paper is estimation of limited dependent variable (LDV) panel data models with unobserved individual specific effects and endogenous explanatory variables. The paper contributes to this literature by extending three existing cross-sectional limited dependent variable (LDV) estimators, that allow for endogenous regressors, to a panel data model. Focus is on estimation of effects of time-invariant endogenous regressors, since, to our knowledge, besides doing joint maximum likelihood estimation (MLE), no alternative consistent parametric estimators than the ones suggested here exist. We conduct a Monte Carlo study in a probit model where we compare the performance of these estimators to a host of alternative estimators, focusing on estimation of the marginal effect on the probability of observing a one, since this statistic is often of largest interest from an applied point of view.

In empirical studies the specification of a model where the outcome of interest depends on unobserved time invariant individual specific effects often comes directly from economic theory, e.g. as the marginal utility of wealth in life-cycle models in Heckman and Macurdy (1980). On a less structural level, the allowance for individual effects is a parsimonious way of specifying individual heterogeneity, when aiming at estimating the causal effect of a regressor on a given outcome, allowing for common unobserved determinants of both. The estimators considered in this paper reflect different aspects of empirical interests in this type of modelling.

There is a large literature that struggles with the difficulties in estimating non-linear models with unobserved individual specific effects. When the correlation between individual specific effects and regressors is unrestricted, most existing estimators rely on time variation in the regressor of interest to remove the impact from the unobserved individual effect. Examples include the linear within (fixed effect) estimator, the conditional logit estimator (e.g. Andersen (1970)), the minimum distance probit estimator suggested by Chamberlain (1980), the maximum score estimator developed by Manski (1987), and the semiparametric estimators suggested by Honore (1992), Kyriazidou (1997) and Lee (1999). These estimators can not identify the effects of time-invariant variables. Therefore, the effects of variables (on relevant outcomes) such as education, gender, race, birthweight, height and any variable that is determined at a given point in time can not be identified using these methods. One could also imagine certain variables that vary over time for some reason are time-invariant in specific samples, e.g. number of children for elderly, minimum wages in given regions in given periods etc. In a linear regression model, this can be solved using

the approach due to Hausman and Taylor (1981). We show that in LDV models, when the endogenous time-invariant variable is continuous, the problem of estimating their effects can be solved by extending different cross-sectional LDV estimators to the panel data case. The cross-sectional estimators are the two-stage IV (2SIV) method suggested by Lee (1981), the two-stage conditional maximum likelihood (2SCML) estimator suggested by Rivers and Vuong (1988) and the two-stage estimator suggested by Amemiya (1978) (AGLS) and elaborated by Newey (1987). Our extensions essentially boil down to doing a random effects estimation in the second stage rather than a cross-sectional LDV estimation. This is potentially very important in LDV models, since heteroscedasticity bias may be severe (see e.g. Horowitz (1993)) and is transmitted to all estimated parameters.

In contrast to this literature, it has been argued that if the aim is to estimate causal marginal effects on the outcome of interest, rather than structural parameters, the focus on structural LDV models unnecessarily complicates estimation. In relation to this, we ask a simple question. Given that with a LDV model at hand, many applied researchers focus on the marginal effects, e.g. on the probability of observing a one in a probit model, rather than on the probit parameter per se, how well does simpler linear models approximate this marginal effect? Linear models might be superior as they rely less on distributional assumptions. The question is related to issues raised by Angrist (2001) and the accompanying discussion in a recent issue of *Journal of Business and Economic Statistics* (JBES). The answer to the question may support applied researchers, especially when doing exploratory studies, but, from how we read the discussion in JBES, should not guide the choice of estimator and model. The latter choice depends more crucially on what the analysis is developed to answer, whether there is a need for extrapolation and in the end beliefs about credibility of extrapolation from structural models as opposed to e.g. local average treatment effects based on a natural experiment. To look at how the suggested estimators perform in small samples, and to try to answer the question raised above, we conduct a Monte Carlo study.

Although we present the panel models in a general LDV framework, we confine attention to a probit model in the Monte Carlo studies, where we focus on the estimated marginal effects, for reasons mentioned above. There has been some prior Monte Carlo studies in the cross-sectional model on the structural parameters, but no studies on marginal effects. Using a probit model with individual unobserved effects as data generating process, we compare the properties of the cross-section and the panel versions of the 2SIV, the 2SCML and the AGLS estimators with the simple probit, the random effects (RE) probit and the following linear estimators: OLS, 2SLS, Fixed

Effects, and Hausman and Taylors (1981) 2SLS estimator. The Monte Carlo results suggest the expected that with weak instruments, most two-stage estimators perform bad and the worse the endogeneity problem, the larger advantage of two-stage estimators. The results suggest that the two-stage linear estimators produce the worst results in case of weak instruments. Regarding the two-stage LDV models, the AGLS estimators have a large small sample bias when no endogeneity is present, and in general the 2SCML and 2SIV estimators seem to be preferable for small samples. Linear estimators do not perform better than LDV estimators and often perform worse, particular for the types of misspecification considered here. The results suggest that, at least with respect to the designs and sample sizes considered in this paper, our extentions to panel models do not improve small sample properties.

The paper is outlined as follows. In the next section we specify the panel LDV model and present the panel estimators. In section 3 we write up alternative linear model estimators, in section 4 the Monte Carlo designs are described, in section 5 the results are presented and section 6 concludes.

2. THE MODEL

The LDV model we consider is given by the equations:

$$\begin{aligned}
 (1), (2) \quad & y_{it}^* = X_{1it}\beta + \gamma z_i + \varepsilon_{it} \\
 & y_{it} = \tau(y_{it}^*, \psi) \\
 & i = 1, \dots, n \quad t=1, \dots, T
 \end{aligned}$$

where X_{1it} are regressors of individual i in period t , y_{it}^* is an unobserved latent variable, z_i is a continuous time-invariant regressor, and ε_{it} is an error term. We observe y_{it} according to some known function of y_{it}^* , τ , which may depend on nuisance parameters, ψ . This general model includes e.g. censored regression models, some duration models, some transformation models and quantal response models.

The focus of our attention is how to estimate γ when z_i and ε_{it} are correlated. We focus on the reduced form model where z_i is determined as:

$$(3) \quad z_i = X_i \Pi + V_i$$

where X may include X_1 and errors and regressors fulfil:

$$(4) \quad \varepsilon_{it} | X_i, X_{1it} \sim N.I.D(0, \Sigma), \quad E(V_i | X) = 0$$

that is, the latent variable errors are jointly normal distributed for a given individual and errors in (1) and (3) have mean zero conditional on the covariates, X . The covariance between V_i and ε_{it} is thus left unrestricted and z_i is endogenous if this covariance is non-zero¹. With linear index specifications of y^* and z_i , we need at least one variable in X that is excluded from the y -equation to identify γ . Let X_2 be such excluded variables. Given the panel structure of the model we can specify an error component structure:

$$(5) \quad \varepsilon_{it} = \alpha_i + v_{it}$$

where α_i is a time-invariant individual effect and v_{it} is a random error. We restrict attention to the case where endogeneity of z_i arises through correlation between V_i and α_i . A variety of estimators have been suggested in the mentioned papers considering the cross-sectional model. We focus on three of these, which are all based on a first stage linear estimation of (3). The estimators can be applied in our panel model, basically by applying a random effects estimator in a second stage estimation, as we will show now.

A two-stage procedure suggested by Lee, summarized in Lee (1981), replaces z_i by its predicted value from the first stage linear estimation. We can do the same in the panel model, yielding the following second stage equation:

$$(6) \quad \begin{aligned} y_{it}^* &= X_{1it}\beta + \gamma(X_i\hat{\Pi}) + \alpha_i + \tilde{v}_{1it} \\ \tilde{v}_{1it} &= \gamma(z_i - \hat{z}_i) + v_{it} \end{aligned}$$

where a ‘^’ denotes estimates from the first stage. The difference to the cross-sectional model is that after inserting fitted values to take care of endogeneity of z_i , the individual specific effect is

¹ Removing time subscripts, this is the cross-sectional model considered by Newey (1987), which is closely related to models considered in Lee (1981) and Blundell and Smith (1989). A more general system is considered by Blundell and Smith (1994) and, for the probit model, by Heckman (1978) and Amemiya (1978), where feedback from the latent variable to the endogenous regressor is allowed (structural shifts). The probit version of the system consisting of (1), (2) and (3) was considered by Rivers and Vuong (1988) and the special case of the Tobit model was considered by Nelson and Olsen (1978), Amemiya (1979) and Blundell and Smith (1986). None of these consider extensions to panel models.

still present but is now uncorrelated with regressors, i.e. it is a random effect. The cross-sectional estimator is often referred to as the two-stage instrumental variable probit (2SIV). We therefore refer to this as the 2SIVR estimator, where R stands for random effects.

Rivers and Vuong (1988) suggested a control function estimator, using an estimate of $E(\varepsilon_{it}|V_i)$ to correct for endogeneity. Their estimator is called the two-stage conditional maximum likelihood estimator (2SCML). We assume, as they do, that the conditional mean is linear in V , implying the following²:

$$(7) \quad \begin{aligned} E(\varepsilon_{it} | V_i) &= E(\alpha_i | V_i) = \eta V_i, \quad \eta = \frac{\sigma_{\alpha V}}{\sigma_V^2} \\ \Rightarrow \alpha_i &= \eta V_i + w_i \end{aligned}$$

where η is the population regression parameter of regressing α_i on V_i . Therefore w_i is orthogonal to z_i by construction. Using (7) to replace α_i by V_i and w_i , and replacing V_i by its estimate from the first stage estimation, the equation for y^* becomes:

$$(8) \quad \begin{aligned} y_{it}^* &= X_{it}\beta + \gamma z_i + \eta \hat{V}_i + w_i + \tilde{v}_{2it} \\ \tilde{v}_{2it} &= \gamma(V_i - \hat{V}_i) + v_{it} \end{aligned}$$

Again, the presence of w_i makes this second stage a random effects model, hence we refer to this estimator as the 2SCMLR estimator³.

² Note that even though this can be justified by a joint normality assumption on (α_i, V_i) , this is not required. In general, the two-stage conditional maximum likelihood procedure is based on the conditional likelihood: $f(y_i|X_{1i}, z_i, \theta_1, \theta_2)f(z_i|X_i, \theta_2)$ and first maximizes $f(z_i|X_i, \theta_2)$ and then, given the estimate θ_2^* , maximizes $f(y_i|X_{1i}, z_i, \theta_1, \theta_2^*)$. Vuong (1984) shows that this procedure yields an asymptotic normal distributed estimate of θ_1 given standard regularity conditions. It is required that the distribution of z_i does not depend on θ_1 , that a consistent estimate of θ_2 can be obtained and that the conditional distribution of $y|z$ is correctly specified. In particular, when the first stage is a linear regression, the distribution of z can be left unspecified.

³ This is related to the panel model suggested by Vella and Verbeek (1999), section 3, but they focus on time-varying endogenous regressors without the linear index assumption. They explicitly

Finally, an estimator suggested by Newey (1987) replaces z_i in (8) by the reduced form equation (3). In the panel version the second stage equation then becomes:

$$(9) \quad \begin{aligned} y_{it}^* &= X_{1it}\alpha_1 + X_{2it}\alpha_2 + \lambda\hat{V}_i + w_i + \tilde{v}_{3it} \\ \tilde{v}_{3it} &= \lambda(V_i - \hat{V}_i) + v_{it} \end{aligned}$$

where $\lambda = \gamma + \eta$ and the α parameters are related to the structural parameters β and γ in the following way:

$$(10) \quad \alpha_1 = \beta + \gamma\Pi_1, \quad \alpha_2 = \gamma\Pi_2$$

with Π is splitted into Π_1 and Π_2 according to the division of X into X_1 and X_2 . Therefore, if there are more than one variable in X_2 , γ is overidentified and γ and β can be estimated as suggested by Amemiya (1978), using GLS on the latter equations, yielding:

$$(11) \quad \hat{\delta}_A = (\hat{D}'W\hat{D})^{-1}\hat{D}'W\hat{\alpha}$$

where $\delta = (\beta, \gamma)$, $D = [\Pi, J]$, with J defined as the matrix s.t. $X_1 = XJ$ and W is a weighting matrix⁴. Again, the difference to the cross-sectional estimator is to add random effects, w_i , to the second stage where α_1 and α_2 are estimated. We refer to this as Amemiyas GLS estimator with random effects (henceforth AGLSR). In the following, when referring to any of these estimators in plural we mean both the cross-sectional and the panel version, i.e. the 2SIV estimators refer to the 2SIVR and the 2SIV estimator.

From an applied point of view, we note that all the panel estimators require estimation of a random effects LDV model in the second stage. With normal distributed individual effects this can be done for instance by using Gaussian quadrature, see Butler and Moffitt (1982). Note also that the AGLS estimators require one additional estimation compared to the 2SIV and 2SCML estimators. An attractive feature of the 2SCML estimators is that they provide an estimate of η , which often is of theoretical interest, and the t-test that η is zero is an easy accessible test for exogeneity of z_i . We

mention that with their procedure they can not test for endogeneity of time-invariant variables like education in their empirical application.

⁴ In the just identified case, δ can be obtained as $D^{-1}\alpha$.

also note that our panel specification allows the use of instrumental variables (X_2 variables) generated from X_1 as suggested by Hausman and Taylor (1981)⁵.

Asymptotic normality holds for all the estimators, as seen e.g. by applying the proof in Newey (1987) or in Vuong (1984), replacing individual likelihood contributions with joint likelihoods over time for each individual. Newey shows that under joint normality of the error terms, when W is a consistent estimator of the asymptotic covariance matrix of the estimate of $(\alpha - D\delta)$, the AGLS estimator is asymptotically efficient in the class of GLS estimators which includes the 2SIV estimator and the 2SCML estimator. Rivers and Vuong (1988) note that in the just-identified case, under joint normality, 2SCML is identical to joint MLE, and thus efficient. This is also the case for our panel estimator.

The small sample behavior of the structural cross-sectional LDV estimates has been investigated in a few number of studies. Rivers and Vuong (1988) conduct a small Monte Carlo study based on 100 observations, where they compare the performance of 2SIV, 2SCML and G2SP cross-sectional estimates in the probit model, where G2SP stands for Generalized Two-stage Simultaneous Probit, which is Amemiyas estimator that inspired Newey (1987) to suggest the AGLS⁶. They conclude that the G2SP and the 2SIV never outperform the 2SCML estimator, and the 2SCML performs better in most simulations. Alvarez and Glasgow (2000) conduct Monte Carlo studies comparing simple probit with 2SIV and 2SCML estimates. They find that simple probit estimators may be substantially biased when endogeneity is present and ignored. Both two-stage estimators perform better and, in samples of size 10.000, the 2SCML performs much better than the 2SIV, but in samples of size 300, the reverse holds⁷. Evaluating changes in predicted probabilities, the 2SCML

⁵ I.e. using individual means over time and deviation from means as instruments. See also extensions by Breusch, Mizon and Schmidt (1989).

⁶ The difference between G2SP and AGLS is that G2SP does not include the residual from equation (3) as regressor, and thus is unconditional on V . This implies that it is not always at least as efficient than 2SCML as AGLS is, as shown by Rivers and Vuong (1988).

⁷ The latter seemingly contradicts the result found by Rivers and Vuong (1988). This must be due to a minor difference in the designs, Alvarez and Glasgow using two instruments independent of

and the 2SIV estimator perform equally well with a low bias compared to simple probits. Mroz and Guilkey (1992) conduct a Monte Carlo study, where they among others compare the 2SIV estimator with joint normal MLE⁸. The bias of the 2SIV estimator relative to the joint normal ML estimator is only eight percent with 1000 observations under normality, whereas the bias of the joint ML is more than twice the size of the bias of the 2SIV estimator when the error in the first stage is skewed. Finally, Bollen et al. (1995) cite the results of a large Monte Carlo Study, comparing simple probit, 2SCML, joint MLE and a GMM estimator. They stress the intuitively appealing result that the behavior of the 2SCML estimator depends crucially on the explanatory power in the first stage estimation. We note that for all three panel estimators, to obtain consistent standard errors corrections are needed to account for the first-stage estimation. Newey describes how to do this for the AGLS estimator, and a similar approach is feasible here. For the 2SCMLR estimator, the expressions in Vuong (1984) can be used, but the presence of random effects complicates the correct asymptotic variance considerably.

3. ALTERNATIVE LINEAR ESTIMATORS

Estimation of LDV models requires a lot of technical knowledge and, as they are not yet standard in statistical packages, some computational work. A relevant question is to what extent it pays off? In answering this question, one may like to know e.g. how well-behaved the LDV estimators are in finite samples, how robust they are to misspecification, and whether simpler alternatives exist. The latter has been pursued by Angrist (2001), where he argues that much of the difficulties arising in limited dependent variable models can be avoided, if the aim is to estimate causal marginal effects, rather than a specific structural form. Following this thought we line up alternative linear models that could be of interest in relation to our specific model. Before doing this, we stress that in many

the single exogenous regressor in the main equation, whereas Rivers and Vuong use two exogenous regressors and one or two instruments, all variables drawn from a joint normal distribution.

⁸ Mroz and Guilkey go a step further and compare the performance of the estimators when joint normality does not hold, using discrete factor approximation to estimate the joint distribution of error terms.

cases, the LDV parameters are the main aim. This is e.g. the case if only the sign of the effect is needed (then, in some models, the LDV parameter is sufficient), if a structural model is specified and the parameter has theoretical content or the estimate may be needed in other estimations.

A linear panel model with fixed effects of the LDV outcome is given by the following:

$$(12) \quad y_{it} = X_{it}\beta^l + \gamma^l z_i + \alpha_i^l + v_{it}^l$$

where the superscript l indicates that the parameters and errors are different in this model than in the LDV model. As is well-known, the parameters in this model describe the marginal effects on the mean of observed outcomes if covariates are conditionally mean independent of the error term. This marginal effect is often what the researcher is after in the first place rather than a structural LDV coefficient. Note, however, that consistency of these estimates requires correct model specification, i.e. linearity of effects of observed and unobserved variables, which is clearly unrealistic in many cases exactly because of the LDV nature of y . In some cases this may be a correct way to model y , e.g. when all regressors are dummy variables. It could be that even when this is not the case, linear estimates may provide a reasonable approximation to mean marginal effects or, as is common in LDV models, marginal effects evaluated at means of explanatory variables.

When z_i is correlated with individual effects α_i^l , 2SLS estimation, where z_i is instrumented, may give consistent estimates of unknown parameters in (12)⁹. As noted by Hausman (1978), 2SLS estimation is algebraically equivalent to OLS on either one of the following equations:

$$(13), (14) \quad \begin{aligned} y_{it} &= X_{it}\beta^l + \gamma^l z_i + \theta \widehat{V}_i + \xi_{1it} \\ y_{it} &= X_{it}\beta^l + \gamma^l \widehat{z}_i + \xi_{2it} \end{aligned}$$

⁹ In the case of a dummy endogenous regressor (a treatment indicator) and a dummy instrument, Abadie (1999) finds the best linear (least squares) approximation to the average causal effect for compliers. Compliers are those who obtain treatment because they are affected by the instrument. Abadie shows that this best approximation is equal to the 2SLS estimator in a model with regressors, when the probability of being affected by the instrument is linear in the regressors, but in general 2SLS is not the best linear approximation.

where a ‘^’ again denotes estimates from the first stage estimation of (3) and the ξ_i ’s are error terms with mean zero. As seen, these are the linear versions of the 2SIV and 2SCML estimators, which therefore in this special case are identical. If other covariates than z_i are correlated with α_i , this 2SLS estimator is not consistent. Hausman and Taylor (1981) suggested a consistent estimator of γ , removing the influence of the correlation between other regressors and α_i by first estimating β consistently by the within estimator obtained as:

$$(15) \quad Qy = QX_1\beta + Qv^l$$

where Q is the within transformation, $(Qy)_{it} = y_{it} - y_{i\cdot}$, where $y_{i\cdot}$ is the mean over time of y_{it} . Let $\hat{\beta}_w$ be the within estimate. They then apply OLS to the following equation:

$$(16) \quad y_{it} - X_{it}\hat{\beta}_w = \gamma^l z_i + \eta^l \hat{V}_i + \xi_{3i}$$

we refer to this as the 2SLSF estimator, F for fixed effects. Without inclusion of the estimated residual, we refer to this as the fixed effect (FE) estimator, which is consistent if z_i is exogenous, but allows X_1 to be correlated with α^l .

4. MONTE CARLO DESIGNS

In this section we describe the Monte Carlo designs used to investigate the properties of the estimators specified in the last sections. In particular, we will focus on a binomial response probit model with one time-varying and one time-invariant regressor:

$$(17) \quad y = \tau(y^*, \psi) = 1(\beta_0 + x_{it}\beta_1 + \gamma z_i + \alpha_i + \varepsilon_{it} > 0)$$

We define a benchmark design, where we let $(\beta_0, \beta_1, \gamma) = (1, 1, 1)$ and $\alpha_i, \varepsilon_{it}, x_{i1}$ be i.i.d. standard normal distributed and $x_{it} = x_{it-1} + 1$. Furthermore we generate an instrumental variable w_i , also standard normal and generate z_i as follows:

$$(18) \quad z_i = \rho w_i + \lambda \alpha_i + v_i$$

where v_i is standard normal. Therefore we can inspect how the different estimators behave when we let λ vary, which determines the degree to which z_i and y_{it} are determined by a common

component¹⁰. We can also let ρ vary, which determines the correlation between the instrument and z_i . This benchmark design is compared against designs with different sources of misspecification. First it may be of interest to infer whether one of the LDV estimators are more robust against misspecification of the distribution of α_i , so in a second design we draw α_i from a chi-squared distribution with one degree of freedom which is a right skewed distribution. Finally we also compare with a third design where we let z_i depend on α_i and its square, to infer the impact on the AGLS and 2SCML estimators which are based on an assumption that $\alpha_i|V_i$ has a mean that is linear in V_i .

4.1 Monte Carlo Designs with Real Data.

It is often found that Monte Carlo simulations depend on the distribution of the regressors, such that the use of artificial regressors may produce too nice results, see e.g. the chapter on Monte Carlo simulations in Davidson and MacKinnon (1993). We therefore also consider the behavior of the estimators using real data regressors.

We use a two-period representative panel of Danish workers interviewed in 1990 and 1995 (The Danish National Work Environment Cohort Study (WECS)). The specific sample we use is described in Arendt (2001). It was the empirical analysis in the latter paper that motivated the 2SCMLR panel estimator suggested in this paper. We used the 2SCMLR estimator to estimate education effects on health, modelled in an ordered quantal response model. There is a large literature within economics concerned with the interpretation of education effects on health, summarized in Grossman and Kaestner (1997). The main concern is that the correlation reflects common unobserved components, such as time preferences, determined prior to educational attainment. The outcome could be any other variable than health, where we would suspect that education effects are partly due to selection. We use 500 observations on women from this sample. Education is measured in years of education. We limit the number of regressors to a year dummy, a dummy for being a white collar worker and age. Instrumental variables for education include means over time of age and of the white collar dummy, in spirit of the suggestion by Hausman and Taylor (1981). In addition to these instruments, we use two Danish school reforms, in 1958 and

¹⁰ With this linear set-up, $V=\alpha\lambda+v$, st. when α , v and w are independent and have variance one: $\eta = \text{Cov}(\alpha, V)/\text{Var}(V)=\lambda/(1+\lambda^2)$. For λ equal to .1, this is equal to 0.099, and when λ is .5, this is 0.4.

1975, to instrument education. The reforms and the validity of these reforms as instruments for education are also discussed in Arendt (2001). We generate two dummy instrumental variables, indicating whether individuals are affected or not by the reforms.

Artificial health variables using these data are constructed in the following way. First we estimate the parameters in equation (3) and form predicted education. Then we draw i.i.d. standard normal errors α_i and v_{it} and generate education as in (18), replacing ρw by predicted education. Finally, the latent outcome is generated using the real regressors with prefixed values of β and γ and adding the generated errors α_i and v_{it} . See the appendix for details and descriptive statistics on the data.

5. MONTE CARLO RESULTS

This section contains the results from the Monte Carlo simulations¹¹. We focus on estimation of marginal effects on the probability that y is one, but note that in panel data models with unobserved heterogeneity, several marginal effects can be calculated, see Lillard and Willis (1978), section 3, for a discussion. Since in practice, the individual specific effect is not observed, calculation of individual probabilities is not feasible. A second approach is to let the individual effect vary and take a mean across its distribution, while a third approach evaluates at mean effects equal to zero. Since there is a tradition in applied econometrics to evaluate marginal effects at means of observable covariates we will do this, but will average out the unobservable individual effect for the panel estimators¹². With standard normal errors and variables, the true marginal effect on the

¹¹ The simulations are conducted using GaussTM vrs. 3.2.4., and the Gauss “rndn” random normal number generator. We use the Probit, OLS and Maxlik procedures to estimate the different models. We integrate out individual effects by Hermite integration formulas with three points, see Butler and Moffitt (1982) and the appendix, using a procedure written in GaussTM by Paul Fackler, adapted from Press et al., Numerical Recipes in Fortran, 2nd ed. The random effects probit likelihood is given in the appendix.

¹² To be specific, marginal effects are calculated as follows:

$$\frac{\partial P(y=1 | X, z)}{\partial z} = \gamma \phi(-\beta_0 - \bar{x}\beta_1 - \bar{z}\gamma),$$

probability that y equals one is 0.1607¹³. We report results from eight different probit models (probit, 2SCML, AGLS, 2SIV with and without random effects) as well as for the linear OLS, 2SLS, FE, 2SLSF models. We will refer to all estimators that account for endogeneity of the time-invariant variable as two-stage estimators, and the others as one-stage estimators (including the FE although it is estimated in two stages). In all designs we report the bias and mean squared error of estimated marginal effects¹⁴. 100 Monte Carlo simulations are conducted for each experiment.

In table 1 we report results using a data set with 200 observations and two time periods, on the benchmark design, i.e. all regressors and errors are normal, with different values of ρ and λ . The value of λ determines the correlation between z_i and individual effects, and ρ determines the correlation between the instrument and z_i . In the simulations we use λ equal to zero (i.e. no common individual effect, so z_i is exogenous, hence two-stage procedures are not needed), 0.1 and 0.5, and ρ equal to 0.05, 0.2 and 0.5. The case with ρ equal to 0.05 is supposed to illustrate a case with a weak instrument. Note that in all cases, random effects are present in the true model.

Starting with the top left corner of table 1, ρ is 0.05. In the bottom row we report the partial R-squared of w , the instrument, from the first stage estimation of z_i on x_1 and w , to indicate whether we have a weak instrument. As seen, with ρ equal to 0.05, this is quite low, around 0.007. Values of this size and even lower, have been observed e.g. with IV-estimation based on “natural experiments”, see e.g. Bound et al.(1995). As a consequence, for all values of λ ,

for the simple probit estimator, where bars denote empirical means and ϕ is the standard normal density function, and as:

$$\frac{\partial P(y = 1 | X, z)}{\partial z} = \gamma \int \phi(-\beta_0 - \bar{x}\beta_1 - z\gamma - \alpha) d\alpha$$

for the random effects probit estimators, using Hermite integration for calculation of the integral.

¹³ This is the integral with respect to α over $\phi(-1-E(X)-\alpha)\phi(\alpha)$, where the mean of X is a half, since $X_2=X_1+1$. We have simulated this as the mean over 60,000 draws of $\phi(-1.5-\alpha)$ where α is drawn from a standard normal.

¹⁴ The bias of an estimate $\hat{\theta}$ of θ is $E(\hat{\theta} - \theta)$ and the MSE is $E(\hat{\theta} - \theta)^2$. We calculate this using the empirical mean of the sample of estimates obtained from the Monte Carlo Simulations.

estimators, except perhaps the 2SIV, have a very high MSE. In the first column, λ is 0 so z_i is exogenous, and therefore the RE probit estimator is the correct estimator to use. The table shows that the RE probit estimator does estimate the marginal effect precisely, both regarding bias and MSE, although e.g. the probit performs even better. The 2SLSF has the highest MSE, followed by the AGLS estimators which in addition have the highest bias. Increasing λ to 0.1, in column (3) and (4) of the top of table 1, reduces the bias of the AGLS estimators and the estimator with the lowest bias is now the 2SCMLR, although all two-stage estimators, except the 2SIV, still have a high MSE. With λ equal to 0.5, the bias of all estimators increases (except the 2SLSF), and the 2SIV, OLS and the FE estimators have the smallest bias. The MSE is particular high for the 2SLS and the 2SLSF estimators. These results illustrate that two-stage estimators, linear as well as non-linear, are very imprecise as expected when the instruments used are weak. The only exception is the 2SIV estimator without random effects. We note that for the 2SCMLR estimator there is a small gain, both regarding bias and MSE, in taking account of the individual effects, i.e. when compared to the 2SCML estimator, whereas this is not the case for the AGLS and the 2SIV estimators.

In the next block, ρ is 0.2, which yields a partial R-squared for the instrument around 0.04. This is, we think, still likely to be a realistic level in many situations with individual data. When λ is 0, most estimators have a low bias, with the exception of the AGLS and the 2SLSF estimators. With λ equal to 0.1, the bias and MSE of the AGLS estimators are again reduced dramatically. The bias and MSE are still higher for most two-stage estimators than for one-stage estimators. Results change somewhat when λ is 0.5, where the bias of one-stage estimators increase beyond that of most two-stage estimators, but note that no clear ordering between the estimators that account for heterogeneity and their corresponding cross-sectional estimators can be made.

When ρ is increased further to 0.5, the partial R-squared is around 0.2, which is more rare in practice with individual data, and the MSE of the two-step estimators in particular diminishes further. Notable changes occurs for the linear 2SLSF estimator for high values of λ . Again, AGLS is a little worse than other estimators in the case of exogeneity, and the biases rise as before when λ is increased from 0.1 to 0.5, most for the probit and the RE probit.

In table 2 we increase the number of observations to 1000. The results are not that different from the case with 200 observations. Thus with a weak instrument, the 2SLSF is the worst in terms of bias, followed by the AGLS estimators under exogeneity (when λ is zero). We note though that compared to the case with 200 observations, the MSE of the two-stage estimators decreases in most

cases. We also see that with this amount of observations, with λ being different from zero, the advantage of taking account of endogeneity is higher, even when the instruments are weak (with 2SIVR for $(\lambda, \rho) = (0.5, 0.05)$ and AGLSR for $(\lambda, \rho) = (0.5, 0.2)$ and $(0.5, 0.5)$ being exceptions). The 2SCML estimators seem to perform particular well.

Table 3 show results with simulations similar to those in table 2, except that we have increased the number of periods to five¹⁵. Although the asymptotic results only depend on the size of n , the number of individuals, it could be that in small samples the number of time periods may affect the results. Table 3 shows that this is the case. With weak instruments, both bias and MSE increase for most estimators. The AGLSR estimators now perform particularly bad, even with good instruments and a high degree of endogeneity. The 2SCML and the 2SIV estimators are preferred to probit and RE probit when endogeneity is severe ($\lambda = 0.5$) and reasonable instruments are available ($\rho > 0.05$), and perform just as well as 2SLS when $\rho > 0.05$.

So far we have considered the small sample behavior of the different estimators under normality assumptions. In table 4 we look at the behavior when individual effects are distributed as chi-squared with one degree of freedom, minus one to obtain mean zero. We still draw z_i as a linear function of individual effects, such that the assumption used by the 2SCML and AGLS estimators, that $E(\alpha_i | V_i)$ is linear in V_i , is correct. One might expect that linear estimators provide a better approximation to marginal effects than LDV estimators because they do not rely on distributional assumptions regarding the individual effect. The true marginal effect under this scenario is now 0.2088¹⁶.

This larger effect reflects that the distribution of α_i is right skewed but enters the marginal effect with a minus, and is thus putting largest weight on individuals with marginal effects near plus one (the density of $-\alpha - 1$ has a vertical asymptote at one). The general tendency therefore is to underestimate the marginal effect, because cross-sectional estimators do not integrate out individual

¹⁵ We generate x_{i1} as standard normal, but let $x_{it} = x_{it-1} + 0.25$ implying that $E(X_i) = 0.2*0 + 0.2*0.25 + 0.2*0.5 + 0.2*0.75 + 0.2*1 = 0.5$, to obtain the same true marginal effect: the integral over $\phi(-1.5-\alpha)\phi(\alpha) = 0.1607$.

¹⁶ This is simulated as the mean over 60,000 draws of $\phi(-1.5-\alpha)$, where α is drawn from a squared normal, subtracted by one to make the mean equal to zero.

effects, and the panel estimators use a weight function (the normal distribution) that places too much weight on individuals with low marginal effects. We see that the AGLS estimators and the 2SIV may be substantially biased. The 2SCML estimators unexpectedly perform very well. It is seen that none of the linear estimators provide a better approximation to marginal effects than the 2SCML estimators.

In table 5 we allow normal distributed individual effects to affect z_i quadratically:

$$z_i = \rho w_i + \lambda \alpha_i - \lambda^2 \alpha_i^2 + v_i$$

implying that the control function $E(\alpha|V)$, incorporated by the 2SCML and the AGLS estimators, is misspecified. The true marginal effect is 0.1607 as in the benchmark scenario. The results found in the benchmark scenario for λ less than 0.5 apply here as well. For λ equal to 0.5 most estimators are severely biased as expected, but when ρ is larger than 0.05, this is not worse for all the 2SCML and AGLS estimators than other two-stage estimators, as one might expect. It is noteworthy that the OLS estimator performs reasonably well in all cases, and the 2SLS has a larger bias than most non-linear estimators when λ is 0.5.

In order to obtain some robustness against the assumed normal distribution of regressors, we conduct a final set of simulations using a design with real data as described above. Table 6 reports the results. The top of the table contains results where both school reforms and Hausman-Taylor type of instruments are used, and in the bottom, only school reforms are used. Because the endogenous variable, education, is simulated, the partial R-squared is very high (0.44-0.5) in the top part of the table where all instruments are used, whereas in the bottom table, the partial R-squared is much lower; only 0.03. Therefore, as might be expected, in the top of the table two-stage estimators perform better than estimators assuming exogeneity of education. It is seen that the 2SCML/-R and 2SLS estimators perform really well in particular, whereas the 2SLSF, the FE and the 2SIVR have large biases. Even in the case with low partial R-squared several two-stage estimators perform better than OLS, probit and RE probit.

6. CONCLUSION

We have extended three cross-sectional LDV model estimators to a panel model with individual specific effects. To our knowledge, besides joint MLE, these are the only existing consistent and asymptotic normal estimators for the effect of a time-invariant endogenous continuous regressor in

a LDV model with unobserved individual effects. However, these models are time-consuming to apply. It is therefore of great interest how well they perform in small samples. In addition, when the outcome of interest is the marginal effect on the LDV outcome rather than LDV parameters per se, comparison against much simpler linear approximations can be conducted. We conducted Monte Carlo simulations of the probit model, paying special attention to the estimated marginal effect on the probability that the binomial variable equals one.

Using the design with artificial normal regressors and errors, we make the following general remarks¹⁷. When there is only weak suspicion of endogeneity (λ low), it might be preferable to use the simple probit, and AGLS seems to be the worst solution. The 2SIV estimator seems to be less affected by the problem of weak instruments than other estimators accounting for endogeneity, and 2SLS and 2SLSF seem to be the worst estimators with weak instruments. If a reasonable instrument is available and endogeneity is present ($\rho > 0.05, \lambda > 0$), two-stage estimators are often preferable in terms of bias, but not in terms of MSE. The extension of the cross-sectional estimators to the panel model, for designs considered here, does not seem to give any small sample improvements.

Increasing the number of individuals from 200 to 1000 showed a relative benefit in favor of two-stage estimators, whereas increasing the number of time periods introduced further imprecision for the AGLS estimators in particular.

For practical purposes, it is relevant to note that the 2SCML and 2SLS estimators perform reasonably well in many scenarios, the latter despite of the DGP being a LDV model. The 2SCML estimators were relatively robust against the sources of misspecification considered here. Since endogeneity is easily tested in these models, it seems that they constitute good starting points, perhaps compared to the OLS, Probit or the 2SIV estimators for robustness. We showed that linear estimators are subject to the same problems as LDV estimators, and do not perform better than these under misspecification. It should be stressed that the relative good performance of linear estimators in the benchmark design is not expected to hold if we had not confined attention to marginal effects evaluated at mean values of the regressors, stemming from the fact that a linear model yields marginal effects that are constant across individuals, which is impossible to hold for

¹⁷ We should have in mind that the results might be subject to some uncertainty, especially with respect to biases for those estimators with large MSE, since the number of Monte Carlo loops has been kept relatively low at the moment.

all individuals. This is not the case in LDV models, although the functional form of the marginal effects is given by the distributional assumption. However, the latter can and should be tested in practice.

We wrap up with some suggestions for future work. In general we know little about the small sample behavior of many existing LDV estimators. In our Monte Carlo study we focused on the binomial probit model, but it would be interesting to see how the estimators behave in other LDV models. Furthermore, a version of the 2SCMLR estimator has been considered in the case when the causal variable is discrete by Orme (1996). This could be compared to recent causal effects estimators suggested by Mullahy (1997) and Abadie (2000). Finally, as found in our study, parametric models are likely to be biased under misspecification, so the framework could be extended to semiparametric models. A first step could be to estimate the distribution of the individual effect by discrete factor approximation, as done by Mroz and Guilkey (1992) in cross-sectional models, and a second step could either be where the second stage estimation does not involve parametric assumptions, e.g. inspired by approaches reviewed by Blundell and Powell (2000), or where attention is directed at non-parametric estimation of marginal effects. Because of the curse-of-dimensionality of the latter a combination might be beneficial.

Appendix A: The Likelihood in the Random Effects Probit Model

For all the panel LDV estimators we need to estimate the random effects probit model. We therefore write down the likelihood for this model. The contribution for individual i to the likelihood for the random effects probit model is:

$$f(y_i | X_i, z_i) = \int \prod_{t=1}^T \Phi(c_{it})^{1-y_{it}} (1 - \Phi(c_{it}))^{y_{it}} \phi(w_i) dw_i$$

where e.g. in the 2SCMLR model, $c_{it} = X_{1it}\beta + \gamma z_i + \eta V_i + w_i$ and X_{1i} is the vector $(X_{1i1}, \dots, X_{1iT})$, and ϕ and Φ are standard normal probability and cumulative distribution functions. Gaussian-Hermite quadrature can be used to approximate this integral. The quadrature of order n approximates integrals involving e^{-x^2} by a sum:

$$\int e^{-z^2} f(z) dz \approx \sum_{i=1}^n w_i f(x_i)$$

where w and x are specific values, depending on the number n . Then for the normal density we have:

$$\int \phi(z) f(z) dz \approx \sum_{i=1}^n \frac{w_i}{\sqrt{\pi}} f(\sqrt{2}x_i)$$

We use $n=3$, which gives $x = (-1.2247, 0, 1.2247)$, $w = (0.2954, 1.1826, 0.2954)$. For applied researchers we note that the 2SIVR and the 2SCMLR models can be estimated without messing with the quadrature themselves. One needs a linear regression to form the residuals and predicted values and a random effects LDV estimator. In StataTM the latter is available for some models with the procedure `xtgee`. The GLLAMM package by Rabe-Hesketh includes other random effect models (see <http://www.iop.kcl.ac.uk/IoP/Departments/BioComp/programs/gllamm.html>). The AGLS estimator is implemented by J. B. Gelbach for the probit (see <http://www.glue.umd.edu/~gelbach/ado/>) and can be modified to other LDV models using the `xtgee` procedures. A problem is that standard errors must be corrected. Besides Gelbach's latest version of the cross-sectional AGLS estimator, I have no knowledge of procedures that does this at the moment. Newey (1987) and Vuong (1984) outline how to obtain correct standard errors, although the results become particularly cumbersome for the 2SCMLR estimator. Bootstrapping might be another solution. Bollen et al. (1995) cite a Monte Carlo study, p.115, showing that standard error correction does not seem to matter for the 2SCML probit estimator.

Appendix B: Specification of Design using Real Data

First we present some descriptive statistics on the used data:

TABLE A. Descriptive Statistics for the WECS Data.

Variable	Mean	Std.dev	Min	Max
Age	37.642	9.251	18	59
White Collar	0.818	0.387	0	1
Education	13.48	2.33	7	18
Predicted Educ.	13.48	0.997	10.737	14.207
Mean White Coll.	0.808	0.331	0	1
Reform 58	0.586	0.493	0	1
Reform 75	0.196	0.397	0	1

Notes: 500 observations on Women from the WECS data. 1990. Mean White Coll. is the mean over 1990 and 1995 of the White Collar dummy.

The Monte Carlo simulations are conducted using the following specification:

$$y_{it} = 1(\beta_0 + \beta_1 \tilde{E}_i + \beta_2 A_{it} + \beta_3 WC_{it} + \beta_4 D_t + \alpha_i + \varepsilon_{it} > 0)$$

$$(\beta_0, \beta_1, \beta_2, \beta_3, \beta_4) = (-0.5, -0.1, 0.02, -0.2, 0.05)$$

$$i = 1, \dots, 500 \quad t = 1990, 1995$$

where A_{it} is age, WC_{it} is a dummy for being a white collar worker and D_t is a year dummy. E_i -tilde is simulated education of individual i , constructed as follows. First we estimate a linear regression of years of education on A_{it} , W_{it} , the means over time for these, D_t and two school reform instruments. The result, with p-values below parameter estimates is:

$$\tilde{E}_i = 9.274 - 0.324A_{i1} + 0.087WC_{i1} + 0.328MA_i + 2.831MWC_i + 0.554R58_i + 0.473R75_i$$

$$(0.206) (0.915) (0.845) (0.914) (0.000) (0.141) (0.456)$$

$$Adj.R^2 = 0.172, N = 500$$

where MX_i is the individual mean of X in 1990 and 1995, $R58_i$ is a dummy for whether individual i was affected by a 1958 reform, but not by a 1975 reform, and $R75_i$ is a dummy for whether individual i was affected by the 1975 reform.

References

Abadie, A. (1999). Semiparametric Estimation of Instrumental Variable Models for Causal Effects, Mimeo, Massachusetts Institute of Technology.

Alvarez, M. R. and G. Glasgow (2000). Two-Stage Estimation of Non-Recursive Choice Models, *Political Analysis* 8 (2), 147-165.

Amemiya, T. (1978). The Estimation of a Simultaneous Equation Generalized Probit Model, *Econometrica* 46 (5), 1193-1205.

Amemiya, T. (1979). The Estimation of a Simultaneous Equation Generalized Tobit Model, *International Economic Review* 20, 169-181.

Andersen, E. B. (1970). Asymptotic Properties of Conditional Maximum Likelihood Estimators, *Journal of the Royal Statistical Society, Series B*, 32, 283-301.

Angrist, J. D. (2001). Estimation of Limited Dependent Variable Models with Dummy Endogenous Regressors: Simple Strategies for Empirical Practice, *Journal of Business & Economic Statistics* 19 (1), 2-16.

Arendt, J. N. (2001). Education effects on Health. A Panel Data Analysis using School Reforms for Identification, Mimeo, Institute of Economics, University of Copenhagen.

Blundell, R. and J. Powell (2000). Endogeneity in Nonparametric and Semiparametric Regression Models, Mimeo Prepared for the Econometric Society World Meetings Seattle, August 2000.

Blundell, R. and R. Smith (1986). An Exogeneity Test for a Simultaneous Equation Tobit Model with an Application to Labor Supply, *Econometrica* 54 (3), 679-85.

Blundell, R. and R. Smith (1989). Estimation in a class of Simultaneous Equation Limited Dependent Variable Models, *Review of Economic Studies* 56 (1), 37-57.

Blundell, R. and R. Smith (1994). Coherency and Estimation in Simultaneous Models with Censored or Qualitative Dependent Variables, *Journal of Econometrics* 64, 355-373.

Bollen, K. A., Guilkey, D. K. and T. A. Mroz (1995). Binary Outcomes and Endogenous Explanatory Variables: Tests and Solutions with an Application to the Demand for Contraceptive Use in Tunisia, *Demography* 32 (1), 111-131.

Bound, J., Jaeger, D. A. and R. M. Baker (1995). Problems with Instrumental Estimation When the Correlation Between the Instruments and the Endogenous Explanatory Variable is Weak, *Journal of the American Statistical Association* 90 (430), 443-450.

Breusch, T. S., Mizon, G. E. and P. Schmidt (1989). Efficient Estimation Using Panel Data, *Econometrica* 57 (3), 695-700.

Buttler, J. and R. Moffitt (1982). A Computationally Efficient Quadrature Procedure for the One-Factor Multinomial Probit Model, *Econometrica* 50 (3), 761-764.

Chamberlain, G. (1980). Analysis of Covariance with Qualitative Data, *Review of Economic Studies* 47 (1), 225-238.

Davidson, R. and J. G. MacKinnon (1993). *Estimation and Inference in Econometrics*, New York: Oxford University Press.

Grossman, M. and R. Kaestner (1997). Effects of Education on Health, in Behrman, J. and N. Stancey (Eds.), *The Social Benefits of Education*, Ann Arbor, The University of Michigan Press, 69-123.

Hausman, J. A. (1978). Specification Tests in Econometrics, *Econometrica* 46 (6), 1251-1271.

Hausman, J. A. and W. E. Taylor (1981). Panel Data and Unobservable Individual Effects, *Econometrica* 49 (6), 343-414.

Heckman, J. J. (1978). Dummy Endogenous Variables in a Simultaneous Equation System, *Econometrica* 46 (5), 931-959.

Heckman, J. J. and T. E. MaCurdy (1980). A Life Cycle Model of Female Labour Supply, *Review of Economic Studies* 47 (1), 47-74.

Honore, B. E. (1992). Trimmed LAD and Least Squares Estimation of Truncated and Censored Regression Models with Fixed Effects, *Econometrica* 60 (3), 533-565.

Horowitz, J. L. (1993). Semiparametric and Nonparametric Estimation of Quantal Response Models, in G. S. Maddala, C. R. Rao and H. D. Vinod (Eds.), *Handbook of Statistics* 11, Elsevier Science, Amsterdam (1993), 45-72.

Kyriazidou, E. (1997). Estimation of a Panel Data Sample Selection Model, *Econometrica* 65 (6), 1335-64.

Lee, L. F. (1981). Simultaneous Equation Models with Discrete and Censored Dependent Variables, in Manski, C. and D. McFadden (Eds.), *Structural Analysis of Discrete Data with Economic Applications*, Cambridge, MA: MIT Press.

Lee, M. (1999). A Root-n Consistent Semiparametric Estimator for Related-Effect Binary Response Panel Data, *Econometrica* 67 (2), 427-433.

Lillard, L. A. and R. J. Willis (1978). Dynamic Aspects of Earnings Mobility, *Econometrica* 46 (5), 985-1012.

Manski, C. F. (1987). Semiparametric Analysis of Random Effects Linear Models from Binary Panel Data, *Econometrica* 55 (2), 357-362.

Mroz, T. A. and D. K. Guilkey (1992). Discrete Factor Approximations for Use in Simultaneous Equation Models with Both Continuous and Discrete Endogenous Variables, Carolina Population Center Papers 92-03, University of North Carolina at Chapel Hill.

Nelson, F. and L. Olsen (1978). Specification and Estimation of a Simultaneous Equation Model with Limited Dependent Variables, *International Economic Review* 19, 659-705.

Newey, W. K. (1987). Efficient Estimation of Limited Dependent Variable Models with Endogenous Explanatory Variables, *Journal of Econometrics* 36, 231-250.

Orme, C. (1996). The Initial Conditions Problem and Two-Step Estimation in Discrete Panel Data Models, University of Manchester, Working Paper No. 9633.

Rivers, D. and Vuong, Q. H. (1988). Limited Information Estimators and Exogeneity Tests for Simultaneous Probit Models, *Journal of Econometrics* 39, 347-366.

Vella, F. and M. Verbeek (1999). Two-Step Estimation of Panel Data Models with Censored Endogenous Variables and Selection Bias, *Journal of Econometrics* 90 (2), 239-263.

Vuong, Q. H. (1984). Two-Stage Conditional Maximum Likelihood Estimation of Econometric Models, Social Science Working Paper 538, California Institute of Technology.

Table 1. Monte Carlo Results for Estimation of Marginal Effects of Time Invariant Regressor in a Panel, 200 Observations.

$\rho=0.05$	$\lambda=0$		$\lambda=0.1$		$\lambda=0.5$	
	Bias	MSE	Bias	MSE	Bias	MSE
Estimator						
OLS	-0.0104	0.0006	-0.0070	0.0006	0.0354	0.0017
FE	-0.0072	0.0039	-0.0062	0.0040	0.0329	0.0036
2SLS	0.0305	0.1869	-0.0070	0.1997	0.1543	2.9370
2SLSF	0.2080	23.5234	0.5133	19.1671	0.4600	53.1792
Probit	0.0006	0.0006	0.0064	0.0008	0.0641	0.0049
RE Probit	-0.0097	0.0007	-0.0048	0.0008	0.0484	0.0032
2SCML	0.0350	0.2031	0.0260	0.2327	0.0771	0.6397
2SCMLR	0.0291	0.1829	0.0004	0.1958	0.0709	0.5967
AGLS	0.2409	1.7077	0.0053	1.7315	0.0964	0.5444
AGLSR	0.3247	2.2576	0.0209	1.8326	0.0586	0.4103
2SIV	-0.0081	0.0051	-0.0192	0.0054	-0.0330	0.0051
2SIVR	-0.0045	0.2123	0.0359	0.4016	0.1105	0.1060
Partial R2	0.0078		0.0071		0.0066	

$\rho=0.2$	$\lambda=0$		$\lambda=0.1$		$\lambda=0.5$	
	Bias	MSE	Bias	MSE	Bias	MSE
Estimator						
OLS	-0.0121	0.0007	-0.0008	0.0005	0.0319	0.0015
FE	-0.0180	0.0041	0.0055	0.0033	0.0383	0.0041
2SLS	-0.0115	0.0189	-0.0348	0.0227	-0.0060	0.0181
2SLSF	-0.0304	2.5576	0.2962	24.3198	0.1417	1.5810
Probit	0.0010	0.0008	0.0114	0.0007	0.0215	0.0065
RE Probit	-0.0096	0.0008	0.0000	0.0006	0.0429	0.0026
2SCML	0.0020	0.0188	-0.0296	0.0204	0.0114	0.0217
2SCMLR	-0.0076	0.0153	-0.0382	0.0184	-0.0008	0.0185
AGLS	0.1333	2.8409	0.0022	0.0310	-0.0128	0.0165
AGLSR	0.1463	2.1161	-0.0140	0.0254	-0.0557	0.0096
2SIV	0.0007	0.0015	-0.0129	0.0018	-0.0173	0.0022
2SIVR	-0.0129	0.0084	0.0275	0.0292	-0.0001	0.0159
Partial R2	0.0431		0.0431		0.0385	

$\rho=0.5$	$\lambda=0$		$\lambda=0.1$		$\lambda=0.5$	
	Bias	MSE	Bias	MSE	Bias	MSE
Estimator						
OLS	-0.0131	0.0008	-0.0033	0.0005	0.0210	0.0008
FE	-0.0133	0.0044	0.0003	0.0031	0.0254	0.0034
2SLS	-0.0160	0.0021	-0.0196	0.0023	-0.0305	0.0031
2SLSF	-0.0223	0.0649	-0.0088	0.0540	0.0031	0.0775
Probit	-0.0003	0.0006	0.0109	0.0008	0.0499	0.0034
RE Probit	-0.0111	0.0006	-0.0004	0.0007	0.0359	0.0021
2SCML	-0.0050	0.0017	-0.0076	0.0023	-0.0093	0.0028
2SCMLR	-0.0169	0.0018	-0.0172	0.0024	-0.0202	0.0028
AGLS	0.0236	0.1012	-0.0038	0.0025	-0.0121	0.0033
AGLSR	0.0323	0.0886	-0.0227	0.0027	-0.0639	0.0054
2SIV	-0.0168	0.0027	-0.0014	0.0010	-0.0122	0.0012
2SIVR	0.0617	0.1182	-0.0060	0.0020	-0.0136	0.0023
Partial R2	0.2063		0.2096		0.1671	

Notes: Bias and mean squared error of marginal effect on probability of positive outcome of continuous time-invariant variable. All are calculated from 100 simulations of the benchmark design on 200 obs. with 2 time periods, see the text. λ determines the correlation between z and the individual effect and ρ determines the correlation between z and the instrument.

Table 2. Monte Carlo Results for Estimation of Marginal Effects of Time Invariant Regressor in a Panel, 1000 observations.

$\rho=0.05$ Estimator	$\lambda=0$		$\lambda=0.1$		$\lambda=0.5$	
	Bias	MSE	Bias	MSE	Bias	MSE
OLS	-0.0121	0.0003	0.0015	0.0002	0.0342	0.0013
FE	-0.0129	0.0008	0.0042	0.0009	0.0340	0.0019
2SLS	-0.0043	0.2289	-0.0231	0.1710	-0.0212	0.0544
2SLSF	0.8743	121.3321	-1.2385	80.0597	0.4463	58.8585
Probit	-0.0009	0.0001	0.0168	0.0004	0.0621	0.0040
RE Probit	-0.0117	0.0003	0.0050	0.0002	0.0458	0.0022
2SCML	-0.0202	0.2638	0.0051	0.1842	0.0099	0.0564
2SCMLR	-0.0241	0.2360	-0.0082	0.1591	-0.0023	0.0483
AGLS	0.2013	4.2889	0.0236	0.1012	0.0364	0.0950
AGLSR	0.2593	4.1352	0.0323	0.0886	-0.0095	0.0374
2SIV	-0.0140	0.0035	-0.0168	0.0029	-0.0237	0.0018
2SIVR	0.0072	0.2082	0.0617	0.1182	0.0613	0.0623
Partial R2	0.0034		0.0036		0.0029	

$\rho=0.2$ Estimator	$\lambda=0$		$\lambda=0.1$		$\lambda=0.5$	
	Bias	MSE	Bias	MSE	Bias	MSE
OLS	-0.0132	0.0003	0.0005	0.0001	0.0307	0.0010
FE	-0.0123	0.0010	0.0015	0.0007	0.0286	0.0013
2SLS	-0.0099	0.0020	-0.0151	0.0025	-0.0223	0.0021
2SLSF	-0.0031	0.3608	-0.0016	0.2942	-0.0656	0.4734
Probit	-0.0018	0.0002	0.0152	0.0003	0.0593	0.0037
RE Probit	-0.0125	0.0002	0.0034	0.0000	0.0434	0.0020
2SCML	0.0048	0.0022	0.0018	0.0027	-0.0018	0.0019
2SCMLR	-0.0063	0.0020	-0.0104	0.0027	-0.0162	0.0018
AGLS	-0.0070	0.0043	-0.0011	0.0037	-0.0053	0.0025
AGLSR	-0.0068	0.0023	-0.0176	0.0021	-0.0613	0.0047
2SIV	-0.0026	0.0003	-0.0048	0.0004	-0.0132	0.0023
2SIVR	0.0003	0.0023	0.0001	0.0025	-0.0107	0.0021
Partial R2	0.0391		0.0396		0.0317	

$\rho=0.5$ Estimator	$\lambda=0$		$\lambda=0.1$		$\lambda=0.5$	
	Bias	MSE	Bias	MSE	Bias	MSE
OLS	-0.0144	0.0003	-0.0037	0.0001	0.0216	0.0005
FE	-0.0176	0.0008	-0.0037	0.0006	0.0201	0.0009
2SLS	-0.0125	0.0004	-0.0153	0.0006	-0.0229	0.0010
2SLSF	-0.0280	0.0084	-0.0164	0.0085	-0.0313	0.0111
Probit	0.0007	0.0001	0.0130	0.0004	0.0513	0.0027
RE Probit	-0.0102	0.0002	0.0013	0.0002	0.0359	0.0014
2SCML	0.0024	0.0003	-0.0003	0.0004	-0.0021	0.0005
2SCMLR	-0.0081	0.0004	-0.0120	0.0005	-0.0138	0.0006
AGLS	-0.0007	0.0004	0.0030	0.0004	-0.0029	0.0005
AGLSR	-0.0101	0.0007	-0.0211	0.0008	-0.0607	0.0040
2SIV	-0.0014	0.0002	-0.0037	0.0002	-0.0105	0.0003
2SIVR	-0.0064	0.0005	-0.0032	0.0004	-0.0096	0.0004
Partial R2	0.2010		0.1993		0.1681	

Notes: Same as table 1, now with 1000 observations.

Table 3. Monte Carlo Results for Estimation of Marginal Effects of Time Invariant Regressor in a Panel, 200 observations, 5 Periods.

$\rho=0.05$ Estimator	$\lambda=0$		$\lambda=0.1$		$\lambda=0.5$	
	Bias	MSE	Bias	MSE	Bias	MSE
OLS	-0.0114	0.0006	0.0058	0.0005	0.0372	0.0019
FE	0.2143	0.0614	0.2399	0.0709	0.3406	0.1280
2SLS	0.0226	13.0908	0.0712	1.2282	0.0995	0.6987
2SLSF	2.1494	161.5025	-0.8271	1.6321	1.2527	70.1212
Probit	-0.0651	0.0048	-0.0863	0.0080	-0.1317	0.0180
RE Probit	-0.0063	0.0006	0.0141	0.0008	0.0558	0.0037
2SCML	0.0777	18.9919	0.2603	2.4740	0.1046	0.3913
2SCMLR	-0.0409	14.3164	0.2766	2.7638	0.1711	0.2221
AGLS	0.1620	23.3020	1.78201	171.8078	1.6155	60.8675
AGLSR	0.2601	23.8158	2.0224	182.0584	3.0163	208.7492
2SIV	-0.0318	0.0065	-0.0428	0.0063	-0.0409	0.0054
2SIVR	0.1309	2.1682	0.1338	2.1775	0.0274	0.2891
Partial R2	0.0085		0.0071		0.0077	

$\rho=0.2$ Estimator	$\lambda=0$		$\lambda=0.1$		$\lambda=0.5$	
	Bias	MSE	Bias	MSE	Bias	MSE
OLS	-0.0086	0.0006	0.0029	0.0006	0.0370	0.0017
FE	0.2154	0.0602	0.2413	0.0704	0.3291	0.1209
2SLS	0.0132	0.0281	0.0011	0.0144	-0.0215	0.0633
2SLSF	-0.1214	13.5223	-0.0440	13.2076	0.1740	13.6416
Probit	-0.0691	0.0053	-0.0834	0.0076	-0.1322	0.0182
RE Probit	-0.0019	0.0005	0.0111	0.0007	0.0570	0.0039
2SCML	0.0332	0.0329	0.0189	0.0179	0.0048	0.12788
2SCMLR	0.0224	0.0334	0.0186	0.0179	0.0121	0.1401
AGLS	1.4013	12.1650	1.5276	30.6576	1.1509	10.6518
AGLSR	1.4605	11.6859	1.8418	33.2036	1.8802	24.6452
2SIV	-0.0214	0.0020	-0.0190	0.0017	-0.0185	0.0022
2SIVR	0.0235	0.0304	0.0161	0.0151	-0.0110	0.0674
Partial R2	0.0366		0.0386		0.0397	

$\rho=0.5$ Estimator	$\lambda=0$		$\lambda=0.1$		$\lambda=0.5$	
	Bias	MSE	Bias	MSE	Bias	MSE
OLS	-0.0090	0.0005	-0.0040	0.0004	0.0573	0.0036
FE	0.2197	0.0582	0.2165	0.0615	0.3375	0.1206
2SLS	-0.0097	0.0018	-0.0106	0.0016	0.0126	0.0018
2SLSF	0.1986	0.3205	0.1492	0.3897	0.2443	0.3474
Probit	-0.0700	0.0054	-0.0762	0.0062	-0.0887	0.0084
RE Probit	-0.0012	0.0005	0.0042	0.0004	0.0750	0.0062
2SCML	0.0108	0.0021	0.0071	0.0014	0.0365	0.0034
2SCMLR	0.0039	0.0021	-0.0034	0.0014	0.0228	0.0025
AGLS	0.3739	1.3059	0.6150	2.1674	0.6338	2.2537
AGLSR	0.3491	1.2150	0.6662	2.4876	0.5559	2.4031
2SIV	-0.0101	0.0007	-0.0114	0.0007	0.0141	0.0011
2SIVR	-0.0014	0.0019	-0.0022	0.0014	0.0220	0.0024
Partial R2	0.2033		0.2072		0.1719	

Notes: As table 1, except now with 5 time periods.

Table 4. Monte Carlo Results for Estimation of Marginal Effects of Time Invariant Regressor in a Panel Probit Model, 200 observations, with Individual Effects being Chi-squared distributed.

$\rho=0,05$	$\lambda=0$		$\lambda=0,1$		$\lambda=0,5$	
Estimator	Bias	MSE	Bias	MSE	Bias	MSE
OLS	-0.0462	0.0025	-0.0313	0.0014	-0.0324	0.0014
FE	-0.0526	0.0051	-0.0295	0.0028	-0.0280	0.0018
2SLS	0.0112	0.4327	-0.0459	0.1919	-0.0431	0.1063
2SLSF	-0.6460	24.1736	0.0949	16.8893	0.4268	10.2847
Probit	-0.0234	0.0014	-0.0033	0.0012	0.0347	0.0030
RE Probit	-0.0328	0.0020	-0.0136	0.0013	0.0219	0.0024
2SCML	0.0154	0.4029	-0.0289	0.2028	0.0104	0.1378
2SCMLR	0.0062	0.3241	-0.0301	0.1967	0.0011	0.1212
AGLS	0.2663	6,1707	-0.1401	0.8534	-0.2722	6.6217
AGLSR	0.3478	7,3456	-0.1500	0.4653	-0.5151	13.8617
2SIV	-0.0890	0.0129	-0.1068	0.0155	-0.1229	0.0164
2SIVR	-0.1180	0.3693	-0.0787	0.5593	-0.0463	0.3012
Partial R2	0.0072		0.0079		0.0067	

$\rho=0,2$	$\lambda=0$		$\lambda=0,1$		$\lambda=0,5$	
Estimator	Bias	MSE	Bias	MSE	Bias	MSE
OLS	-0.0524	0.0031	-0.0304	0.0013	-0.0334	0.0015
FE	-0.0540	0.0049	-0.0361	0.0038	-0.0302	0.0019
2SLS	-0.0286	0.0395	-0.0442	0.0138	-0.0628	0.0171
2SLSF	-0.2168	1.6984	-0.1071	1.3348	0.1465	1,4181
Probit	-0.0308	0.0017	-0.0007	0.0011	0.0362	0.0032
RE Probit	-0.0402	0.0024	-0.0113	0.0012	0.0231	0.0025
2SCML	-0.0178	0.0389	-0.0122	0.0160	-0.0183	0.0200
2SCMLR	-0.0288	0.0352	-0.0235	0.0145	-0.0267	0.0165
AGLS	-0.0321	0.0142	-0.0368	0.0160	-0.0097	0.0281
AGLSR	-0.0599	0.0135	-0.0779	0.0126	-0.1034	0.0176
2SIV	-0.0823	0.0078	-0.0826	0.0085	-0.1112	0.0132
2SIVR	-0.0217	0.0122	-0.0315	0.0141	-0.0164	0.0150
Partial R2	0.0395		0.0469		0.0333	

$\rho=0,5$	$\lambda=0$		$\lambda=0,1$		$\lambda=0,5$	
Estimator	Bias	MSE	Bias	MSE	Bias	MSE
OLS	-0.0499	0.0028	-0.0410	0.0021	-0.0347	0.0015
FE	-0.0543	0.0046	-0.0438	0.0032	-0.0332	0.0020
2SLS	-0.0443	0.0038	-0.0517	0.0042	-0.0570	0.0051
2SLSF	-0.0720	0.0492	-0.0712	0.0434	-0.0504	0.0676
Probit	-0.0257	0.0014	-0.0132	0.0012	0.0379	0.0032
RE Probit	-0.0337	0.0019	-0.0226	0.0015	0.0258	0.0025
2SCML	-0.0200	0.0029	-0.0231	0.0034	-0.0162	0.0045
2SCMLR	-0.0284	0.0032	-0.0322	0.0037	-0.0265	0.0047
AGLS	-0.0330	0.0035	-0.0174	0.0034	-0.0138	0.0043
AGLSR	-0.0722	0.0068	-0.0731	0.0067	-0.1068	0.0128
2SIV	-0.0750	0.0065	-0.0870	0.0081	-0.1135	0.0133
2SIVR	-0.0404	0.0040	-0.0285	0.0032	-0.0241	0.0036
Partial R2	0.2028		0.2015		0.1387	

Notes: As table 1, except that individual effects are Chi-squared(1) distributed.

Table 4. Monte Carlo Results for Estimation of Marginal Effects of Time Invariant Regressor in a Panel Probit Model, 200 observations, with Individual Effects being Chi-squared distributed.

$\rho=0,05$	$\lambda=0$		$\lambda=0,1$		$\lambda=0,5$	
Estimator	Bias	MSE	Bias	MSE	Bias	MSE
OLS	-0.0462	0.0025	-0.0313	0.0014	-0.0324	0.0014
FE	-0.0526	0.0051	-0.0295	0.0028	-0.0280	0.0018
2SLS	0.0112	0.4327	-0.0459	0.1919	-0.0431	0.1063
2SLSF	-0.6460	24.1736	0.0949	16.8893	0.4268	10.2847
Probit	-0.0234	0.0014	-0.0033	0.0012	0.0347	0.0030
RE Probit	-0.0328	0.0020	-0.0136	0.0013	0.0219	0.0024
2SCML	0.0154	0.4029	-0.0289	0.2028	0.0104	0.1378
2SCMLR	0.0062	0.3241	-0.0301	0.1967	0.0011	0.1212
AGLS	0.2663	6,1707	-0.1401	0.8534	-0.2722	6.6217
AGLSR	0.3478	7,3456	-0.1500	0.4653	-0.5151	13.8617
2SIV	-0.0890	0.0129	-0.1068	0.0155	-0.1229	0.0164
2SIVR	-0.1180	0.3693	-0.0787	0.5593	-0.0463	0.3012
Partial R2	0.0072		0.0079		0.0067	

$\rho=0,2$	$\lambda=0$		$\lambda=0,1$		$\lambda=0,5$	
Estimator	Bias	MSE	Bias	MSE	Bias	MSE
OLS	-0.0524	0.0031	-0.0304	0.0013	-0.0334	0.0015
FE	-0.0540	0.0049	-0.0361	0.0038	-0.0302	0.0019
2SLS	-0.0286	0.0395	-0.0442	0.0138	-0.0628	0.0171
2SLSF	-0.2168	1.6984	-0.1071	1.3348	0.1465	1,4181
Probit	-0.0308	0.0017	-0.0007	0.0011	0.0362	0.0032
RE Probit	-0.0402	0.0024	-0.0113	0.0012	0.0231	0.0025
2SCML	-0.0178	0.0389	-0.0122	0.0160	-0.0183	0.0200
2SCMLR	-0.0288	0.0352	-0.0235	0.0145	-0.0267	0.0165
AGLS	-0.0321	0.0142	-0.0368	0.0160	-0.0097	0.0281
AGLSR	-0.0599	0.0135	-0.0779	0.0126	-0.1034	0.0176
2SIV	-0.0823	0.0078	-0.0826	0.0085	-0.1112	0.0132
2SIVR	-0.0217	0.0122	-0.0315	0.0141	-0.0164	0.0150
Partial R2	0.0395		0.0469		0.0333	

$\rho=0,5$	$\lambda=0$		$\lambda=0,1$		$\lambda=0,5$	
Estimator	Bias	MSE	Bias	MSE	Bias	MSE
OLS	-0.0499	0.0028	-0.0410	0.0021	-0.0347	0.0015
FE	-0.0543	0.0046	-0.0438	0.0032	-0.0332	0.0020
2SLS	-0.0443	0.0038	-0.0517	0.0042	-0.0570	0.0051
2SLSF	-0.0720	0.0492	-0.0712	0.0434	-0.0504	0.0676
Probit	-0.0257	0.0014	-0.0132	0.0012	0.0379	0.0032
RE Probit	-0.0337	0.0019	-0.0226	0.0015	0.0258	0.0025
2SCML	-0.0200	0.0029	-0.0231	0.0034	-0.0162	0.0045
2SCMLR	-0.0284	0.0032	-0.0322	0.0037	-0.0265	0.0047
AGLS	-0.0330	0.0035	-0.0174	0.0034	-0.0138	0.0043
AGLSR	-0.0722	0.0068	-0.0731	0.0067	-0.1068	0.0128
2SIV	-0.0750	0.0065	-0.0870	0.0081	-0.1135	0.0133
2SIVR	-0.0404	0.0040	-0.0285	0.0032	-0.0241	0.0036
Partial R2	0.2028		0.2015		0.1387	

Notes: As table 1, except that individual effects are Chi-squared(1) distributed.

Table 5. Monte Carlo Results for Estimation of Marginal Effects of Time Invariant Regressor in a Panel, 200 observations. Individual Effects depending quadratically on Z-residual.

$\rho=0.05$	$\lambda=0$		$\lambda=0.1$		$\lambda=0.5$	
Estimator	Bias	MSE	Bias	MSE	Bias	MSE
OLS	-0.0096	0.0006	0.0003	0.0005	-0.0114	0.0009
FE	-0.0114	0.0045	0.0008	0.0045	-0.1344	0.0212
2SLS	0.0697	0.3243	0.0079	0.1916	-0.2263	3.5122
2SLSF	0.1524	22.9050	-0.3943	13.8899	-2.3906	6.5399
Probit	0.0002	0.0007	0.0147	0.0009	0.0427	0.0027
RE Probit	-0.0106	0.0007	0.0032	0.0007	0.0302	0.0017
2SCML	0.0708	0.2968	0.0116	0.1671	-0.1871	4.6373
2SCMLR	0.0553	0.2728	0.0048	0.1499	-0.1939	-0.1522
AGLS	0.2349	3.0041	-0.0261	2.1213	-0.0008	0.0030
AGLSR	0.2111	2.1373	-0.0746	2.0797	-0.0099	0.0025
2SIV	-0.0320	0.0055	-0.0238	0.0051	-0.0063	0.0009
2SIVR	0.2115	2.2518	-0.0042	0.2736	-0.0011	0.0020
Partial R2	0.0067		0.0083		0.0065	

$\rho=0.2$	$\lambda=0$		$\lambda=0.1$		$\lambda=0.5$	
Estimator	Bias	MSE	Bias	MSE	Bias	MSE
OLS	-0.0100	0.0006	-0.0056	0.0007	-0.0077	0.0007
FE	-0.0092	0.0033	-0.0170	0.0050	-0.1298	0.0201
2SLS	-0.0213	0.0154	-0.0221	0.0148	-0.0835	0.0222
2SLSF	0.0338	1.5530	-0.1511	2.1272	-1.5668	2.6561
Probit	0.0013	0.0006	0.0086	0.0007	0.0474	0.0032
RE Probit	-0.0094	0.0006	-0.0027	0.0006	0.0351	0.0022
2SCML	-0.0113	0.0138	-0.0074	0.0173	-0.0423	0.0241
2SCMLR	-0.0188	0.0130	-0.0210	0.0166	-0.0499	0.0233
AGLS	-0.0054	0.0159	-0.0146	0.0237	-0.0523	0.0253
AGLSR	-0.0081	0.0147	-0.0311	0.0170	-0.1214	0.0223
2SIV	-0.0063	0.0020	-0.0070	0.0018	-0.0442	0.0034
2SIVR	-0.0025	0.0158	-0.0135	0.0186	-0.0561	0.0197
Partial R2	0.0444		0.0447		0.0321	

$\rho=0.5$	$\lambda=0$		$\lambda=0.1$		$\lambda=0.5$	
Estimator	Bias	MSE	Bias	MSE	Bias	MSE
OLS	-0.0131	0.0007	-0.0029	0.0004	-0.0196	0.0010
FE	-0.0146	0.0030	-0.0023	0.0034	-0.1146	0.0156
2SLS	-0.0124	0.0020	-0.0193	0.0023	-0.0741	0.0084
2SLSF	-0.0228	0.0532	-0.0097	0.0561	-0.6144	0.4351
Probit	0.0015	0.0006	0.0130	0.0008	0.0379	0.0026
RE Probit	-0.0094	0.0006	0.0013	0.0006	0.0252	0.0018
2SCML	0.0018	0.0019	-0.0051	0.0023	-0.0267	0.0048
2SCMLR	-0.0094	0.0018	-0.0158	0.0023	-0.0365	0.0051
AGLS	-0.0008	0.0030	-0.0101	0.0023	-0.0487	0.0051
AGLSR	-0.0099	0.0025	-0.0336	0.0029	-0.1246	0.0168
2SIV	-0.0063	0.0009	-0.0031	0.0008	-0.0336	0.0026
2SIVR	-0.0011	0.0020	-0.0134	0.0019	-0.0693	0.0068
Partial R2	0.2024		0.2050		0.1473	

Notes: As table 1, except that individual effects affect z quadratically.

Table 6. Monte Carlo Results for Estimation of Marginal Effects of Time Invariant Regressor in a Panel, 500 observations, Real Data.

Estimator	$\lambda=0$		$\lambda=0.1$		$\lambda=0.5$	
	Bias	MSE	Bias	MSE	Bias	MSE
OLS	-0.0027	0.0002	0.0119	0.0003	0.0563	0.0033
FE	0.0264	0.0066	0.0295	0.0086	0.0161	0.0066
2SLS	-0.0038	0.0002	-0.0011	0.0003	0.0004	0.0003
2SLSF	0.0266	0.0066	0.0295	0.0087	0.0155	0.0066
Probit	-0.0098	0.0003	-0.0245	0.0007	-0.0697	0.0050
RE Probit	-0.0043	0.0002	0.0119	0.0003	0.0584	0.0036
2SCML	-0.0034	0.0002	-0.0006	0.0003	0.0005	0.0003
2SCMLR	-0.0053	0.0003	-0.0020	0.0003	-0.0010	0.0004
AGLS	0.0185	0.0031	0.0121	0.0063	0.0280	0.0045
AGLSR	-0.0077	0.0034	-0.0174	0.0044	-0.0125	0.0034
2SIV	0.0102	0.0002	0.0090	0.0002	0.0108	0.0002
2SIVR	0.0763	0.0574	0.0160	0.0590	-0.0033	0.0605
Partial R2	0.5079		0.4986		0.4446	
True Marg.Eff.	-0.0252		-0.0252		-0.0252	
Share Positive	0.2999		0.3003		0.2948	

Estimator	$\lambda=0$		$\lambda=0.1$		$\lambda=0.5$	
	Bias	MSE	Bias	MSE	Bias	MSE
OLS	-0.0012	0.0002	0.0125	0.0003	0.0548	0.0031
FE	0.0232	0.0068	0.0123	0.0055	0.0083	0.0051
2SLS	-0.0055	0.0035	-0.0001	0.0034	0.0033	0.0049
2SLSF	0.0235	0.0069	0.0123	0.0057	0.0078	0.0052
Probit	-0.0114	0.0003	-0.0252	0.0008	-0.0681	0.0048
RE Probit	-0.0022	0.0002	0.0126	0.0003	0.0567	0.0034
2SCML	-0.0050	0.0035	0.0006	0.0035	0.0041	0.0051
2SCMLR	-0.0135	0.0049	-0.0050	0.0048	-0.0007	0.0069
AGLS	0.0010	0.0005	0.0065	0.0001	0.0034	0.0004
AGLSR	0.0009	0.0004	0.0052	0.0003	0.0039	0.0003
2SIV	0.0095	0.0001	0.0109	0.0002	0.0104	0.0001
2SIVR	0.0743	0.7411	-0.0123	0.8179	-0.0773	1.2074
Partial R2	0.0368		0.0398		0.0320	
True Marg.Eff.	-0.0252		-0.0252		-0.0252	
Share Positive	0.2966		0.29596		0.2898	

Notes: As table 1, but now with real regressors from data for 500 working women. z is years of education and exogenous regressors are age and a dummy for being a white collar worker. In the top table the instruments are individual means over time of exogenous regressors and indicators for whether individuals are affected by 2 school reforms. In the bottom table only school reforms are used. See text for description and specific design.