



Chalmers Publication Library

Focus on magnetoplasmonics

This document has been downloaded from Chalmers Publication Library (CPL). It is the author's version of a work that was accepted for publication in:

New Journal of Physics (ISSN: 1367-2630)

Citation for the published paper: Armelles, G. ; Dmitriev, A. (2014) "Focus on magnetoplasmonics". New Journal of Physics, vol. 16

http://dx.doi.org/10.1088/1367-2630/16/4/045012

Downloaded from: http://publications.lib.chalmers.se/publication/199170

Notice: Changes introduced as a result of publishing processes such as copy-editing and formatting may not be reflected in this document. For a definitive version of this work, please refer to the published source. Please note that access to the published version might require a subscription.

Chalmers Publication Library (CPL) offers the possibility of retrieving research publications produced at Chalmers University of Technology. It covers all types of publications: articles, dissertations, licentiate theses, masters theses, conference papers, reports etc. Since 2006 it is the official tool for Chalmers official publication statistics. To ensure that Chalmers research results are disseminated as widely as possible, an Open Access Policy has been adopted. The CPL service is administrated and maintained by Chalmers Library.

New Journal of Physics

The open access journal at the forefront of physics

Deutsche Physikalische Gesellschaft $igoplus DPG \mid {f IOP}$ Institute of Physics

Editorial

Focus on magnetoplasmonics

Gaspar Armelles¹ and **Alexandre Dmitriev**²

 ¹ IMM-Instituto de Microelectrónica de Madrid (CNM-CSIC), Isaac Newton 8, PTM, E-28760, Tres Cantos, Madrid, Spain
 ² Department of Applied Physics, Chalmers University of Technology, Göteborg 42196, Sweden

E-mail: gaspar@imm.cnm.csic.es and alexd@chalmers.se

Received 26 February 2014 Accepted for publication 26 February 2014 Published 15 April 2014 *New Journal of Physics* **16** (2014) 045012 doi:10.1088/1367-2630/16/4/045012

Abstract

In this 'focus on' collection, a snapshot of the new, fast emerging field of magnetoplasmonics is presented. The research in the field deals with the combination of plasmonics and magnetism to elucidate the fundamentals of spin–plasmon interactions and reach new functionalities such as the enhancement of magneto-optical activity in various materials, active control of plasmons with weak magnetic fields, magnetoplasmonics-based bio- and chemical sensing, magnetophotonic and magnetoplasmonic crystals as modulators of light transmission and reflection, and many others.

Keywords: magnetoplasmonics, magneto-optics, nanoplasmonics

In the last few years, much effort has been devoted to the study of the properties of collective electromagnetic excitations with a formidable ability to couple free-space electromagnetic radiation to the nanoscale, and to strongly localize and enhance the corresponding electromagnetic fields. These excitations, known as surface plasmons, play a very prominent role in the optical response of the large range of nanoscale material systems, from metals to graphene. Their relevance to light–matter interactions was recognized more than a century ago, but it was not until the strong recent development of nanotechnology and our ability to control

Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

materials at the nanoscale that plasmonics has become a burgeoning research field, with applications spanning energy harvesting, telecoms and sensing [1].

To achieve new functionalities, the combination of plasmons with other material properties has become increasingly appealing. In particular, the field dubbed 'magnetoplasmonics' combines magnetism and plasmonics to develop several novel concepts in materials science [2]. To name a few, these are: the fundamental studies of plasmon–spin interactions in various spectral ranges; the enhancement of magneto-optical activity in materials, including diamagnetics, graphene and others; active control of plasmons with weak magnetic fields; magnetoplasmonics-based bio- and chemical sensing; and magnetophotonic and magnetoplasmonic crystals as modulators of light transmission and reflection.

One of the mentioned lines of research in magnetoplasmonics concerns the studies of new ways to control the properties of surface plasmons using magnetic fields (figure 1(a)). Its effect was first analyzed in the early 1970s on structures made from highly doped semiconductors, where surface plasmons occur in the far-infrared region. It was shown that depending on the relative orientation of the surface plasmon wave vector and the applied magnetic field, the magnetic field could control properties of surface plasmons such as, for example, propagation or localization. This versatility makes the magnetic field very attractive for the development of new active devices, but, unfortunately, for pure noble-metal-based structures, the magnetic field required to achieve proper control of surface plasmon properties is too high for application purposes. To decrease this magnetic field, the use of complex structures made from ferromagnetic materials and noble metals has been proposed. These systems exhibit simultaneously magnetic and plasmonic properties. By an adequate choice of the internal structure of the system, the plasmon properties can be controlled by a (low) magnetic field. The plasmon enhancement of magneto-optical (MO) activity towards optical isolation, acoustomagnetoplasmonics or plasmon-mediated MO transparency are explored [3]. There is a move towards ultrafast magnetoplasmonics with such hybrid plasmon-ferromagnet thin films systems for the eventual on-chip control of light [4].

Another facet of magnetoplasmonics is the focus on localized plasmons in magnetic nanostructures or in the combined magnetic–nanoplasmonic architectures (figure 1(b)). Here, in contrast to heavily damped propagating surface plasmons in magnetic metals, strong effects on magneto-optics are present due to plasmon-induced phase modification of the reflected or transmitted light. As a consequence, the enhancement of MO effects [5] and the large tunability of the magneto-optical response [6] are achieved. In practice, magnetoplasmonics start to play a prominent role in the design of next-generation memory storage technology. The hard disk drive industry is facing a major challenge in continuing to provide increased areal density, driven by the ever-increasing data storage requirements. The heat-assisted magnetic recording approach provides a combination of high coercive field magnetic materials with local heating by a plasmon nanoantenna [7]. This approach currently allows up to record-breaking 1 Tb inch⁻² storage densities.

In this 'focus on' collection, a snapshot of the field of magnetoplasmonics is presented. The contributions cover both theoretical and experimental investigations in this fast developing field. In particular, the research areas discussed include: the optical response of subwavelength magnetoplasmonic nanowire gratings [8], theoretical and experimental evaluation of plasmonic and magnetoplasmonic interferometers as refractometric-based sensors [9], the magnetic-field-induced wave-vector modulation and field distribution of surface plasmon polaritons in dielectric/metal and dielectric/metal/dielectric systems [10], and the manipulation of the



Figure 1. (a) Modulation of a propagating surface plasmon by the applied magnetic field when the ferromagnetic layer (gray) is adjacent to plasmon-supporting noble metal (light brown); (b) tunable Kerr and Faraday effects in nanoferromagnets with localized plasmons—the scattering scanning near-field optical microscopy (s-SNOM) experimentally mapped enhanced electromagnetic fields of the excited localized plasmon resonance in the Ni nanodisk are depicted by the colored landscape.

transverse magneto-optical Kerr effect in periodic metal-dielectric hybrid structuresmagnetoplasmonic crystals [11].

References

- [1] For recent reviews—see Stockman M I 2011 Opt. Express 19 22029
 Atwater H et al 2010 Nat. Mater. 9 205
 Konstantatos G et al 2010 Nat. Nanotechnol. 5 391
- [2] Armelles G et al 2013 Adv. Opt. Mater. 1 10
- [3] Temnov V V et al 2010 Nat. Photon. 4 107
 Belotelov V I et al 2011 Nat. Nanotechol. 6 370
 Temnov V V et al 2013 Nat. Commun. 4 1468
 Chin J Y et al 2013 Nat. Commun. 4 1599
 Belotelov V I et al 2013 Nat. Commun. 4 2128
- [4] Temnov V V 2012 Nat. Photon. 6 728
- [5] Gonzalez-Diaz J B et al 2008 Small 4 202
 Banthi J C et al 2012 Adv. Mater. 24 OP36
- [6] Bonanni V et al 2011 Nano Lett. 11 5333
 Maccaferri N et al 2013 Opt. Express 21 9875
 Maccaferri N et al 2013 Phys. Rev. Lett. 111 167401
- [7] Stipe B C *et al* 2010 *Nat. Photon.* **4** 484
 Challener W A *et al* 2009 *Nat. Photon.* **3** 220
- [8] Marinchio H et al 2014 New J. Phys. 16 015007
- [9] Martín-Becerra D et al 2013 New J. Phys. 15 085021
- [10] Torrado J F et al 2013 New J. Phys. 15 075025
- [11] Pohl M et al 2013 New J. Phys. 15 075024