

Predicted multiply imaged X-ray AGNs in the XXL survey

F. Finet,^{1,2,3★} A. Elyiv,^{2,4,5} O. Melnyk,^{2,6} O. Wertz,² C. Horellou⁷ and J. Surdej^{2†}

¹Aryabhata Research Institute of Observational Sciences (ARIES), Manora Peak, Nainital 263 129, Uttarakhand, India

²Extragalactic Astrophysics and Space Observations (AEOS), University of Liège, Allée du 6 Août, 17 (Sart Tilman, Bât. B5c), B-4000 Liège, Belgium

³National Astronomical Observatory of Japan (NAOJ), 650 N. A'Ohoku place, Hilo, HI 96720, USA

⁴Main Astronomical Observatory, Academy of Sciences of Ukraine, 27 Akademika Zabolotmoho St., UA-03680 Kyiv, Ukraine

⁵Dipartimento di Fisica e Astronomia, Università di Bologna, Viale Berti Pichat 6/2, I-40127 Bologna, Italy

⁶Astronomical Observatory, Kyiv National University, 3 Observatorna St., UA-04053 Kyiv, Ukraine

⁷Department of Earth & Space Sciences, Chalmers University of Technology, Onsala Space Observatory, SE-439 92 Onsala, Sweden

Accepted 2015 June 23. Received 2015 June 22; in original form 2014 March 20

ABSTRACT

We estimate the incidence of multiply imaged active galactic nuclei (AGNs) among the optical counterparts of X-ray selected point-like sources in the XXL field. We also derive the expected statistical properties of this sample, such as the redshift distribution of the lensed sources and of the deflectors that lead to the formation of multiple images, modelling the deflectors using both spherical and ellipsoidal singular isothermal mass distributions. We further assume that the XXL survey sample has the same overall properties as the smaller *XMM*-COSMOS sample restricted to the same flux limits and taking into account the detection probability of the XXL survey. Among the X-ray sources with a flux in the [0.5–2] keV band larger than 3.0×10^{-15} erg cm⁻² s⁻¹ and with optical counterparts brighter than an *r*-band magnitude of 25, we expect ~ 20 multiply imaged sources. Out of these, ~ 16 should be detected if the search is made among the seeing-limited images of the X-ray AGN optical counterparts and only one of them should be composed of more than two lensed images. Finally, we study the impact of the cosmological model on the expected fraction of lensed sources.

Key words: gravitational lensing: strong – galaxies: active – cosmological parameters – X-rays: galaxies.

1 INTRODUCTION

The XXL survey,¹ carried out by the space-based X-ray observatory *XMM-Newton*, spans over $\sim 2 \times 25$ square degrees with near 10 ks exposure in each field and is expected to lead to the detection of $\sim 25\,000$ active galactic nuclei (AGNs) down to a limiting flux 10^{-15} erg cm⁻² s⁻¹ in the [0.5–2] keV soft X-ray band (Pierre et al., in preparation). These X-ray data are complemented by multi-wavelength data obtained with the Canada–France–Hawaii Telescope Legacy Survey (CFHTLS) and with the Blanco telescope (Blanco/South Pole Telescope Cosmology Survey) in the (near-) optical *u'*, *g*, *r*, *i* and *z* bands, down to a limiting AB magnitude of ~ 25 . Besides the multi-band imaging of the XXL fields, there is a very large on-going effort to obtain optical spectra of XXL sources, through either the matching of existing survey catalogues or dedicated spectroscopic surveys. Among these spectroscopic data acquisition programmes, the VIMOS Public Extragalactic Redshift Survey (Guzzo & Le Fèvre 2010) covers most of the northern field, the southern field being covered using the AAOmega multi-object spec-

trograph on the Anglo-Australian Telescope, an instrument used for the Galaxy And Mass Assembly project (Driver et al. 2009).

The completeness of this multi-wavelength data base over the entire XXL field provides a unique sample to search for multiply imaged AGNs. We have thus initiated such a search among the optical counterparts of point-like sources in the soft X-ray band. Besides the scientific interest provided by each multiply imaged source, the goal of this project is to construct a statistically clean sample of lensed sources that will be used, in combination with samples of multiply imaged sources from other recent surveys, to independently constrain the cosmological model.

The choice of the soft X-ray point-like sources is motivated by the higher sensitivity of *XMM-Newton* in this band. Furthermore, this spectral band should contain a larger fraction of type-I AGNs than the hard X-ray. On average, type-I AGNs with a detectable optical counterpart are expected to have a higher redshift than type-II AGNs (more absorbed in the visible and thus more difficult to detect in the optical at high redshift). As higher redshift sources have a higher probability of being lensed, this population is more likely to undergo gravitational lensing with the formation of multiple images. The better angular resolution achievable in the optical domain will allow us to unravel the multiply imaged sources.

A search for gravitational lenses among the optical counterparts of X-ray sources has already been carried out for a subset of the

* E-mail: finet@astro.ulg.ac.be

† Also, Directeur de Recherche honoraire du F.R.S.-FNRS.

¹ <http://ifru.cea.fr/xxl>

XXL field, the *XMM-Newton* Large-Scale Structure (XMM-LSS) field (Elyiv et al. 2013). For this smaller field, visual inspection of all optical counterparts has been done in order to identify the multiply imaged AGN candidates that are now awaiting spectroscopic confirmation. The search for lensed sources in the larger XXL field is in progress.

In this paper, we present a prospective analysis of the lensed AGN population detectable within the XXL survey, as well as a study of their expected statistical properties. In order to perform this analysis, we have reformulated the mathematical formalism to study the statistical aspects of gravitational lensing, basing the statistical formalism on the observables of the source population.

This paper is structured as follows. We present the mathematical approach in Section 2. Namely, we derive the expression allowing us to calculate the probability for a source to be lensed with the formation of multiple images, modelling the deflector population by means of a spherical mass distribution and then taking into account the internal ellipticity of the deflector mass distribution. We explain how this expression may be averaged over the entire population of sources detected in the survey, thanks to the source joint probability density, with which we derive the expression of the expected fraction of lensed sources in a survey, as well as the expected redshift distribution of the lensed sources and that of the deflectors. Our simulations also account for the inability of the ground-based CFHT and Blanco telescopes to resolve multiple images with too small angular separations.

In Section 3, we present the observational constraints used to estimate the expected properties of the XXL population in the X-ray and optical domains: these were inferred from the deeper (but smaller) XMM-COSMOS field (Brusa et al. 2010). Finally, in Section 4 we present our results, i.e. the expected number of multiply imaged sources in the XXL and the XMM-LSS fields, as well as the expected statistical properties of these lensed sources. We also investigate the fraction of lensing events as a function of their number of lensed images and we investigate how the fraction of multiply imaged sources changes as a function of the cosmological mass density parameter, Ω_m .

2 MATHEMATICAL FORMALISM

2.1 Lensing optical depth

Multiple images due to gravitational lensing occur when light rays emitted from a background source are deviated towards the observer by a foreground deflector located near the line of sight. In our case, the amplified lensed sources have to be above the survey flux limit in both the X-ray and the optical domains, and the lensed images have to be resolved in the latter. The probability for a source to be lensed depends on its redshift, its X-ray flux and r -band magnitude. In this section, we derive an expression to calculate the probability for a source to undergo a gravitational lensing event as a function of its redshift and its X-ray flux only, considering the minimal angular separation resolvable in the r band. We will analyse the validity of this simplification in Section 5 by formally including the r magnitude in the calculations.

Let us consider a source with redshift z_s , with an observed flux f_X in the X-ray band, and a lens with a mass distribution characterized by a set of parameters M' , located at an intermediate position along the line of sight at redshift z_d . In the lens plane perpendicular to the line of sight, we can define an area $\Sigma(z_s, z_d, f_X, M')$ centred on the source projected on the lens plane, called the lensing cross-section in which the presence of a deflector leads to the detection of multiple images by the observer (where the multiple images are

resolved in the r -band CCD frames). The lensing cross-section is a function of the redshifts of the source and the deflector, the X-ray flux f_X of the source and the deflector mass distribution parameters M' (some mass distributions are more efficient at deflecting light rays and thus have a larger lensing cross-section).

The probability $d\tau(z_s, z_d, f_X, M')$ for this source to be multiply imaged due to the presence of a deflector in the redshift range $[z_d, z_d + dz_d]$ is given by (Turner, Ostriker & Gott 1984)

$$d\tau = (1 + z_d)^3 n_D(z_d, M') \frac{cdt}{dz_d} \Sigma(z_s, z_d, f_X, M') dz_d, \quad (1)$$

where $n_D(z_d, M')$ is the volume density in the comoving reference frame of deflectors characterized by the mass parameters M' . The quantity cdt/dz_d represents the infinitesimal light-distance element at redshift z_d per deflector redshift unit, which, in a flat expanding Friedmann-Lemaître-Robertson-Walker (FLRW) universe, is given by (e.g. Peebles 1993)

$$\frac{cdt}{dz_d} = \frac{c}{H_0(1 + z_d)} [(1 + z_d)^3 \Omega_m + (1 - \Omega_m)]^{-1/2}, \quad (2)$$

where Ω_m is the present-day value of the cosmological mass density parameter.

The envelope of the lensing cross-sections at different deflector redshifts z_d defines the *lensing volume* in which the presence of a deflector leads to the detection of multiple lensed images of the background source. The probability $\tau(z_s, f_X, M')$ for a source to be lensed with the formation of multiple images can be calculated by integrating equation (1) over all values of the deflector redshift z_d , which leads to

$$\begin{aligned} \tau(z_s, f_X, M') &= \int_0^{z_s} (1 + z_d)^3 n_D(z_d, M') \frac{cdt}{dz_d} \Sigma(z_s, z_d, f_X, M') dz_d. \end{aligned} \quad (3)$$

The definition of $\tau(z_s, f_X, M')$ in equation (3) corresponds to an optical depth, which for small values can be assimilated to a probability. For this reason, we refer to $\tau(z_s, f_X, M')$ as the source *lensing optical depth* or *lensing probability*, without distinction.

In equation (3), the comoving density $n_D(z_d, M')$ of deflectors assumes one type of deflectors with similar characteristics defined by the parameters M' . Considering the mass distribution to be characterized by means of deflectors with a central velocity dispersion σ , in the range $[\sigma, \sigma + d\sigma]$, $n_D(z_d, \sigma)$ may be expressed by means of the velocity dispersion function (VDF) of galaxies $\Phi_\sigma(\sigma, z_d)$,

$$n_D(z_d, \sigma) = \Phi_\sigma(\sigma, z_d) d\sigma. \quad (4)$$

To take into account the contribution of all galaxies with different central velocity dispersions, equation (3) has to be integrated over σ . The VDF in equation (4) can be either measured directly or inferred from the luminosity function (LF) of the deflecting galaxies, using the mean Faber–Jackson or Tully–Fisher relationship, depending on the type of galaxies considered. However, Sheth et al. (2003) have shown that neglecting the dispersion of the Faber–Jackson relationship leads to a wrong estimate of the VDF. We thus use the VDF determined directly through observations (thereby following Sheth et al. 2003; Mitchell et al. 2005; Choi, Park & Vogeley 2007; Chae 2010). Sheth et al. (2003) and Choi et al. (2007) have shown that the VDF of early- and late-type galaxies is well fitted by the modified Schechter function

$$\Phi_\sigma(\sigma, z_d) d\sigma = \Phi_* \left(\frac{\sigma}{\sigma_*} \right)^\alpha \exp \left(- \left(\frac{\sigma}{\sigma_*} \right)^\beta \right) \frac{\beta}{\Gamma(\alpha/\beta)} \frac{d\sigma}{\sigma}, \quad (5)$$

where Φ_* and σ_* are the characteristic number density and central velocity dispersion, respectively, α and β are the slope coefficients of the VDF for low and high values of σ , respectively, and where $\Gamma(x)$ is the Gamma function.

Although less numerous, early-type galaxies are much more efficient deflectors than late-type ones, which tend to form multiple images with smaller angular separation. Late-type galaxies contribute by less than 10 per cent to the gravitational lensing events in a typical sample of lensed AGNs selected in the optical or near-infrared (Fukugita & Turner 1991; Maoz & Rix 1993; Keeton, Kochanek & Falco 1998; Kochanek et al. 2000), although in radio-selected samples, thanks to the better angular resolution of the survey, the fraction of lensing events formed by late-type deflectors may be higher (e.g. the CLASS survey where at least 5 of 22 deflectors are late-type galaxies; see Browne et al. 2003). In the present work, because the multiple images will be searched for in the SDSS r band, we consider the deflector population to be only composed of early-type galaxies, and we will study in Section 4 the validity of this assumption.

Constraints from strong-lensing statistics on the evolution of the VDF of early-type galaxies show very little evolution or are consistent with a no-evolution assumption (see e.g. Chae 2003, 2010; Ofek, Rix & Maoz 2003; Oguri et al. 2012). Consequently, throughout this work we assume the deflector VDF to be constant with redshift in the comoving reference frame, or in other words, that there is no impact of the evolution of the deflector population on the VDF and we use the value of the VDF parameters determined in the local Universe by Choi et al. (2007), i.e.

$$[\Phi_*, \sigma_*, \alpha, \beta] = [8 \times 10^{-3} h^3 \text{Mpc}^{-3}, 161 \text{ km s}^{-1}, 2.32, 2.67].$$

The lensing cross-section in equation (1) depends on the deflector mass distribution parameters M' . For a deflector mass distribution modelled as a singular isothermal sphere (SIS; a spherical mass distribution with a volume density scaling as r^{-2}), the mass distribution is characterized by the line-of-sight velocity dispersion σ and the lensing cross-section $\Sigma(z_s, z_d, f_X, \sigma)$ can be defined as [see Turner et al. (1984) for definition and Claeskens (1999) for the formalism followed in this paper]

$$\Sigma(z_s, z_d, f_X, \sigma) = b_0^2(z_s, z_d, \sigma) \iint_{S_y} \frac{N_{f_X}(f_X/A(\mathbf{y}))}{N_{f_X}(f_X)} d\mathbf{y}, \quad (6)$$

where we have introduced the source vector $\mathbf{y} = (y_1, y_2)$ which Cartesian coordinates are projected on the deflector plane, and normalized to the scale factor b_0 (i.e. the Einstein radius in the deflector plane). $A(\mathbf{y})$ is the total amplification of the lensed images formed for a source located at the position \mathbf{y} , i.e. the sum of the amplification moduli of each lensed image and $N_{f_X}(f_X)$ is the differential number counts function (DNCF) as a function of the source flux f_X in the X-ray band. For an SIS deflector, the scale factor is given by (see e.g. Claeskens 1999)

$$b_0(\sigma, z_d, z_s, \text{UM}) = 4\pi \left(\frac{\sigma}{c}\right)^2 \frac{D_{\text{OD}} D_{\text{DS}}}{D_{\text{OS}}}, \quad (7)$$

where D_{OD} , D_{DS} and D_{OS} represent the different angular diameter distances between the observer, the deflector and the source, c is the speed of light and $\text{UM} = (\Omega_m, H_0)$ is a set of parameter values characterizing the universe model as a flat expanding FLRW one.

The ratio $N_{f_X}(f_X/A(\mathbf{y}))/N_{f_X}(f_X)$ in equation (6) is known as the *amplification bias* (Turner 1980; Turner et al. 1984; Fukugita & Turner 1991). It is introduced to take into account a favourable bias when estimating the lensing probability in a flux-limited sample induced by the amplification phenomenon. The amplification

may lead to the inclusion of sources in a flux-limited sample that are intrinsically fainter than the flux limit but have undergone a gravitational lensing amplification. Since the ratio of the DNCF in equation (6) at two different X-ray flux levels (i.e. f_X and $f_X/A(\mathbf{y})$) is likely to be slightly dependent on the source redshift, it would certainly be more accurate to consider the redshift dependence of the DNCF. However, at the time of this work, the XXL redshifts are not yet available and we do not have access to a source sample large enough to constrain the redshift dependence of the DNCF (the source sample used is described in Section 3). Consequently, we consider the value of the DNCF ratios averaged over the whole redshift range. Because of the presence of $A(\mathbf{y})$ in the amplification bias, the expression of $\Sigma(z_s, z_d, f_X, \sigma)$ is intrinsically linked to the amplification map of the deflector defining the amplification as a function of the source position, itself determined by the mass distribution of the deflector. It is out of the scope of this paper to explain this dependence in detail and the reader may refer to e.g. Hezaveh & Holder (2011) for some further description. Nevertheless, because of the presence of the total amplification A in its argument, the calculation of the amplification bias in equation (6) necessitates the knowledge of $N_{f_X}(f_X)$ for sources fainter than the survey limiting flux. This may be estimated either by extrapolating $N_{f_X}(f_X)$ or by considering a parent source sample accessing fainter fluxes. In this work, we will consider a deeper source sample described in Section 3.

Finally, S_y in equation (6) represents the area in the y plane, centred on the deflector, in which the presence of a source leads to a *lensing event*, i.e. the lensing cross-section, normalized to the scale factor.

The *lensing event* may be defined in different ways. It may refer to the formation of multiple images or to the formation of a given number of lensed images, or the formation of lensed images with an angular separation sufficiently large to be detected in a survey. Depending on the definition adopted for the lensing event, the integration domain S_y differs and, consequently, this leads to a different definition of the lensing cross-section Σ in equation (6) and of the lensing optical depth in equation (3). In most cases, the integration in equation (6) must be performed numerically.

When modelling the deflector mass distribution by means of an SIS profile, the lensing events can lead to the formation of two images at maximum. We define $\tau_{\text{SIS}}(z_s, f_X)$ as the probability for a source to be multiply imaged when the deflectors are modelled by an SIS mass distribution. $\tau_{\text{SIS}}(z_s, f_X)$ is thus calculated by inserting equations (4)–(7) into equation (3) and by performing the integration over σ . The integration over σ can be performed analytically, leading to the following expression (Turner et al. 1984; Mitchell et al. 2005 for derivation using the VDF)

$$\tau_{\text{SIS}}(z_s, f_X) = \Phi_* \frac{\Gamma((\alpha + 4)/\beta)}{\Gamma(\alpha/\beta)} \int_0^{z_s} (1 + z_d)^3 \frac{cdt}{dz_d} \Sigma_{\text{SIS}}(z_s, z_d, f_X, \sigma_*) dz_d. \quad (8)$$

$\Sigma_{\text{SIS}}(z_s, z_d, f_X, \sigma_*)$ represents the lensing cross-section defined by equation (6), for $\sigma = \sigma_*$, when considering the area S_y for the case of an SIS deflector. The integration in equation (8) must be performed numerically.

The introduction of an internal ellipticity in the mass distribution used to model the deflectors allows us to account for the formation of more than two lensed images. This is the reason why the singular isothermal ellipsoid (SIE) mass distribution has been introduced (Kormann, Schneider & Bartelmann 1994). The ellipticity parameter in the SIE profile is the axial ratio q of the deflector mass

projected on the deflector plane and the mass distribution parameters M' are now σ and q . As for the case of the SIS deflector, we can define the probability $\tau_{\text{SIE}}(z_s, f_X)$ for a source to be lensed with the formation of multiple images, irrespectively of the number of the lensed images, when modelling the deflector population with SIE mass distributions (Huterer, Keeton & Ma 2005). On the other hand, we may also define the probability $\tau_{\text{SIE},i}(z_s, f_X)$ for a source to be lensed with the formation of i images (with $i = 2, 3$ or 4). As the lensing cross-section now depends on the axial ratio q of the deflector (through the dependence of S_y in equation 6), in order to calculate $\tau_{\text{SIE}}(z_s, f_X)$ we must also integrate equation (1) over the axial ratio q , using an appropriate probability distribution. Furthermore, the deflector density function in equations (3) and (4) must take into account how the deflector population is distributed as a function of both σ and q , and thus we have to introduce the dependence of n_D on the axial ratio q . In equation (4), the density $n_D(\sigma, q)$ of deflectors with a central velocity dispersion in the range $[\sigma, \sigma + d\sigma]$ and an axial ratio in the range $[q, q + dq]$ may be expressed as

$$\begin{aligned} n_D(\sigma, q) &= \Phi_\sigma(\sigma) d_{q|\sigma}(\sigma, q) d\sigma \\ &= \Phi_\sigma(\sigma) d_q(q) d\sigma dq, \end{aligned} \quad (9)$$

where $d_{q|\sigma}(\sigma, q)$ represents the normalized axial ratio distribution for the deflectors with a central velocity dispersion σ and $d_q(q)$ is the marginal normalized distribution as a function of q . The last equality arises if we assume that the deflector distributions as a function of σ and q are mutually independent.² $\Phi_\sigma(\sigma)$ is given by equation (5), as in the case of the SIS mass model. There are pieces of strong evidence from the study of various gravitational lens samples, that elliptical galaxy isophotes and the mass distribution ellipticities are aligned and have well-correlated values (see Koopmans et al. 2006; Sluse et al. 2012 for independent confirmations). Furthermore, as previously mentioned, there is no evidence for strong evolution effects in the VDF of early-type galaxies from lensing surveys. Thanks to these two observational facts, $d_q(q)$ can be estimated from the distribution of the isophotes of early-type galaxies as measured in the local Universe. We therefore use the axial ratio distribution measured by Choi et al. (2007) from a sample of elliptical galaxies in the local Universe.

$\tau_{\text{SIE}}(z_s, f_X)$ can thus be calculated by inserting equations (5) and (9) into equation (1), and by integrating over σ and q . Using equations (6) and (7), and performing the analytical integration over σ , τ_{SIE} can thus be expressed as

$$\begin{aligned} \tau_{\text{SIE}}(z_s, f_X) &= \Phi_* \frac{\Gamma((\alpha + 4)/\beta)}{\Gamma(\alpha/\beta)} \\ &\int_0^{z_s} (1 + z_d)^3 \frac{cdt}{dz_d} \int_0^1 d_q(q) \Sigma_{\text{SIE}}(z_s, z_d, f_X, \sigma_*, q) dq dz_d, \end{aligned} \quad (10)$$

where $\Sigma_{\text{SIE}}(z_s, z_d, f_X, \sigma_*, q)$ represents the lensing cross-section calculated through numerical integration of equation (6), when considering an SIE deflector with an axial ratio q and a central velocity dispersion σ_* . Similarly, the probability $\tau_{\text{SIE},i}(z_s, f_X)$ for a source to be lensed with the formation of i images, when modelling the deflectors with SIE profiles, can be calculated using $\Sigma_{\text{SIE},i}$ in the previous relation and considering in S_y only the area in which a

source should be located in order to lead to the formation of i lensed images.

We have developed *Matlab* toolboxes and libraries to perform the numerical integration in the expressions of $\tau_{\text{SIS}}(z_s, f_X)$ and $\tau_{\text{SIE}}(z_s, f_X)$ from equations (8) and (10), as well as for the calculation of $\tau_{\text{SIE},i}(z_s, f_X)$, taking into account that lensed images angularly too close to each other cannot be resolved in the survey. The numerical integrations are made in two steps. First, we create a data base of the lensing cross-sections in equation (6), considering $b_0 = 1$, ranging over all possible values of f_X and over the ratio $\theta_{\text{mis}}/\theta_E$, where θ_{mis} represents the smallest angular separation for which point-like images of same brightness are resolved in the survey, and also over q for the SIE case. The integration of the double integral in equation (6) is performed by plain Monte Carlo integration, where we randomly generate $\sim 10^6$ source positions \mathbf{y} , and calculate the position and amplification of the lensed images, which contribute to the integral only if the multiple images can be resolved and brighter than the X-ray limiting flux of the survey. In the second step, the lensing optical depths are calculated by integrating equations (8) and (10) using trapezoidal integration and the data base of lensing cross-sections. We have thus all the tools needed for the calculation of the lensing optical depth of a source with known redshift and apparent X-ray flux, when considering a population of deflectors modelled with SIS or SIE mass distributions.

Similar expressions for the lensing optical depths such as $\tau_{\text{SIS}}(z_s, f_X)$, $\tau_{\text{SIE}}(z_s, f_X)$ and $\tau_{\text{SIE},i}(z_s, f_X)$ (equations 8 and 10, respectively), as well as their differential contribution as a function of the deflector redshift $d\tau/dz_d$, have already been derived and used for the analysis of statistical samples of lensed sources, either to constrain the cosmological model (e.g. Turner et al. 1984; Fukugita, Futamase & Kasai 1990; Turner 1990; Fukugita & Turner 1991; Kochanek 1992; Maoz & Rix 1993; Surdej et al. 1993; Cen et al. 1994; Chae et al. 2002; Keeton 2002; Chae 2003; Oguri et al. 2012) or to study the population of deflectors (Keeton 1998; Keeton & Kochanek 1998; Keeton et al. 1998; Kochanek et al. 2000; Chae 2003, 2010; Ofek et al. 2003), some of the work having considered the ellipticity of the mass distribution as well as the effect of external shear on the statistics (Huterer et al. 2005; Oguri & Marshall 2010; Oguri et al. 2012).

The various expressions derived for the optical depth τ and its differential contribution $d\tau/dz_d$ are considered for a single source. This may be averaged over the whole detected population to derive the mean optical depth through the sample as well as the expected redshift distributions of the lensed sources and deflectors. To average over the population of sources, some previous works have made use of the source LF, and integrate over the absolute magnitude of the source population (e.g. Oguri & Marshall 2010; Oguri et al. 2012) which, for the integration process, necessitates the choice of a universe model.

In the next subsection, we propose a slightly different formulation that allows us to average any function over the entire population of sources detected in a survey, where the averaging is done based on the observables, i.e. the distribution in the (z_s, f_X) plane of the detected sources. Because the formalism is based directly on the observed distribution of sources, it naturally takes into account the detection biases of the sources. We apply this method to derive useful expressions such as the average lensing optical depth in a sample, the redshift distributions of the lensed sources and of the deflectors effectively leading to the formation of multiple lensed images.

² Although not strictly justified, this assumption is made because of the lack of observational constraints for the distribution of early-type galaxies in the (σ, q) plane, as well as for calculation time consideration.

2.2 Joint source probability density and fraction of multiply imaged sources

Let us consider a survey characterized by its limiting flux, different biases in the source detection procedure and its angular coverage. Each detected source is characterized by its redshift z_s and its flux f_X in the selected spectral band (in the present case, the [0.5–2] keV band). The detection of a source within the survey may be considered as a random event with respect to the continuous random variables associated with the source redshift and the X-ray flux, respectively.

We can define a probability $P(z_s, f_X)$ that a source detected in the survey is characterized by an observed redshift and flux in the range $[z_s, z_s + dz_s]$ and $[f_X, f_X + df_X]$, respectively. We may then define the joint probability density $d_{\text{obs}}(z_s, f_X)$ spanning over the (z_s, f_X) plane, associated with this random event. $P(z_s, f_X)$ and $d_{\text{obs}}(z_s, f_X)$ are related through

$$P(z_s, f_X) = d_{\text{obs}}(z_s, f_X) dz_s df_X. \quad (11)$$

The random variables z_s and f_X associated with a detected source follow the joint distribution described by $d_{\text{obs}}(z_s, f_X)$. This function contains all the information about the survey and implicitly takes into account the detection biases. For a sufficiently large number of detected sources, the joint probability density $d_{\text{obs}}(z_s, f_X)$ may be directly estimated from the detected source population, by calculating a smoothed histogram of the source distribution in the (z_s, f_X) plane, normalized by the total number N_{AGN} of sources detected within the survey. Ideally, we would have liked to define the source joint probability density $d_{\text{obs}}(z_s, f_X, r)$ in the (z_s, f_X, r) space, where r represents the SDSS r magnitude of the optical counterpart(s) of the X-ray sources. However, because of the small number of sources observed in our reference sample (see Section 3), the quantity $d_{\text{obs}}(z_s, f_X, r)$ could hardly be accurately determined. When analysing the XXL sample however, the number of detected sources should be sufficient to constrain d_{obs} in the 3D space. We shall therefore postpone such a more detailed study until all optical counterparts of the XXL X-ray sources have been identified.

The normalized marginal probability density distributions associated with the random variables z_s and f_X are closely related to the observations. Indeed, the marginal density distribution obtained by integrating $d_{\text{obs}}(z_s, f_X)$ over z_s or f_X represents the normalized source distribution as a function of the flux $N_{f_X}(f_X)/N_{\text{AGN}}$ and the redshift $N_{z_s}(z_s)/N_{\text{AGN}}$, respectively.

The use of $d_{\text{obs}}(z_s, f_X)$ as a weighing function when performing the integration over the entire population of detected sources allows us to calculate the expected mean value of any function of the random variables z_s and f_X . The mathematical expectation $\langle \tau \rangle$ of the lensing optical depth, i.e. the fraction of sources gravitationally lensed within the detected population, can be calculated by integrating $\tau(z_s, f_X)$ over the (z_s, f_X) plane, weighing with $d_{\text{obs}}(z_s, f_X)$, i.e.

$$\langle \tau \rangle = \iint \tau(z_s, f_X) d_{\text{obs}}(z_s, f_X) dz_s df_X. \quad (12)$$

Using the expression of $\tau_{\text{SIS}}(z_s, f_X)$, $\tau_{\text{SIE}}(z_s, f_X)$ or $\tau_{\text{SIE}, i}(z_s, f_X)$ given by equations (8) and (10) for the calculation of the optical depth $\tau(z_s, f_X)$ in equation (12), we are able to calculate the expected fraction of multiply imaged sources, considering a population of deflectors modelled with either the SIS or SIE mass profile. In the latter case, we can also calculate the expected fraction of lensed sources as a function of the number i of lensed images.

Oguri et al. (2008) have also derived an expression for the number of multiply imaged sources using a binning of the redshift–magnitude space (equation 12 in Oguri et al. 2008), and with the number of sources in the bins as a weighing factor. Their expression corresponds to the discrete equivalent of equation (12), integrating in the redshift–magnitude space, rather than the (z_s, f_X) plane. The essential difference between the approach used in these previous works and ours is the use of the source distribution in the (f_X, z_s) rather than using the source LF as a weighing factor in the absolute magnitude–redshift space. Our approach allows us to account directly for the detection bias of the sources and does not necessitate any assumption of a universe model for the calculation of the weighing factor (which is necessary when using absolute magnitudes and the source LF).

2.3 Normalized redshift distributions

It is now straightforward to establish the normalized distribution $w_{z_d}(z_d)$ of the deflector redshifts expected in the XXL field

$$w_{z_d}(z_d) = \frac{1}{\langle \tau \rangle} \iint \left\{ \frac{d\tau}{dz_d}(z_s, z_d, f_X) d_{\text{obs}}(z_s, f_X) \right\} dz_s df_X, \quad (13)$$

where the differential contribution $d\tau/dz_d$ of the redshift z_d to a source lensing optical depth is given by equation (1).

Similarly, the normalized redshift distribution $w_{z_s}(z_s)$ of the lensed sources is given by

$$w_{z_s}(z_s) = \frac{1}{\langle \tau \rangle} \int \tau(z_s, f_X) d_{\text{obs}}(z_s, f_X) df_X. \quad (14)$$

Oguri et al. (2012) have derived by different means an expression for $w_{z_d}(z_d)$ and applied it to the sample of the SDSSQLS (see equation 23 in Oguri et al. 2012). This estimation was done by binning the redshift–magnitude plane and using the number of sources in the different bins as a weighing factor. Their relation corresponds to the discrete equivalent of equation (13), where the integration runs over the redshift–magnitude plane, rather than the (z_s, f_X) one.

In Oguri & Marshall (2010), the authors have derived a different expression for the redshift distribution of the lensed sources (equation 7 in Oguri & Marshall 2010), equivalent to equation (14), except that the integration runs over the absolute magnitude and the weighing function used in the integration corresponds to the expression of the joint probability density expressed in terms of the source LF.

Finally, let us note that Mitchell et al. (2005) have also derived an estimation for the redshift distribution of the lensed sources in the CLASS survey, assuming the DNCF of the sources to be expressed as a single power-law expression, i.e. the amplification bias being thus constant for each source. However, this assumption is very restrictive as QSOs usually show a DNCF that presents a break at a critical magnitude, and needs to be modelled by a double power-law expression.

The use of $d_{\text{obs}}(z_s, f_X)$ allows an easy calculation of $\langle \tau \rangle$ through the survey, and the calculation of the expected normalized distributions $w_{z_s}(z_s)$ and $w_{z_d}(z_d)$, as a function of the redshift of the lensed sources and of the deflectors, respectively. These distributions are calculated without any assumption about the source population, as $d_{\text{obs}}(z_s, f_X)$ may be directly estimated from the observed data, naturally including the observational biases.

In the next section, we present the observational constraints used for the estimation of the joint probability density of the XXL survey.




Figure 1. The left-hand panel displays the distribution in the (z_s, f_X) plane of the XMM-COSMOS source population (Brusa et al. 2010), restricted to $F_{[0.5-2]\text{keV}} > 3 \times 10^{-15} \text{ erg cm}^{-2} \text{ s}^{-1}$ in the [0.5–2] keV band and with an optical counterpart brighter than $r = 25$, for which a redshift estimate is available. This X-ray band flux limit corresponds to a detection probability of 0.5 in the XMM-LSS field, as defined in Elyiv et al. (2012). The X-ray flux is shown on a logarithmic scale. The right-hand panel displays the joint probability density function $d_{\text{obs}}(z_s, f_X)$ in the (z_s, f_X) plane, corresponding to the XMM-COSMOS source population (Brusa et al. 2010), restricted to the X-ray band flux limit of the XMM-LSS fields.

3 OBSERVATIONAL CONSTRAINTS AND d_{obs} DETERMINATION

Ideally, the joint probability density of the XXL sources should be derived by constructing a histogram of the XXL sources in the redshift–X-ray flux plane. However, the cross-correlation between the X-ray and (near-)optical data and the identification of the source type have not yet been performed. The XXL X-ray sources and their associated multi-wavelength data have characteristics similar to those of one of its sub-fields, the XMM-Large Scale Structure (XMM-LSS), covering 10.9 out of the 44.2 square degrees effectively covered by the XXL survey. The XMM-LSS X-ray sources and their associated multi-wavelength data, presented in Chiappetti et al. (2013), have optical counterparts taken from the CFHTLS W1 catalogue, down to the limiting magnitudes $i' \simeq 25$, $r' \simeq 25$, before correction for Galactic extinction.³ The X-ray source classification and properties of the XMM-LSS field have already been determined (Melnik et al. 2013). The survey limiting flux in the soft band is $F_{[0.5-2]\text{keV}} \simeq 3 \times 10^{-15} \text{ erg s}^{-1} \text{ cm}^{-2}$ (with a detection probability of 0.5, as defined in Elyiv et al. 2012). Nevertheless, at the time of this work, the redshift estimate in the XMM-LSS sample is still ongoing.

Therefore, to retrieve the properties of the expected population to be detected within the XXL survey and their optical counterparts, we used data from a deeper field (~ 60 ks exposure, compared to ~ 10 ks for XMM-LSS): the XMM-COSMOS field, to which we apply the flux cuts of the XXL in the X-ray and the optical, that we assume to be similar to those of the XMM-LSS. The XMM-COSMOS survey, covering a contiguous field of 2 square degrees, is described in Brusa et al. (2010), and has a limiting flux of $F_{[0.5-2]\text{keV}} \simeq 5 \times 10^{-16} \text{ erg s}^{-1} \text{ cm}^{-2}$ ($\simeq 1.4 \times 10^{-15}$ considering the flux with 50 per cent effective area coverage; see fig. 6 in Cappelluti et al. 2007). The flux limit at 50 per cent effective area coverage for the XMM-LSS and that for the COSMOS are in good agreement with the exposure time ratios as $3 \times 10^{-15} / 1.4 \times 10^{-15} \sim \sqrt{60 \text{ ks} / 10 \text{ ks}}$.

³ After correction, the limiting magnitudes of the CFHTLS W1 catalogue are (expressed in AB magnitudes and considering a 5σ signal-to-noise ratio) $r \simeq 24.8$ and $i \simeq 24.5$. See <http://xxlmultiwave.pbworks.com/w/page/54613008/Optical> and http://terapix.iap.fr/rubrique.php?id_rubrique=268 for details.

Their optical catalogue with which the cross-correlation was performed contains sources detected in at least one of the Subaru bands (b, v, g, r, i, z) down to an AB magnitude limit of ~ 27 . The COSMOS sample is almost complete down to $F_{[0.5-2]\text{keV}} \simeq 3 \times 10^{-15} \text{ erg s}^{-1} \text{ cm}^{-2}$ (X-ray detection probability of ≥ 0.98 ; cf. Cappelluti et al. 2007, fig. 6), and 98 per cent of the X-ray sources have an optical counterpart.

The joint probability density of the XXL survey is expected to be quite similar to that of the XMM-COSMOS field if we apply to this survey the same X-ray and optical flux cuts and take into account the different probability of detection for sources fainter than $2 \times 10^{-14} \text{ erg cm}^{-2} \text{ s}^{-1}$. Consequently, we first estimate the COSMOS joint probability density $d_{\text{COSMOS}}(z_s, f_X)$ from a smoothed histogram of the sources from the XMM-COSMOS in the (z_s, f_X) plane, where we apply the X-ray and r -band cutoffs of the XXL. We then take into account the XXL detection probability to determine $d_{\text{obs}}(z_s, f_X)$.

In the left-hand panel of Fig. 1, we have represented the XMM-COSMOS sources in the (z_s, f_X) plane, with fluxes presented along a logarithmic scale. We have only considered sources with a flux larger than $F_{[0.5-2]\text{keV}} = 3 \times 10^{-15} \text{ erg cm}^{-2} \text{ s}^{-1}$ in the [0.5–2] keV band and with an optical counterpart brighter than $r = 25$ in the SDSS r band. We have rejected all X-ray sources for which there was no redshift estimate. Whenever available, spectroscopic redshifts were preferred to photometric ones. The XMM-COSMOS catalogue presented in Brusa et al. (2010) contains 1797 sources in the 2 square degrees. After applying the X-ray and r -band cutoff (and excluding sources with no available r magnitude), there are 630 sources of which 6 are excluded because of no available redshift. The final restricted COSMOS source sample contains 624 sources. The density of sources in the (z_s, f_X) plane is larger for the fainter fluxes, near redshift $z \sim 1$. At any given redshift, there are more sources with a lower flux. Finally, let us also point out the absence of sources with a high flux at high redshift.

In order to estimate the joint probability density $d_{\text{COSMOS}}(z_s, f_X)$ of the COSMOS sources, we have calculated a smoothed histogram of the COSMOS source distribution with the XMM cutoff in the X-ray and r band displayed in the left-hand panel of Fig. 1. For the convenience of the developed software, $d_{\text{COSMOS}}(z_s, f_X)$ has been derived using a logarithmic scale for the X-ray fluxes. We have considered redshift intervals of 0.375 with bin centres separated

Table 1. Estimate of (1) the number of sources with an X-ray flux greater than 3×10^{-15} erg cm $^{-2}$ s $^{-1}$ in the [0.5–2] keV range and $r < 25$ and (2) the number of detected multiply imaged sources in the XMM-COSMOS, XMM-LSS and XXL surveys. The estimates of the number of sources are extrapolated from the XMM-COSMOS catalogue taking into account the angular coverage of the different surveys. For the estimate of the number of multiply imaged sources, the numbers in parentheses correspond to the SIS case. Observational data are presented in bold.

Survey	Coverage (deg 2)	Number of sources	Lensed sources ($\theta_{\text{mis}} = 0$ arcsec) SIE (SIS)	Lensed sources ($\theta_{\text{mis}} = 0.45$ arcsec) SIE (SIS)	>2 images ($\theta_{\text{mis}} = 0$ arcsec)	>2 images ($\theta_{\text{mis}} = 0.45$ arcsec)
XMM-COSMOS	2	621	–	–	–	–
XMM-COSMOS (with XXL $P_{\text{detection}}$)	2	529.5	–	–	–	–
XMM-LSS	10.9	2885	5 (5)	4 (4)	0	0
XXL	44.2	11 701	20 (21)	16 (17)	1	1
			$\langle \tau \rangle$ ($\theta_{\text{mis}} = 0$ arcsec)	$\langle \tau \rangle$ ($\theta_{\text{mis}} = 0.45$ arcsec)	$\langle \tau_{>2} \rangle / \langle \tau \rangle$ ($\theta_{\text{mis}} = 0$ arcsec)	$\langle \tau_{>2} \rangle / \langle \tau \rangle$ ($\theta_{\text{mis}} = 0.45$ arcsec)
SIE			1.698×10^{-3}	1.384×10^{-3}	0.0718	0.0564
SIS			1.788×10^{-3}	1.489×10^{-3}	–	–
SIS (Late)			5.719×10^{-4}	1.629×10^{-6}	–	–

by 0.0625 and logarithmic magnitude intervals of 0.3 separated by bins of 0.05. The derived COSMOS joint probability density is shown in the right-hand panel of Fig. 1. We have intentionally kept the same axis as in the left-hand panel in order to clearly identify the similarities between the two figures. The grey-scale indicates the values of $d_{\text{COSMOS}}(z_s, f_X)$; the darker the grey, the higher the probability of finding a source. The normalization factor of $d_{\text{COSMOS}}(z_s, f_X)$ is the number of sources (624) detected in the 2 square degrees of the COSMOS field, restricted to the XMM-LSS cutoffs.

To take into account the detection probability of the XXL survey, we multiply the $d_{\text{COSMOS}}(z_s, f_X)$ in the right-hand panel of Fig. 1 by the XMM-LSS detection probability as a function of the flux taken from Elyiv et al. (2012, fig. 10), the obtained distribution being $d_{\text{obs}}(z_s, f_X)$. The normalization factor of this distribution represents the number of sources (592.5) with $F_{[0.5-2]\text{keV}} \gtrsim 3 \times 10^{-15}$ erg cm $^{-2}$ s $^{-1}$ and $r < 25$ that would be detected within the XXL survey in the 2 square degrees of the XMM-COSMOS field. From this nor-

malization factor, we may thus estimate the number of sources to be detected in the XMM-LSS and XXL fields with similar characteristics. The expected numbers of sources in the different surveys are summarized in Table 1. Assuming the final XXL catalogue to have similar characteristics as those of the XMM-LSS, this estimate of $d_{\text{obs}}(z_s, f_X)$ is assumed to be valid for both the XMM-LSS and the XXL fields.

As a reliability test of the derived joint probability density $d_{\text{obs}}(z_s, f_X)$, we verify its ability to represent the properties of the observed population of AGNs. In the left-hand panel of Fig. 2, we have represented as a continuous grey line the observed cumulative distribution as a function of the redshift of the XMM-COSMOS sources with an X-ray flux larger than $F_{[0.5-2]\text{keV}} = 3 \times 10^{-15}$ erg cm $^{-2}$ s $^{-1}$ in the [0.5–2] keV band, with $r < 25$ and a redshift estimation.

In the same figure, we have represented with a dashed dark grey line the cumulative source redshift distribution derived from the COSMOS joint probability distribution when not considering the detection probability of the XXL survey. The cumulative redshift




Figure 2. Left-hand panel: cumulative function of the number density $N(z_s)$ of the sources as a function of the redshift for the XMM-COSMOS sources with $F_{[0.5-2]\text{keV}} \gtrsim 3 \times 10^{-15}$ erg cm $^{-2}$ s $^{-1}$ and $r < 25$. We have also represented the cumulative redshift distribution inferred from the marginal distribution of the joint probability density function $d_{\text{COSMOS}}(z_s, f_X)$ and $d_{\text{obs}}(z_s, f_X)$, with and without considering the XXL detection probability (dashed and continuous black curves, respectively). Right-hand panel: differential number counts of the XMM-COSMOS sources as a function of the flux in the soft X-ray band, for all the COSMOS sources, which are used for the amplification bias calculation. We have only represented the X-ray flux range accessible to the XXL survey. We have also represented the marginal distribution obtained from the joint probability density function $d_{\text{COSMOS}}(z_s, f_X)$, as well as the DNCF fit used for the calculation of the amplification bias.

distribution is obtained by integrating $d_{\text{COSMOS}}(z_s, f_X)$ over f_X and by multiplying the number of AGNs detected per square degree in the COSMOS field, where we have applied the XXL cutoffs.

The cumulative redshift distribution inferred from the distribution $d_{\text{COSMOS}}(z_s, f_X)$ reproduces very well the observed cumulative source redshift distribution.

In the same figure, we have represented with a dark continuous line the cumulative redshift distribution derived from the final $d_{\text{obs}}(z_s, f_X)$ when considering the detection probability of the XMM-LSS survey. We remark very good agreement at low redshift between this distribution and the two previous ones, and a significant deviation above redshift ~ 1 due to the fact that the XXL survey does not detect all of the fainter sources, more numerous at higher redshifts.

In the right-hand panel of Fig. 2, we have represented the DDCF of the XMM-COSMOS sources as a function of their X-ray soft band flux. The flux is shown along a logarithmic scale and the error bars are estimated considering a Poisson noise. We have here considered all the COSMOS sources brighter than $F_{[0.5-2]keV} = 1 \times 10^{-15} \text{ erg cm}^{-2} \text{ s}^{-1}$. This DDCF is used to estimate $N_{f_X}(f_X)$ needed when calculating the amplification bias in equation (6). As previously mentioned, the calculation of the lensing optical depth necessitates the knowledge of the DDCF for fluxes fainter than the survey detection limit. We have therefore fitted the observed XMM-COSMOS DDCF $\log_{10}(n(f_X))$ with a third-order polynomial in the flux range covered by the XMM-COSMOS and extrapolated the data linearly outside this flux range. The fit is shown as a continuous dark grey line in the right-hand panel of Fig. 2. This polynomial form of the DDCF is used for the calculation of the amplification bias in our simulations.

In the same figure, we have represented with a dashed line the DDCF per unit of solid angle derived from the joint probability density function $d_{\text{obs}}(z_s, f_X)$, where we have restricted the COSMOS sources to the XXL cutoffs and taken into account the detection probability of the survey. This is obtained by integrating $d_{\text{obs}}(z_s, f_X)$ over z_s and by multiplying by the number N_{AGN}/Ω of AGNs per square degree. This curve is matching fairly well the fitted curve for the brighter sources and is lower for the fainter sources, where the XXL survey only detects part of the sources and where the fraction of excluded sources with $r > 25$ is higher.

The small size of the source sample in the XMM-COSMOS field at low redshift (due to the small survey angular coverage) does not permit a better estimate of $d_{\text{obs}}(z_s, f_X)$ at these low redshift values, because of a large scatter in the observed data at low redshift, especially for the brighter sources. Nevertheless, as these very bright and low-redshift sources are very rare in the survey and as their lensing probability is very small (because of their very low redshift), this does not have a large impact on our simulations. These problems will be reduced in the XXL survey, for which the angular coverage is ~ 20 times larger than that of the XMM-COSMOS field. This confirms, as mentioned earlier, that the COSMOS sample is not large enough to determine the distribution $d_{\text{obs}}(z_s, f_X, r)$ of the sources in the 3D space (z_s, f_X, r) .

4 RESULTS

4.1 Mean lensing optical depth

Using the joint probability density $d_{\text{obs}}(z_s, f_X)$ described in Section 3, we have computed the mean lensing optical depth $\langle \tau \rangle$ for the XXL survey, integrating numerically equation (12). As $d_{\text{obs}}(z_s, f_X)$ is

assumed to be identical for the XXL and the XMM-LSS surveys, the results are valid for both surveys.

To calculate the lensing optical depth, we modelled the deflector population with SIS and SIE mass distributions (equations 8 and 10). We have computed the average optical depth for different values of the minimum image separation θ_{mis} resolvable at optical wavelengths. When calculating the cross-section, we have considered the detection of the lensed images to be achievable down to an angular separation θ_{mis} (independently of their relative amplification). Although this is not strictly accurate, we have made this assumption for the following reason. The regions contributing the most to the lensing cross-section are those where the source is located close to the caustic curves, as these are the most amplified and thus benefit the most from the amplification bias. For these configurations, the lensed images that are the closest to each other (and also the brightest) are those formed on each side of the tangential critical curve. These lensed images have a very similar amplification (Kormann et al. 1994). Consequently, for the detection of these lensed images, the critical parameter is the smallest angular distance under which these point-like images cannot be resolved, independently of their brightness. In practice however, for lensed images close to each other, the minimum angular distance at which the point-like images can be disentangled is dependent on the flux difference between the images, especially at very small angular distances. For the analysis of the final XXL sample, the angular selection function (characterizing the smallest angular distance detectable as a function of the relative amplification of the lensed images) will have to be determined precisely.

Fig. 3 displays the behaviour of the average optical depth $\langle \tau \rangle$ as a function of the minimal image separation θ_{mis} resolvable in the optical survey. We display the results when modelling the deflector population with either SIE or SIS mass profiles. In the former case, we have calculated the total probability of having a multiply imaged source, independently of the number of the lensed images.

For a perfect instrument (i.e. $\theta_{\text{mis}} = 0$ arcsec), when considering deflectors modelled with SIE mass distributions, we find an average lensing probability $\langle \tau_{\text{SIE}} \rangle = 1.698 \times 10^{-3}$. When modelling the




Figure 3. Average lensing probability $\langle \tau \rangle$ calculated for the XXL and XMM-LSS fields as a function of the minimum angular separation θ_{mis} resolvable by the survey in the optical domain, when modelling the deflector population with the SIS and SIE mass distributions. For comparison, we have also computed the case of the late-type galaxies, modelled by means of SIS mass distributions.

deflectors with an SIS mass distribution, the mean lensing probability is $\langle \tau_{\text{SIS}} \rangle = 1.788 \times 10^{-3}$. Modelling the deflectors with SIS mass distributions thus leads to a slightly larger average lensing probability by ~ 5 per cent. The reason for this is that the SIE mass distribution was introduced in order to preserve the projected mass inside the same area but does not conserve the geometrical cross-section nor the amplification probability distribution. In other words, the area inside the caustic curve of an SIS deflector is always larger than or equal to that included inside the caustic curves of an SIE deflector. When averaging the SIE lensing cross-section over the deflector ellipticity distribution, this leads to an equivalent SIE lensing cross-section smaller than that of the SIS case. Furthermore, for these two types of deflectors, the probability of producing a multiply imaged source with a given total amplification slightly differs. The impact of the amplification bias is thus different for the two deflector models. Consequently, the slight differences between $\langle \tau_{\text{SIS}} \rangle$ and $\langle \tau_{\text{SIE}} \rangle$ depend on the DNCF as a function of f_X for the source population, which varies from one survey to another. The overestimate of $\langle \tau_{\text{SIS}} \rangle$ thus has to be estimated independently for each survey. This boost of the average lensing optical depth in the SIS model was studied by Huterer et al. (2005) who concluded that the ellipticity in the deflector mass distribution decreases the mean lensing optical depth. A boost of the SIS model by a few per cent may be expected in a survey with sources showing a steep DNCF.

Both $\langle \tau_{\text{SIE}} \rangle$ and $\langle \tau_{\text{SIS}} \rangle$ decrease for increasing values of the parameter θ_{mis} . When considering a finite resolution of the instrument, some of the lensed images formed are angularly too close to each other and are detected as a single point-like object. Consequently, the probability of *detecting* the lensed images decreases as θ_{mis} increases.

In the present case, the multiple lensed images will be searched for among the optical counterparts of the point-like X-ray sources, because of the better angular resolution in the optical domain. The ground-based observations are limited by the atmospheric seeing. In the northern XXL fields, the CFHT in the r band has a typical seeing of 0.7 arcsec (Salmon et al. 2009) and for the Southern hemisphere the typical seeing with the Blanco telescope is ~ 0.9 arcsec (Desai et al. 2012). We thus considered our full width at half-maximum of the point spread function (PSF) to be homogeneous over the entire sample and equal to ~ 0.9 arcsec.

Thanks to PSF fitting techniques, we can hope to resolve multiple point-like lensed images down to half the full width at half-maximum for lensed images with the same amplification, which constitute the configurations contributing the most to the lensing cross-section as previously explained. Consequently, the typical θ_{mis} value achievable is expected to be $\theta_{\text{mis}} \sim 0.45$ arcsec (in practice, as stressed previously, θ_{mis} depends on the relative amplification of the lensed images, which will have to be taken into account when analysing the final XXL sample). For this value, we have $\langle \tau_{\text{SIE}} \rangle = 1.384 \times 10^{-3}$ and $\langle \tau_{\text{SIS}} \rangle = 1.489 \times 10^{-3}$. The slight overestimate of the SIS mean lensing value relatively to that of the SIE model thus increases with θ_{mis} and reaches ~ 10 per cent for $\theta_{\text{mis}} = 0.45$ arcsec.

For comparison, we have computed the evolution of the mean optical depth as a function of θ_{mis} for the population of late-type galaxies modelled by means of an SIS mass distribution. As for the case of the early-type galaxies, we have considered the comoving density of late-type galaxies to be constant with the redshift and we have used the VDF parameters determined by Chae (2010) in the local Universe, i.e.

$$[\Phi_*, \sigma_*, \alpha, \beta] = [66 \times 10^{-3} h^3 \text{Mpc}^{-3}, 91.5 \text{ km s}^{-1}, 0.69, 2.10].$$

The results are displayed in Fig. 3. When considering a perfect instrument ($\theta_{\text{mis}} = 0$ arcsec), the average lensing optical depth associated with the late-type galaxies is about a third of that of the early-type ones (i.e. $\langle \tau \rangle = 5.719 \times 10^{-4}$). Nevertheless, the decrease of $\langle \tau \rangle$ with θ_{mis} is steeper than that in the early-type galaxy case. Indeed, late-type galaxies are less massive and lead to smaller typical angular separations of the multiple lensed images, which are not disentangled in the seeing-limited images. For $\theta_{\text{mis}} = 0.45$ arcsec, the average lensing optical depth due to late-type galaxies is found to be $\langle \tau \rangle = 1.629 \times 10^{-6}$, three orders of magnitude lower than that of the early-type galaxies. The expected contribution of the late-type galaxies in our sample of gravitationally lensed sources is thus negligible, which validates our assumption of only considering the population of early-type galaxies as the deflectors for the XXL lensed source sample.

Our estimation of the contribution of late-type galaxies to the lensed sources is surprisingly low compared to the observed fraction of late-type lenses in existing samples. Furthermore, it suggests that late-type lenses with image separations larger than 0.5 arcsec are extremely rare, which is also in contradiction with observed samples. Indeed, 2 out of the 13 CLASS lenses from the statistical sample are likely to be produced by late-type galaxies (B0218 and B1933; see Browne et al. 2003) and out of the 26 lensed QSOs of the SDSSQLS statistical sample, one is possibly due to a late-type galaxy (J1313) and has an angular separation larger than 1 arcsec. We do not fully understand the reason for these discrepancies. A possible cause of error is the effect of the lens environment, not considered in this work, which may lead to an additional gravitational shear. Huterer et al. (2005) have shown that the external shear broadens the distribution of angular separation between the lensed images without changing its average value. It may therefore increase the fraction of events with an angular separation larger than 1 arcsec. In the CLASS and SDSSQLS statistical samples, all the lenses produced by late-type galaxies with an angular separation larger than 1 arcsec required external shear to accurately model the position and relative amplification of the lensed images (see Sluse et al. 2012; Suyu et al. 2012 and references therein). Another possible source of discrepancy is the VDF used for late-type galaxies (from Chae 2010) which does not come from direct measurements (as in the case of the early-type VDF; Choi et al. 2007). It is inferred from the local LF of late-type galaxies using the Tully–Fisher relationship (taking into account its dispersion) and assuming a conversion between the circular velocity v_c and σ to be that of an SIS profile (i.e. $\sigma = v_c/\sqrt{2}$). Chae (2010) stresses that lensing statistics of late-type galaxies might necessitate to consider the circular velocity function (rather than the VDF) and consider more realistic mass distribution models. Furthermore, type-specific LF (and VDF) are still potentially biased by misclassification of the galaxy types and, according to Park & Choi (2005), the work of Chae (2010) has an ~ 10 per cent classification mismatch. We therefore advise to use with caution the results concerning the late-type population statistics.

Table 1 summarizes the number of multiply imaged sources expected in the different surveys as well as the expected number of detected events. Assuming that the sources detected in the XXL survey will have the same properties as those in the XMM-COSMOS to which we applied the same flux limits in the X-ray and optical bands (except when estimating the amplification bias) and when accounting for the detection probability of the XXL, we expect the detection of 11 701 sources with $F_{[0.5-2]\text{keV}} > 3 \times 10^{-15} \text{ erg cm}^{-2} \text{ s}^{-1}$ and $r < 25$ in the 44.2 square degrees of the survey. Among these sources, we expect 21 (20 in the SIE case) to be




Figure 4. Fraction $\langle \tau_{\text{SIE}, i} \rangle / \langle \tau_{\text{SIE}} \rangle$ of the lensing events with a given number $i = 2, 3$ or 4 of images as a function of the survey minimum angular separation θ_{mis} , when modelling the deflector population with the SIE mass distribution. Three-image configurations are quads with two blended point-like images.

multiply imaged, out of which 17 (16 in the SIE case) should be detected assuming $\theta_{\text{mis}} = 0.45$ arcsec.

When calculating the lensing probability of a source to be lensed through equation (10), considering a population of deflectors modelled with SIE profiles, we may also calculate the probability $\tau_{\text{SIE}, i}$ of a source to be lensed with the formation of a given number i of lensed images. We have calculated the average value of $\langle \tau_{\text{SIE}, i} \rangle$ for the population of sources to be detected within the XXL and XMM-LSS fields using equation (12), for different values of the θ_{mis} parameter. The results are displayed in Fig. 4, where we have plotted as a function of the value of the θ_{mis} parameter the fraction of lensing events composed of i lensed images $\langle \tau_{\text{SIE}, i} \rangle / \langle \tau_{\text{SIE}} \rangle$ relatively to the total average lensing probability $\langle \tau_{\text{SIE}} \rangle$.

The fraction of lensing events with the formation of two images is always the highest. This is a consequence of the scarcity of very elliptical deflectors (see Choi et al. 2007, fig. 13 for the axial ratio distribution of early-type galaxies). In the case of a perfect instrument (i.e. $\theta_{\text{mis}} = 0$ arcsec), we find that 93 per cent of the lensed sources are composed of two images and this fraction increases with the value of the θ_{mis} parameter.

For a perfect instrument, the lensed sources with formation of more than two images are composed of quads (i.e. four-image configurations). As θ_{mis} increases, some of the four-lensed-image configurations, due to their too small angular separation, have only three point-like images detected (quads with two blended point-like images). Out of the 20 lensed sources formed in the XXL population, only one is expected to be detected with more than two images.

The fraction of lensing systems with more than two images is roughly consistent with the results of Oguri & Marshall (2010) who have calculated the expected number of gravitationally lensed quasars in wide-field optical surveys. For a survey with a limiting magnitude $i = 25$, these authors find a fraction of a little more than ~ 10 per cent of quads (formation of four images) against ~ 7 per cent in our simulations. The slight difference is most likely due to the different ellipticity distribution of the deflectors considered [Oguri & Marshall (2010) consider a combination of oblate and prolate three-dimensional deflectors with a Gaussian distribution of their ellipticity] as well as their consideration of an additional external shear due to the lens environment. Huterer et al. (2005) showed




Figure 5. Normalized redshift distributions of the deflectors, the lensed sources and all AGNs. For each distribution, we have indicated the median redshift value.

that the external shear increases the fraction of quads in a sample of lensed sources.

4.2 Redshift distributions

Using the joint probability density $d_{\text{obs}}(z_s, f_X)$ described in Section 3, we have computed the mathematical expectation of the normalized redshift distribution of the deflectors $w_{z_d}(z_d)$ and of the lensed sources $w_{z_s}(z_s)$, numerically integrating equations (13) and (14), respectively. We have considered $\theta_{\text{mis}} = 0.45$ arcsec and we have modelled the deflectors with SIS mass distributions. The normalized redshift distributions $w_{z_d}(z_d)$ and $w_{z_s}(z_s)$ are shown as a function of the redshift in Fig. 5. For comparison, we have represented the marginal distribution as a function of the redshift of the joint probability density, i.e. the normalized distribution as a function of the redshift of all the sources (independently of the fact that they are being lensed).

In this figure, we have also illustrated the median value of the observed redshift for the different distributions. The redshift distribution of the lensed sources is shifted towards a higher redshift compared to that for all the sources. As the source redshift increases, so does its geometrical lensing volume (i.e. the volume in which the presence of a deflector leads to the formation of multiple lensed images). Consequently, sources with a higher redshift tend to have a higher lensing probability. The mathematical expectation for the redshift moves from $\langle z_{\text{AGN}} \rangle \simeq 1.11$ for the entire source population to $\langle z_s \rangle \simeq 1.8$ for the lensed sources.

For redshifts larger than $\langle z_{\text{AGN}} \rangle$, the distribution of the lensed sources does not appear as a smooth function of the redshift, and we clearly see the presence of bumps or redshift ranges with a probability excess compared to a smooth decreasing function of the redshift. These probability overdensities of lensed source detections correspond to redshift ranges in which strong emission lines of the AGNs enter in the optical SDSS r band, in which the optical counterparts are searched for; they are thus a consequence of a selection bias. The presence of the emission lines in the r band increases their probability of being detected, compared to that of a source with only a continuum-like spectrum. For example, the Mg II line at 279.8 nm enters the r -band filter in the redshift range 2–2.42, for which we see an overdensity in the lensed source redshift distribution.

The redshift distribution of the lensed sources corresponds to the probability density from which the detected lensed source redshift may be considered as a random event, thanks to the fact that, in this case, all the lensed sources are detected and may have their redshift estimated. Given a large enough sample of lensed sources, this distribution could be retrieved from a normalized histogram of the lensed sources, as a function of their redshift.

In Fig. 5, we have also represented the normalized distribution ω_{z_d} , as a function of the redshift, for the deflectors involved in the formation of multiply imaged sources. The deflector redshift median value is $\langle z_d \rangle \sim 0.54$ and the most probable value is $z \lesssim 0.5$. The contribution of deflectors with $z \gtrsim 1$ is very small. This comforts our assumption of a non-evolving deflector population in the calculation of the lensing optical depth, as the population involved is mainly located at low redshift.

In the case of the deflector distribution, the observed redshift distribution of the deflectors involved in the formation of multiple images of a source will be highly biased as most of them are not bright enough to be detected.

Oguri & Marshall (2010) have also estimated the expected lens redshift distribution for lensed quasars detected in optical imaging surveys. The deflector redshift distributions are marginally consistent, although the distribution found in this work peaks at lower redshifts [$z_{\max} \sim 0.6$ for Oguri & Marshall (2010) and $z_{\max} \sim 0.5$ in the present work]. The expected source redshift distribution is also shifted towards lower redshift in our study. This difference in the lensed source and deflector redshift distributions comes from the difference in the source distribution: in the present work, the rather bright X-ray flux cutoff tends to reject sources at high redshift, which are included in the sample of Oguri & Marshall (2010).

4.3 Influence of the cosmological model

The probability for a source with a known apparent flux and redshift to be gravitationally lensed with the formation of multiple images depends on the cosmological model. This may be seen for instance through the dependence of the infinitesimal light-distance element cdt/dz on the cosmological mass density parameter Ω_m in equation (2), as well as through the dependence of the lensing cross-section on Ω_m (see equations 6 and 7) via the definition of the angular diameter distances. If we consider a flat expanding FLRW universe model, the only dependence of the lensing probability on the universe model is made through the cosmological mass density parameter Ω_m . Indeed, although a flat expanding FLRW is totally characterized by Ω_m and H_0 , an increase in the value of H_0 will only act as a scaling factor (decreasing the lensing volume while increasing the density of deflectors). So far, we have considered an FLRW flat universe with $\Omega_m = 0.3$ and $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$. In Fig. 6, we display the behaviour of the average lensing probability $\langle \tau \rangle$ for the XXL sources as a function of Ω_m , for a flat universe with $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$. We have computed both $\langle \tau_{\text{SIE}} \rangle$ and $\langle \tau_{\text{SIS}} \rangle$. Both models lead to the same behaviour of the mean lensing probability as a function of Ω_m and, here as well, we observe that the SIS mass distribution leads to a slight overestimate of the average lensing probability when compared to that corresponding to the SIE model, whatever the value of Ω_m .

There is a very strong dependence of the expected fraction of lensed sources in the survey on Ω_m . For this reason, the statistics of gravitational lensing in a well-defined sample of sources has been widely used to probe the value of Ω_m and test dark energy models (Turner et al. 1984; Fukugita et al. 1990; Turner 1990;




Figure 6. Average lensing probability as a function of the cosmological matter density parameter Ω_m .

Surdej et al. 1993; Keeton 1998, 2002; Chae et al. 2002; Ofek et al. 2003; Mitchell et al. 2005; Cao et al. 2012).

The XXL sample on its own will not allow better constraints on the value of Ω_m than recent lens surveys such as the SDSS-LQS. Oguri et al. (2012) for instance constrained Ω_{Lambda} to $\Omega_{\text{Lambda}} = 0.79^{+0.06(\text{stat.})}_{-0.07} \pm 0.06(\text{syst.})$ on the basis of 19 lenses from the SDSS-LQS statistical sample. The XXL lensed source sample will thus be combined with other recent surveys (including the SDSS-LQS) to better constrain the cosmological parameters.

5 CONSIDERING THE *r*-BAND CUTOFF

In the previous section, when calculating the amplification bias in equation (6), we have only considered the distribution of the sources as a function of their flux f_X in the X-ray band and we have considered the lensing optical depth $\tau(z_s, f_X)$ to be only a function of the source redshift and its flux in the X-ray band. Ideally, the source r magnitude in the optical domain should also be considered for the calculation of $\tau(z_s, f_X, r)$ and $d_{\text{obs}}(z_s, f_X, r)$ and the calculation of the mean lensing optical depth $\langle \tau \rangle$ in equation (12) should include the integration over r .

The joint probability density $d_{\text{obs}}(z_s, f_X, r)$ may be decomposed as

$$d_{\text{obs}}(z_s, f_X, r) = d_{\text{obs}}(z_s, f_X) d(r|z_s, f_X), \quad (15)$$

where $d(r|z_s, f_X)$ is the normalized distribution as a function of r for sources with a redshift and an X-ray flux in the ranges $[z_s, z_s + dz_s]$ and $[f_X, f_X + df_X]$, respectively, and equation (12) remains correct if we consider the optical depth $\tau_{(r)}(z_s, f_X)$ averaged over the r magnitudes

$$\tau_{(r)}(z_s, f_X) = \int \tau(z_s, f_X, r) d(r|z_s, f_X) dr. \quad (16)$$

Equivalently, $\tau_{(r)}(z_s, f_X)$ can be calculated through equation (3) if we consider an average lensing cross-section $\Sigma_{(r)}$

$$\Sigma_{(r)} = b_0^2 \iint_{S_y} B_{(r)} dy, \quad (17)$$

where we have defined the amplification bias $B_{(r)}$ averaged over the r magnitudes

$$B_{(r)} = \int \frac{N(z_s, f_X/A_X, r + 2.5 \log(A_r))}{N(z_s, f_X, r)} d(r|z_s, f_X) dr, \quad (18)$$

where $N(z_s, f_X, r)$ represents the density of sources with a redshift z_s , having an observed flux f_X in the X-ray and an r -band magnitude r , A_X and A_r are the amplifications due to the lensing event in the X-ray and r band, respectively. If the source is point-like, we may




Figure 7. Left: distribution of the COSMOS sources in the (f_X, r) plane. We have represented the X-ray fluxes in terms of the X-ray magnitudes $m_x = -2.5 \log f_X$, and have represented the X-ray and r -band cutoff of the XXL survey. We have also represented the contour plot of the COSMOS source density in the (f_X, r) plane. Right: amplification bias as a function of the total amplification of the lensing event when considering only the DNCF as a function of the X-ray flux f_X (continuous dark grey line) and when averaging over the r -band magnitude distribution of the sources.

then assume that $A_X = A_r$. If not, the ratio A_X/A_r will depend on the size of regions in the AGN emitting the X-ray and the optical fluxes, respectively, as well as the positions of these regions with respect to the caustic curves. The expectation of the ratio A_X/A_r will thus necessitate heavy simulations to be performed. As a first approximation we consider point-like sources.

In this work, the amplification bias $B_{X\text{-ray}}$ has been calculated so far uniquely on the basis of the X-ray flux through the relation

$$B_{X\text{-ray}} = \frac{N(f_X/A_X)}{N(f_X)}, \quad (19)$$

where we have neglected the redshift dependence of the sources as a function of their redshift because of a too small source sample to correctly characterize this possible redshift dependence. In the following, we will compare the amplification bias obtained through equations (18) and (19).

To calculate rigorously the amplification bias through equation (18), we thus need to determine the distribution of the source population in the (z_s, f_X, r) space. Our observational data from the COSMOS sample are not large enough to estimate this distribution over the whole area probed by the survey. Consequently, we assume the distributions $N(z_s, f_X, r)$ and $d(r|z_s, f_X)$ to be independent of the redshift.

Let us now estimate the distribution $N(f_X, r)$ on the basis of the distribution of the COSMOS sample in the (f_X, r) plane, represented in the left-hand panel of Fig. 7. We have represented all the COSMOS sources for which the soft X-ray flux and r -band magnitude are given in the sample of Brusa et al. (2010). As 98 per cent of the detected X-ray sources have an optical counterpart in the r band for X-ray sources brighter than the XXL 0.5 detection probability cut, we can consider this sample to be a complete sample of X-ray sources brighter than the X-ray cut, for which we know the optical counterpart.

To estimate $N(f_X, r)$, we have proceeded as follows. We bin in terms of the $m_x = -2.5 \log f_X$ magnitude, with a bin width of $dm_x = 0.5$ and bin centres ranging from 33.5 to 37. In each m_x bin, we construct the histogram of the sources as a function of the r -band magnitude, using bins with a width of $dr = 0.5$ and bin centres ranging from 17 to 28 with step of 0.5. Each histogram as a function of r is fitted by means of a Gaussian profile. The number of sources in each bin is then divided by $dm_x dr$.

We thus obtain the evolution of the Gaussian fit parameters (i.e. the amplitude A_G , the average position r_G and the standard deviation σ_G) as a function of the m_x bin and we fit the dependence of these parameters as a function of m_x by a linear law in the m_x range 33.5–37. We model the density of the sources in the (f_X, r) plane by

$$N(f_X, r) = A_G(m_x) \exp\left(-\left(\frac{r - r_G(m_x)}{2\sigma_G(m_x)}\right)^2\right). \quad (20)$$

The COSMOS data do not have enough bright sources to constrain $N(f_X, r)$ for m_x brighter than 33.5. We have represented the isodensity contours of the calculated function $N(f_X, r)$ in the left-hand panel of Fig. 7, showing very good agreement between the observed distribution of the COSMOS sources and the modelled density function.

We now use the modelled function $N(f_X, r)$ to calculate the amplification bias in equation (18), for sources with different X-ray-to-optical flux ratios. Assuming that the amplification induced by the lensing event in the X-ray and that in the optical are identical, when amplified, a source is displaced in the (m_x, r) parallel to the dashed line shown in the left-hand panel of Fig. 7. Consequently, the amplification bias is calculated thanks to the evolution of $N(f_X, r)$ along this trajectory.

In the right-hand panel, we represent the behaviour of the amplification bias as a function of the total amplification of the lensing event, for four sources with apparent X-ray fluxes equal to the XXL X-ray limiting flux $f_{XXL\text{-cut}}$, and 5, 10 and 15 times the value of the $f_{XXL\text{-cut}}$. The dashed light grey curve represents the amplification bias $B_{X\text{-ray}}$ obtained when only considering the DNCF of the XXL sources as a function the X-ray flux, calculated through equation (19). The continuous black curve corresponds to the average bias $B_{(r)}$ calculated thanks to equation (18). We see perfect agreement between these curves and the amplification bias calculated only considering the DNCF as a function of the X-ray flux f_X . Consequently, the amplification bias calculated through equation (6) perfectly corresponds to the amplification bias averaged over the r -band magnitudes.

Thus, assuming a point-like source, we may calculate the amplification bias in the combined X-ray/optical data by considering uniquely the X-ray distribution of a complete and deeper sample, which validates the method introduced in Section 2. Nevertheless,

in the analysis of the final XXL sample, we will have to also consider the redshift dependence of the amplification bias, which will be made possible thanks to the much larger size of the sample.

6 CONCLUSIONS

We have calculated the expected statistical properties of the multiply imaged sources to be detected among the optical counterparts of the XXL point-like sources, modelling the deflectors successively with SIS and SIE mass profiles. We find the following.

(i) One expects the formation of 20 (21 using the SIS model) multiply imaged AGNs out of which 16 (17 for the SIS case) should be detected among the optical counterparts with an angular resolution of 0.45 arcsec, and we only expect the detection of one gravitational lens system composed of more than two lensed images.

(ii) The expectations are consistent when modelling deflectors with SIE and SIS mass distributions, although the SIS model leads to a slight overestimate of the mean lensing probability. This overestimate is a function of the amplification bias and is thus different for each survey.

(iii) The late-type galaxy population should not contribute to the lensed sources to be detected.

(iv) Although the detection is done simultaneously in the X-ray and in the optical domain, the amplification bias may be estimated from the X-ray flux distribution, as long as we consider a complete X-ray sample from a deeper survey and for point-like sources.

In this work, we have considered isothermal profiles to model the deflectors. This has allowed us to get first good estimates of the expected number of lensed AGN in the XXL survey. However, more detailed calculations ought to be carried out. Indeed, Auger et al. (2010) and Koopmans et al. (2009), through the analysis of massive early-type deflectors from the Sloan Lens ACS Survey (SLACS) survey, have found a slight deviation from the isothermal profile, with a steeper slope parameter ($\gamma = 2.078$ and 2.085 , respectively, where the mass distribution evolves as $r^{-\gamma}$). If we consider two mass distributions with the same total mass, a steeper profile would lead to a higher fraction of the lens in the centre, which will increase the Einstein angular radius, therefore leading to an increase in the lensing cross-section of the deflector. This increase in the lensing cross-section with the steepness of the radial mass profile has been put in evidence in a series of papers (Mandelbaum, van de Ven & Keeton 2009; van de Ven, Mandelbaum & Keeton 2009), where the authors studied the impact of galaxy shape and density profile on the selection biases in surveys for the detection of strong lenses. Because isothermal profiles are singular, the authors analysed more realistic profiles in order to define the total mass.

Furthermore, Sonnenfeld et al. (2013) have shown from the study of the CFHTLS-Strong Lensing Legacy Survey (SL2S) galaxy-scale lens sample (with deflectors in the range $0.2 < z < 0.8$) that the mass density profile of early-type galaxies depends on the redshift, lower galaxies showing a steeper average profile. As deflectors with a steeper profile tend to have a higher lensing cross-section, this would favour the deflector redshift distribution to be shifted towards lower redshift.

These effects, along with the redshift dependence of the amplification bias, will have to be considered in the analysis of the final XXL sample.

ACKNOWLEDGEMENTS

The authors would like to acknowledge the Communauté Française de Belgique and the Actions de Recherche Concertées de

l'Académie universitaire Wallonie-Europe for their funding during the present research.

REFERENCES

- Auger M. W., Treu T., Bolton A. S., Gavazzi R., Koopmans L. V. E., Marshall P. J., Moustakas L. A., Burles S., 2010, *ApJ*, 724, 511
- Browne I. W. A. et al., 2003, *MNRAS*, 341, 13
- Brusa M. et al., 2010, *ApJ*, 716, 348
- Cao S., Pan Y., Biesiada M., Godlowski W., Zhu Z.-H., 2012, *J. Cosmol. Astropart. Phys.*, 3, 16
- Cappelluti N. et al., 2007, *ApJS*, 172, 341
- Cen R., Gott J. R., III, Ostriker J. P., Turner E. L., 1994, *ApJ*, 423, 1
- Chae K.-H., 2003, *MNRAS*, 346, 746
- Chae K.-H., 2010, *MNRAS*, 402, 2031
- Chae K.-H. et al., 2002, *Phys. Rev. Lett.*, 89, 151301
- Chiappetti L. et al., 2013, *MNRAS*, 429, 1652
- Choi Y.-Y., Park C., Vogeley M. S., 2007, *ApJ*, 658, 884
- Claeskens J.-F., 1999, PhD thesis, Université de Liège, Belgium
- Desai S. et al., 2012, *ApJ*, 757, 83
- Driver S. P. et al., 2009, *Astron. Geophys.*, 50, 050000
- Elyiv A. et al., 2012, *A&A*, 537, A131
- Elyiv A., Melnyk O., Finet F., Pospieszalska-Surdej A., Chiappetti L., Pierre M., Sadibekova T., Surdej J., 2013, *MNRAS*, 434, 3305
- Fukugita M., Turner E. L., 1991, *MNRAS*, 253, 99
- Fukugita M., Futamase T., Kasai M., 1990, *MNRAS*, 246, 24p
- Guzzo L., Le Fèvre O. eds 2010, *AIP Conf. Proc. Vol. 1241, Invisible Universe. Am. Inst. Phys., New York*, p. 39
- Hezaveh Y. D., Holder G. P., 2011, *ApJ*, 734, 52
- Huterer D., Keeton C. R., Ma C.-P., 2005, *ApJ*, 624, 34
- Keeton C. R., II 1998, PhD thesis, Harvard University
- Keeton C. R., 2002, *ApJ*, 575, L1
- Keeton C. R., Kochanek C. S., 1998, *ApJ*, 495, 157
- Keeton C. R., Kochanek C. S., Falco E. E., 1998, *ApJ*, 509, 561
- Kochanek C. S., 1992, *ApJ*, 384, 1
- Kochanek C. S. et al., 2000, *ApJ*, 543, 131
- Koopmans L. V. E., Treu T., Bolton A. S., Burles S., Moustakas L. A., 2006, *ApJ*, 649, 599
- Koopmans L. V. E. et al., 2009, *ApJ*, 703, L51
- Kormann R., Schneider P., Bartelmann M., 1994, *A&A*, 284, 285
- Mandelbaum R., van de Ven G., Keeton C. R., 2009, *MNRAS*, 398, 635
- Maoz D., Rix H.-W., 1993, *ApJ*, 416, 425
- Melnyk O. et al., 2013, *A&A*, 557, A81
- Mitchell J. L., Keeton C. R., Frieman J. A., Sheth R. K., 2005, *ApJ*, 622, 81
- Ofek E. O., Rix H.-W., Maoz D., 2003, *MNRAS*, 343, 639
- Oguri M., Marshall P. J., 2010, *MNRAS*, 405, 2579
- Oguri M. et al., 2008, *AJ*, 135, 512
- Oguri M. et al., 2012, *AJ*, 143, 120
- Park C., Choi Y.-Y., 2005, *ApJ*, 635, 29
- Peebles P., 1993, *Principles of Physical Cosmology*. Princeton Univ. Press, Princeton, NJ
- Salmon D., Cuillandre J.-C., Barrick G., Thomas J., Ho K., Matsushige G., Benedict T., Racine R., 2009, *PASP*, 121, 905
- Sheth R. K. et al., 2003, *ApJ*, 594, 225
- Sluse D., Chantry V., Magain P., Courbin F., Meylan G., 2012, *A&A*, 538, A99
- Sonnenfeld A., Treu T., Gavazzi R., Suyu S. H., Marshall P. J., Auger M. W., Nipoti C., 2013, *ApJ*, 777, 98
- Surdej J. et al., 1993, *AJ*, 105, 2064
- Suyu S. H. et al., 2012, *ApJ*, 750, 10
- Turner E. L., 1980, *ApJ*, 242, L135
- Turner E. L., 1990, *ApJ*, 365, L43
- Turner E. L., Ostriker J. P., Gott J. R., 1984, *ApJ*, 284, 1
- van de Ven G., Mandelbaum R., Keeton C. R., 2009, *MNRAS*, 398, 607

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.