

EPJ Web of Conferences **93**,01047 (2015)
 DOI: 10.1051/epjconf/20159301047
 © Owned by the authors, published by EDP Sciences, 2015

Mixed-symmetry octupole and hexadecapole excitations in $N=52$ isotones

Andreas Hennig^{1,a}, Mark Spieker¹, Volker Werner^{2,3}, Tan Ahn^{2,4}, Vassia Anagnostatou^{2,5}, Nathan Cooper², Vera Derya¹, Michael Elvers^{1,2}, Janis Endres¹, Phil Goddard^{2,5}, Andreas Heinz^{2,6}, Richard O. Hughes^{2,7}, Gabriela Ilie^{2,8}, Milena N. Mineva⁹, Simon G. Pickstone¹, Pavel Petkov^{1,9}, Norbert Pietralla³, Desirée Radeck^{1,2}, Tim J. Ross^{5,7}, Deniz Savran^{10,11}, and Andreas Zilges¹

¹ Institut für Kernphysik, Universität zu Köln, D-50937 Köln, Germany

² Wright Nuclear Structure Laboratory, Yale University, New Haven, Connecticut 06520, USA

³ Institut für Kernphysik, Technische Universität Darmstadt, D-64289 Darmstadt, Germany

⁴ National Superconducting Cyclotron Laboratory NSCL, Michigan State University, East Lansing, Michigan 48824, USA

⁵ Department of Physics, University of Surrey, Guildford, GU2 7XH, UK

⁶ Fundamental Fysik, Chalmers Tekniska Högskola, SE-41296 Göteborg, Sweden

⁷ University of Richmond, Richmond, Virginia 23173, USA

⁸ National Institute for Physics and Nuclear Engineering, Bucharest-Magurele, RO-77125, Romania

⁹ Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, BG-1784 Sofia, Bulgaria

¹⁰ ExtreMe Matter Institute EMMI and Research Division, GSI, D-64291 Darmstadt, Germany

¹¹ Frankfurt Institute for Advanced Studies FIAS, D-60438 Frankfurt a.M., Germany

Abstract. In addition to the well-established quadrupole mixed-symmetry states, octupole and hexadecapole excitations with mixed-symmetry character have been recently proposed for the $N = 52$ isotones ^{92}Zr and ^{94}Mo . We performed two inelastic proton-scattering experiments to study this kind of excitations in the heaviest stable $N = 52$ isotope ^{96}Ru . From the combined experimental data of both experiments absolute transition strengths were extracted.

1 Introduction

Isovector excitations of valence-shell nucleons are usually denoted as mixed-symmetry states (MSS) [1]. They are predicted in the proton-neutron version of the Interacting Boson Model (IBM-2) [2–4] and can be distinguished from fully-symmetric states (FSS) by their F -spin quantum number [5]. As an experimental signature for MSS, the IBM-2 predicts strong $M1$ transitions to their symmetric counterparts with transition matrix elements in the order of $1 \mu_N$. The collective structure of low-lying states in near-spherical, vibrational nuclei is dominated by the quadrupole degree of freedom. By now, mixed-symmetry quadrupole excitations in vibrational nuclei are well established as collective features near closed shells [6]. In addition to the quadrupole degree of freedom, mixed-symmetry excitations of octupole and hexadecapole character have been proposed in the $N = 52$ isotones ^{92}Zr and ^{94}Mo [7–9]. The identification is based on remarkably strong $M1$ transitions between the lowest-lying 3^- and 4^+ states. Recently, the strong $M1$ transition between the lowest-lying 4^+ states in ^{94}Mo was successfully described by including g -boson excitations in IBM-2 calculations [9], suggesting FS and MS one-phonon hexadecapole admixtures in the 4_1^+ and 4_2^+ states, respectively. It

is the purpose of the present work to study possible mixed-symmetry octupole and hexadecapole states in the heaviest stable $N = 52$ isotope ^{96}Ru .

2 Experiments

The determination of absolute transition strengths requires the measurement of spins and parities of excited states, γ -decay branching ratios, multipole mixing ratios, and nuclear level lifetimes. For this purpose, two inelastic proton-scattering experiments were performed. In a first experiment, performed at the Wright Nuclear Structure Laboratory (WNSL) at Yale University, USA, a proton beam with an energy of $E_p = 8.4$ MeV impinged on a $106 \mu\text{g}/\text{cm}^2$ enriched ^{96}Ru target, supported by a ^{12}C backing with a thickness of $14 \mu\text{g}/\text{cm}^2$. The scattered protons were detected in coincidence with de-exciting γ -rays using five silicon particle detectors and eight BGO-shielded Clover-type HPGe detectors, respectively. From the acquired $p\gamma$ coincidence data γ -decay branching ratios were extracted, while the additionally acquired $\gamma\gamma$ coincidence data were used to determine spins and multipole mixing ratios by means of a $\gamma\gamma$ angular correlation analysis.

In order to extract nuclear level lifetimes in the fs range, we performed a second proton scattering experiment at the Institute for Nuclear Physics at the Univer-

^ae-mail: hennig@ikp.uni-koeln.de




Figure 1. (Color online) Excerpt of the experimental level scheme of ^{96}Ru . $M1$, $E1$, and $E2$ transitions are indicated with black, blue, and red arrows, respectively. The width of the arrows is proportional to the γ -decay branching ratios. Along with the $2_{ms}^+ (\equiv 2_3^+) \rightarrow 2_1^+$ transition [13], $M1$ transitions with sizeable strengths were observed between the lowest-lying 3^- and 4^+ states.

sity of Cologne, Germany. The same target as used for the experiment at WNSL was bombarded with a beam of 7.0 MeV protons. The scattered protons were detected with the new particle detector array SONIC embedded within the γ -spectrometer HORUS to allow for a coincident detection of scattered protons and de-exciting γ -rays. Nuclear level lifetimes were extracted by means of the Doppler-shift attenuation method (DSAM) [10] from the $p\gamma$ coincidence data. Since the initial direction and velocity of the recoil nucleus, as well as its excitation energy can be extracted from the energy of the scattered proton, the $p\gamma$ coincidence yields several advantages for the DSAM measurement [11]:

- The angle θ_γ between the direction of the γ -ray emission and the direction of motion of the recoil nucleus can be extracted on an event-by-event basis.
- Feeding from higher-lying states is eliminated by gating on the excitation energy.
- Peak centroids can be extracted from proton-gated γ -ray spectra, yielding an increased peak-to-background ratio.

The slowing-down process of the ^{96}Ru recoil nuclei in the target and stopper materials was modeled by means of the Monte-Carlo simulation program DSTOP96 [12]. A comparison of the calculated Doppler-shift attenuation factor with the experimentally determined value finally yields the nuclear level lifetime.

3 Experimental results

From the combined experimental data of both experiments absolute transition strengths were calculated. The results concerning one-phonon mixed-symmetry states in ^{96}Ru are shown in Figure 1, pointing out $M1$ transitions with sizeable strengths of $0.14(4) \mu_N^2$ and $0.90(18) \mu_N^2$ between the low-lying 3^- and 4^+ states, respectively. Based on

their absolute $M1$ transition strengths, the $3_2^{(-)}$ state at $E_x = 3077$ keV and the 4_2^+ state at $E_x = 2462$ keV are likely candidates to show mixed-symmetry one-phonon octupole and hexadecapole contributions, respectively.

Acknowledgments

This work is supported by the DFG under grant Nos. (ZI-510/4-2) and SFB 634, the U.S. Department of Energy Grant No. DE-FG02-01ER40609, and the BMBF Grant No. 05P12RDFN8. P.P. is grateful for the financial support of the Bulgarian Science Fund under contract DFNI-E 01/2. D.S. acknowledges support by the Alliance Program of the Helmholtz Association (HA216/EMMI). D.R. and D.S. acknowledge the German Academic Exchange Service (DAAD) for financial support. S.G.P. and M.S. are supported by the Bonn-Cologne Graduate School of Physics and Astronomy.

References

- [1] F. Iachello, Phys. Rev. Lett. **53**, 1427 (1984)
- [2] A. Arima, and F. Iachello, Phys. Rev. Lett. **35**, 1069 (1975)
- [3] A. Arima, T. Otsuka, F. Iachello, and I. Talmi, Phys. Lett. B **66**, 205 (1977)
- [4] T. Otsuka, A. Arima, and F. Iachello, Nucl. Phys. A **309**, 1 (1978)
- [5] P. van Isacker, K. Heyde, J. Jolie, and A. Sevrin, Ann. Phys. **171**, 253 (1986)
- [6] N. Pietralla, P. von Brentano, and A. F. Lisetskiy, Prog. Part. Nucl. Phys. **60**, 225 (2008)
- [7] C. Fransen, N. Pietralla, Z. Ammar, D. Bandyopadhyay, N. Boukharouba, P. von Brentano, A. Dewald, J. Gableske, A. Gade, J. Jolie, U. Kneissel, S. R. Leshner, A. F. Lisetskiy, M. T. McEllistrem, M. Merrik, H. H. Pitz, N. Warr, V. Werner, and S. W. Yates, Phys. Rev. C **67**, 024307 (2003)
- [8] M. Scheck, P. A. Butler, C. Fransen, V. Werner, and S. W. Yates, Phys. Rev. C **81**, 064305 (2010)
- [9] R. J. Casperson, V. Werner, and S. Heinze, Phys. Lett. B **721**, 51 (2013)
- [10] T. K. Alexander and J. S. Foster, Adv. Nucl. Phys. **10**, 197 (1978)
- [11] G. G. Seaman, N. Benczer-Koller, M. C. Bertin, and J. R. MacDonald, Phys. Rev. **188**, 1706 (1969)
- [12] P. Petkov, J. Gableske, O. Vogel, A. Dewald, P. von Brentano, R. Krücken, R. Peusquens, N. Nicloay, A. Gizon, J. Gizon, D. Bazzacco, C. Rossi-Alvarez, S. Lunardi, P. Pavan, D. R. Napoli, W. Andrejtscheff, and R. V. Jolos, Nucl. Phys. A **640**, 293 (1998)
- [13] N. Pietralla, C. J. Barton, R. Krücken, C. W. Beausang, M. A. Caprio, R. F. Casten, J. R. Copper, A. A. Hecht, H. Newman, J. R. Novak, and N. V. Zamfir, Phys. Rev. C **64**, 031301(R) (2001)