



**CIRANO**  
Centre interuniversitaire de recherche  
en analyse des organisations

**Série Scientifique**  
*Scientific Series*

**97s-06**

**GARCH for Irregularly  
Spaced Financial Data:  
The ACD-GARCH Model**

*Eric Ghysels, Joanna Jasiak*

Montréal  
Février 1997

## **CIRANO**

Le CIRANO est une corporation privée à but non lucratif constituée en vertu de la Loi des compagnies du Québec. Le financement de son infrastructure et de ses activités de recherche provient des cotisations de ses organisations-membres, d'une subvention d'infrastructure du ministère de l'Industrie, du Commerce, de la Science et de la Technologie, de même que des subventions et mandats obtenus par ses équipes de recherche. La *Série Scientifique* est la réalisation d'une des missions que s'est données le CIRANO, soit de développer l'analyse scientifique des organisations et des comportements stratégiques.

*CIRANO is a private non-profit organization incorporated under the Québec Companies Act. Its infrastructure and research activities are funded through fees paid by member organizations, an infrastructure grant from the Ministère de l'Industrie, du Commerce, de la Science et de la Technologie, and grants and research mandates obtained by its research teams. The Scientific Series fulfils one of the missions of CIRANO: to develop the scientific analysis of organizations and strategic behaviour.*

## **Les organisations-partenaires / The Partner Organizations**

- École des Hautes Études Commerciales.
- École Polytechnique.
- McGill University.
- Université de Montréal.
- Université du Québec à Montréal.
- Université Laval.
- MEQ.
- MICST.
- Avenor.
- Banque Nationale du Canada.
- Bell Québec.
- Fédération des caisses populaires de Montréal et de l'Ouest-du-Québec.
- Hydro-Québec.
- La Caisse de dépôt et de placement du Québec.
- Raymond, Chabot, Martin, Paré.
- Société d'électrolyse et de chimie Alcan Ltée.
- Télélobe Canada.
- Ville de Montréal.

Ce document est publié dans l'intention de rendre accessibles les résultats préliminaires de la recherche effectuée au CIRANO, afin de susciter des échanges et des suggestions. Les idées et les opinions émises sont sous l'unique responsabilité des auteurs, et ne représentent pas nécessairement les positions du CIRANO ou de ses partenaires.

*This paper presents preliminary research carried out at CIRANO and aims to encourage discussion and comment. The observations and viewpoints expressed are the sole responsibility of the authors. They do not necessarily represent positions of CIRANO or its partners.*

# GARCH for Irregularly Spaced Financial Data: The ACD-GARCH Model\*

*Eric Ghysels<sup>†</sup>, Joanna Jasiak<sup>‡</sup>*

## Résumé / Abstract

Nous développons une classe de modèles ARCH pour les séries temporelles échantillonnées à intervalles inégaux comme des observations liées à des transactions de marché. Notre approche est fondée sur la méthode d'agrégation temporelle pour les modèles ARCH de Drost et Nijman (1993) et de Drost et Werker (1994), et d'autre part sur le modèle autorégressif des moyennes conditionnelles des durées entre les transactions financières de Engle et Russell (1996). La classe de modèles présentée ici est nommée ACD-GARCH. Ce type de modèles peut être défini comme un GARCH aux coefficients aléatoires où la durée entre les transactions détermine la dynamique des paramètres. Le ACD-GARCH devient un modèle bivarié quand sa formation admet les interactions entre les volatilités des rendements passés et les durées et vice-versa. Sinon, la série de durées est considérée exogène par rapport au processus de volatilité. Cette condition est exigée dans l'estimation du modèle ACD-GARCH en deux étapes. La spécification bivariée nous permet de tester l'existence de la causalité de type Granger entre les volatilités et les durées. Sous conditions générales, diverses procédures d'estimation par la méthode de moments généralisés sont considérées, dont certaines fournissent les estimateurs, à la fois de type GMM et de type QMLE. Pour ce qui est des applications, nous présentons une étude empirique basée sur les données de transactions du titre IBM en 1993. Nos résultats indiquent que la volatilité des rendements sur les prix d'actions de IBM causent, au sens de Granger, les durées entre les transactions. Nous observons aussi que la persistance du processus GARCH diminue fortement quand on introduit les durées dans la formulation du modèle.

*We develop a class of ARCH models for series sampled at unequal time intervals set by trade or quote arrivals. Our approach combines insights from the temporal aggregation for GARCH models discussed by Drost and Nijman (1993) and Drost and Werker (1994), and the autoregressive conditional duration model of Engle and Russell (1996) proposed to model the spacing between consecutive financial transactions. The class of models we introduce*

---

\* Correspondence Address: Eric Ghysels, CIRANO, 2020 University Street, 25th floor, Montréal, Qc, Canada H3A 2A5 Tel: (514) 985-4000 Fax: (514) 985-4039 e-mail: ghyselse@cirano.umontreal.ca  
The first author would like to acknowledge the financial support of the Social Sciences and Humanities Research Council of Canada. We would also like to thank Robert Engle, Nour Meddahi and the participants of the Financial Econometrics Workshops at Tilburg University, the RIBES and Tinbergen Institute seminars in Rotterdam and the Universidad Carlos III in Madrid for their comments.

<sup>†</sup> Pennsylvania State University and CIRANO

<sup>‡</sup> York University

*here will be called ACD-GARCH. It can be described as a random coefficient GARCH, or doubly stochastic GARCH, where the durations between transactions determine the parameter dynamics. The ACD-GARCH model becomes genuinely bivariate when past asset return volatilities are allowed to affect transaction durations and vice versa. Otherwise the spacings between trades are considered exogenous to the volatility dynamics. This assumption is required in a two-step estimation procedure. The bivariate setup enables us to test for Granger causality between volatility and intra-trade durations. Under general conditions we propose several GMM estimation procedures, some having a QMLE interpretation. As illustration we present an empirical study of the IBM tick-by-tick data. We find that volatility of IBM stock prices Granger causes intra-trade durations. We also find that the persistence in GARCH drops dramatically once intra-trade durations are taken into account.*

**Mots Clés :** Hétéroscedasticité, activité de marché, volatilité, causalité, modèles de durées

**Keywords :** Heteroskedasticity, Market Activity, Tick-by-Tick Data, Volatility, Causality, Duration Models

JEL : C22, C32, D41, G1

# 1 Introduction

The Autoregressive Conditional Heteroskedastic (ARCH) class of models introduced by Engle (1982) is widely used to capture the temporal dynamics of asset return volatility. Several recent papers, including Bera and Higgins (1995), Bollerslev, Chou and Kroner (1992), Bollerslev, Engle and Nelson (1994), Diebold and Lopez (1995) and Palm (1996) survey the extensive number of applications to financial time series and the impressive theoretical developments which took place over the last decade. In their generic form ARCH processes model the conditional variance as a measurable function of past returns. The volatility dynamics is explicitly modelled in discrete time and while the sampling frequency is in general left unspecified, data are assumed to be equally spaced in time. This setup is adequate for the many applications involving financial time series sampled weekly, daily, or intradaily at hourly or finer time intervals. There is a growing interest in the modelling of financial time series sampled at the transaction-based frequency. Such data became more widely available over the last years due to the implementation of the electronic trading systems on major financial markets.

In this paper we introduce a class of ARCH models for return series sampled at time intervals set by the trade arrivals. Any general formulation of GARCH for irregularly spaced financial data is extremely complex. We propose a class of processes which provides an *approximation* to the analytically and computationally intractable general case. Our approach combines insights from the temporal aggregation procedure for GARCH models proposed by Drost and Nijman (1993) and Drost and Werker (1996) and the autoregressive conditional duration model (ACD) of Engle and Russell (1995) for intra-trade duration sequences. The class of models introduced here will be called ACD-GARCH, although strictly speaking they do not belong to the GARCH class of models. Indeed, the ACD-GARCH models are intrinsically bivariate as they involve past returns and the time intervals between past transactions.<sup>1</sup> They can also be viewed as a time deformed GARCH model and therefore there is some legitimacy to call it GARCH. An ACD-GARCH model is in fact a random coefficient GARCH model, or doubly stochastic GARCH, where the duration between transactions determine the dynamics of the entire parameter vector. The parameter behavior is described by the temporal aggregation formulas of weak GARCH. In this sense our specification differs from the recently proposed ISAR-ARCH model of Pai, Polasek

---

<sup>1</sup>Hence, the conditional volatility is not exclusively a measurable function of past returns, a characteristic attributed to the definition of the class of ARCH models, see e.g. Bollerslev, Engle and Nelson (1994).

and Kozumi (1995) where the unequal spacing in data is accommodated by ad hoc varying autoregressive coefficients in the conditional mean of returns and the coefficients on lagged squared errors in the conditional variance equation.

Since the ACD-GARCH model is genuinely bivariate, past volatilities may determine the transaction durations and vice-versa. In this context, causal relationships between the volatility and durations arise as an interesting issue to investigate. In the univariate ACD model discussed by Engle and Russell (1996), the spacings between trades is not a function of the volatility process. We propose GMM as an estimation method applicable under general conditions. A sequential procedure can be adopted when we assume that the spacing of transactions in time is weakly exogenous in the sense of Engle, Hendry and Richard (1983). In our empirical illustration we use tick-by-tick data on IBM stock extracted from the ISSM (Institute for the Study of Security Markets) 1993 data set. We estimate ACD-GARCH models and introduce tests for causality between IBM stock volatility and intra-trade durations. We find such causality to be significant.

The paper is organized as follows: The general structure is presented in section 2. Next, we discuss the estimation methods in section 3. The empirical study IBM tick-by-tick data is presented in section 4. Conclusions appear in section 5.

## 2 The ACD-GARCH class of models

We consider asset prices recorded on a real time basis. A simple example would be transaction prices of a particular stock where each transaction has a time stamp. We assume that observations are represented by the pairs  $(y_{t_i}, t_i)_{i=0}^N$  where  $y_{t_i} = \ln p_{t_i}$ , the log of the transaction price recorded at time  $t_i$ . The data are unequally spaced and tied to the point process of transaction events. In many cases the arrival times are modelled as a Poisson process, possibly non-homogenous with the intensity determined by a set of covariates.<sup>2</sup> Engle and Russell (1995) argued that the class of Poisson processes does not fit well the financial tick-by-tick data as the occurrence of market transactions shows a strong temporal dependence. They suggested an autoregressive conditional durations model, henceforth denoted ACD, very similar to ARCH

---

<sup>2</sup>This is the so-called proportional hazard model. See e.g. Lancaster (1990) for a detailed discussion. Gouriéroux, Jasiak and Le Fol (1996) propose nonparametric methods for hazard models and their associated duration and survival functions applied to French tick-by-tick data from the Bourse de Paris.

to accommodate the persistence in intra-trade durations. We will focus almost exclusively our attention on the sequential trading process taking each tick into account. The framework we develop, however, is generic and general with respect to the transaction event. For instance, the ticks may be restricted to large block trades instead of all trades, or may be the time that elapses before a predetermined number of shares is traded (say 10000 shares for instance).

In a first subsection we present the structure of the models. Next we provide a time deformation interpretation to ACD-GARCH models. A third subsection covers some special cases and the final subsection discusses causality between trading and volatility.

## 2.1 The Model Structure

Following the approach of Engle and Russell we call a duration the difference between subsequent transaction times  $t_{i+1} - t_i, i = 0 \dots N$ . The conditional expected intra-trade time  $\psi_{i+1} = E \left[ (t_{i+1} - t_i) \mid (t_j - t_{j-1})_{j \leq i} \right]$  can be represented as a measurable function of past durations. In analogy to the GARCH (1,1)

$$\psi_{i+1} = \omega^d + \alpha^d (t_i - t_{i-1}) + \beta^d \psi_i, \quad (2.1)$$

where the parameter vector  $\theta^d = (\omega^d, \alpha^d, \beta^d)$  is indexed by  $d$  to indicate that the parameters pertain to the durations dynamics. Dividing the durations by their conditional means yields a sequence of i.i.d. variables:

$$(t_{i+1} - t_i) / \psi_{i+1} \sim g(\psi_{i+1}; \theta^d), \quad i = 0, \dots, N \quad (2.2)$$

where  $g$  denotes the assumed distribution function, like for example an exponential or Weibull [see Engle and Russell (1995)]. Eventually, the conditional mean duration equation (2.1) can be extended from ACD (1,1) to a more general setup:

$$E \left[ (t_{i+1} - t_i) \mid (t_j - t_{j-1})_{j \leq i}, (y_{t_j} - y_{t_{j-1}})_{j \leq i}, (x_{t_j})_{j \leq i} \right] = \psi_{i+1} \equiv \psi \left[ (t_j - t_{j-1})_{j=i}^{i-p+1}, (\psi_j)_{j=i}^{i-q+1}, (y_{t_j} - y_{t_{j-1}})_{j=i}^{i-r+1}, (x_{t_j})_{j=i}^{i-s+1}; \theta^d \right] \quad (2.3)$$

where  $x_{t_i}$  represents a state variable process which may include deterministic components or other variables such as the trading volume of the transactions occurring at or before  $t_i$ . Whenever past returns or

especially past volatilities appear in the above equation, we allow other market dependent variables to affect the trading process beyond the dynamics captured by the autoregressive structure in (2.1). This suggests of course testing for causality, an issue which will be discussed later.

Consider now the asset price process  $\{y_{t_i}\}, i = 0, \dots, N$ . The return series is defined as a sequence of differences  $\{(y_{t_i} - y_{t_{i-1}})\}$ , indexed by the transaction numbers  $i = 1, \dots, N$ . Note that the current return value is expected to be observed over the period  $\psi_{i+1}$  before the next trade arrives.

Ultimately we would like to construct the expected volatility at the next tick (or large block trade or sale of the next 10000 share, depending on the definition of events), more precisely:

$$\bar{\sigma}_{i+1}^2 \equiv E \left[ (y_{t_{i+1}} - y_{t_i})^2 \mid (t_j - t_{j-1})_{j \leq i}, (y_{t_j} - y_{t_{j-1}})_{j \leq i}, (x_{t_j})_{j \leq i} \right]$$

where we assume that  $(y_{i+1} - y_i)$  has conditional mean zero. This will be difficult to accomplish, however, if we aim for analytic expressions and stay within the class of ARCH models. Indeed, the ARCH class of models, as originally defined by Engle (1982) does not temporally aggregate. Since we deal with unequally spaced data temporal aggregation is crucial. Hence, a first step is to weaken the definition of ARCH such that temporal aggregation is possible. This was developed by Drost and Nijman (1993). In particular, we assume that the first differences  $\{(y_{t_i} - y_{t_{i-1}}), i = 1, \dots, N\}$  have finite fourth moments and form a stationary sequence satisfying the weak GARCH (1,1) definition [see Drost and Nijman, 1993]. Especially we assume that  $\{(y_{t_i} - y_{t_{i-1}}), i = 1, \dots, N\}$  consists of selected observations drawn from a underlying data generating process evolving on a time grid set by equal, very small time increments.<sup>3</sup> Hence, for the moment we will remain in a discrete time setting which will cause us some difficulty. Some of these difficulties will be solved by adopting a continuous time framework to which we turn our attention later in this section. Drost and Nijman (1993) have shown that the class of weak GARCH (1,1) processes is closed under temporal aggregation. We will rely on this result to model the dynamics of the volatility process over unequal intra-trade intervals.

Let us denote  $\varepsilon_i = y_{t_i} - y_{t_{i-1}}$  and fix the next trade  $t_{i+1} = t_i + \delta$ , i.e. when time is measured in seconds the intra-trade duration will be  $\delta$ . For any *given*  $\delta$  the weak GARCH (1,1) models provides the best linear projection

---

<sup>3</sup>In practice, we will consider as very small time unit, the accuracy limit of the records which usually is one second.



$$\sigma_{i+1}^2(\delta) \equiv P_\delta \left[ \varepsilon_{i+1}^2 | 1, (\varepsilon_j)_{j \leq 1}, (\varepsilon_j^2)_{j \leq i} \right] \quad (2.4)$$

where  $P_\delta$  denotes the best linear projection. The temporal aggregation formula of Drost and Nijman (1983), which will be presented shortly, enable us to compute this linear projection for any given  $\delta$ . There is an important technical issue emerging here which we cannot accommodate properly in this framework. Indeed, the temporal aggregation formula provide a mapping between equally spaced representations with different sampling frequencies and do not strictly speaking apply to unequally spaced data. We will ignore this issue until we adopt a continuous time framework later in this section. The continuous time framework is more suitable to deal with this, but for the moment we will remain with the discrete time setup. Since the ACD model provides us with a conditional duration distribution for  $\delta$ . It is worth noting that  $P_\delta$  is only a function of past returns, a feature which is proper to ARCH models. At this stage, however, we can introduce information about intra-trade durations, as the ACD model allows to compute:

$$\tilde{\sigma}_{i+1}^2 = \int \sigma_{i+1}^2(\delta) g(\delta; \psi_{i+1}, \theta^d) d\delta \quad (2.5)$$

which is the expected conditional volatility based on the distribution function  $g$  appearing in (2.2) with conditional mean  $\psi_{i+1}$ . While the integration in (2.5) is in principle feasible it is analytically intractable and can only be accomplished numerically. To side step this difficulty we propose to use  $\sigma_{i+1}^2(\psi_{i+1})$  which yields the structure for the class of ACD-GARCH models we introduce in this paper. Using the temporal aggregation formula of Drost and Nijman (1993) we can write the ACD-GARCH class of models more explicitly as:

$$\sigma_{i+1}^2 = \omega_i + \alpha_i \varepsilon_i^2 + \beta_i \sigma_i^2 \quad (2.6)$$

where:

$$\omega_i = \psi_{i+1} \omega \frac{1 + (\alpha + \beta)^{\psi_{i+1}}}{1 - (\alpha + \beta)}, \quad (2.7)$$

$$\alpha_i = (\alpha + \beta)^{\psi_{i+1}} - \beta_i, \quad (2.8)$$

$$\beta_i / (1 + \beta_i^2) = \left[ a(\alpha, \kappa, \psi_{i+1}) (\alpha + \beta)^{\psi_{i+1}} - b(\alpha, \beta, \psi_{i+1}) \right] / \left[ a(\alpha, \kappa, \psi_{i+1}) \left\{ 1 + (\alpha + \beta)^{2\psi_{i+1}} \right\} - 2b(\alpha, \beta, \psi_{i+1}) \right], \quad (2.9)$$

$$\begin{aligned}
a(\alpha, \beta, \kappa, \psi_{i+1}) &= \psi_{i+1} (1 - \beta)^2 + 2\psi_{i+1} (\psi_{i+1} - 1) (1 - \alpha - \beta)^2 \times \\
&\quad \left\{ 1 - (\alpha + \beta)^2 + \alpha^2 \right\} / \left[ (\kappa - 1) \left( 1 - (\alpha + \beta)^2 \right) \right] \\
&\quad + 4c(\alpha, \beta, \psi_{i+1}) / \left( 1 - (\alpha + \beta)^2 \right),
\end{aligned} \tag{2.10}$$

$$\begin{aligned}
b(\alpha, \beta, \psi_{i+1}) &= \left\{ \alpha \left( 1 - (\alpha + \beta)^2 \right) + \alpha^2 (\alpha + \beta) \right\} \left( 1 - (\alpha + \beta)^{2\psi_{i+1}} \right) \\
&\quad / \left( 1 - (\alpha + \beta)^2 \right),
\end{aligned} \tag{2.11}$$

$$\begin{aligned}
c(\alpha, \beta, \psi_{i+1}) &= \left\{ \psi_{i+1} (1 - \alpha - \beta) - 1 + (\alpha + \beta)^{\psi_{i+1}} \right\} \times \\
&\quad \left\{ \alpha \left( 1 - (\alpha + \beta)^2 \right) - \alpha^2 (\alpha + \beta) \right\}.
\end{aligned} \tag{2.12}$$

The constant parameters  $\omega$ ,  $\alpha$ ,  $\beta$  in (2.7) - (2.12) characterize the conditional volatility of the underlying process which is defined at the smallest time interval. The parameter  $k$  in (2.9) and (2.10) denotes the kurtosis. For the purpose of parameter identifications we will also use the relationship  $\alpha\psi_{i+1} + \beta\psi_{i+1} = (\alpha + \beta)^{\psi_{i+1}}$ .

When we define an error term  $\eta_i = \varepsilon_i^2 - \sigma_i^2$  such that

$$\varepsilon_i^2 = \omega_i + (\alpha_i + \beta_i) \varepsilon_{i-1}^2 + \eta_i + \beta_{i+1} \eta_{i-1} \tag{2.13}$$

we obtain an alternative representation for the model. It reveals that there are two sources of error in the  $\eta_i$  process, one related to the innovation in the duration process  $(t_{i+1} - t_i) - \psi_{i+1}$  and the other related to the usual GARCH volatility innovation. This mixture of errors suggests that the  $\eta_i$  process is most likely fat-tailed.

Finally, the ACD-GARCH can be written as a system of equations

$$\psi_{i+1} = \omega^d + \alpha^d (t_i - t_{i-1}) + \beta^d \psi_i \tag{2.14}$$

$$\sigma_i^2 = \omega_i + \alpha_i \varepsilon_{i-1}^2 + \beta_i \sigma_{i-1}^2. \tag{2.15}$$

## 2.2 Time deformation

As noted in the introduction, the ACD-GARCH model does not satisfy the weak GARCH, nor the ARCH definitions in the sense that it does not represent the conditional variance as a measurable function defined on the filtration of past returns. In some way the ACD-GARCH model is in fact a stochastic volatility model. It was also noted that the GARCH aggregation formula apply in principle to equally spaced data with different sampling frequencies; a technical issue we ignored when dealing with irregularly spaced series. In this section we address some of these issues by noting that the class of ACD-GARCH models could also be viewed as a time deformed GARCH diffusion.<sup>4</sup> Following Drost and Werker (1996) one can define a diffusion:

$$dy(t) = \sigma(t)dW_1(t) \quad (2.16)$$

$$d\sigma(t)^2 = \delta \left( \omega - \sigma(t)^2 \right) dt + \sqrt{2\lambda\delta} \sigma(t)^2 dW_2(t) \quad (2.17)$$

where  $W_1(t)$  and  $W_2(t)$  are two independent Brownian motions with  $\omega > 0$ ,  $\delta > 0$ , and  $\lambda \in (0, 1)$ . Using the results of Drost and Werker (1996) one interprets the ACD-GARCH model as a time deformed GARCH diffusion with time deformation  $\psi_{i+1}$  yielding:

$$\omega_i = \psi_{i+1} \omega \{1 - \exp(-\psi_{i+1}\delta)\} \quad (2.18)$$

$$\alpha_i = \exp(-\psi_{i+1}\delta) - \beta_i \quad (2.19)$$

$$\beta_i = (\beta_i^2 + 1) (c_i) (\exp(-\psi_{i+1}\delta) - 1) / [c_i \{1 + \exp(-2\psi_{i+1}\delta)\} - 2] \quad (2.20)$$

where

$$c_i = \frac{[4 \{ \exp(-\psi_{i+1}\delta) - 1 + \psi_{i+1}\delta \} + 2\psi_{i+1}\delta \{1 + \psi_{i+1}\delta(1 - \lambda) / \lambda \}]}{[1 - \exp(-2\psi_{i+1}\delta)]} \quad (2.21)$$

---

<sup>4</sup>For more details on time deformation and its applications to financial data, see Ghysels and Jasiak (1994) and Ghysels, Gouriéroux and Jasiak (1996a, b).

Equations (2.18) through (2.21) yield the ACD-GARCH appearing in (2.6) as an *exact* discretization of the diffusion (2.16-17) with the ACD conditional mean  $\varphi_{i+1}$  as the directing process. The time deformation is purely exogenous to volatility when we restrict our attention to pure ACD models. However, in the more general case where past volatility affects expected trade durations we may have a partially endogenous directing process.<sup>5</sup> This time deformation interpretation clarifies, as noted earlier, that we are in fact dealing with a class of stochastic volatility models.

### 2.3 Special cases

The ACD-GARCH provides a convenient framework encompassing some well-known models. We discuss two special cases which arise when additional restrictions are imposed on the duration dynamics.

The first is the equally spaced traditional GARCH(1,1) model which is obtained by setting  $\psi_{i+1} \equiv \psi \forall i$ . It applies to *unequally* spaced data with a constant conditional duration distribution. Such a situation would occur when the point process governing transactions is in fact a time homogeneous Poisson process. We noted in the introduction that the event process is not exclusively tied to every transactions tick. For instance, the intra-trade durations may reflect to occurrence of large block trades or other specific trades. If these arrive at a Poisson rate we can simply use standard GARCH processes to predict the volatility of returns despite the fact that such trades are irregularly spaced.

Another special case worth mentioning is the situation where  $\psi_i$  follows a deterministic periodic pattern, i.e.  $\psi_i = \psi_{i-S}$  for  $\forall i$  and given  $S$ . Then, one obtains a periodic GARCH model, introduced by Bollerslev and Ghysels (1996), for unequally spaced data. Such a situation occurs if transactions arrive as a Poisson process which is time non-homogeneous with a periodic pattern in the hazard rates.

### 2.4 Causality

An interesting issue to investigate in the context of ACD-GARCH is the causality from the volatility process to the intra-trade duration sequence.

---

<sup>5</sup>This distinction has consequences with respect to estimation. If we were to estimate directly the diffusion instead of the discrete time weak GARCH we could in principle use the methods proposed by Conley et al. (1996) when the directing process is exogenous. Otherwise one needs to adopt the methods discussed in Ghysels and Jasiak (1996).

More explicitly, the question is whether past returns or past volatilities should appear on the right hand side of the (2.3).

There are various ways to test the exclusion restrictions associated with the absence of returns and/or volatility in the intra-trade durations process. The most appealing, though not the most optimal, is to estimate the ACD model without any consideration for volatility, i.e. the univariate process of intra-trade durations as in Engle and Russell (1996). The ACD-GARCH is estimated next using the return data and tests are applied to the bivariate process of residuals generated by the two univariate models. Such a procedure is quite similar to recently proposed causality tests in conditional variances across various financial asset price movements. Cheung and Ng (1996) developed such a procedure to test causality in conditional variances based on the early work by Haugh (1976) and McLeod and Li (1983) based on estimating the univariate ARCH models and using the cross-correlation function (CCF) of the squared standardized residuals. In theory two issues can be investigated, namely the traditional causality between the conditional mean of returns and the intra-trade durations as well as the causality between the volatility and durations, i.e. the conditional variance-mean causality.

The first problem can be addressed by applying the CCF-based Pierce-Haugh test to the standardized residuals, i.e. the i.i.d sequence  $d_i$  and the return innovation process  $\varepsilon_i$ :

$$d_i = (t_{i+1} - t_1) / \psi_{i+1}, \quad (2.22)$$

$$e_i = (y_i - y_{i-1}) / \sigma_i, \quad (2.23)$$

For independent sequences  $\{d_i\}$  and  $\{e_i\}$ , the existence of their second moments implies that:

$$\begin{pmatrix} \sqrt{N} \hat{c}_{de}(k) \\ \sqrt{N} \hat{c}_{de}(k') \end{pmatrix} \rightarrow AN \left( \begin{bmatrix} 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right), k \neq k' \quad (2.24)$$

where  $\hat{c}_{de}(k)$  is the sample cross-correlation at lag  $k$ ,

$$\hat{c}_{de}(k) = \hat{c}v_{de}(k) / \sqrt{\hat{c}v_{dd}(0) \hat{c}v_{ee}(0)}, \quad (2.25)$$

and  $\hat{c}v_{de}(k)$  is the  $k$ th lag sample cross covariance [see Hannan (1970), Cheng and Ng (1996)]:

$$\hat{c}v_d(k) = N^{-1} \sum (d_i - \bar{d})(e_{i-k}).$$

Causality at lag  $k$  can be tested by comparing  $\sqrt{N}\hat{c}_{de}(k)$  to a critical value from the standard normal distribution. Alternatively, a chi-square test statistic based on:

$$S_k = N \sum_{i=j}^k \hat{c}_{de}(i)^2 \quad (2.26)$$

can be computed. It is chi-square distributed with  $(k - j + 1)$  degrees of freedom and can be used to test the null hypothesis of no causality from lag  $j$  to lag  $k$ .

The test of causality between the volatility process and the intra-trade durations involves the conditional mean of durations and the conditional variance of returns. Hence the  $k$ th lag sample cross-correlation of the squared standardized residuals  $e_i^2$ , denoted by  $E_i$ , and  $d_i$  are computed:

$$\hat{c}_{dE}(k) = \hat{c}v_{dE}(k) / \sqrt{\hat{c}v_{dd}(0) \hat{c}v_{EE}(0)}, \quad (2.27)$$

where

$$\hat{c}v_{dE}(k) = N^{-1} \sum (d_i - \bar{d})(E_{i-k} - \bar{E}).$$

For  $d_i$  and  $e_i^2$  defined by (2.22) and (2.23) we show in the appendix that under suitable regularity conditions  $\sqrt{N}(\hat{c}_{dE}(k_1), \dots, \hat{c}_{dE}(k_m))$  converges to  $N(0, I_m)$  as  $N \rightarrow \infty$ , where  $k_1, \dots, k_m$  are  $m$  different integers.<sup>6</sup>

It should be recalled that the residual-based causality tests are conservative tests since they are constructed from univariate time series processes. In the context of high-frequency data this negative feature of such tests is probably not of great concern. More elaborate tests based on the genuine bivariate representation of the trade and volatility process can be considered as well. They will be briefly covered in the next section. To conclude it is important to stress the importance of causality testing. Indeed, when trading and volatility are not mutually exogenous we really have to deal with the joint process of  $d_i$  and  $e_i^2$ . Most microstructure models suggest such a bivariate setup (see e.g. O'Hara (1995) for a survey of the literature).

---

<sup>6</sup>For the above results to hold, the conditional mean of the returns has to be included in the model and consistently estimated. We need to remain cautious however, as the eventual causality in mean violates the independence condition and affects the size of the mean-variance causality test [see Cheung and Ng (1996)].

### 3 Estimation

It is assumed that the econometrician has a high-frequency data set of financial time series consisting of observations  $\left( (y_{t_i}, t_i, x_{t_i})_{i=0}^N, Y_o, T_o, X_o \right)$  where  $Y_o, T_o, X_o$  are sets of initial conditions of  $y_{t_i}, t_i$  and  $x_{t_i}$  which contain the  $p, r, s$  lagged presample observations (see equation (2.3)). As noted in section 2, the  $x_{t_i}$  process may include deterministic components as well as other variables included in the ACD-GARCH model, trading volume or past returns being a particularly relevant examples.

It would be tempting to formulate a maximum likelihood estimator (henceforth MLE) for ACD-GARCH models since taken separately both ARCH and ACD models are typically estimated using MLE. However, in estimating the two components simultaneously we encounter a major obstacle to maintaining a fully specified distributional framework. Indeed, Drost and Nijman (1993) argued that the temporal aggregation results hold for a weaker than the standard (strong) definition of ARCH and this requires an estimation via a set of moment conditions or via quasi-MLE or QMLE. The necessary moment conditions are provided by the projection equations which define weak GARCH models. Alternatively, the theoretical likelihood of GARCH could serve as a distributional form for the QMLE. Under certain circumstances, related to the causality issues discussed in section 2.3, we could even consider maintaining a ML estimation approach for the ACD model. Indeed, a two-step procedure can be considered, starting first with the estimation of the ACD parameters followed by the estimation of the conditional variance model for the tick-by-tick data. In a MLE framework such a two-step procedure would amount to a situation where the spacing of transactions in time is weakly exogenous, in the sense of Engle, Hendry and Richard (1983). We first cover the general joint estimation framework in section 3.1 and discuss the two-step procedures in section 3.2. In section 3.1 we focus on the GMM estimation and show a particular GMM estimator having a QMLE interpretation.

#### 3.1 Joint ACD and GARCH estimation

We assume that the parameter vector  $\theta$  has two components, namely (1)  $\theta^d$  a subvector describing the parameters of the ACD model and (2)  $\theta^r$  a subvector which contains the parameters of the conditional mean and variance of tick-by-tick returns. This setup is the same as in section 2.3. When we assume that  $x_{t_i}$  contains at least a constant then the dimension of  $\theta^d$  equals  $p + q + r + s$ , using the notation of (2.4) with  $s \geq 1$ . The vector  $\theta^r$  can also be divided into two components, namely

$\mu^r$  which governs the conditional mean, and may contain as components the constant drift or seasonal dummies and the vector  $\zeta \equiv (\omega, \alpha, \beta)$  which determines equations (2.7) through (2.12). To set the scene for the discussion of the estimation procedure we consider a generic GMM setup based on a set of moment conditions:

$$E f_i(\theta) = 0 \tag{3.1}$$

where  $f_i(\theta) \equiv \left( f_i^d(\theta)', f_i^r(\theta)' \right)'$  and

$$f_i^d(\theta) \equiv f^d \left( (t_j - t_{j-1})_{j=i+1}^{i-p}, (\psi_j)_{j=i}^{i-q}, (y_{t_j} - y_{t_{j-1}})_{j=i}^{i-r}, (x_{t_j})_{j=i}^{i-s}; \theta \right) \tag{3.2}$$

$$f_i^r(\theta) \equiv f^r \left( (t_j - t_{j-1})_{j=i}^{i-p+1}, (\psi)_{j=i}^{i-q+1}, (y_{t_j} - y_{t_{j-1}})_{j=i+1}^{i-r+1}, (x_{t_j})_{j=i}^{i-s+1}; \theta \right) \tag{3.3}$$

It should be noted that  $t_{i+1} - t_i$  enters  $f_i^d(\theta)$  while  $y_{t_{i+1}} - y_{t_i}$  enters  $f_i^r(\theta)$ . Hence, the first set of moment conditions pertains to the durations between transactions while the second set of moment conditions draws on projection equations of the conditional mean and variance of returns. Moreover, within this generic general framework the entire parameter vector  $\theta$  enters both sets of moment conditions. Hence, as we discussed in section 2.3, past volatility  $\sigma_i$  could possibly help to determine expected future durations such as  $\psi_{i+1}$ . This can be accommodated in (2.3) by letting  $x_{t_i} \equiv (1, \sigma_i)$ .<sup>7</sup> In the sequel we consider this general framework. Later, in section 3.2 we examine the more restrictive setting where  $f_i^d(\theta^d)$  does not involve  $\theta^r$ . While in principle a wide variety of choices for  $f_i^d$  and  $f_i^r$  exist there is a rather natural choice which emerges from the discussion in section 2. Although, we cannot consider MLE for the joint duration/conditional volatility model it would be logical to start with distributional properties of the normalized durations, i.e.  $(t_i - t_{i-1})/\psi_i$  in (2.2), to construct  $f_i^d(\theta)$ . It has the advantage that the two-step procedures examined in the next section are special cases of the procedures considered here. For the exponential version of the ACD model the log likelihood function can be written:

---

<sup>7</sup>Needless to say that such a situation could become numerically quite involved. Indeed, with  $q = p = 1, r = 0$  and  $x_{t_i} \equiv (1, \sigma_i)$  in (2.3) it would require recursive calculations of  $\psi_{i+1}$  given  $\sigma_i$  and  $\sigma_{i+1}$  given  $\psi_{i+1}$  and  $\sigma_i$  starting with  $\sigma_0$ .



$$L^E(\theta) \equiv - \sum_{i=1}^N l_i^d(\theta) \equiv - \sum_{i=1}^N [\ln(\psi_i) + (t_i - t_{i-1}) / \psi_i],$$

resulting in the score moment condition:

$$f_i^d(\theta) = \partial l_{i+1}^d(\theta) / \partial \theta^d = [(t_{i+1} - t_i) / \psi_{i+1}(\theta) - 1] / \psi_{i+1}(\theta) \times \partial \psi_{i+1}(\theta) / \partial \theta^d. \quad (3.4)$$

The score function of the ACD model, required in this approach is its derivative with respect to  $\theta^d$ . The dependence on the entire vector  $\theta$  is revealed by making  $\psi_i$  an explicit function of  $\theta$ . The computation of this moment condition is relatively straightforward. As indicated by Engle and Russell, it involves recursive computations of  $\partial \psi_i / \partial \theta^d$  which are quite similar to those encountered in score functions of GARCH models. Engle and Russell also consider a Weibull distributed ACD requiring an appropriate expression for  $\partial l_{i+1}^d(\theta) / \partial \theta^d$ .<sup>8</sup>

We turn our attention now to the second set of moment conditions  $f_i^r(\theta)$ . There are two particular choices for  $f_i^r(\theta)$  that would naturally emerge from the discussion in section 2. The first choice is computationally simple, while the second is more involved but has the advantage of yielding a QMLE interpretation of the resulting GMM estimator. The moment conditions for the first approach are obtained from the projection equations resulting from (2.4) and (2.6). In particular consider the instrument set  $Z_{mi} \equiv \left[ 1, (t_j - t_{j-1})_{j=i}^{i-p+1}, (y_{t_j} - y_{t_{j-1}})_{j=1}^{i-r+1}, (x_{t_j})_{j=i}^{i-r+1} \right]$ , then for  $\varepsilon_{i+1}$  defined in (2.4) we have :

$$E \varepsilon_{i+1} Z'_{mi} = 0. \quad (3.5)$$

For the conditional variance we can use:

$$E [\varepsilon_{i+1}^2 - \omega_{i+1} - (\alpha_{i+1} + \beta_{i+1}) \varepsilon_i^2] Z'_{Vi-1} = 0, \quad (3.6)$$

---

<sup>8</sup>In fact, we could entertain the possibility to model normalized durations via SNP density, [see Gallant and Tauchen (1989)], and use the SNP score in (3.4).

where  $Z_{Vi} \equiv [1, \varepsilon_i, \dots, \varepsilon_{i-l+1}, \varepsilon_i^2, \dots, \varepsilon_{i-k+1}^2]$ . The combination of (3.5) and (3.6) yields:

$$f_i^r(\theta) \equiv \left( \left( \varepsilon_{i+1} Z_{mi}' \right)', \left( [\varepsilon_{i+1} - \omega_{i+1} - (\alpha_{i+1} + \beta_{i+1}) \varepsilon_i^2] Z_{Vi-1}' \right)' \right)' \quad (3.7)$$

which amounts to  $r + p + s + k + l + 2$  moment restrictions. The sample equivalent of (3.1) forms the basis of the following GMM estimation procedure:

$$\hat{\theta}_N^G \equiv \theta \text{Arg min} \left( N^{-1} \sum_{i=1}^N f_i(\theta) \right)' \Omega_N^{-1} \left( N^{-1} \sum_{i=1}^N f_i(\theta) \right), \quad (3.8)$$

where  $\Omega_N^{-1}$  is the optimal weighting matrix. Hansen (1982) showed that matrix should be estimated from:

$$\Omega \equiv E \begin{bmatrix} f_i^d(\theta)' f_i^d(\theta) & f_i^d(\theta)' f_i^r(\theta) \\ f_i^r(\theta)' f_i^d(\theta) & f_i^r(\theta)' f_i^r(\theta) \end{bmatrix}. \quad (3.9)$$

Equation (3.9) emphasizes the fact that crossproducts between  $f_i^d(\theta)$  and  $f_i^r(\theta)$  are involved in the computation of the optimal GMM estimator defined in (3.8). A priori we do not expect these crossproducts to be zero. Indeed, it was shown in (2.13) that the error process  $\eta_i$  is related to the innovation in the duration process error  $(t_{i+1} - t_i) - \psi_{i+1}$  which makes the first element of  $f_i^r(\theta)$  nonorthogonal to  $f_i^d(\theta)$ . It is important to emphasize this feature at this stage as the two-step procedure presented in the next section yields an estimator inefficient relative to  $\hat{\theta}_N^G$  because the off-diagonal elements in (3.9) are set equal to zero.

We denoted the GMM estimator in (3.8) by  $\hat{\theta}_N^G$  to distinguish it from the second GMM estimator, denoted  $\hat{\theta}_N^Q$  where the index  $Q$  refers to a QMLE interpretation of the estimator. We mentioned before that maintaining a fully specified distribution of normalized returns, i.e.  $(y_{t_i} - y_{t_{i-1}}) / \sigma_i \equiv Z_i$ , is not compatible with the temporal aggregation condition and therefore MLE of ACD-GARCH is ruled out. Yet, as stressed by Drost and Nijman, it is possible to provide a QMLE interpretation to the usual ARCH-type MLE procedures. This suggests to construct the moment

condition from a score function using a QMLE argument for  $\theta^r$ . Indeed, the efficiency of  $\hat{\theta}_N^G$  could be improved by using instruments other than  $Z_{mi}$  and  $Z_{Vi}$ . In fact, Meddahi and Renault (1995) examine QMLE estimation of weak GARCH processes and provide an optimal instrument GMM interpretation. We replace  $f_i^r(\theta)$  specified in (3.7) by:

$$f_i^r(\theta) = \partial l_{i+1}^r(\theta) / \partial \theta^r \quad (3.10)$$

where the quasi-log likelihood function  $l_i^r$  for return  $y_{t_i} - y_{t_{i-1}}$ , assuming  $Z_i$  is i.i.d.  $N(0, 1)$  for all  $i$ , is:

$$l_i^r(\theta) = (2\pi)^{-1/2} \exp \left\{ -0.5 \left[ (y_{t_i} - y_{t_{i-1}}) / \sigma_i(\theta) \right]^2 \right\} \\ - \ln |J_i(\theta)| - 0.5 \ln (\sigma_i^2(\theta)) \quad (3.11)$$

with  $J_i(\theta) = \partial \sigma_i(\theta) / \partial \psi_{i+1} | \psi_{i+1}$ . Since the score function in (3.10) involves the derivative with respect to  $\theta^r$  we obtain an exactly identified system of equations when the scores  $\partial l_i^d(\theta) / \partial \theta^d$  and  $\partial l_i^r(\theta) / \partial \theta^r$  are stacked. This results in a GMM estimator which is independent of its weighting matrix and therefore yields a QMLE interpretation to the joint duration/volatility estimator  $\hat{\theta}_N^Q$ . The fact that we need to compute the score of (3.11) may be computationally more intensive compared to the GMM estimator based on the projection equations defined in (3.5) and (3.6). Hence there is some advantage in using  $\hat{\theta}_N^G$  instead of  $\hat{\theta}_N^Q$  in terms of computer time.

Several concluding comments can be made before turning our attention to the two-step procedure. Consider first the asymptotic distribution. Under suitable regularity conditions, which may or may not be easy to verify or establish (see Bollerslev, Engle and Nelson (1994) for further discussion and references), the estimators have standard normal asymptotic distributions (see e.g. Hansen (1982)). It is worth noting that the MLE of Engle and Russell will be inconsistent, even under proper distributional assumptions when  $\partial \psi_i(\theta) / \partial \theta^r \neq 0$  as is the case with  $x_{t_i} \equiv (1, \sigma_i)$ , since it would require joint estimation of transaction durations and volatilities. It was already noted that in principle (3.8) may involve over identifying restrictions under the  $\hat{\theta}_N^G$  specification when  $\dim(\theta) < 2(1 + p + r + s) + k + l + q$ . In such cases one must choose an estimator for  $\hat{\Omega}_N$ , an issue covered in detail by Newey and West (1988),

and Andrews and Monahan (1992), among others. Finally, a standard J-statistic could be applied as model diagnostic.

### 3.2 Two-step estimators

Engle, Hendry and Richard (1984) drew attention to properties of likelihood functions which allow for separate estimation of  $\theta^d$  and  $\theta^r$ . As noted in the previous section we do not really operate in the context of MLE to straightforwardly apply the concepts of weak and strong exogeneity introduced by Engle and al. (1984). Yet, it is only the weakening of the GARCH structure that prevents us from using MLE. Hence, in principle we can focus on the properties of the duration process between transactions and discuss its ML estimation separately. This raises the possibility of introducing two-step procedures.

If we assume that the spacing between transactions is weakly exogenous, in the sense of Engle, Hendry and Richard, with respect to the return process then the parameter vector  $\theta^d$  can be estimated first and separately. This amounts to assuming that the score function  $\partial l_{i+1}^d(\theta) / \partial \theta^d$  is independent of  $\theta^r$  or put differently  $l_{i+1}^d(\theta)$  is a function of  $\theta^d$  only. With this restriction, we can estimate the ACD-GARCH model in two steps: First we estimate the ACD model following the procedure described in Engle and Russell (1995). Under the correctly specified distributional assumptions this estimator will in fact be the most efficient by virtue of MLE. This estimator, denoted  $\hat{\theta}_1^d$  space can be used to construct an expected durations sequence  $\psi_i(\hat{\theta}_1^d)$  using (2.1) in the ACD(1,1) case or a more general recursion formula for ACDX( $p, q$ ) models. The sequence of expected durations  $\psi_i(\hat{\theta}_1^d)$  can be used to estimate the weak GARCH model. As noted in the previous section one may choose a different instrument setup. Either using projection equation arguments as in (3.7) or a score function with a Gaussian QMLE specification as in (3.10). Other possibilities could be considered as well of course. The former will be denoted  $\hat{\theta}_{NG}^r$  while the latter will be denoted  $\hat{\theta}_{NQ}^r$ .

## 4 Empirical illustration: IBM stock prices

In this section we present an application of the ACD-GARCH to intraday observations on IBM prices and intra-trade durations recorded in November 1993. The sample originally consisted of 43328 observations. Prior to estimation the durations between the market closure and the

next day's opening were deleted. We also removed observations corresponding to the market openings and trades occurring simultaneously at the same price, yielding zero returns at zero length durations. The sample reduced to 13038 observations. These data were adjusted for seasonal effects in the duration series and higher order autocorrelation in the returns. The data exhibited intra-day seasonality in the mean durations. Table 4.1 shows the average durations throughout the eight hour trading day.

**Table 4.1: Hourly Averages of Durations**

Time	Mean
9 <sup>00</sup> - 10 <sup>00</sup>	33.0949
10 <sup>00</sup> - 11 <sup>00</sup>	34.3788
11 <sup>00</sup> - 12 <sup>00</sup>	36.2974
12 <sup>00</sup> - 13 <sup>00</sup>	39.3115
13 <sup>00</sup> - 14 <sup>00</sup>	34.9770
14 <sup>00</sup> - 15 <sup>00</sup>	33.4801
15 <sup>00</sup> - 16 <sup>00</sup>	38.5714

These intra-daily components were removed by using the splines to compute the deterministic means conditioned on the time of day and dividing each observation by this value [see Engle and Russell (1995)]. No daily or weekly seasonal components were detected.

The transformed series has mean 0.9999 and standard deviation 0.4469, i.e. features an underdispersion compared to the exponentially distributed series with  $\lambda = 1$ . It is also characterized by a significant degree of autocorrelation as indicated by the value of the portmanteau statistic  $Q(20)=523.5944$ .

No significant predictable patterns were detected in the returns on IBM stock. The original series exhibited some persistence in the conditional mean and was replaced by residuals of an AR(3) process with mean 0.0000, variance 0.0578 and kurtosis 5.4939. The portmanteau statistics computed for the levels and squares of the transformed process are  $Q(20)=14.3650$  and 264.4364 consecutively.

The first estimation method is the two-stage approach under the duration exogeneity assumption. In the first stage we estimated the ACD (1,1) model by Maximum Likelihood based on an exponential density function with intensity parameter  $\lambda = 1$ . The parameter estimates are presented in Table 4.3.

**Table 4.3: Maximum Likelihood Estimates of the ACD (1,1) Model**

parameter	estimate	std.error	p.value
$\omega^d$	0.0028	0.0055	0.3041
$\alpha^d$	0.0128	0.0064	0.0231
$\beta^d$	0.9843	0.0089	0.0000

We can easily observe that the sum of the  $\alpha^d$  and  $\beta^d$  parameters is close to unity revealing the possible presence of a long memory pattern.<sup>9</sup> The residuals of the ACD(1,1) feature a high degree of autocorrelation (Q(20)=169.5615) and the null hypothesis of a white noise process is rejected. To improve the fit, we increase the lag length and estimate an ACD(2,2) model summarized in Table 4.4.

**Table 4.4: Maximum Likelihood Estimates of the ACD(2,2) Model**

parameter	estimate	std.error	p.value
$\omega^d$	0.0009	0.0026	0.3689
$\alpha_1^d$	0.0760	0.0218	0.0002
$\alpha_2^d$	-0.0710	0.0208	0.0003
$\beta_1^d$	1.3805	0.4241	0.0006
$\beta_2^d$	-0.3863	0.4187	0.1781

This specification turns out to be more successful in accommodating the persistence in durations. The portmanteau statistic associated with the residuals is Q(20)=20.0516 and the white noise hypothesis can not be rejected at level 0.05.

The sequence of predicted duration  $\{\psi_i\}$  was used in the second step of the estimation procedure to evaluate the time varying parameters  $\alpha_i$  and  $\beta_i$  appearing in (2.6) through (2.12). We relied on a normal approximation of the density of returns yielding QMLE estimates of the parameters  $\omega$  and  $\alpha$  of the underlying process reported in Table 4.5. The coefficient  $\beta$  was computed from the relation appearing in (2.8). Given the large number of observations we computed the average over all  $\psi$  in the sample. In principle one could obtain standard errors but as this is quite computationally involved we did not calculate them for  $\beta$ . The duration-dependent parameters  $\alpha_i$  and  $\beta_i$  have means 0.1578 and

<sup>9</sup>The issue of long memory in intra-trade durations is discussed in Ghysels and Jasiak (1996) were supporting evidence for the long memory hypothesis is in fact found.

0.0004. They take values within the intervals  $[0., 0.2212]$  and  $[0., 0.0141]$  respectively. The expected intra-trade durations range from 0.8082 and 2.2995 with a mean one by construction. The results in Table 4.5 are quite interesting as they reveal that the persistence in volatility has completely disappeared once the intra-trade duration persistence is driving the volatility process. Indeed the well known IGARCH phenomenon does not appear in the underlying process as the sum of  $\alpha$  and  $\beta$  is very small. Moreover, the random coefficients  $\alpha_i$  and  $\beta_i$  also take values indicating low persistence. The setup of equations (2.6) and (2.7) reveals perhaps why this is the case. Indeed the drift  $\omega_i$  depends directly on  $\psi_{i+1}$ , i.e. the expected duration which absorbs all the persistence. Since intra-trade durations take account of the persistence it is interesting to investigate whether they are exogenous in the sense of Granger causality. The series of rescaled ACD-GARCH and ACD (2,2) residuals were used to test causality as discussed in section 2.4. We investigated the impact of past returns (levels) on the durations and of past volatilities (squares) on the durations as well. We proceeded by computing the crosscorrelation functions at various lags  $k$  reported in the Table 4.6.

**Table 4.5: Quasi - Maximum Likelihood Estimates of the ACD-GARCH (1,1)**

parameter	estimate	std.error	p.value
$\omega$	0.0503	0.0003	0.0000
$\alpha$	0.0729	0.0083	0.0000
$\beta$	0.0838	—	—

The first column in Table 4.6 contains the lags  $k$  corresponding to the number of ticks. The second one shows the values of the crosscorrelation function between the levels of returns and durations ( $r_{de}$ ) while the third contains the crosscorrelations between past squared returns and duration ( $r_{dE}$ ). The null hypothesis of independence cannot be rejected at any lag in the levels. We find however evidence confirming the existence of Granger causality between past volatilities and durations at lags 4 and 10 at the 5% significance level. The joint hypothesis of independence at lags 1 though 10 cannot be rejected at 5% for both ( $r_{de}$ ) and ( $r_{dE}$ ).

**Table 4.6: Cross-correlations in levels and squares of standardized residuals**

lag $k$	levels $r_{de}$	squares $r_{dE}$
1	-0.0025	0.0081
2	0.0003	0.0079
3	0.0087	0.0036
4	0.0064	*0.0209
5	-0.0042	0.0019
6	-0.0123	-0.0048
7	-0.0073	0.0035
8	0.0115	-0.0109
9	-0.0082	0.0019
10	0.0077	*0.0198

Q(10) = 7.9255      Q(10) = 14.8523

Note:  $\chi^2(10) = 18.3$ . “\*” indicates significance at the 5% level.

## 5 Conclusion

In this paper we introduced ACD-GARCH models which is a class of ARCH models for asset return series sampled at time intervals set by the trade arrivals. Our approach combines insights from the temporal aggregation procedure for GARCH models and ACD for intra-trade duration sequences. Since the ACD-GARCH models are intrinsically bivariate we also tested for Granger causal relationships between the volatility and intra-trade durations. Several interesting results emerged from our analysis of IBM data. First of all we find that once the persistence in intra-trade durations are taken into account the volatility dynamics show very little temporal dependence. However, intra-trade durations are not exogenous in a Granger causality sense. Indeed, the results based on tick-by-tick data on IBM stock show that volatility and trading are interdependent.

We also propose a GMM estimation method applicable under general conditions. A sequential procedure can also be adopted when we assume that the spacing of transactions in time is weakly exogenous. Admittedly, the methods we propose are still computationally complex to be implemented on a real time basis in a tick-by-tick trading basis. As we noted throughout the paper it is straightforward to consider events other than the next trading tick. One may for instance consider the durations



between the sale of a total of 10000 shares and predict price volatility that is associated with this event. Another example is that of large block trades, like the sale of 10000 shares in one single trade. Both these events are less frequent and therefore make the use of ACD-GARCH easier in the context of real time trading.

## Appendix A

In principle the causality from durations to volatility and vice versa can be investigated. Although, in the framework of the ACD-GARCH model we restrict our attention to causality from volatility to intra-trade durations we will consider the general case in the proof of Proposition 1. The demonstration follows arguments also used in Cheung and Ng (1996). We will use the following notation:

$$\begin{aligned}\frac{[(y_i - y_{i-1})]^2}{\sigma_i^2} &= \frac{\Delta y_i^2}{\sigma_i^2} = E_i, \\ \frac{t_i - t_{i-1}}{\psi_i} &= \frac{\Delta t_i}{\psi_i} = d_i,\end{aligned}$$

where  $d_i$  has mean 1 and  $\varepsilon_i$  has mean 0 and variance 1. We adopt a general specification of the conditional mean of durations:

$$\psi_i = \varphi_0^d + \sum_{l=1}^{\infty} \varphi_l(\theta^d) \{ \Delta t_{i-l} - \varphi_0^d \},$$

where  $\varphi_l(\theta^d)$  are uniquely defined functions of  $\theta^d$ , the subvector of  $\theta$  which parameterizes the intra-duration process. We assume that the above specification satisfies stationarity conditions. We first show that:

$$\sqrt{N} \frac{\partial \text{cov}_{dE}(k)}{\partial \theta_i} \Big|_{\theta=\theta^0} = O(1), \quad \forall \theta_i \in \theta. \quad (1)$$

Consider  $\theta_j \in \theta^d$ .

$$\frac{\partial \text{cov}_{dE}(k)}{\partial \theta_j^d} \Big|_{\theta=\theta^0} = N^{-1} \sum_i p_i q_{i-k}, \quad (2)$$

where

$$p_i = -\frac{d_i}{\psi_i} \sum_{l=1}^{\infty} \frac{\partial \varphi_l(\theta^d)}{\partial \theta_j^d} (\Delta t_{i-l} - \varphi_0^d) \quad (3)$$

$$q_{i-k} = E_{i-k} - 1 \quad (4)$$

The highest order of  $d_i$  in  $p_i$  is 2. Under the null hypothesis of independence  $p_i$  and  $q_{i-k}$  are independent. Under the null the same result holds for  $\theta_j \in \theta^r$ , the parameter vector of the return process rather than  $\theta^d$  for the duration process, under the following specification of the conditional variance of returns:

$$\sigma_i^2 = \varphi_0^r + \sum_{l=1}^{\infty} \varphi_l(\theta^r, \bar{\delta}_i) \{ \Delta y_{i-l}^2 - \varphi_0^r \},$$

where  $\varphi_l(\theta^r, \bar{\delta}_i)$  are functions of  $\theta^r$  and a fixed value of the duration  $\bar{\delta}$  and satisfy stationarity assumptions. We then have:

$$\left. \frac{\partial cov_{dE}(k)}{\partial \theta_j^r} \right|_{\theta=\theta^0} = N^{-1} \sum_i p_i q_{i-k},$$

where  $p_i = d_i - 1$  and:

$$q_{i-k} = -\frac{E_{i-k}}{\sigma_{i-k}} \sum_{l=1}^{\infty} \frac{\partial \varphi_l(\theta^r, \bar{\delta})}{\partial \theta_j^r} (\Delta y_{i-k-l}^2 - \varphi_0^r)$$

The highest order of  $\varepsilon_{i-k} q_{i-k}$  is 4.

The stationarity and existence of the fourth moment of  $d_i$  and the fourth moment of  $E_i$  imply that:

$$\sqrt{N} \frac{\sum p_i q_{i-k}}{N} \xrightarrow{N \rightarrow \infty} N(0, \sigma^2), \quad (5)$$

[Hannan (1970), theorem 14, p.232]. Hence for any  $\theta_j$  *epsilon*  $\theta$ :

$$\sqrt{N} \left. \frac{\partial cov_{dE}(k)}{\partial \theta_j} \right|_{\theta=\theta^0} = O(1). \quad (6)$$

We now expand  $c\hat{ov}_{dE}(k)$  around the true parameter  $\theta^0$ ,

$$\begin{aligned} \sqrt{N} c\hat{ov}_{dE}(k) &= \sqrt{N} cov_{dE}(k) + \sqrt{N} (\hat{\theta} - \theta_0) \left. \frac{\partial cov_{dE}(k)}{\partial \theta} \right|_{\theta=\theta^0} \\ &\quad + \sqrt{N} (\hat{\theta} - \theta^0)' \left. \frac{\partial^2 cov_{dE}(k)}{\partial \theta \partial \theta'} \right|_{\theta=\theta^*} (\hat{\theta} - \theta^0), \quad (7) \end{aligned}$$

where  $\|\theta^0 - \theta^*\| \leq \|\theta^0 - \hat{\theta}\|$ . Since by assumption  $\hat{\theta}$  is consistent,

$$(\hat{\theta} - \theta^0) = O(1). \quad (8)$$

By this result and if we assume that for all  $\theta$  in an open convex neighborhood of  $(\theta^0)$  and for all  $N$ ,  $\sqrt{N} \partial^2 c\hat{ov}_{dE}(k) / \partial \theta_i \partial \theta_j$  exists and is bounded in probability for  $\theta_i, \theta_j \in \theta$ , where  $\theta = (\theta^d, \theta^r)$  are the parameters of the duration and return processes and  $\theta^0 = (\theta^{d^0}, \theta^{r^0})$  is the true parameter vector, then we have that:

$$\sqrt{N}\hat{c}v_{dE}(k) = \sqrt{N}cov_{dE}(k) + \sqrt{N}(\hat{\theta} - \theta^0) \left. \frac{\partial cov_{dE}(k)}{\partial \theta} \right|_{\theta=\theta^0} + o(1). \quad (9)$$

The second term is  $o(1)$  by (1) and (8). Hence

$$\sqrt{N}\hat{c}v_{dE}(k) = \sqrt{N}cov_{dE}(k) + o(1). \quad (10)$$

For  $k \neq k'$  we have

$$\begin{aligned} N\hat{c}v_{dE}(k) \hat{c}v_{dE}(k') &= (\sqrt{N}cov_{dE}(k) + o(1)) (\sqrt{N}cov_{dE}(k') + o(1)) \\ &= N cov_{dE}(k) cov_{dE}(k') + o(1) \end{aligned} \quad (11)$$

By a similar argument

$$\hat{c}v_{dd}(\theta) = cov_{dd}(\theta) + o(1).$$

and

$$\hat{c}v_{EE}(\theta) = cov_{EE}(\theta) + o(1).$$

Applying the Slutsky and Cramer-Rao theorems imply that  $\hat{c}_{dE}(k)$  and  $c_{dE}(k)$  have the same asymptotic distribution and

$$\sqrt{N}(\hat{c}_{dE}(k_1), \dots, \hat{c}_{dE}(k_m)) \rightarrow N(0, I_m).$$

## References

- [1] Andrews, D.W.K. and J.C. Monahan, "An Improved Heteroskedasticity and Autocorrelation Consistent Covariance Matrix Estimator," *Econometrica* 60, 953-966.
- [2] Bera, A.K. and M.L. Higgins (1995), "On ARCH models : properties, estimation and testing", In L. Exley, D.A.R. George, C.J. Roberts and S. Sawyer (eds), *Surveys in Econometrics*. (Basil Blackwell : Oxford).
- [3] Bollerslev, T., Y.C. Chou and K. Kroner (1992), "ARCH Modelling in Finance : A Selective Review of the Theory and Empirical Evidence", *Journal of Econometrics* 52, 201-224.
- [4] Bollerslev, T., R. Engle and D. Nelson (1994), "ARCH Models", in R.F. Engle and D. McFadden (eds.), *Handbook of Econometrics*, Volume IV, (North-Holland, Amsterdam).
- [5] Bollerslev, T. and E. Ghysels (1996), "On Periodic Autoregression Conditional Heteroskedasticity", *Journal of Business and Economic Statistics*.
- [6] Cheung, Y.W. and L.K. Ng, (1996), "A causality-in-variance test and its applications to financial market prices", *Journal of Econometrics* 72, 33-48.
- [7] Diebold, F.X. and J.A. Lopez (1995), "Modelling Volatility Dynamics", in K. Hoover (ed), *Macroeconomics : Developments, Tensions and Prospects*.
- [8] Drost, F.C. and T.E. Nijman (1993), "Temporal Aggregation of GARCH Processes", *Econometrica* 61, 909-927.
- [9] Drost, F.C. and B.J.M. Werker (1996), "Closing the GARCH Gap: Continuous Time GARCH Modelling", *Journal of Economics*.
- [10] Engle, R.F. (1982), "Autoregressive Conditional Heteroskedasticity with Estimates of the Variance of United Kingdom Inflation", *Econometrica* 50, 987-1007.
- [11] Engle, R.F. and J.R. Russell (1995), "Forecasting Transaction Rates: The Autoregressive Conditional Duration Model", Discussion paper U.C.S.D.
- [12] Engle, R.F., D. Hendry and J.F. Richard (1983), "Exogeneity", *Econometrica*.
- [13] Gallant, A.R. and G. Tauchen (1989), "Semiparametric Estimation of Conditionally Constrained Heterogeneous Processes: Asset Pricing applications", *Econometrica* 57, 1091-1120.
- [14] Ghysels, E., C. Gouriéroux and J. Jasiak (1996a), "Market Time and Asset Price Movements: Theory and Estimation", Discussion paper CIRANO and C.R.D.E., Université de Montréal.
- [15] Ghysels, E., C. Gouriéroux and J. Jasiak (1996b), "Trading Patterns, Time Deformation and Stochastic Volatility in Foreign Exchange Markets", Paper presented at the HFDF Conference, Zürich.

- [16] Ghysels, E. and J. Jasiak (1994), “Stochastic Volatility and Time Deformation: An Application of Trading Volume and Leverage Effects”, Paper presented at the Western Finance Association Meeting, Santa Fe.
- [17] Ghysels, E. and J. Jasiak (1996), “Long term dependence in Trading”, Discussion Paper.
- [18] Hannan, E.J. (1970), “Multiple Time Series”, (Wiley, New York, N.Y.).
- [19] Hansen, L.P. (1982), “Large Sample Properties of Generalized Method of Moments Estimators”, *Econometrica* 50, 1029-1054.
- [20] Haugh, L.D. (1976), “Checking the Independence of Two Covariance Stationary Time Series: A Univariate Residual Cross-Correlation Approach”, *Journal of the American Statistical Association*, 379-385.
- [21] Lancaster, T. (1990), “The Econometric Analysis of Transition Data”, (Cambridge University Press Cambridge).
- [22] Meddahi, N. and E. Renault (1995), “Linear Statistical Inference for ARCH-type Processes”, Discussion Paper, GREMAQ.
- [23] Newey, W.K. and K. West (1987), “A Simple, Positive Semi-Definite, Heteroskedasticity and Autocorrelation Consistent Covariance Matrix”, *Econometrica* 55, 703-708.
- [24] Nijman, T., and F. Palm (1990), “Parameter identification in ARIMA processes in the presence of regular but incomplete sampling”, *Journal of Time Series Analysis*, 11, 239-248.
- [25] Nijman, T. and F. Palm (1990), “Predictive Accuracy Gain from Disaggregate Sampling in ARIMA Models”, *Journal of Business and Economic Statistics*, 8, 405-412.
- [26] Palm, F.C. (1996), “GARCH Models of Volatility”, in G.S. Maddala and C.R. Rao (eds.), *Handbook of Statistics*, Volume 14, (North-Holland, Amsterdam).
- [27] Palm, F.C. and Nijman, Th. (1984), “Missing Observations in the Dynamic Regression Model”, *Econometrica* 52, 1415-1435.

## Liste des publications au CIRANO

### Cahiers CIRANO / *CIRANO Papers* (ISSN 1198-8169)

- 96c-1 Peut-on créer des emplois en réglementant le temps de travail ? / par Robert Lacroix
- 95c-2 Anomalies de marché et sélection des titres au Canada / par Richard Guay, Jean-François L'Her et Jean-Marc Suret
- 95c-1 La réglementation incitative / par Marcel Boyer
- 94c-3 L'importance relative des gouvernements : causes, conséquences et organisations alternative / par Claude Montmarquette
- 94c-2 Commercial Bankruptcy and Financial Reorganization in Canada / par Jocelyn Martel
- 94c-1 Faire ou faire faire : La perspective de l'économie des organisations / par Michel Patry

### Série Scientifique / *Scientific Series* (ISSN 1198-8177)

- 97s-08 Living on a Noisy and Dusty Street: Implications for Environmental Evaluation / Tagreed Boules, Robert Gagné et Paul Lanoie
- 97s-07 The Location of Comparative Advantages on the Basis of Fundamentals Only / Thijs ten Raai et Pierre Mohnen
- 97s-06 GARCH for Irregularly Spaced Financial Data: The ACD-GARCH Model / Eric Ghysels, Joanna Jasiak
- 97s-05 Can Capital Markets Create Incentives for Pollution Control? / Paul Lanoie, Benoît Laplante et Maité Roy
- 97s-04 La régie des services informatiques : Le rôle de la mesure et des compétences dans les décisions d'impartition / Benoit A. Aubert, Suzanne Rivard et Michel Patry
- 97s-03 Competition and Access in Telecoms: ECPR, Global Price Caps, and Auctions / Marcel Boyer
- 97s-02 L'économie des organisations : Mythes et réalités / Marcel Boyer
- 97s-01 Fixed Wages, Piece Rates, and Intertemporal Productivity: A Study of Tree Planters in British Columbia / Harry J. Paarsch et Bruce Shearer
- 96s-34 Structural Change and Asset Pricing in Emerging Markets / René Garcia et Eric Ghysels
- 96s-33 Monitoring of Pollution Regulation: Do Local Conditions Matter? / Catherine Dion, Paul Lanoie et Benoit Laplante
- 96s-32 Technological Learning and Organizational Context: Fit and Performance in SMEs / Élisabeth Lefebvre, Louis A. Lefebvre et Lise Préfontaine
- 96s-31 Piece Rates, Fixed Wages, and Incentive Effects: Statistical Evidence from Payroll Records / Harry J. Paarsch et Bruce Shearer
- 96s-30 R&D Spillover and Location Choice under Cournot Rivalry / Ngo Van Long et Antoine Soubeyran
- 96s-29 Efficient Income Redistribution in a Growing Economy / Gerhard Sorger

---

· Vous pouvez consulter la liste complète des publications du CIRANO et les publications elles-mêmes sur notre site World Wide Web à l'adresse suivante :

<http://www.cirano.umontreal.ca/publication/page1.html>