
STROBE-X: X-ray Timing & Spectroscopy on Dynamical Timescales from Microseconds to Years



Colleen A. Wilson-Hodge (NASA/MSFC), Paul Ray (NRL), Keith Gendreau (NASA/GSFC), Deepto Chakrabarty (MIT), Marco Feroci (INAF-IASF/INFN), Tom Maccarone (TTU), Zaven Arzoumanian (CRESST/GSFC), Ron Remillard (MIT), Kent Wood (Praxis/NRL), Christopher Griffith (NRC/NRL), Peter Jenke (UAH)

Spectroscopic Time-Resolving Observatory for Broadband Energy X-rays (STROBE-X)



STROBE-X Instruments

X-ray Concentrator Array

- Concentrator optics and detectors, scaled up from NICER
- Energy resolution: 85-175 eV FWHM
- Effective area @ 1.5 keV: 3.4 m²





Large Area Detector


- SDDs and micropore collimators from ESA's LOFT M3 & M4
- Energy resolution: 200-240 eV FWHM
- Effective Area @ 10 keV 7.6 m²

Wide Field Monitor

- LOFT SDDs and mask
- Energy resolution: 300 eV FWHM
- Instantaneous FoV: 1/3 of sky


STROBE-X Black Hole Science

- Probes stationary spacetimes near black holes to explore effects of strong gravity
- Complementary to gravitational wave measurements that probe dynamic spacetimes




STROBE-X Black Hole Science


- Measures BH spin using QPOs
- Exploits X-ray
 reverberation mapping
 of BH accretion flows
 across all mass scales
 from stellar mass to
 supermassive




STROBE-X Science



- Fully determines the ultradense matter equation of state using > 20 pulsars
- Explores cosmic chemical evolution for ~100 high redshift clusters
- Continuously surveys the dynamic X-ray sky with large duty cycle and high spectral and time resolution



Summary



- STROBE-X is a probe class (<\$1B) observatory designed for X-ray timing and spectroscopy in the 0.2-30 keV band
- STROBE-X has huge collecting area and good spectral resolution
- It is optimized for the study of matter in the most extreme conditions in the universe