

Elsevier Editorial System(tm) for Sensors & Actuators: B. Chemical
Manuscript Draft

Manuscript Number: SNB-D-15-00099R1

Title: Magnet bioreporter device for ecological toxicity assessment on heavy metal contamination of coal cinder sites

Article Type: Research Paper

Keywords: whole-cell bioreporter; magnetic nanoparticles (MNPs); magnet bioreporter device; heavy metal; toxicity

Corresponding Author: Dr. Dayi Zhang, Ph.D.

Corresponding Author's Institution: Lancaster University

First Author:

Order of Authors: Jianli Jia; Hanbing Li; Shuang Zong; Bo Jiang; Guanghe Li; Odafe Ejenavi; Jingrong Zhu

Abstract: A novel magnet bioreporter device was developed in this research for soil toxicity assessment, via magnetic nanoparticles functionalized whole-cell bioreporters. The whole-cell bioreporter ADPWH_recA kept response capability to DNA damage after magnetic nanoparticles (MNPs) functionalization, and could be harvested from soil samples by permanent magnet to reduce the soil particle disturbance. Compared to conventional treatments applying bioreporter directly in soil-water mixture (SW-M treatment) or supernatant (SW-S treatment), MNPs functionalized bioreporter via the magnet device (MFB) treatment achieved high sensitivity to evaluate the toxicity and bioavailability of chromium contamination in soils from 10 mg/kg to 5,000 mg/kg soil dry weight. The MNPs functionalized bioreporter also achieved high reproducibility with pH value from 5.0 to 9.0, salinity from 0% to 5% and temperature from 20°C to 40°C. A case study was carried out on the ecological toxicity assessment of heavy metal contamination at the coal cinder site via the magnet bioreporter device. The heavy metal toxicity declined with the increasing distance to the coal cinder point, and a significant accumulation of heavy metal toxicity was observed along the vertical distribution. No direct link was found between the pollution load index (PLI) and heavy metal toxicity, and the results suggested the bioreporter test monitored the toxicity of heavy metals in soils and was an important approach for ecological risk assessment. Magnet bioreporter device also offered the high throughput biological measurement and was feasible for in situ monitoring.



To:

Editor of Sensor and Actuators B: Chemical

Dr Dayi Zhang

Lecturer

Lancaster Environment Centre

Lancaster University

Lancaster, LA1 4YQ

30 Dec 2014

Telephone: +44 (0) 1524 510288**Fax:** +44 (0)1524 510082**Email:** d.zhang@lancaster.ac.uk

Dear Editor

I would like to submit this manuscript, entitled "*Magnet bioreporter device for ecological toxicity assessment on heavy metal contamination of coal cinder sites*", for the consideration in *Sensor and Actuators B: Chemical*. The heavy metals leaching from coal cinder have severely contaminated soils in China and its toxicity was hard to be evaluated on the field by traditional chemical analysis. This paper published a novel concept of magnetic bioreporter device to improve the sensitivity and reliability of whole-cell bioreporter in soil measurement. Functionalized with the magnetic nanoparticles, the whole-cell bioreporter could sense the toxicity of soils in situ and subsequently isolated by magnetic field for bioluminescent signal detection. The magnetic bioreporter device was further applied for the toxicity assessment on a coal cinder sites in China. The results illustrated the horizontal/vertical toxicity distribution and its correlation with pollution load index. This novel magnetic device benefitted the biosensor application on the field, particularly contributing to the soil measurement, with high feasibility in practical environmental monitoring.

No conflict of interest exists in the submission of this manuscript, and the manuscript has approved by all authors for publication. The authors would like to declare that the work described is original research that has not been published previously, and is not under consideration for publication elsewhere, in whole or in part.

Thanks for your consideration. If you have any questions, please feel free to contact with me.

Yours sincerely

Dr Dayi Zhang

Reviewer #1: The paper describes magnetic bioreporter device for ecological toxicity assessment on heavy metal contamination of coal cinder sites. However, before it could be publish, some points need to be addressed as follows:

3. Results and discussions

No discussion on optimization parameter of magnetic whole-cell bioreceptor based, such as pH, temperature, medium etc. Please give comment on this issue.

Answer:

Thanks for the comments and the author has carried out more research work on the impacts of pH and temperature. All these data are included in the reproducibility part.

3.1. Higher sensitivity ...

Please give a mechanism of whole-cell bioreceptor toward heavy metals. In this subtitle, it could be better if the analytical characteristic is summarized in Table, in term of linear range, sensitivity, LOD, reproducibility etc. What was the life time of this magnetic bioreporter cells?

Answer:

The mechanisms of whole-cell bioreporter's response to heavy metal are the SOS response activated by the DNA damage. The author has added more description and discussion on this issue. The additional table (Table 1) is added to summarize the main features of the magnetic bioreporter.

3.2. Heavy metal ...

The heavy metal profile in Table 1 needs to added with information regarding the instrumentation used for the determination of heavy metal listed.

Answer:

The analytical instrument for Hg is DMA-80 Hg analyzer (Milestone S.r.L., Italy) and the other elements were analysed by ICP-MS (X series II, Thermo Fischer Scientific, USA). The information has been listed with detailed detection procedure in the

materials and methods section. The author has added in the note of the table for better description in accordance with the comments.

Other comment

It could be better if the performance of the proposed magnetic bioreporter cells could be compared with other toxicity assessment on heavy metals.

Answer:

Thanks for this good comments and the performance of ADPWH_recA has been tested and compared to other toxicity assessment in our previous research (Y.Z. Song, G.H. Li, S.F. Thornton, I.P. Thompson, S.A. Banwart, D.N. Lerner, et al., Optimization of Bacterial Whole Cell Bioreporters for Toxicity Assay of Environmental Samples, *Environ Sci Technol*, 43(2009) 7931-7938). Sharing the same mechanisms of SOS response, the author believes the magnetic bioreporter has the same performance comparing to other toxicity assay. The authors have added some more discussions in the revised manuscript.

Reviewer #2: This manuscript presented a magnet bioreporter device for soil toxicity assessment by magnetic nanoparticles functionalized whole-cell bioreporters. The whole-cell bioreporters with magnetic nanoparticles functionalization could be separated from soil samples to reduce the soil particle interference. Although this work is innovate, major revision should be performed before this manuscript can be accepted for publication.

1. Why did the author choose the whole-cell *Acinetobacter* ADP1_recA reporter for toxicity assessment? And why did the author choose chromium as the target analyte? Did the reporter represent any specificity to chromium? Did other heavy metals have genotoxic effects on the reporter?

Answer:

Various whole-cell bioreporter can be applied for toxicity test, like *E. coli*, but they are not suitable for soil detection mainly due to two reasons as induction condition and response stability. First of all, the induction condition for *E. coli* is 37 degree,

which is much higher than natural temperature in soils. *Acinetobacter* can survive and response to carcinogens from 15 degree to 37 degree, allowing the toxicity assessment at the real soil temperature. Secondly, the reporter gene in *E. coli* is on the plasmid, requiring antibiotics to maintain the plasmid copies and responsive sensitivity. The reporter gene in *Acinetobacter* is on the chromosome, which is single copy and stable with no need of antibiotic cultivation. From these reasons, the author suggests that *Acinetobacter* is suitable candidate as bioreporter strain for environmental monitoring and assessment.

From the chemical analysis of heavy metal profiles in the contaminated soils, chromium is the major contaminants and the highest pollutant level compared to other heavy metals. For this reason, chromium is chosen as the targeting analyte to investigate the biological response and the feasibility that the magnetic biosensor device can be applied for heavy metal contamination evaluation. The description was added in the materials and methods section.

From our previous work, ADPWH_recA responds to various carcinogens, including mitomycin C, UV light, ethidium bromide and H₂O₂ (Y.Z. Song, G.H. Li, S.F. Thornton, I.P. Thompson, S.A. Banwart, D.N. Lerner, et al., Optimization of Bacterial Whole Cell Bioreporters for Toxicity Assay of Environmental Samples, *Environ Sci Technol*, 43(2009) 7931-7938). From the SOS response mechanisms, it is not a specific bioreporter to sense targeting chemical molecules, but responding to the generic SOS activation of *recA* gene. It is obvious that other heavy metals might also cause the bioluminescent response, which is undergoing. Current work shows that ADPWH_recA only represents positive response to Zn and Be. Further work is still undergoing and the outcome might be discussed in future publication. The author thanks reviewers' comments and added more discussions in the revised manuscript addressing the details shown above.

2. When detecting real soil samples by the whole-cell bioreporter, no extra preprocessing procedures were introduced. Is it possible that some substances in samples affect the bioreporter during the incubation and further interfere the toxicity evaluation? How did you solve the interference problem?

Answer:

From our previous work on toxicity assessment on soils (Y.Z. Song, G.H. Li, S.F. Thornton, I.P. Thompson, S.A. Banwart, D.N. Lerner, et al., Optimization of Bacterial Whole Cell Bioreporters for Toxicity Assay of Environmental Samples, *Environ Sci Technol*, 43(2009) 7931-7938; B. Jiang, D. Zhu, Y. Song, D. Zhang, Z. Liu, X. Zhang, et al., Use of a whole-cell bioreporter, *Acinetobacter baylyi*, to estimate the genotoxicity and bioavailability of chromium(VI)-contaminated soils, *Biotechnol Lett*, 37(2015) 343-348), some substances (organic matters, other heavy metals, etc.) in soils significantly affected the bioavailability of heavy metals and they are part of the soil meso-environment determining the ecological risks of heavy metal contamination. As for other environmental factors, like pH, temperature and salinity, are further discussed with new experiment in the revised version. From the data, all these factors will not influence the bioreporter response within the soil properties. The author thinks that the interference will not affect the toxicity assessment by whole-cell bioreporter.

3. How did the reporter attach to the magnetic nanoparticles? Is the magnetic nanoparticles coupled with any chemical groups?

Answer:

The magnetic nanoparticles were attached on the membrane of whole-cell bioreporter by electrostatic attraction between Fe-OO- and amino-groups. More discussions are made in the revised version.

4. In figure 4, the ordinate is the "bioluminescence response ratio", and in figure 5, the ordinate is the "relative bioluminescence response ratio". In the text, the two words were also used alternatively. Did these two words have different meanings or not?

Answer:

Thanks for the comments and the author misrepresented the phrase. It should be "relative bioluminescence response ratio" (as defined in the Materials section) throughout the manuscript. The author has corrected all the mistakes.

5. Why the MFB present positive bioluminescent response to the low chromium concentration in soil and negative response to high chromium concentration? Why are the simulation curves parabolic shaped? And how could the toxicity of the contaminated soils be predicted by the curves? According to the curve, even under the treatment to two samples in different chromium concentration, the reporter may present same bioluminescent response. The author should explain it in detail.

Answer:

Chromium has both cytotoxicity and genotoxicity on living microorganisms. At lower concentration, the cytotoxicity is not obvious and would not affect bacterial growth, and the positive relationship between chromium amount and bioluminescent response was therefore observed. At higher concentration (>100 mg/kg dry soil weight in this study), the cytotoxicity of chromium became dominant and significantly affected cell growth and activity. The bacterial growth curve was significantly suppressed by the high chromium concentration. The author has added the growth curve against different concentration of chromium (Fig. S1). The results showed that no growth change was observed when the chromium concentration was less than 100 mg/kg and this is the range for positive relationship between bioluminescent response and chromium contamination. At higher level, the cell growth and activity was inhibited, causing the dramatic loss of bioluminescent signal. The synergetic effects of cytotoxicity and genotoxicity were then the main reason causing the parabolic curve.

To determine the real toxicity, both relative bioluminescent response ratio and growth curve were considered. If the growth was not affected, the bioluminescence data belonged to the positive relationship range, and it was on the opposite when the growth was inhibited. The author has added more detailed discussion in the revised version.

6. What is the detection limit and linear range of the magnet bioreporter device?

Answer:

From our previous publication (Jiang, B., Zhu, D., Song, Y., Zhang, D., Liu, Z., Zhang, X., Huang, W.E. and Li, G.*, (2015) Use of a whole-cell bioreporter, *Acinetobacter baylyi*, to estimate the genotoxicity and bioavailability of chromium(VI)-contaminated soils, *Biotechnology Letters*, 37(2), 343-348), the limit of detection is 130 mg/kg soil and the linear range is 130-5200 mg/kg (ultrasonic treatment). Without pretreatment, the response of ADPWH_recA to chromium contaminated soils is too weak to be quantified, similar as our research results. The test of this magnet bioreporter work demonstrated a higher sensitivity that the limit of detection is 1 mg/kg soil and the linear range is 1-100 mg/kg. Compared with the bioreporter growth data, the detection range of the magnetic device can achieve 1-5000 mg/kg. The author has added the discussion and summarized the limited of detection and other key parameters in Table 1.

7. In figure 4, why did the response ratio not fit well with the model prediction of 10% chromium bioavailability when chromium concentration was below 100 mg/kg soil dry weight?

Answer:

The bioavailability of chromium changes with the contamination level. With lower chromium in the soils, different bioavailability will be observed due to different adsorption by soil particles. The author thought it is the main reason causing the unexpected bioluminescent response ratio when chromium concentration was <100 mg/kg soil. Some further discussion was addressed on this issue in the revised version.

8. In figure 5, the authors concluded that "the toxicity of heavy metals declined with the increasing distance to the coal cinder point" because "Except for 0 m point, the bioluminescence response ratio dropped from 1.47 (10 m) to 1.10 (150 m) in the surface soil, 1.34 (10 m) to 1.16 (150 m) in the middle soil, and 1.26 (0 m) to 0.58 (150 m) in the bottom soil." However, did lower bioluminescence response ratio really mean lower toxicity? Please combine the results obtained from figure 4 to explain the correlations between toxicity and bioluminescence response ratio.

Answer:

Thanks for the comments. As discussed for the bioreporter growth curve, whether the growth is inhibited can be the indicator to determine the toxicity range of soil samples. From Fig. S4, all the bioreporter growth curve behaved similar without any inhibition, and the results indicated that no strong cytotoxicity was found in the heavy metal contaminated soils. Located in the linear relationship range, higher relative bioluminescent response ratio represented higher ecological toxicity. The author has revised the discussion in the new manuscript.

9. What about the reproducibility of magnetic nanoparticles functionalized whole-cell bioreporters?

Answer:

Thanks for the comments and it is the authors' mistake missing the description for the reproducibility test. In our experiment, three biological replicates were carried out for each test, and five individual tests were repeated for the same sample sets to evaluate the reproducibility. The data and error bar on each graph were the mean and standard error of five individual tests. The author has corrected the sentence in the materials and methods section.

10. Since the soil ecological risks were not associated with neither the load of PC1 nor PC2, the circles in figure 6 is needless and should be deleted.

Answer:

Thanks for the comments and the author has moved Figure 6 to Supplementary Material as supporting information for the manuscript.

Reviewer #3: This paper reported a magnet biosensor for soil toxicity monitoring. It was stated that this biosensor "offered the high throughput biological measurement and was feasible for in situ monitoring".

This reviewer would like to suggest some revisions below.

1. Replicates and errors. No related data were given in this draft. The authors

should perform some efforts focusing on replicates and sensing errors.

Answer:

For all the tests, three biological replicates were carried out. All the data in the graph were the average of replicates with error bars. The information was in Section 2.2.

2. What was the repeatability of this biosensor. What was the testing error between biosensors of different lots?

Answer:

Thanks for the comments and it is the authors' mistake missing the description for the reproducibility test. Actually, three biological replicates were carried out for each sample, and five individual tests were repeated for the same sample sets to evaluate the reproducibility. The data and error bar on each graph were the mean and standard error of five individual tests. The author has corrected the sentence in the materials and methods section.

3. Did the authors considered the influence from other competing ions, pollutions and moisture on your biosensor?

Answer:

The author has added the influence of salinity on the bioreporter's performance and the results indicated <4% salinity did not affect its response ratio. During the bioluminescence measurement, the soils were saturated and the original moisture of soils will not affect the assessment results.

From the mechanisms of SOS response and ADPWH_recA bioreporter, it responded to all the DNA damage. Thus, any other carcinogen will also cause the positive response of bioreporter. However, the chromium was just chosen as the target analyte since it is the main heavy metal contamination at the sites. The MNPs functionalized bioreporter actually evaluate the synergetic toxicity of all the contaminants in the environmental samples. The author has added more discussion in the revised manuscript.

4. The authors should try more practical samples to testify the accuracy and practicability of this biosensor.

Answer:

Thanks for the comment. The author wanted to collect more samples in different direction and at different distance and depth, to further test the reproducibility and reliability of the magnetic biosensor by more sample testing. However, on the north and west to the cinder sites, nearly all the surface is the hard surface (within the main chemical production area) and no appropriate soils can be collected. The east and south soils were near to the road and the soils might be contaminated by the coal transportation activities. Furthermore, the local company did not allow us to take samples below 50 cm to protect the underground pipeline. From local investigation, we have tried out best to choose the most appropriate sampling line and sampled the maximal samples for the work, as illustrated in the figure.

More tests were carried out according to reviewers' suggestions, and the impacts of pH, salinity and temperature were comprehensively discussed in the revised version.

To further testify the accuracy and practicability, the authors have more samples collected from different cinder sites around China (about 70 samples from 5 companies), and further work will focus on the relationship between heavy metal profiles, biosensor results and microbial community, to deeper understand how the magnetic biosensor device can be applied to assess the toxicity and how the toxicity has affected indigenous microorganisms. In this paper, our main goal is to prove the feasibility of magnetic bioreporter device at one site, and it will be used as a technical tool for more samples to illustrate the ecological impacts of coal cinder in the next step.

18 **Abstract**

19 A novel magnet bioreporter device was developed in this research for soil toxicity
20 assessment, via magnetic nanoparticles functionalized whole-cell bioreporters. The
21 whole-cell bioreporter ADPWH_recA kept response capability to DNA damage after
22 magnetic nanoparticles (MNPs) functionalization, and could be harvested from soil
23 samples by permanent magnet to reduce the soil particle disturbance. Compared to
24 conventional treatments applying bioreporter directly in soil-water mixture (SW-M
25 treatment) or supernatant (SW-S treatment), MNPs functionalized bioreporter via the
26 magnet device (MFB) treatment achieved high sensitivity to evaluate the toxicity and
27 bioavailability of chromium contamination in soils from 10 mg/kg to 5,000 mg/kg soil
28 dry weight. The MNPs functionalized bioreporter also achieved high reproducibility
29 with pH value from 5.0 to 9.0, salinity from 0% to 5% and temperature from 20°C to
30 40°C. A case study was carried out on the ecological toxicity assessment of heavy
31 metal contamination at the coal cinder site via the magnet bioreporter device. The
32 heavy metal toxicity declined with the increasing distance to the coal cinder point, and
33 a significant accumulation of heavy metal toxicity was observed along the vertical
34 distribution. No direct link was found between the pollution load index (*PLI*) and
35 heavy metal toxicity, and the results suggested the bioreporter test monitored the
36 toxicity of heavy metals in soils and was an important approach for ecological risk
37 assessment. Magnet bioreporter device also offered the high throughput biological
38 measurement and was feasible for *in situ* monitoring.

39 **Key words:** whole-cell bioreporter, magnetic nanoparticles (MNPs), magnet
40 bioreporter device, heavy metal, toxicity

41

42

43 **1. Introduction**

44 Heavy metals are the key anthropogenic environmental contaminants, mainly caused
45 by industrial activities [1, 2]. All around the world and particularly in China,
46 numerous heavy metal contaminated sites are found due to the improper disposal of
47 various chemical wastes [3], including coal cinders [4], and the key pollutants include
48 chromium, mercury, arsenic, lead, cadmium, manganese, cobalt, copper, nickel and
49 zinc. They have high mobility through the leachate and further contaminate the
50 biospheric soils [5, 6], with respective carcinogenic, teratogenic and mutagenic effects
51 [7]. The high level of heavy metal in soils threatens the ecological system [8], poses
52 potential risks to human health [9] and draws attention on early warning for potential
53 cancer induction [10]. Due to the complex composition and synergetic effects in soils,
54 traditional chemical and physical analysis only provides the amount information [3],
55 but the toxicity and bioavailability of heavy metal contamination from coal cinder are
56 hard to be evaluated.

57 Recently, whole-cell bioreporter has become initiative and legislative tool for
58 environmental monitoring, with capability to sense the bioavailability and toxicity of
59 contaminated water and soil samples [11]. With genetically engineered bacteria, yeast,
60 fungi, or animal cells, the biological signals of whole-cell bioreporter are initiated by
61 phenotypic color (*lacZ*), fluorescent (*gfp/yfp*) or bioluminescent (*luc/lux*) genes [12-
62 14]. It offers highly sensitive, rapidly analytic, easy operation and cost-effective
63 feasibility for *in situ* pollutants assessment [15]. Some whole-cell bioreporter
64 specifically senses the heavy metal molecules [16, 17] or their
65 cytotoxicity/genotoxicity [18].

66 Though the application of whole-cell bioreporter in water sample is successful, it
67 suffers from the heterogeneous features of soils [19]. Exposed to whole-cell
68 bioreporter, the soil particles will absorb the bioluminescent signal (*lux* or *luc*) or give
69 strong fluorescent interference (*gfp*). Some recent work has assessed the
70 bioavailability and toxicity of copper [20], cobalt and nickel [21] via direct exposing
71 the whole-cell bioreporter to the soils [22], but the biological sensitivity and
72 specificity are significantly reduced. Some pre-treatments, like water extraction or
73 ultrasonication, are therefore applied to transfer contaminants into aqueous phase for
74 biological analysis [23]. Particularly for heavy metal, the aqueous extraction has been
75 used for whole-cell bioreporters to sense the bioavailability of chromium [24],

76 mercury [16], lead and cadmium [25, 26] in soils. Nevertheless, the main drawback is
77 the neglect of the real occurrence of pollutants in the porous soil [17]. Technically, a
78 new type of bioreporter device is required to sense the soil contaminants *in situ* and
79 effectively separate the living reporter cells from the soil particles for biological
80 signal detection. Magnetic nanoparticles (MNPs) functionalization offers the
81 feasibility of magnetic remote control and is biocompatible for whole cell bioreporter
82 [27]. Its equipping and portability for *in situ* monitoring is still under development
83 and required further research.

84 In this work, a novel magnet bioreporter device was developed and optimized for
85 effective monitoring and assessment of coal cinder contaminated soils. With whole-
86 cell *Acinetobacter* ADP1_recA reporter [28], the magnet device effectively reduced
87 the impacts of soil particles and improved the sensitivity and reproducibility,
88 comparing to the direct exposure of bioreporters to the soils. The MNPs
89 functionalized bioreporter was able to evaluate the ecological toxicity of heavy metal
90 contamination, via the high throughput and easy operation magnet device. This work
91 showed the feasibility and potential of *in situ* environmental risk assessment via
92 whole-cell bioreporter for the coal contaminated sites.

93 **2. Material and methods**

94 *2.1 Bioreporter strain and incubation*

95 In this research, the *Acinetobacter baylyi* ADPWH_recA whole-cell bioreporter was
96 introduced for environmental ecological toxicity evaluation [28, 29]. Compared to
97 other plasmid based or *Escherichia coli* hosted toxicity bioreporter, the reporter gene
98 was located on the chromosomal with high stability and *Acinetobacter* was soil
99 bacterium to tolerate the ambient soil environment and achieve high sensitivity. After
100 cultivation in Luria-Bertani (LB) medium overnight at 30°C, the 10.0 mL
101 ADPWH_recA cells were harvested by 3,000 rpm centrifugation for 10 minutes. The
102 bioreporter pellets were further washed by deionized water and resuspended in 10 mL
103 deionized water for magnetic nanoparticles functionalization or 10 mL fresh MMS
104 medium for toxicity measurement. The 1.0 litre MMS medium contained 1.0 g
105 (NH₄)SO₄, 2.5 g KH₂PO₄, 0.1 g MgSO₄·7H₂O, 0.005 g FeSO₄·7H₂O, 0.25 g
106 nitrilotriacetic acid (NTA), 0.55 g NaOH, 3.24 g sodium succinate (20 mM) and 1 mL
107 Bauchop and Elsdon solution.

108 *2.2 Direct toxicity measurement on soil samples*

109 The direct toxicity measurement of soil samples were conducted for the supernatant of
110 soil/water mixtures (SW-S) and the bulk soil/water mixtures (SW-M) respectively.
111 For SW-S treatment, the 100 mg soil sample was suspended in 1 mL deionized water
112 and homologized by 150 rpm shaking for 10 min. The 20 μ L supernatant and 180 μ L
113 bioreporter suspensions were added into each well of a black clear-bottom 96-well
114 microplate. For SW-M treatment, the 1.0 mL bioreporter suspensions were added into
115 100 mg soil sample, and the mixture was directly transferred into the microplate. The
116 incubation and induction was conducted at 30°C for 4 hours, in the Spectra M5 Plate
117 Reader (Molecular Devices, USA). Three biological replicates were carried out for
118 each sample and the bioluminescent signal was measured every 10 minutes with 5
119 seconds shaking before each reading.

120 *2.3 Magnetic nanoparticles synthesis and functionalization*

121 All the chemicals in this study were analytical grade and purchased from Sigma
122 Aldrich (UK) without specific statement. Magnetic nanoparticles synthesis followed
123 chemical deposition method [27]. The 12.5 mL NaOH (1.5 M) was added dropwisely
124 into the mixture of 1.0 mL FeCl₃ (2 M in 2 M HCl) and 0.5 mL FeCl₂ (1 M in 2 M
125 HCl) with 40k Hz ultrasonic homologization, until the appearance of dark iron oxide
126 precipitates. The precipitates were further separated by permanent magnet and washed
127 with deionized water until the supernatant reached pH=7.0.

128 For bioreporter functionalization, 10 mL bioreporter suspensions (in deionized water)
129 were mixed with 1 mL MNPs suspension, incubated at room temperature for 10
130 minutes with 150 rpm shaking. The MNPs-bioreporter was subsequently harvested by
131 a permanent magnet and washed twice by deionized water. The magnetized
132 bioreporter was finally suspended in 10 mL MMS medium for soil toxicity
133 assessment via the magnet bioreporter device.

134 *2.4 Magnet bioreporter device and operation*

135 The magnet bioreporter device contained the magnet probe assay and plastic cover for
136 bioreporter strain transfer, as illustrated in Fig. 1a. The magnet probe assay was
137 assembled by 96 magnet probes (1 cm length and 3 mm id), fixed on the plastic base
138 and patterned (20.66 mm distance between each magnet probe) for the high
139 throughput bioreporter measurement on the 96-well microplate. For MNPs

140 functionalized bioreporter via the magnet device (MFB treatment), the 1.0 mL
141 bioreporter suspension was mixed with 100 mg soil samples and transferred into each
142 well of the 96-well microplate (Fig. 1 b-1). For the determination of the best
143 cultivation time before magnetic separation, the bioreporter cells were magnetically
144 harvest at 0, 15, 30, 45, 60, 75, 90, 105 and 120 min. The recovered MNPs
145 functionalized bioreporter was counted by plate count and the bioluminescent
146 response was also measured.

147 After incubation at 30°C for 1 hour as the optimal cultivation condition, the magnet
148 probe assay (with plastic cover) was emerged into the reaction system for 30 seconds
149 (Fig. 1 b-2). The bioreporter cells were then separated from the soil suspension and
150 attached on the plastic cover by magnetic field. The magnet device was transferred
151 and emerged in another 96-well microplate, supplemented with 200 μ L fresh MMS
152 medium (Fig. 1 b-3). Removing the magnet probe, the plastic cover and microplate
153 was incubated at 30°C for 5 minutes with 150 rpm shaking (Fig. 1 b-4). The
154 bioreporter cells were resuspended in the fresh MMS medium and the
155 bioluminescence was further measured on the Spectra M5 Plate Reader (Molecular
156 Devices, USA). The detection and data analysis followed the same instruction for the
157 direct toxicity measurement on soil samples.

158 For reproducibility test, MNPs-bioreporter was applied to sense the toxicity of 100
159 mg/kg chromium contaminated soils in the medium with different pH values and salt
160 contents. The pH value in the induction medium was adjusted by 1.0 M HCl or 1.0 M
161 NaOH solution as 4.0, 5.0, 6.0, 7.0, 8.0, 9.0 and 10.0. The series of salty medium was
162 prepared by adding NaCl into the MMS medium with the final concentration of 1%,
163 2%, 3%, 4%, 5% and 10%. As for the temperature influence, the temperature during
164 induction period was controlled by the Spectra M5 Plate Reader at 10°C, 15°C, 20°C,
165 30°C, 37°C, 40°C and 45°C. To evaluate the life-time of MNPs-bioreporters, the
166 bioreporter suspension was stored at 4°C and taken out for direct toxicity
167 measurement without any pre-treatment.

168 *2.5 Toxicity model for heavy metal contaminated soils*

169 For the calibration of heavy metal toxicity in soils, chromium was chosen at the target
170 analyte since it was the main heavy metal contamination at the coal cinder site. The
171 artificial chromium contaminated soils were prepared by mixing 1.0 g

172 uncontaminated soil with 1 mL potassium bichromate solution with the concentration
 173 of 0, 0.1, 0.2, 0.5, 1, 2, 5, 10, 20, 50, 100, 200, 500, 1000, 2000 and 5,000 mg/L. The
 174 soil slurry was air dried in the laminar hood. The chromium contamination in soils
 175 was 0, 0.1, 0.2, 0.5, 1, 2, 5, 10, 20, 50, 100, 200, 500, 1000, 2000 and 5,000 mg/kg
 176 soil dry weight, respectively.

177 The genotoxicity of hexavalent chromium in soil was identified as several
 178 mechanistic processes, including mutagenic effects on DNA and direct inductive
 179 immunological responses [30]. Such genotoxic effects consequently caused the
 180 accumulation of single stranded DNA (ssDNA) and the activation of SOS response
 181 for damaged DNA repair. The SOS response (bioluminescent response of
 182 ADPWH_recA) is the function of LexA-like SOS repressor (LSR , cell^{-1}) and can be
 183 simulated by the previous gene expression model [31, 32], as shown in Equation (1).

$$184 \quad SOS_{r,s} = 1 + \left(\frac{k_{dSLR}}{2 \cdot (1 + k_{ssDNA})} \cdot [LSR] \right) \cdot \frac{[Cr]}{(K_{mLSR} \cdot K_{Cr} \cdot k_{ssDNA} \cdot k_{Cr})^{-1} + [Cr]} \quad (1)$$

185 Here, $SOS_{r,s}$ (SOS response ratio) is dependent on the hexavalent chromium
 186 contamination level in soils ($[Cr]$, mg/kg). K_{Cr} is the isotherm equilibrium of
 187 chromium-DNA adduct (DNA phosphodiester backbone with chromium) and k_{Cr}
 188 represents ssDNA generation constant from the chromium-DNA adduct ($\text{L}/(\text{cell} \cdot \text{g})$)
 189 chromium). k_{dSLR} represents the equilibrium coefficient of LSR dimer ($dSLR$, cell^{-1})
 190 and monomer ($mLSR$, cell^{-1}) and k_{ssDNA} represents the cleavage reaction constant of
 191 LSR dimer. K_{mLSR} is the dynamic gene expression (SOS response) level activated by
 192 LSR monomer.

193 2.6 Sites description

194 A total of 16 soil samples were taken from the methanol plant of Yulin Energy and
 195 Chemical Industry, Yanzhou Coal Corporation, China (Fig. 2). The site (698,000 m^2)
 196 was located in Yulin Shaanxi Province (N38°34'41.9'', E109°55'50.4''), in the
 197 junction of Maowusu Sandy Land and the Loess Plateau. The annual coal
 198 consumption was 31,200 tonnes and the soils have been seriously contaminated by the
 199 coal with high heavy metal content. The sampling sites were designed along the
 200 leeward direction of the cinder heap, with the distance of 0, 10, 50, 80 and 150 m. The
 201 uncontaminated soil sample was collected in the living area of the plant, 500 m away
 202 from the heap. At each point, the soils were sampled at different depth of 0-20 cm

203 (surface soil), 20-35 cm (middle soil) and 35-50 cm (bottom soil) to evaluate the
204 toxicity profiles caused by the trace metal transportation.

205 *2.7 Chemical analysis*

206 Before chemical analysis, all of soil samples were seized by 200 mesh. Mercury was
207 determined by DMA-80 Hg analyzer (Milestone S.r.L., Italy). For other trace
208 elements, the samples were digested in an UltraCLAVE microwave high pressure
209 reactor (Milestone S.r.L., Italy), containing 330 mL distilled H₂O, 30 mL 30% H₂O₂
210 and 2 mL 98% H₂SO₄ as the digestion solution [33]. With 50 bars initial nitrogen
211 pressure, the microwave digestion program was listed in Table S1. Further digestion
212 for 50 mg soil sample was conducted in 5 mL 40% HF, 2 mL 65% HNO₃, and 1 mL
213 30% H₂O₂ [34]. The inductively coupled plasma mass spectrometry (ICP-MS, X
214 series II, Thermo Fischer Scientific, USA) was used for the determination of the trace
215 elements in a pulse counting mode (three points per peak). In this study, the multi-
216 element standards (Inorganic Ventures, CCS-1, CCS-4, CCS-5, and CCS-6) were
217 referenced for the calibration of trace element concentrations. As and Se were
218 determined by ICP-MS with collision cell technology (CCT) due to their volatility
219 [35]. Polyfluoroalkoxy volumetric flasks were used without drying on electric hot
220 plate to avoid As/Se volatile loss. With the 1 µg/L tuning solution, the torch position
221 and ion lenses were optimized before real sample measurement. The optimal
222 parameters of the ICP-CCT-MS and calibration curves of As/Se were listed in Table
223 S2 and S3.

224 *2.8 Data analysis*

225 The bioluminescence response was calculated by averaging the bioluminescent signal
226 from the 7 time points between 180 and 240 minutes for each well. The relative
227 bioluminescence response ratio was the specific value of the bioluminescence
228 response of contaminated soil samples to that of the uncontaminated soils. The heavy
229 metal profiles in soil samples were statistically analysed by SPSS software (Version
230 15.0 for Windows) via Principal Component Analysis (PCA). The equality and
231 normality of data were tested by Brown-Forsythe and Shapiro-Wilk test respectively,
232 and the null hypothesis was rejected for $p < 0.05$.

233 Contamination factor (*CF*) is defined as the ratio of the heavy metal concentration in
234 the sample soil to the baseline concentration in background soil, as shown in Equation

235 (2) [36]. Pollution load index (*PLI*) is determined as the n^{th} root of the n CF in
236 Equation (3) [36]. The *CF* and *PLI* are empirical indices to evaluate the level of heavy
237 metal contamination, and the higher values indicate heavier contamination of
238 individual and multiple heavy metals respectively.

$$239 \quad CF = \frac{[\text{Heavy metal in sample soils}]}{[\text{Heavy metal in background soils}]} \quad (2)$$

$$240 \quad PLI = (CF_1 \times CF_2 \times \dots \times CF_n)^{1/n} \quad (3)$$

241 **3. Results and discussions**

242 *3.1 Higher sensitivity and reproducibility of magnet bioreporter device*

243 The MNPs functionalized bioreporter could be magnetic remote controlled for
244 effectively separation from the soil particles. The MNPs were biocompatible, and the
245 viability and bioluminescent signal of whole-cell bioreporter remained over 99%
246 comparing to the native bioreporter cells [27]. With the strong electrostatic attraction
247 between the negative iron oxide (Fe-OO^-) and positive amino-groups ($-\text{NH}_3^+$) on
248 bacterial membrane, the separation effectiveness by magnetic field was above 99.6%
249 and the synthesized MNPs had neither cytotoxicity nor genotoxicity on bacterial
250 bioreporter cells [37]. MNPs functionalized whole-cell bioreporter therefore had the
251 feasibility to sense the toxicity of soil samples *in situ* and subsequently isolated for
252 bioluminescent signal measurement.

253 Due to the cell division, the MNPs functionalized bioreporter gradually lost their
254 magnetic capacity [38]. Though longer incubation with soil samples could improve
255 the chemical uptake by bioreporter cells for higher responsive ratio, the less recovery
256 rate consequently resulted in lower bioluminescent signal and lower sensitivity. Fig. 3
257 illustrated that, within 45 minutes incubation, over 90% living bioreporter cells were
258 isolated from the soil/water mixture based on plate count. The results fitted with
259 previous study that about 12% free bioreporter cells were observed after 120 min
260 cultivation in rich medium [27]. As for the bioluminescence and relative response
261 ratios, the bioluminescent signals were stable from 3600 RLU to 3800 RLU when the
262 incubation time was less than 75 minutes, and the response ratio ranged from 1.90 to
263 2.00. The results suggested a highly reliable responsive period between 30 and 70
264 minutes. The optimal incubation time for sufficient bioreporter cell recovery and high

265 response sensitivity was identified as 60 minutes, and it was further applied in the
266 following work on calibration curve and real soil sample assessment.

267 The summarized features of the magnetic ADPWH_recA whole-cell bioreporter were
268 listed in Table 1 from the reproducibility test. After 1 hour pre-incubation of the
269 MNPs functionalized bioreporter, the cells were captured by permanent magnet and
270 resuspended in fresh medium without soil disturbance for another 4 hours. As a soil
271 bacterium, ADPWH_recA had strong tolerance to the environmental variations and
272 maintained high reproducibility under different pH, salinity and temperature condition.
273 The relative bioluminescent response ratio maintained stable (1.44 to 1.51) when pH
274 value ranged from 5.0 to 9.0, dramatically dropping to 1.25 at pH=4.0 and 1.12 at
275 pH=10.0 (Fig. 4a). The results were similar to previous research on the pH influence
276 on *Acinetobacter baylyi* ADP1 that *Acinetobacter* based bioreporter could tolerate
277 large pH variation [39]. Fig. 4b also illustrated the good responsive performance of
278 MNPs functionalized ADPWH_recA at 20°C (relative bioluminescent response
279 ratio=1.47), 30°C (relative bioluminescent response ratio=1.50) and 37°C (relative
280 bioluminescent response ratio=1.49). The tiny reduction of bioluminescent response
281 at 15°C and 40°C attributed to the less bacterial activities at inappropriate
282 temperatures, and the response was very weak under even lower (10°C) or higher
283 (45°C) temperature conditions. Salinity did not significantly affect the reproducibility
284 of ADPWH_recA and the relative bioluminescent response ratios were above 1.45
285 when the salinity was no higher than 3%, but were gradually suppressed at higher
286 salinity level (Fig. 4c). Therefore, the MNPs functionalized bioreporters had high
287 reproducibility under the normal pH value, salinity and temperature conditions of
288 natural soils and no specific pre-treatment was required for real soil sample
289 assessment. High activity and responsive sensitivity of MNPs functionalized whole-
290 cell bioreporters was also observed after 30 days storage at 4°C (Fig. 4d). Without
291 any pre-treatment, the stored bioreporter cells could be directly applied for soil
292 assessment and the relative bioluminescent response ratio was above 1.45 for
293 chromium contaminated soils of 100 mg/kg soil dry weight. The life-time of MNPs
294 functionalized bioreporter was the same to the original *Acinetobacter* based
295 bioreporters [28, 40], indicating that MNPs functionalization had minimal impacts on
296 the bacterial activities and was an appropriate approach to expand its application in
297 soil contamination assessment.

298 From the calibration curve of soil/water supernatant (SW-S), soil/water mixture (SW-
299 M) and MNPs functionalized bioreporter (MFB) (Fig. 5), magnet device had the
300 highest responsive sensitivity and illustrated the chromium bioavailability in
301 contaminated soils. In SW-S and SW-M treatments, ADPWH_recA bioreporter did
302 not show any positive response to the chromium due to the strong light adsorption by
303 soil particles. The negative bioluminescent response was observed when chromium
304 concentration was above 100 mg/kg soil dry weight for both treatments. Significant
305 positive response was only found in MFB treatment and the limit of detection was 1
306 mg/kg soil dry weight (Fig. 5 and Table 1). From 1 mg/kg to 100 mg/kg chromium
307 contamination in dry soils, the relative bioluminescence response ratio showed a
308 linear relationship to quantify the toxicity and bioavailability of chromium in soil
309 samples, ranging from 1.05 to 1.60. Above 500 mg/kg soil dry weight, chromium
310 predominantly behaved the cytotoxicity effects and all the three treatments had similar
311 inhibited bioluminescent signal. From the whole-cell bioreporter growth curve (Fig.
312 S1), there was no significant growth difference when the chromium concentration was
313 less than 500 mg/kg soil dry weight, in which range that the relative bioluminescent
314 response ratio was positively correlated with chromium. It therefore explained the
315 decreasing bioluminescent response ratio at higher chromium level that strong
316 cytotoxicity of chromium inhibited bioreporter growth and activities.

317 Given the model simulation of bioreporter's response to chromium with different
318 bioavailability in aqueous phase in Fig. 5 [41], the results further revealed the
319 bacteria-contaminant interaction within the porous soils and its impacts on bioreporter
320 response. With lower chromium bioavailability, the calibration curve shifted towards
321 higher chromium values. The SOS response coefficient ($K_{SLR} \cdot K_{Cr} \cdot k_{ssDNA} \cdot k_{MO}$,
322 3.8) and genotoxicity coefficient ($\frac{k_{dsSLR}}{2 \cdot (1 + k_{ssDNA})} \cdot [LSR]_{total}$, 1.724 L/mg) kept stable,
323 similar to previous research [41]. Referring to the synergetic efficiency through the
324 SOS repair process (including genotoxin DNA damage, ssDNA recognition and SOS
325 box activation), the similar SOS response coefficients indicated the same SOS
326 mechanism of bioreporter's responsive to chromium genotoxicity and cytotoxicity in
327 the soils via the magnet bioreporter device [32]. Since the bioluminescent signal of
328 ADPWH_recA was regulated by the SOS process, all the carcinogens causing DNA
329 damage would activate its response, including mitomycin C, UV light, ethidium
330 bromide and H₂O₂ [22]. The bioreporter therefore did not respond to a particular

331 heavy metal (like chromium), but evaluated the synergistic toxicity of all the
332 carcinogens in environmental samples.

333 From the parabolic curve of MNPs functionalized bioreporter to hexavalent chromium,
334 both the relative bioluminescent response ratio and growth curve (Fig. S1) were
335 considered to evaluate the toxicity of unknown environmental samples. In absence of
336 growth inhibition, the sample had low cytotoxicity and its bioluminescent response
337 belonged to the positive relationship range, oppositely in presence of growth
338 inhibition. Only the MFB treatment had the positive bioluminescence response when
339 chromium concentration was less than 200 mg/kg soil dry weight, and the response
340 ratio fitted well with the model prediction of 10% chromium bioavailability when
341 chromium concentration was above 100 mg/kg soil dry weight. At lower chromium
342 contamination level, chromium bioavailability changed due to the complex adsorption
343 effects of soil particles and the irregular bioluminescent response ratio represented the
344 changing bioavailable fraction. Given heavy chromium contamination level (>500
345 mg/kg soil dry weight), MFB and SW-M treatments had similar responsive results,
346 significantly higher than SW-S treatment. Since the whole-cell bioreporter only
347 sensed the water soluble chromium in the supernatant of soil-water mixture in SW-S
348 treatment, it measured the chromium toxicity in the unbound water phase. The
349 dominant fraction of chromium existed in the bound water or was absorbed on the soil
350 particles, and their carcinogenic effects was only assessable by the direct-contact
351 bioreporter assay [24]. The results indicated magnetic functionalized bioreporter
352 could effectively evaluate the real toxic effects of chromium in soils, by directly
353 contacting soil particles and seeking for chromium in unbound/bound water or
354 adsorbed on particle surface. The portable magnet bioreporter device provided the
355 ready-to-use and nature-inspired technique for soil *in situ* measurement by optimizing
356 the operation procedure and enhancing the bioluminescent signal [42].

357 *3.2 Heavy metal contamination profiles in soils*

358 The heavy metal profiles of the 16 investigated soil samples and the coal cinder were
359 listed in Table 2. The chromium was 23 times enriched in the rough cinder (from
360 38.18 mg/kg to 920.82 mg/kg), followed by nickel (5.5 times enrichment). The
361 enrichment of other heavy metals ranged from 1.8 to 3.7 times, except for zinc (no
362 significant change) and mercury (loss due to volatile during the combustion process)
363 [43]. Similar to previous research on coal combustion residues [44, 45], chromium

364 and nickel were mostly found concentrated in bottom ash or cinder as the dominant
365 toxic heavy metal residues, due to their low volatility and high stability.

366 As for heavy metal contamination in soils, the investigated soils only showed
367 significant contamination of chromium from 448.66 mg/kg to 802.77 mg/kg soil dry
368 weight, with the *CF* values from 1.12 to 2.01 (398.94 mg/kg soil dry weight in
369 background). The levels of the other heavy metals were similar to the background
370 soils (p -value>0.05), as the *CF* value of beryllium ranged from 0.80 to 1.19, nickel
371 from 0.40 to 0.95, copper from 0.50 to 1.34, zinc from 0.43 to 0.78, arsenic from 0.41
372 to 0.88, selenium from 0.33 to 1.37, cadmium from 0.50 to 0.96, lead from 0.83 to
373 1.40, uranium from 0.38 to 1.28 and mercury from 0.13 to 4.13. The results indicated
374 that the key ecological toxicity was attributed to chromium, same as revealed by many
375 previous research on mutagens in heavy metal contaminated soils [46], and its toxicity
376 in soils could be effectively evaluated by various biological assays [47, 48]. The
377 magnet bioreporter device in this study was therefore feasible to enhance the toxicity
378 test sensitive by directly exposing MNPs-functionalized whole-cell bioreporter cells
379 and diagnosing their bioluminescent response signal.

380 For soils nearer to the coal cinder point (0 m and 10 m), there was higher heavy metal
381 contamination in the upper layer soil. The chromium contamination declined from
382 745.15 mg/kg (0-20 cm, 0 m) to 505.60 mg/kg (35-50 cm, 0 m), and from 802.77
383 mg/kg (0-20 cm, 10 m) to 525.79 mg/kg (35-50 cm, 10 m), respectively. Except for
384 Be, Pb and Hg, all the other heavy metal elements (Ni, Cu, Zn, As, Se, Cd, and U) has
385 the same vertical decreasing distribution. Comparing to the heavy metal composition
386 in the cinders, chromium was also the key carcinogenic heavy metal in soils. Given
387 the sequence of the exchangeable fractions of heavy metals as $Cd > Zn > Cu > Ni >$
388 $Pb > Cr$ [5], chromium was further concentrated in top soils with less mobility. The
389 results further suggested that the main source of heavy metal contamination in soils
390 was the leachates from the cinders and their residues were dependant on the
391 transportation process in soils.

392 *3.3 Ecological risk profiles of heavy metal contaminated soils*

393 In the ecological toxicity evaluation by directly applying whole-cell bioreporters to
394 the soil (SW-M, Fig. S2 in Supplementary Materials) and soil-water supernatant (SW-
395 S, Fig. S3 in Supplementary Materials), ADPWH_recA only behaved negative

396 (relative bioluminescence response ratio < 1.0) or neutral signal (relative
397 bioluminescence response ratio = 1.0) and was not suitable to quantify the toxicity
398 impacts of heavy metal contamination *in situ*. Fig. 6 illustrated the ecological toxicity
399 profiles of the soil samples by the magnetic bioreporter device (MFB treatment), and
400 the toxicity of heavy metals declined with the increasing distance to the coal cinder
401 point. From the whole-cell bioreporter growth curve for the soil samples (Fig. S4), all
402 the heavy metal contaminated soils did not show inhibition effects on bacterial growth,
403 indicating all the bioluminescent signals were within the linear response range and the
404 relative bioluminescence response ratio had positive relationship with the ecological
405 toxicity in soils. Except for 0 m point, the relative bioluminescence response ratio
406 dropped from 1.47 (10 m) to 1.10 (150 m) in the surface soil, 1.34 (10 m) to 1.16 (150
407 m) in the middle soil, and 1.26 (0 m) to 0.58 (150 m) in the bottom soil. At the 0 m
408 point, the low bioluminescence signal of surface soil was caused by the high
409 cytotoxicity effects of chromium (745.15 mg/kg soil dry weigh) and the growth of
410 ADPWH_recA bioreporter was inhibited. The soil sample at 0 m point was therefore
411 characterized with the highest ecological risk.

412 From the toxicity vertical distribution, the ecological risks had a significant decline in
413 deeper soils. Attributing to the heavy metals leachates from the coal cinders, the
414 ecological risk distribution fitted well with chemical analysis and previous studies.
415 The high ecological risk at the surface soils than bottom soils suggested the leakage
416 and vertical transportation chromium in soils [49]. Comparing to the horizontal
417 ecological risk distribution, the results further identified the main toxicity sources as
418 the heavy metals from the coal cinders.

419 *3.4 Correlation between soil heavy metal profiles and ecological risk*

420 The Principle Component Analysis (PCA) illustrated the main factors causing the
421 ecological risks in soil samples (Fig. S5a). More precisely, the principle component 1
422 (PC1) was the heavy metal contamination level, accounting for 60.5% of the total
423 variance. At the sampling points nearer to the coal cinder site (0 m and 10 m), the
424 surface and middle soils were heavily contaminated and therefore recognized as
425 isolated square (red) and circle (blue) to the higher value of PC1-axis. For the rest
426 soils, they gathered due to similar contamination level (*PLI*). PC1 was therefore
427 derived from the external heavy metal sources, leaching from the coal cinder for the
428 surface soil (0-20 cm) and heavy metal vertical transportation for middle soil (20-35

429 cm). The soil depth was the principle component 2 (PC2), contributing to 13.3% of
430 the total variance). Heavy metals distribution and mobility were reported to depend on
431 soil properties and depth [50], and their spatial distribution in different depths of soils
432 also affected the mobility and bioavailability [51]. Nevertheless, the soil ecological
433 risks (illustrated as the area of each symbol) were associated with neither the load of
434 PC1 nor PC2, suggesting that they were complicatedly affected by both heavy metal
435 profiles and soil features.

436 There was also no significant correlation between heavy metal pollution load index
437 (*PLI*) and ecological risk (p -value>0.05) (Fig. S5b). Higher *PLI* indicated high heavy
438 metal contamination level, but did not fit with the ecological risk distribution.
439 Previous research had shown the positive correlation between heavy metal content
440 and ecological toxicity at the contaminated sites with individual heavy metal pollutant,
441 like chromium residues [24] or copper contaminated agricultural soils [52]. The
442 ecological toxicity was only affected by the individual *EF* value and bioavailability in
443 soil. At the coal cinder contaminated sites, we found the existence of multiple heavy
444 metals and their synergic/antagonistic effects consequently resulted in complicated
445 ecological toxicity [53]. Many evidences had revealed that the toxicity of individual
446 or multiple heavy metals behaved antagonistic or additive effects, dependent on the
447 composition and soil features, like organic matters or pH value [54, 55]. In this case,
448 *PLI* was an empirical indicator evaluating the multiple heavy metal contamination
449 level, but suffered from identifying and characterizing the interaction between various
450 heavy metal molecules and their association with soil particles. From the mechanisms
451 of ADPWH_recA to sense all the carcinogens activating SOS process, the response of
452 whole-cell bioreporter effectively represented the synergic/antagonistic effects of
453 multiple heavy metals. By directly exposing the living bioreporter cells to the
454 contaminated samples *in situ*, the MNPs functionalized bioreporter had its feasibility
455 as an important approach, supplementary to chemical analysis, in ecological risk
456 assessment and environmental risk management.

457 **4. Conclusion**

458 This work developed a novel magnet bioreporter device for soil toxicity assessment,
459 via magnetic nanoparticles functionalized whole-cell bioreporters. The living
460 magnetic bioreporter cells could sense the carcinogenic chemicals in the soil and
461 effectively separated from the soil-water slurry in the bioluminescence detection step

462 to avoid the disturbance of soil particles. Comparing to the conventional treatments
463 directly applying bioreporter in soil-water mixture or supernatant, the magnet
464 bioreporter device achieved high sensitivity and reproducibility under soil pH, salinity
465 and temperature conditions. The dose-toxicity calibration curve revealed the impacts
466 of chromium bioavailability on its ecological risk in soils, where strong genotoxicity
467 was identified when chromium concentration was from 1 mg/kg to 500 mg/kg soil dry
468 weight and the cytotoxic inhibition was found at chromium over 500 mg/kg soil dry
469 weight. For the first time, the ecological toxicity of heavy metal contaminated soils
470 was evaluated by the whole-cell bioreporter at the coal cinder site. Though the
471 existence of heavy metal contamination contributed to the main ecological risks at the
472 site, the pollution load index (*PLI*) had no significantly relationship with the
473 ecological toxicity distribution. The synergic and antagonistic effects of soil multiple
474 heavy metal contamination brought the challenges for environmental risk assessment
475 by chemical analysis. The magnetic bioreporter device behaved as an alternative
476 approach for the high throughput biological measurement and was feasible for *in situ*
477 monitoring.

478 **Acknowledgement**

479 The authors would like to thank National Basic Research Program of China (973
480 Program, No. 2014CB238906) and National Natural Science Foundation of China
481 (No. 41301331) for financial support. Prof Shifeng Dai (China University of Mining
482 & Technology, Beijing) helped in ICP-MS analysis of heavy metal components in the
483 soil samples.

484

485

486 **Reference**

- 487 [1] X.D. Li, O.W.H. Wai, Y.S. Li, B.J. Coles, M.H. Ramsey, I. Thornton, Heavy
488 metal distribution in sediment profiles of the Pearl River estuary, South China, Appl
489 Geochem, 15(2000) 567-81.
- 490 [2] K. Loska, D. Wiechula, I. Korus, Metal contamination of farming soils affected by
491 industry, Environ Int, 30(2004) 159-65.
- 492 [3] S.R. Smith, A critical review of the bioavailability and impacts of heavy metals in
493 municipal solid waste composts compared to sewage sludge, Environ Int, 35(2009)
494 142-56.
- 495 [4] H. Alizai, D. Preston, S.S. Meng, M. Kolkas, Heavy metal contamination in the
496 coastal sediments of the Lower New York Bay, Northeastern Geology and
497 Environmental Sciences, 25(2003) 226-31.
- 498 [5] P.J. He, Z. Xiao, L.M. Shao, J.Y. Yu, D.J. Lee, In situ distributions and
499 characteristics of heavy metals in full-scale landfill layers, J Hazard Mater, 137(2006)
500 1385-94.
- 501 [6] Z. Dang, C.Q. Liu, M.J. Haigh, Mobility of heavy metals associated with the
502 natural weathering of coal mine spoils, Environ Pollut, 118(2002) 419-26.
- 503 [7] M. Valko, H. Morris, M.T.D. Cronin, Metals, toxicity and oxidative stress, Curr
504 Med Chem, 12(2005) 1161-208.
- 505 [8] K.E. Giller, E. Witter, S.P. McGrath, Toxicity of heavy metals to microorganisms
506 and microbial processes in agricultural soils: A review, Soil Biol Biochem, 30(1998)
507 1389-414.
- 508 [9] L. Jarup, Hazards of heavy metal contamination, Br Med Bull, 68(2003) 167-82.
- 509 [10] M.L. Farre, R. Brix, D. Barcelo, Screening water for pollutants using biological
510 techniques under European Union funding during the last 10 years, Trends in
511 Analytical Chemistry, 24(2005) 532-45.
- 512 [11] S. Belkin, Microbial whole-cell sensing systems of environmental pollutants,
513 Curr Opin Microbiol, 6(2003) 206-12.
- 514 [12] T.K. Van Dyk, E.J. DeRose, G.E. Gonye, LuxArray, a high-density, genomewide
515 transcription analysis of *Escherichia coli* using bioluminescent reporter strains, J
516 Bacteriol, 183(2001) 5496-505.
- 517 [13] E.A. Meighen, Genetics of bacterial bioluminescence, Annu Rev Genet, 28(1994)
518 117-39.

- 519 [14] J. Sanseverino, R.K. Gupta, A.C. Layton, S.S. Patterson, S.A. Ripp, L. Saidak, et
520 al., Use of *Saccharomyces cerevisiae* BLYES expressing bacterial bioluminescence
521 for rapid, sensitive detection of estrogenic compounds, *Appl Environ Microbiol*,
522 71(2005) 4455-60.
- 523 [15] S.F. D'Souza, *Microbial Sensors*, *Biosens Bioelectron*, 16(2001) 337-53.
- 524 [16] L.D. Rasmussen, S.J. Sorensen, R.R. Turner, T. Barkay, Application of a mer-lux
525 biosensor for estimating bioavailable mercury in soil, *Soil Biol Biochem*, 32(2000)
526 639-46.
- 527 [17] A. Ivaska, M. Virtab, A. Kahrua, Construction and use of specific luminescent
528 recombinant bacterial sensors for the assessment of bioavailable fraction of cadmium,
529 zinc, mercury and chromium in the soil, *Soil Biol Biochem*, 34(2002) 1439-47.
- 530 [18] S. Rodriguez-Mozaz, M.J. Lopez de Alda, D. Barcelo, Biosensors as useful tools
531 for environmental analysis and monitoring, *Analytical and Bioanalytical Chemistry*,
532 386(2006) 1025-41.
- 533 [19] J.R. van der Meer, S. Belkin, Where microbiology meets microengineering:
534 design and applications of reporter bacteria, *Nat Rev Microbiol*, 8(2010) 511-22.
- 535 [20] P. Corbisier, E. Thiry, L. Diels, Bacterial biosensors for the toxicity assessment
536 of solid wastes, *Environmental Toxicology and Water Quality*, 11(1996) 171-7.
- 537 [21] C. Tibazarwa, P. Corbisier, M. Mench, A. Bossus, P. Solda, M. Mergeay, et al.,
538 A microbial biosensor to predict bioavailable nickel and its transfer to plants, *Environ*
539 *Pollut*, 113(2001) 19-26.
- 540 [22] Y. Song, B. Jiang, S. Tian, H. Tang, Z. Liu, C. Li, et al., A whole-cell bioreporter
541 approach for the genotoxicity assessment of bioavailability of toxic compounds in
542 contaminated soil in China, *Environ Pollut*, 195C(2014) 178-84.
- 543 [23] V.H.C. Liao, M.T. Chien, Y.Y. Tseng, K.L. Ou, Assessment of heavy metal
544 bioavailability in contaminated sediments and soils using green fluorescent protein-
545 based bacterial biosensors, *Environ Pollut*, 142(2006) 17-23.
- 546 [24] B. Jiang, D. Zhu, Y. Song, D. Zhang, Z. Liu, X. Zhang, et al., Use of a whole-cell
547 bioreporter, *Acinetobacter baylyi*, to estimate the genotoxicity and bioavailability of
548 chromium(VI)-contaminated soils, *Biotechnol Lett*, 37(2015) 343-8.
- 549 [25] R. Turpeinen, J. Salminen, T. Kairesalo, Mobility and bioavailability of lead in
550 contaminated boreal forest soil, *Environ Sci Technol*, 34(2000) 5152-6.

551 [26] H. Fritze, J. Perikomaki, T. Petanen, T. Pennanen, M. Romantschuk, M. Karp, et
552 al., A microcosms study on the effects of Cd containing wood ash on the coniferous
553 humus fungal community and the Cd bioavailability, *Soils Sediments*, 1(2001) 146-50.

554 [27] D. Zhang, R.F. Fakhrullin, M. Özmen, H. Wang, J. Wang, V.N. Paunov, et al.,
555 Functionalization of whole-cell bacterial reporters with magnetic nanoparticles,
556 *Microbial Biotech*, 4(2011) 89-97.

557 [28] Y.Z. Song, G.H. Li, S.F. Thornton, I.P. Thompson, S.A. Banwart, D.N. Lerner,
558 et al., Optimization of Bacterial Whole Cell Bioreporters for Toxicity Assay of
559 Environmental Samples, *Environ Sci Technol*, 43(2009) 7931-8.

560 [29] D. Zhang, A. Ding, S. Cui, C. Hu, S.F. Thornton, J. Dou, et al., Whole cell
561 bioreporter application for rapid detection and evaluation of crude oil spill in seawater
562 caused by Dalian oil tank explosion, *Water Res*, 47(2013) 1191-200.

563 [30] K.P. Nickens, S.R. Patierno, S. Ceryak, Chromium genotoxicity: A double-edged
564 sword, *Chem Biol Interact*, 188(2010) 276-88.

565 [31] A.A. Al-Anizi, M.T. Hellyer, D. Zhang, Toxicity assessment and modeling of
566 *Moringa oleifera* seeds in water purification by whole cell bioreporter, *Water Res*,
567 56(2014) 77-87.

568 [32] D. Zhang, Y. Zhao, Y. He, Y. Wang, Y. Zhao, Y. Zheng, et al., Characterization
569 and Modeling of Transcriptional Cross-Regulation in *Acinetobacter baylyi* ADP1,
570 *Acs Synthetic Biology*, 1(2012) 274-83.

571 [33] S. Dai, X. Wang, Y. Zhou, J.C. Hower, D. Li, W. Chen, et al., Chemical and
572 mineralogical compositions of silicic, mafic, and alkali tonsteins in the late Permian
573 coals from the Songzao Coalfield, Chongqing, Southwest China, *Chem Geol*,
574 282(2011) 29-44.

575 [34] S. Dai, X. Wang, V.V. Seregin, J.C. Hower, C.R. Ward, J.M.K. O'Keefe, et al.,
576 Petrology, mineralogy, and geochemistry of the Ge-rich coal from the Wulantuga Ge
577 ore deposit, Inner Mongolia, China: New data and genetic implications, *Int J Coal*
578 *Geol*, 90(2012) 72-99.

579 [35] X. Li, S. Dai, W. Zhang, T. Li, X. Zheng, W. Chen, Determination of As and Se
580 in coal and coal combustion products using closed vessel microwave digestion and
581 collision/reaction cell technology (CCT) of inductively coupled plasma mass
582 spectrometry (ICP-MS), *Int J Coal Geol*, 124(2014) 1-4.

583 [36] M.A.H. Bhuiyan, L. Parvez, M.A. Islam, S.B. Dampare, S. Suzuki, Heavy metal
584 pollution of coal mine-affected agricultural soils in the northern part of Bangladesh, *J*
585 *Hazard Mater*, 173(2010) 384-92.

586 [37] C. Chen, D. Zhang, S.F. Thornton, M. Duan, Y. Luo, A. Ding, et al.,
587 Functionalization and immobilization of whole cell bioreporters for the detection of
588 environmental contamination, *Environmental Engineering and Management Journal*,
589 12(2013) 1417-22.

590 [38] D. Zhang, J.P. Berry, D. Zhu, Y. Wang, Y. Chen, B. Jiang, et al., Magnetic
591 nanoparticle-mediated isolation of functional bacteria in a complex microbial
592 community, *ISME J*, 9(2015) 603-14.

593 [39] C. Li, H. Wang, L. Zhou, Y. Zhang, F. Song, J. Zhang, Quantitative
594 measurement of pH influence on SalR regulated gene expression in *Acinetobacter*
595 *baylyi* ADP1, *J Microbiol Methods*, 79(2009) 8-12.

596 [40] D. Zhang, Y. He, Y. Wang, H. Wang, L. Wu, E. Aries, et al., Whole-cell
597 bacterial bioreporter for actively searching and sensing of alkanes and oil spills,
598 *Microbial Biotech*, 5(2012) 87-97.

599 [41] Y. Xu, J. Zhu, H. Li, Y. Du, D. Zhang, Removal and immobilization of
600 hexavalent chromium contaminated water by magnetic nanoparticles, *Environ Sci*
601 *Technol*, Under review.

602 [42] E. Michelini, L. Cevenini, M.M. Calabretta, S. Spinuzzi, C. Camborata, A. Roda,
603 Field-deployable whole-cell bioluminescent biosensors: so near and yet so far, *Anal*
604 *Bioanal Chem*, 405(2013) 6155-63.

605 [43] R. Meij, L.H.J. Vredendregt, H.T. Winkel, The fate and behavior of mercury in
606 coal-fired power plants, *J Air Waste Manag Assoc*, 52(2002) 912-7.

607 [44] F. Goodarzi, F.E. Huggins, H. Sanei, Assessment of elements, speciation of As,
608 Cr, Ni and emitted Hg for a Canadian power plant burning bituminous coal,
609 *International Journal of Coal Geology*, 74(2008) 1-12.

610 [45] F. Goodarzi, Assessment of elemental content of milled coal, combustion
611 residues, and stack emitted materials: Possible environmental effects for a Canadian
612 pulverized coal-fired power plant, *International Journal of Coal Geology*, 65(2006)
613 17-25.

614 [46] P.A. White, L.D. Claxton, Mutagens in contaminated soil: A review, *Mutat Res-*
615 *Rev Mutat*, 567(2004) 227-345.

- 616 [47] H.Q. Wang, Clastogenicity of chromium contaminated soil samples evaluated by
617 Vicia root-micronucleus assay, Mutation Research-Fundamental and Molecular
618 Mechanisms of Mutagenesis, 426(1999) 147-9.
- 619 [48] J.C. Codina, C. Pereztorrente, A. Perezgarcia, F.M. Cazorla, A. Devicente,
620 Comparison of microbial tests for the detection of heavy metal genotoxicity, Archives
621 of Environmental Contamination and Toxicology, 29(1995) 260-5.
- 622 [49] X. Ma, Z. Lu, J. Cheng, Ecological risk assessment of open coal mine area,
623 Environ Monit Assess, 147(2008) 471-81.
- 624 [50] L. Hernandez, A. Probst, J.L. Probst, E. Ulrich, Heavy metal distribution in some
625 French forest soils: Evidence for atmospheric contamination, Sci Total Environ,
626 312(2003) 195-219.
- 627 [51] K. Nemati, N.K. Abu Bakar, M.R. Abas, E. Sobhanzadeh, Speciation of heavy
628 metals by modified BCR sequential extraction procedure in different depths of
629 sediments from Sungai Buloh, Selangor, Malaysia, J Hazard Mater, 192(2011) 402-10.
- 630 [52] K.K. Brandt, P.E. Holm, O. Nybroe, Bioavailability and toxicity of soil particle-
631 associated copper as determined by two bioluminescent *Pseudomonas fluorescens*
632 biosensor strains, Environ Toxicol Chem, 25(2006) 1738-41.
- 633 [53] M. Holmstrup, A.M. Bindesbol, G.J. Oostingh, A. Duschl, V. Scheil, H.R.
634 Kohler, et al., Interactions between effects of environmental chemicals and natural
635 stressors: A review, Sci Total Environ, 408(2010) 3746-62.
- 636 [54] C.A.M. VanGestel, P.J. Hensbergen, Interaction of Cd and Zn toxicity for
637 *Folsomia candida* Willem (Collembola:Isotomidae) in relation to bioavailability in
638 soil, Environmental Toxicology and Chemistry, 16(1997) 1177-86.
- 639 [55] S. Preston, N. Coad, J. Townend, K. Killham, G.I. Paton, Biosensing the acute
640 toxicity of metal interactions: Are they additive, synergistic, or antagonistic?, Environ
641 Toxicol Chem, 19(2000) 775-80.

642

643

644 **Figure caption**

645 **Fig. 1** Schematic instruction for magnet bioreporter device. (a-1) and (a-2) for
646 magnetic probe assay and the 96-well microplate respectively. (b-1) The 1.0 mL
647 MNPs functionalized bioreporter was mixed with soil samples and further incubated
648 at 30°C for 1 hour; (b-2) separation from soil suspensions via magnetic probe. (b-3)
649 Resuspension in fresh MMS medium; (b-4) incubation and bioluminescence
650 measurement 30°C for 3 hours.


651 **Fig. 2.** Location of research area in Yulin and the sampling sites.

652 **Fig. 3.** The cell recovery rate (A) and bioluminescent response (B) of whole-cell
653 bioreporter against the incubation time with soils. Over 90% of living bioreporter
654 cells were successfully harvested from the soil/water mixture within 45 minutes
655 incubation. The relative bioluminescence response ratio ranged between 1.90 and 2.00
656 when the incubation time was less than 75 minutes. The 60-minute incubation was
657 identified as the optimal time for sufficient bioreporter cell recovery and high
658 response sensitivity.

659 **Fig. 4.** The impacts of pH (a), temperature (b), salt (c) and storage time (d) on
660 magnetic bioreporter's response to artificial chromium contaminated soils. The
661 chromium concentration was 100 mg/kg soil dry weight.

662 **Fig. 5.** The calibration curve for toxicity assessment on artificial chromium
663 contaminated soils. Grey circle refers to magnet bioreporter device (MFB); white
664 diamond represents direct measurement of soil/water supernatant (SW-S); white circle
665 is the direct measurement of soil/water mixture (SW-M). The black line represents the
666 simulation of whole-cell bioreporter's response to chromium toxicity with 100% 100%
667 bioavailability, and a significant bioluminescent response curve shift was found for 50%
668 (red line), 30% (yellow line) and 10% (green line) chromium bioavailability
669 respectively.

670 **Fig. 6.** Ecological toxicity assessment of heavy metal contaminated soils via magnetic
671 bioreporter device.




672

673 **Fig. 1** Schematic instruction for magnet bioreporter device. (a-1) and (a-2) for
 674 magnetic probe assay and the 96-well microplate respectively. (b-1) The 1.0 mL
 675 MNPs functionalized bioreporter was mixed with soil samples and further incubated
 676 at 30°C for 1 hour; (b-2) separation from soil suspensions via magnetic probe. (b-3)
 677 Resuspension in fresh MMS medium; (b-4) incubation and bioluminescence
 678 measurement 30°C for 3 hours.

679

680




681

682 **Fig. 2.** Location of research area in Yulin and the sampling sites.

683


684



685

686

(a)




687

688

(b)


689 **Fig. 3.** The cell recovery rate (a) and bioluminescent response (b) of whole-cell
 690 bioreporter against the incubation time with soils. Over 90% of living bioreporter
 691 cells were successfully harvested from the soil/water mixture within 45 minutes
 692 incubation. The relative bioluminescence response ratio ranged between 1.90 and 2.00
 693 when the incubation time was less than 75 minutes. The 60-minute incubation was
 694 identified as the optimal time for sufficient bioreporter cell recovery and high
 695 response sensitivity.




696

697 **Fig. 4.** The impacts of pH (a), temperature (b), salt (c) and storage time (d) on
 698 magnetic bioreporter's response to artificial chromium contaminated soils. The
 699 chromium concentration was 100 mg/kg soil dry weight.

700





711

712 **Fig. 6.** Ecological toxicity assessment of heavy metal contaminated soils via magnetic
 713 bioreporter device.

714

715

Table

Table 1. Analytical characteristics of magnetic ADPWH_recA whole-cell bioreporter.

Characteristics	Description
Limit of detection	1 mg/kg chromium(VI) in dry soil
Linear range	1-100 mg/kg chromium(VI) in dry soil
Pre-incubation time	1 hour
Detection time	4 hour
Reproducibility	High reproducible when pH value is from 4.0 to 9.0, salinity ranges from 0% to 3%, and temperature is from 20 °C to 37 °C
Storage time	30 days

Table 2. Heavy metal contamination in coal/cinder (mg/kg coal or cinder dry weight) and coal cinder contaminated sites (mg/kg soil dry weight).

Samples	Be	Cr	Ni	Cu	Zn	As	Se	Cd	Pb	U	Hg
Raw coal	0.27	38.18	4.93	6.62	8.55	2.26	0.23	0.05	3.43	0.48	0.006
Rough cinder	0.83	920.82	26.89	22.66	9.01	4.14	0.84	0.12	7.44	1.42	0.001
Background soil	1.33	398.94	30.99	16.22	37.91	6.36	0.87	0.26	21.62	1.20	0.008
0 m											
0-20 cm	1.07	745.15	23.11	17.92	28.78	4.71	0.75	0.25	18.19	1.29	0.003
20-35 cm	1.19	552.97	17.61	10.55	27.72	3.18	0.43	0.22	20.26	0.64	0.005
35-50 cm	1.27	505.60	14.68	10.25	20.97	3.10	0.44	0.16	20.36	0.64	0.003
10 m											
0-20 cm	1.55	802.77	29.38	21.67	23.05	5.26	1.00	0.22	18.05	1.48	0.002
20-35 cm	1.58	620.79	21.71	16.52	29.26	5.60	1.19	0.24	30.31	1.20	0.001
35-50 cm	1.26	525.79	15.80	9.74	17.53	3.00	0.29	0.15	21.44	0.56	0.001
50 m											
0-20 cm	1.16	508.43	13.80	8.93	28.10	3.66	0.36	0.14	20.08	0.54	0.033
20-35 cm	1.12	482.68	12.29	8.09	16.19	2.62	0.41	0.13	19.88	0.45	0.001
35-50 cm	1.12	628.89	15.03	9.35	18.22	3.08	0.46	0.14	20.17	0.53	0.001
80 m											
0-20 cm	1.15	483.48	13.26	9.03	18.61	3.22	0.30	0.16	20.01	0.53	0.001
20-35 cm	1.19	502.34	13.84	9.31	19.76	3.35	0.45	0.18	21.03	0.57	0.001
35-50 cm	1.29	613.57	19.76	15.32	29.44	4.01	0.43	0.25	21.28	0.86	0.001
150 m											
0-20 cm	1.10	494.60	13.56	8.90	19.03	3.24	0.58	0.15	19.99	0.56	0.001
20-35 cm	1.24	474.09	16.46	10.87	24.61	4.48	0.43	0.22	20.51	0.76	0.001
35-50 cm	1.25	448.66	18.50	13.00	28.81	5.08	0.41	0.22	21.70	0.90	0.001

Note: The analytical instrument is Hg analyzer (DMA-80) for Hg and ICP-MS (X series II) for other elements.

Supplementary Material

[Click here to download Supplementary Material: Supplementary Material.docx](#)

Revised manuscript with highlight

[Click here to download Supplementary Material: Manuscript_Highlight.docx](#)

Biographies

Jianli Jia received her PhD degree in environmental science and engineering from Tsinghua University in 2005. Currently, she is an associate professor at the School of Chemical and Environmental Engineering, China University of Mining & Technology (Beijing), China. Her main research area is bioremediation of contaminated land, particularly the heavy metal contaminated sites.

Hanbing Li received his BSc degree in environmental engineering in 2014 from Beijing University of Technology, China. He is currently a PhD research student at Lancaster University, UK. His research topic is the application of magnetic nanoparticles in environmental monitoring and remediation.

Shuang Zong received her BSc degree in environmental engineering in 2014 from China University of Mining & Technology (Beijing), China. She is currently an MSc student at China University of Mining & Technology (Beijing), China. Her research topic is ecological risk assessment at the coal cinder sites.

Bo Jiang received her PhD degree in environmental science and engineering from Tsinghua University in 2015. Her PhD research topic is the construction and application of whole-cell bioreporter in environmental monitoring.

Guanghe Li received his PhD degree in 1992 from China University of Geosciences. He is currently a professor at the School of Environment, Tsinghua University, China. As the senior researcher in contaminated site and groundwater management in China, his specialists are risk assessment and remediation of contaminated land, ecotoxicological characterization of contaminated sites and theories/technologies for water pollution control.

Odafe Ejenavi received his MSc degree in geology in 1999 from University of Benin, Nigeria. He is currently a PhD research student at Lancaster University, UK. His research topic is the impact of crude oil exploration on the fortunes of the Niger Delta and his interests are the application of whole-cell bioreporters for crude oil contamination *in situ* monitoring.

Jingrong Zhu received her MSc degree in environmental science in 2014 from Lancaster University, UK. Her research topic is removal and immobilization of hexavalent chromium contaminated water by magnetic nanoparticles, and her main research interests include magnetic nanoparticles synthesis and functionalization.

Dayi Zhang received his PhD degree in environmental science and engineering from Tsinghua University in 2009. Currently, he is a lecture of environmental science and technology at Lancaster University, UK. His main research interests are whole-cell bioreporter in environmental monitoring, environmental microbiology, synthetic biology, and gene regulation mechanisms.