

CRYSTALLOGRAPHIC  
COMMUNICATIONS

ISSN 2056-9890

Received 25 June 2015

Accepted 10 July 2015

Edited by H. Stoeckli-Evans, University of  
Neuchâtel, Switzerland**Keywords:** crystal structure; piperazinium salts;  
anticancer activity; hydrogen bonding**CCDC references:** 1412124; 1412123**Supporting information:** this article has  
supporting information at [journals.iucr.org/e](http://journals.iucr.org/e)



# Crystal structures of 4-phenylpiperazin-1-ium 6-chloro-5-ethyl-2,4-dioxypyrimidin-1-ide and 4-phenylpiperazin-1-ium 6-chloro-5-isopropyl-2,4- dioxypyrimidin-1-ide

**Monirah A. Al-Alshaikh,<sup>a</sup> Ali A. El-Emam,<sup>b\*</sup> Omar A. Al-Deeb,<sup>b</sup> Mohammed S. M.  
Abdelbaky<sup>c</sup> and Santiago Garcia-Granda<sup>c\*</sup>**<sup>a</sup>Department of Chemistry, College of Sciences, King Saud University, Riyadh 11451, Saudi Arabia, <sup>b</sup>Department of  
Pharmaceutical Chemistry, College of Pharmacy, King Saud University, Riyadh 11451, Saudi Arabia, and <sup>c</sup>Department of  
Physical and Analytical Chemistry, Faculty of Chemistry, Oviedo University-CINN, Oviedo 33006, Spain.\*Correspondence e-mail: [elemam5@hotmail.com](mailto:elemam5@hotmail.com), [sgg@uniovi.es](mailto:sgg@uniovi.es)


The title molecular salts,  $C_{10}H_{15}N_2^+ \cdot C_6H_6ClN_2O_2^-$ , (I), and  $C_{10}H_{15}N_2^+ \cdot C_7H_8ClN_2O_2^-$ , (II), consist of 4-phenylpiperazin-1-ium cations with a 6-chloro-5-ethyl-2,4-dioxypyrimidin-1-ide anion in (I) and a 6-chloro-5-isopropyl-2,4-dioxypyrimidin-1-ide anion in (II). Salt (I) crystallizes with two independent cations and anions in the asymmetric unit. In the crystal structures of both salts, the ions are linked *via* N—H...O and N—H...N hydrogen bonds, forming sheets which are parallel to (100) in (I) and to (001) in (II). In (I), the sheets are linked *via* C—H...Cl hydrogen bonds, forming a three-dimensional framework.

## 1. Chemical context

2,4-Dioxypyrimidine derivatives (uracils) and their related analogues are known for their diverse chemotherapeutic activities including anticancer activity (Ghoshal & Jacob, 1997; Spáčilová *et al.*, 2007; Blokhina *et al.*, 1972), anti-HIV activity (Tanaka *et al.*, 1995; El-Emam *et al.*, 2004) and antibacterial activity (Al-Turkistani *et al.*, 2011). In addition, the piperazine nucleus constitutes the core pharmacophore of several biologically active compounds which display antiviral (Romero *et al.*, 1994, 1996), anticancer (Fytas *et al.*, 2015; Kamal *et al.*, 2015; Arnatt *et al.*, 2014), antitubercular and antibacterial (Nagesh *et al.*, 2014; Peng *et al.*, 2015; Kapić *et al.*, 2011; Wang *et al.*, 2014) and central nervous system activities (Bender *et al.*, 2014; Bali *et al.*, 2010).




(I)




(II)

OPEN  ACCESS



**Figure 1**  
The molecular structure of compound (I), showing the atom labelling. Displacement ellipsoids are drawn at the 50% probability level.



**Figure 2**  
The molecular structure of compound (II), showing the atom labelling. Displacement ellipsoids are drawn at the 50% probability level.

As a result of the relative acidity of 2,4-dioxypyrimidines (Kurinovitch & Lee, 2002; Jang *et al.*, 2001; Nguyen *et al.*, 1998), the title piperazinium salts were isolated as minor byproducts during the reaction of 1-phenylpiperazine with 5-alkyl-6-chlorouracils (Al-Turkistani *et al.*, 2011). In a continuation of our interest in the structures of piperazinium salts (Al-Omary *et al.*, 2014), we report herein on the isolation and crystal structures of these two new piperazinium salts, (I) and (II).

## 2. Structural commentary

The molecular structures of the title salts (I) and (II) are illustrated in Figs. 1 and 2, respectively. Compound (I) crystallizes with two independent 4-phenylpiperazin-1-ium cations (*A* and *B*) and two independent 6-chloro-5-ethyl-2,4-dioxypyrimidin-1-ide anions (*C* and *D*) in the asymmetric unit. In both compounds, the piperazine rings adopt a distorted chair conformation with a positively charged protonated N atom. In compound (I), the mean plane of the piperazine ring makes a

**Table 1**  
Hydrogen-bond geometry (Å, °) for (I).

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N1—H1...O2 <sup>i</sup>	0.86	2.00	2.859 (4)	173
N3—H3A...O1 <sup>ii</sup>	0.89	2.83	3.465 (4)	129
N6—H6A...O4 <sup>iii</sup>	0.89	1.81	2.681 (5)	165
N7—H7...O3 <sup>iv</sup>	0.86	2.02	2.873 (4)	174
N3—H3A...N2 <sup>ii</sup>	0.89	1.92	2.808 (4)	174
N6—H6B...N8 <sup>v</sup>	0.89	1.92	2.798 (5)	169
C10—H10B...O2 <sup>vi</sup>	0.97	2.46	3.355 (5)	154
C26—H26A...O3 <sup>vii</sup>	0.97	2.58	3.444 (5)	147
C16—H16...Cl2 <sup>viii</sup>	0.93	2.80	3.462 (4)	129


Symmetry codes: (i)  $-x+1, -y-1, -z+2$ ; (ii)  $x, y+1, z$ ; (iii)  $x, y-1, z$ ; (iv)  $-x+2, -y+1, -z+2$ ; (v)  $-x+2, y-\frac{1}{2}, -z+\frac{3}{2}$ ; (vi)  $x, -y+\frac{1}{2}, z-\frac{1}{2}$ ; (vii)  $-x+2, -y, -z+2$ ; (viii)  $-x+2, y+\frac{1}{2}, -z+\frac{3}{2}$ .

dihedral angle of  $34.8(2)^\circ$  with the attached phenyl ring in cation *A*, and  $39.7(2)^\circ$  in cation *B*. The equivalent dihedral angle is  $39.61(9)^\circ$  in the cation of compound (II). In the uracil anions, the pyrimidine rings are almost planar with r.m.s. deviations of  $0.008 \text{ \AA}$  in both anions (*C* and *D*) of compound (I), and  $0.024 \text{ \AA}$  in compound (II).

## 3. Supramolecular features

In the crystal of (I), two tetranuclear units are formed, involving cation *A* and anion *C*, and cation *B* and anion *D*, via N—H...O and C—H...O hydrogen bonds. These units are linked via N—H...N hydrogen bonds, forming separate *A/B* and *C/D* sheets parallel to the *bc* plane (Table 1 and Fig. 3). The sheets are linked via C—H...Cl hydrogen bonds, forming a three-dimensional framework (Fig. 3 and Table 1).

In the crystal of (II), the cation and anion are linked by N—H...O and C—H...O hydrogen bonds, forming chains extending along the *b*-axis direction. The chains are linked via



**Figure 3**  
The crystal packing of compound (I), viewed along the *b* axis, showing the most relevant hydrogen bonding (dashed lines; see Table 1).

Table 2

Hydrogen-bond geometry (Å, °) for (II).

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N2—H2N...N4	0.89	1.93	2.813 (2)	174
N2—H3N...O1 <sup>i</sup>	0.89	1.84	2.705 (2)	164
N3—H3...O2 <sup>ii</sup>	0.86	1.98	2.834 (2)	174
C3—H3A...O2 <sup>iii</sup>	0.97	2.54	3.394 (2)	147

Symmetry codes: (i)  $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$ ; (ii)  $-x + \frac{1}{2}, -y + \frac{5}{2}, -z + \frac{1}{2}$ ; (iii)  $x + \frac{1}{2}, -y + 2, z$ .

N—H...N hydrogen bonds, forming sheets lying parallel to the *ac* plane (Table 2 and Fig. 4).

#### 4. Database survey

A search of the Cambridge Structural Database (Version 5.36, last update November 2014; Groom & Allen, 2014) for the anion 6-chloro-5-ethyl-2,4-dioxypyrimidin-1-ide, present in compound (I), gave no hits, while for the anion 6-chloro-5-isopropyl-2,4-dioxypyrimidin-1-ide, present in compound (II), one hit was obtained, with the cation 4-(2-methoxyphenyl)piperazin-1-ium (Al-Omary *et al.*, 2014).

#### 5. Synthesis and crystallization

**Compound (I):** A mixture of 6-chloro-5-ethyluracil (349 mg, 2.0 mmol), 1-phenylpiperazine (325 mg, 2.0 mmol) and anhydrous potassium carbonate (276 mg, 2.0 mmol), in ethanol




Figure 4

The crystal packing of compound (II), viewed along the *b* axis, showing the most relevant hydrogen bonding (dashed lines; see Table 2).

(8 ml), was heated under reflux for 6 h. On cooling, the precipitate, thus formed was separated by filtration to yield 306 mg (51%) of 5-ethyl-6-(4-phenyl-1-piperazinyl)uracil. The filtrate was concentrated by vacuum distillation to 5 ml and allowed to stand at room temperature overnight to yield compound (I) as colourless crystals (m.p.: 459–461 K). <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>, 500.13 MHz): δ 0.93 (*t*, 3H, CH<sub>3</sub>, *J* = 7.0 Hz), 2.35 (*q*, 2H, CH<sub>2</sub>), 3.25 (*s*, 4H, piperazine-H), 3.45 (*s*, 4H, piperazine-H), 6.83–6.95 (*m*, 3H, Ar—H), 7.21 (*d*, 2H, Ar—H, *J* = 6.6 Hz), 8.15–8.17 (*m*, 2H, NH<sub>2</sub>), 10.83 (*s*, 1H, NH).

Table 3

Experimental details.

	(I)	(II)
Crystal data		
Chemical formula	C <sub>10</sub> H <sub>15</sub> N <sub>2</sub> <sup>+</sup> ·C <sub>6</sub> H <sub>6</sub> ClN <sub>2</sub> O <sub>2</sub> <sup>−</sup>	C <sub>10</sub> H <sub>15</sub> N <sub>2</sub> <sup>+</sup> ·C <sub>7</sub> H <sub>8</sub> ClN <sub>2</sub> O <sub>2</sub> <sup>−</sup>
<i>M<sub>r</sub></i>	336.82	350.84
Crystal system, space group	Monoclinic, <i>P</i> 2 <sub>1</sub> / <i>c</i>	Monoclinic, <i>I</i> 2/ <i>a</i>
Temperature (K)	293	101
<i>a</i> , <i>b</i> , <i>c</i> (Å)	21.676 (1), 7.6446 (5), 20.5444 (8)	20.5012 (3), 7.4565 (1), 23.1414 (3)
β (°)	95.065 (5)	90.639 (1)
<i>V</i> (Å <sup>3</sup> )	3391.0 (3)	3537.34 (8)
<i>Z</i>	8	8
Radiation type	Cu Kα	Cu Kα
μ (mm <sup>−1</sup> )	2.12	2.05
Crystal size (mm)	0.17 × 0.08 × 0.06	0.34 × 0.13 × 0.09
Data collection		
Diffractometer	Agilent Xcalibur Ruby Gemini	Agilent Xcalibur Ruby Gemini
Absorption correction	Multi-scan ( <i>CrysAlis PRO</i> ; Agilent, 2014)	Multi-scan ( <i>CrysAlis PRO</i> ; Agilent, 2014)
<i>T<sub>min</sub></i> , <i>T<sub>max</sub></i>	0.809, 0.880	0.760, 0.828
No. of measured, independent and observed [ <i>I</i> > 2σ( <i>I</i> )] reflections	32461, 6532, 3596	13174, 3396, 2926
<i>R<sub>int</sub></i>	0.135	0.069
(sin θ/λ) <sub>max</sub> (Å <sup>−1</sup> )	0.612	0.612
Refinement		
<i>R</i> [ <i>F</i> <sup>2</sup> > 2σ( <i>F</i> <sup>2</sup> )], <i>wR</i> ( <i>F</i> <sup>2</sup> ), <i>S</i>	0.066, 0.185, 1.01	0.044, 0.122, 1.03
No. of reflections	6457	3346
No. of parameters	415	217
H-atom treatment	H-atom parameters constrained	H-atom parameters constrained
Δρ <sub>max</sub> , Δρ <sub>min</sub> (e Å <sup>−3</sup> )	0.42, −0.36	0.55, −0.56

Computer programs: *CrysAlis PRO* (Agilent, 2014), *SIR2011* (Burla *et al.*, 2012), *Mercury* (Macrae *et al.*, 2008), *SHELXL2014* (Sheldrick, 2015), *PLATON* (Spek, 2009) and *publCIF* (Westrip, 2010).

$^{13}\text{C}$  NMR (DMSO- $d_6$ , 125.76 MHz):  $\delta$  13.80 (CH<sub>3</sub>), 19.55 (CH<sub>2</sub>), 44.18, 47.86 (piperazine-C), 116.32, 119.62, 128.44, 150.70 (Ar-C), 108.88, 153.90, 155.94, 164.80 (pyrimidine-C),

**Compound (II):** 6-Chloro-5-isopropyluracil (377 mg, 2.0 mmol), 1-phenylpiperazine (325 mg, 2.0 mmol) and anhydrous potassium carbonate (276 mg, 2.0 mmol), in ethanol (8 ml), was heated under reflux for 6 h. On cooling, the precipitate thus formed was separated by filtration to yield 566 mg (90%) of 5-isopropyl-6-(4-phenyl-1-piperazinyl)uracil. The filtrate was concentrated by vacuum distillation to 5 ml and allowed to stand at room temperature overnight to yield compound (II) as colourless crystals (m.p.: 473–475 K).  $^1\text{H}$  NMR (DMSO- $d_6$ , 500.13 MHz):  $\delta$  1.20 (*d*, 6H, CH<sub>3</sub>, *J* = 7.8 Hz), 2.52–2.56 (*m*, 1H, CH), 3.18 (*s*, 4H, piperazine-H), 3.24 (*s*, 4H, piperazine-H), 6.88–7.02 (*m*, 3H, Ar-H), 7.20–7.22 (*m*, 2H, Ar-H), 8.04–8.08 (*m*, 2H, NH<sub>2</sub>), 11.02 (*s*, 1H, NH).  $^{13}\text{C}$  NMR (DMSO- $d_6$ , 125.76 MHz):  $\delta$  19.98 (CH<sub>3</sub>), 27.0 (CH), 44.50, 47.98 (piperazine-C), 116.16, 119.80, 129.04, 150.0 (Ar-C), 110.82, 152.30, 154.04, 164.06 (pyrimidine-C).

## 6. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 3. The H atoms were included in calculated positions and treated as riding atoms: N–H = 0.86–0.90 Å, C–H = 0.95–1.00 Å with  $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$  for methyl H atoms and  $1.2U_{\text{eq}}(\text{N,C})$  for other H atoms.

## Acknowledgements

The authors are grateful to the Deanship of Scientific Research at King Saud University for funding this study through the research group project No. PRG-1436-23. We also acknowledge financial support from the Spanish Ministerio de Economía y Competitividad (MINECO-13-MAT2013-40950-R, FPI grant BES-2011-046948 to MSMA).

## References

- Agilent (2014). *CrysAlis PRO*. Agilent Technologies UK Ltd, Yarnton, England.
- Al-Omary, F. A. M., Ghabbour, H. A., El-Emam, A. A., Chidan Kumar, C. S. & Fun, H.-K. (2014). *Acta Cryst.* **E70**, o245–o246.
- Al-Turkistani, A. A., Al-Deeb, O. A., El-Brollosy, N. R. & El-Emam, A. A. (2011). *Molecules*, **16**, 4764–4774.
- Arnatt, C. K., Adams, J. L., Zhang, Z., Haney, K. M., Li, G. & Zhang, Y. (2014). *Bioorg. Med. Chem. Lett.* **24**, 2319–2323.
- Bali, A., Sharma, K., Bhalla, A., Bala, S., Reddy, D., Singh, A. & Kumar, A. (2010). *Eur. J. Med. Chem.* **45**, 2656–2662.

- Bender, A. M., Clark, M. J., Agius, M. P., Traynor, J. R. & Mosberg, H. I. (2014). *Bioorg. Med. Chem. Lett.* **24**, 548–551.
- Blokhina, N. G., Vozny, E. K. & Garin, A. M. (1972). *Cancer*, **30**, 390–392.
- Burla, M. C., Caliandro, R., Camalli, M., Carrozzini, B., Cascarano, G. L., Giacovazzo, C., Mallamo, M., Mazzzone, A., Polidori, G. & Spagna, R. (2012). *J. Appl. Cryst.* **45**, 357–361.
- El-Emam, A. A., Massoud, M. A., El-Bendary, E. R. & El-Sayed, M. A. (2004). *Bull. Korean Chem. Soc.* **25**, 991–996.
- Fytas, C., Zoidis, G., Tsotinis, A., Fytas, G., Khan, M. A., Akhtar, S., Rahman, K. M. & Thurston, D. E. (2015). *Eur. J. Med. Chem.* **93**, 281–290.
- Ghoshal, K. & Jacob, S. T. (1997). *Biochem. Pharmacol.* **53**, 1569–1575.
- Groom, C. R. & Allen, F. H. (2014). *Angew. Chem. Int. Ed.* **53**, 662–671.
- Jang, Y. H., Sowers, L. C., Çağın, T. & Goddard, W. A. III (2001). *J. Phys. Chem. A*, **105**, 274–280.
- Kamal, A., Sreekanth, K., Shankaraiah, N., Sathish, M., Nekkanti, S. & Srinivasulu, V. (2015). *Bioorg. Chem.* **59**, 23–30.
- Kapić, S., Paljetak, H. Č., Jakopović, I. P., Fajdetić, A., Ilijaš, M., Štimac, V., Brajša, K., Holmes, D. J., Berge, J. & Alihodžić, S. (2011). *Bioorg. Med. Chem.* **19**, 7281–7298.
- Kurinovich, M. A. & Lee, J. K. (2002). *J. Am. Soc. Mass Spectrom.* **13**, 985–995.
- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
- Nagesh, H. N., Suresh, A., Sairam, S. D. S. S., Sriram, D., Yogeewari, P. & Chandra Sekhar, K. V. G. (2014). *Eur. J. Med. Chem.* **84**, 605–613.
- Nguyen, M. T., Chandra, A. K. & Zeegers-Huyskens, T. (1998). *J. Chem. Soc. Faraday Trans.* **94**, 1277–1280.
- Peng, C.-T., Gao, C., Wang, N.-Y., You, X.-Y., Zhang, L.-D., Zhu, Y.-X., Xu, Y., Zuo, W.-Q., Ran, K., Deng, H.-X., Lei, Q., Xiao, K.-J. & Yu, L.-T. (2015). *Bioorg. Med. Chem. Lett.* **25**, 1373–1376.
- Romero, D. L., Morge, R. A., Biles, C., Berrios-Pena, N., May, P. D., Palmer, J. R., Johnson, P. D., Smith, H. W., Busso, M., Tan, C.-K., Voorman, R. L., Reusser, F., Althaus, I. W., So, A. G., Resnick, L., Tarpley, W. G. & Aristoff, P. A. (1994). *J. Med. Chem.* **37**, 999–1014.
- Romero, D. L., Olmsted, R. A., Poel, T. J., Morge, R. A., Biles, C., Keiser, B. J., Kopta, L. A., Friis, J. M., Hosley, J. D., Stefanski, K. J., Wishka, D. G., Evans, D. B., Morris, J., Stehle, R. G., Sharma, S. K., Yagi, Y., Voorman, R. L., Adams, W. J., Tarpley, W. G. & Thomas, R. C. (1996). *J. Med. Chem.* **39**, 3769–3789.
- Sheldrick, G. M. (2015). *Acta Cryst.* **C71**, 3–8.
- Spáčilová, L., Džubák, P., Hajdúch, M., Krupková, S., Hradil, P. & Hlaváč, J. (2007). *Bioorg. Med. Chem. Lett.* **17**, 6647–6650.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Tanaka, H., Takashima, H., Ubasawa, M., Sekiya, K., Inouye, N., Baba, M., Shigeta, S., Walker, R. T., De Clercq, E. & Miyasaka, T. (1995). *J. Med. Chem.* **38**, 2860–2865.
- Wang, S.-F., Yin, Y., Qiao, F., Wu, X., Sha, S., Zhang, L. & Zhu, H.-L. (2014). *Bioorg. Med. Chem.* **22**, 2409–2415.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

## supporting information

*Acta Cryst.* (2015). E71, 956-959 [https://doi.org/10.1107/S2056989015013298]

## Crystal structures of 4-phenylpiperazin-1-ium 6-chloro-5-ethyl-2,4-dioxypyrimidin-1-ide and 4-phenylpiperazin-1-ium 6-chloro-5-isopropyl-2,4-dioxypyrimidin-1-ide

**Monirah A. Al-Alshaikh, Ali A. El-Emam, Omar A. Al-Deeb, Mohammed S. M. Abdelbaky and Santiago Garcia-Granda**

### Computing details

For both compounds, data collection: *CrysAlis PRO* (Agilent, 2014); cell refinement: *CrysAlis PRO* (Agilent, 2014); data reduction: *CrysAlis PRO* (Agilent, 2014); program(s) used to solve structure: *SIR2011* (Burla *et al.*, 2012); program(s) used to refine structure: *SHELXL2014* (Sheldrick, 2015); molecular graphics: *Mercury* (Macrae *et al.*, 2008); software used to prepare material for publication: *SHELXL2014* (Sheldrick, 2015), *PLATON* (Spek, 2009) and *pubCIF* (Westrip, 2010).

### (I) 4-Phenylpiperazin-1-ium 6-chloro-5-ethyl-2,4-dioxypyrimidin-1-ide

#### Crystal data

$C_{10}H_{15}N_2^+ \cdot C_6H_6ClN_2O_2^-$

$M_r = 336.82$

Monoclinic,  $P2_1/c$

$a = 21.676$  (1) Å

$b = 7.6446$  (5) Å

$c = 20.5444$  (8) Å

$\beta = 95.065$  (5)°

$V = 3391.0$  (3) Å<sup>3</sup>

$Z = 8$

$F(000) = 1424$

$D_x = 1.319$  Mg m<sup>-3</sup>

Cu  $K\alpha$  radiation,  $\lambda = 1.54184$  Å

Cell parameters from 2418 reflections

$\theta = 4.1\text{--}70.3^\circ$

$\mu = 2.12$  mm<sup>-1</sup>

$T = 293$  K

Prism, colourless

$0.17 \times 0.08 \times 0.06$  mm

#### Data collection

Agilent Xcalibur Ruby Gemini  
diffractometer

Radiation source: Enhance (Cu) X-ray Source

Detector resolution: 10.2673 pixels mm<sup>-1</sup>

$\omega$  scans

Absorption correction: multi-scan  
(*CrysAlis PRO*; Agilent, 2014)

$T_{\min} = 0.809$ ,  $T_{\max} = 0.880$

32461 measured reflections

6532 independent reflections

3596 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.135$

$\theta_{\max} = 70.7^\circ$ ,  $\theta_{\min} = 4.1^\circ$

$h = -26 \rightarrow 26$

$k = -9 \rightarrow 9$

$l = -21 \rightarrow 25$

#### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.066$

$wR(F^2) = 0.185$

$S = 1.01$

6457 reflections

415 parameters

0 restraints

Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier map  
 Hydrogen site location: inferred from neighbouring sites  
 H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0563P)^2]$$

where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = 0.001$   
 $\Delta\rho_{\max} = 0.42 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.36 \text{ e } \text{\AA}^{-3}$

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl2	1.13235 (5)	0.04012 (14)	0.82294 (5)	0.0408 (3)
Cl1	0.65035 (5)	-0.09582 (15)	0.82219 (5)	0.0437 (3)
O1	0.51594 (14)	-0.6011 (4)	0.82514 (13)	0.0381 (7)
O2	0.55114 (13)	-0.3192 (4)	1.02030 (13)	0.0344 (7)
O3	1.04166 (14)	0.2995 (4)	1.02077 (13)	0.0378 (7)
N1	0.53525 (15)	-0.4583 (4)	0.92238 (15)	0.0288 (8)
H1	0.5119	-0.5322	0.9401	0.035*
O4	1.02351 (14)	0.5927 (4)	0.82725 (14)	0.0415 (8)
N7	1.03409 (15)	0.4421 (4)	0.92325 (15)	0.0306 (8)
H7	1.0140	0.5236	0.9410	0.037*
N2	0.57984 (15)	-0.3662 (4)	0.82841 (15)	0.0301 (8)
N8	1.07502 (16)	0.3338 (5)	0.82954 (16)	0.0330 (8)
N3	0.56236 (15)	0.6236 (4)	0.69138 (16)	0.0320 (8)
H3A	0.5695	0.6194	0.7347	0.038*
H3B	0.5338	0.7053	0.6817	0.038*
N4	0.64335 (15)	0.3661 (5)	0.64533 (16)	0.0330 (8)
N5	0.86014 (15)	0.0992 (4)	0.85634 (16)	0.0325 (8)
N6	0.93121 (15)	-0.1748 (4)	0.80703 (16)	0.0337 (8)
H6A	0.9565	-0.2655	0.8153	0.040*
H6B	0.9245	-0.1637	0.7639	0.040*
C1	0.54296 (19)	-0.4798 (5)	0.85679 (19)	0.0313 (9)
C18	1.05419 (19)	0.3021 (5)	0.9626 (2)	0.0328 (10)
C17	1.04364 (19)	0.4618 (6)	0.85811 (19)	0.0313 (9)
C4	0.60638 (18)	-0.2374 (5)	0.8663 (2)	0.0315 (9)
C2	0.56213 (18)	-0.3272 (5)	0.9617 (2)	0.0308 (9)
C20	1.09503 (18)	0.1988 (5)	0.8674 (2)	0.0309 (9)
C11	0.68606 (18)	0.2301 (5)	0.6354 (2)	0.0311 (9)
C9	0.6682 (2)	0.5311 (5)	0.6731 (2)	0.0345 (10)
H9A	0.7051	0.5634	0.6524	0.041*
H9B	0.6796	0.5166	0.7195	0.041*
C27	0.82314 (18)	0.2433 (5)	0.8709 (2)	0.0315 (9)
C3	0.60166 (18)	-0.2074 (5)	0.93082 (19)	0.0301 (9)
C25	0.82936 (19)	-0.0520 (5)	0.8253 (2)	0.0341 (10)
H25A	0.8201	-0.0296	0.7789	0.041*

---

H25B	0.7907	-0.0736	0.8443	0.041*
C19	1.08842 (19)	0.1695 (5)	0.93181 (19)	0.0316 (9)
C23	0.96153 (19)	-0.0141 (6)	0.8346 (2)	0.0357 (10)
H23A	0.9982	0.0104	0.8123	0.043*
H23B	0.9742	-0.0317	0.8806	0.043*
C21	1.1136 (2)	0.0116 (6)	0.9694 (2)	0.0377 (10)
H21A	1.1044	-0.0924	0.9433	0.045*
H21B	1.0928	0.0007	1.0091	0.045*
C28	0.7616 (2)	0.2637 (6)	0.8454 (2)	0.0397 (11)
H28	0.7448	0.1857	0.8139	0.048*
C10	0.62039 (19)	0.6734 (5)	0.6625 (2)	0.0336 (10)
H10A	0.6365	0.7809	0.6825	0.040*
H10B	0.6116	0.6941	0.6161	0.040*
C5	0.6318 (2)	-0.0569 (5)	0.9687 (2)	0.0359 (10)
H5A	0.6721	-0.0353	0.9533	0.043*
H5B	0.6381	-0.0893	1.0145	0.043*
C7	0.53832 (19)	0.4524 (6)	0.6672 (2)	0.0378 (10)
H7A	0.5248	0.4610	0.6211	0.045*
H7B	0.5029	0.4196	0.6903	0.045*
C16	0.74788 (19)	0.2396 (6)	0.6587 (2)	0.0357 (10)
H16	0.7618	0.3334	0.6849	0.043*
C15	0.7893 (2)	0.1105 (6)	0.6436 (2)	0.0412 (11)
H15	0.8308	0.1192	0.6592	0.049*
C26	0.87103 (19)	-0.2105 (6)	0.8348 (2)	0.0371 (10)
H26A	0.8784	-0.2368	0.8811	0.044*
H26B	0.8511	-0.3110	0.8132	0.044*
C12	0.6660 (2)	0.0859 (6)	0.5972 (2)	0.0369 (10)
H12	0.6246	0.0765	0.5812	0.044*
C8	0.58805 (18)	0.3138 (5)	0.6776 (2)	0.0344 (10)
H8A	0.5991	0.2979	0.7240	0.041*
H8B	0.5724	0.2033	0.6597	0.041*
C32	0.8473 (2)	0.3663 (5)	0.9167 (2)	0.0374 (10)
H32	0.8884	0.3573	0.9337	0.045*
C24	0.91766 (19)	0.1391 (6)	0.8266 (2)	0.0362 (10)
H24A	0.9370	0.2417	0.8473	0.043*
H24B	0.9083	0.1645	0.7805	0.043*
C29	0.7254 (2)	0.3984 (7)	0.8662 (2)	0.0480 (12)
H29	0.6843	0.4089	0.8493	0.058*
C31	0.8113 (2)	0.4999 (6)	0.9369 (2)	0.0442 (12)
H31	0.8282	0.5796	0.9677	0.053*
C13	0.7077 (2)	-0.0424 (6)	0.5832 (2)	0.0423 (11)
H13	0.6938	-0.1384	0.5582	0.051*
C22	1.1830 (2)	0.0208 (6)	0.9874 (2)	0.0433 (11)
H22A	1.1963	-0.0826	1.0111	0.065*
H22B	1.2040	0.0288	0.9483	0.065*
H22C	1.1924	0.1219	1.0141	0.065*
C14	0.7692 (2)	-0.0311 (6)	0.6054 (2)	0.0450 (12)
H14	0.7970	-0.1173	0.5948	0.054*

C30	0.7503 (2)	0.5173 (6)	0.9120 (2)	0.0487 (13)
H30	0.7261	0.6082	0.9259	0.058*
C6	0.5943 (2)	0.1094 (6)	0.9627 (2)	0.0529 (13)
H6C	0.6158	0.2002	0.9878	0.079*
H6D	0.5887	0.1440	0.9177	0.079*
H6E	0.5546	0.0899	0.9788	0.079*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Cl2	0.0470 (6)	0.0405 (6)	0.0359 (6)	0.0097 (5)	0.0099 (5)	-0.0029 (5)
Cl1	0.0503 (7)	0.0474 (7)	0.0347 (6)	-0.0174 (5)	0.0101 (5)	0.0010 (5)
O1	0.0488 (18)	0.0360 (17)	0.0301 (16)	-0.0110 (14)	0.0063 (14)	-0.0051 (13)
O2	0.0396 (17)	0.0410 (17)	0.0226 (15)	-0.0075 (13)	0.0030 (13)	0.0008 (12)
O3	0.0446 (18)	0.0400 (18)	0.0299 (16)	0.0085 (14)	0.0087 (14)	0.0022 (13)
N1	0.0337 (18)	0.0299 (18)	0.0231 (17)	-0.0075 (15)	0.0036 (14)	0.0018 (14)
O4	0.0462 (19)	0.0444 (19)	0.0349 (17)	0.0130 (15)	0.0087 (14)	0.0079 (14)
N7	0.0323 (18)	0.033 (2)	0.0266 (18)	0.0058 (15)	0.0057 (15)	-0.0034 (15)
N2	0.0301 (18)	0.036 (2)	0.0244 (17)	-0.0008 (15)	0.0033 (14)	0.0004 (15)
N8	0.0331 (19)	0.036 (2)	0.0305 (19)	0.0011 (16)	0.0053 (15)	-0.0007 (16)
N3	0.0349 (19)	0.0336 (19)	0.0280 (18)	0.0064 (15)	0.0052 (15)	-0.0001 (15)
N4	0.0304 (19)	0.033 (2)	0.0355 (19)	-0.0054 (15)	0.0046 (15)	-0.0023 (16)
N5	0.0297 (18)	0.0307 (19)	0.038 (2)	0.0003 (15)	0.0089 (16)	-0.0023 (16)
N6	0.038 (2)	0.035 (2)	0.0287 (19)	0.0004 (16)	0.0042 (16)	-0.0002 (15)
C1	0.032 (2)	0.033 (2)	0.028 (2)	-0.0028 (18)	0.0043 (18)	0.0009 (18)
C18	0.035 (2)	0.032 (2)	0.032 (2)	-0.0007 (18)	0.0079 (19)	0.0012 (18)
C17	0.032 (2)	0.035 (2)	0.028 (2)	0.0040 (18)	0.0048 (18)	0.0007 (18)
C4	0.027 (2)	0.035 (2)	0.033 (2)	-0.0055 (17)	0.0018 (18)	0.0043 (18)
C2	0.029 (2)	0.031 (2)	0.032 (2)	0.0002 (17)	0.0011 (18)	0.0027 (18)
C20	0.028 (2)	0.033 (2)	0.032 (2)	0.0001 (18)	0.0054 (18)	-0.0055 (18)
C11	0.029 (2)	0.033 (2)	0.032 (2)	0.0018 (18)	0.0075 (18)	0.0018 (18)
C9	0.039 (2)	0.029 (2)	0.036 (2)	-0.0052 (19)	0.009 (2)	-0.0006 (19)
C27	0.031 (2)	0.033 (2)	0.031 (2)	-0.0006 (18)	0.0070 (18)	0.0058 (18)
C3	0.031 (2)	0.033 (2)	0.026 (2)	-0.0006 (17)	-0.0008 (17)	0.0012 (17)
C25	0.032 (2)	0.038 (2)	0.032 (2)	-0.0057 (19)	0.0044 (18)	0.0012 (19)
C19	0.034 (2)	0.032 (2)	0.029 (2)	0.0009 (18)	0.0051 (18)	0.0000 (18)
C23	0.031 (2)	0.039 (3)	0.038 (2)	-0.0023 (19)	0.0063 (19)	-0.006 (2)
C21	0.042 (3)	0.038 (2)	0.035 (2)	0.006 (2)	0.010 (2)	0.004 (2)
C28	0.039 (2)	0.044 (3)	0.037 (3)	0.004 (2)	0.006 (2)	-0.001 (2)
C10	0.041 (2)	0.033 (2)	0.027 (2)	0.0013 (19)	0.0050 (19)	0.0011 (18)
C5	0.041 (2)	0.039 (3)	0.028 (2)	-0.009 (2)	0.0027 (19)	0.0014 (19)
C7	0.031 (2)	0.044 (3)	0.038 (2)	0.004 (2)	0.0009 (19)	-0.003 (2)
C16	0.034 (2)	0.038 (2)	0.036 (2)	0.0002 (19)	0.0054 (19)	-0.0014 (19)
C15	0.036 (2)	0.045 (3)	0.043 (3)	0.006 (2)	0.008 (2)	0.004 (2)
C26	0.038 (2)	0.039 (3)	0.034 (2)	-0.006 (2)	0.007 (2)	-0.0022 (19)
C12	0.040 (2)	0.037 (2)	0.034 (2)	0.000 (2)	0.0024 (19)	0.0026 (19)
C8	0.035 (2)	0.031 (2)	0.038 (2)	-0.0043 (19)	0.0045 (19)	-0.0033 (19)
C32	0.039 (2)	0.033 (2)	0.041 (3)	-0.0047 (19)	0.011 (2)	-0.002 (2)



C24	0.032 (2)	0.041 (3)	0.037 (2)	-0.0063 (19)	0.0111 (19)	-0.002 (2)
C29	0.044 (3)	0.055 (3)	0.045 (3)	0.015 (2)	0.006 (2)	0.005 (2)
C31	0.055 (3)	0.037 (3)	0.043 (3)	-0.001 (2)	0.017 (2)	-0.003 (2)
C13	0.058 (3)	0.034 (2)	0.037 (3)	-0.004 (2)	0.014 (2)	-0.006 (2)
C22	0.042 (3)	0.046 (3)	0.043 (3)	0.005 (2)	0.007 (2)	0.009 (2)
C14	0.053 (3)	0.037 (3)	0.047 (3)	0.010 (2)	0.017 (2)	0.002 (2)
C30	0.056 (3)	0.045 (3)	0.047 (3)	0.017 (2)	0.019 (3)	0.005 (2)
C6	0.065 (3)	0.040 (3)	0.051 (3)	0.002 (2)	-0.008 (3)	-0.010 (2)

*Geometric parameters (Å, °)*

C12—C20	1.758 (4)	C19—C21	1.509 (6)
C11—C4	1.747 (4)	C23—C24	1.508 (6)
O1—C1	1.249 (5)	C23—H23A	0.9700
O2—C2	1.248 (5)	C23—H23B	0.9700
O3—C18	1.249 (5)	C21—C22	1.517 (6)
N1—C1	1.382 (5)	C21—H21A	0.9700
N1—C2	1.384 (5)	C21—H21B	0.9700
N1—H1	0.8600	C28—C29	1.386 (6)
O4—C17	1.243 (5)	C28—H28	0.9300
N7—C17	1.380 (5)	C10—H10A	0.9700
N7—C18	1.388 (5)	C10—H10B	0.9700
N7—H7	0.8600	C5—C6	1.508 (6)
N2—C1	1.347 (5)	C5—H5A	0.9700
N2—C4	1.351 (5)	C5—H5B	0.9700
N8—C20	1.342 (5)	C7—C8	1.513 (6)
N8—C17	1.354 (5)	C7—H7A	0.9700
N3—C7	1.478 (5)	C7—H7B	0.9700
N3—C10	1.487 (5)	C16—C15	1.388 (6)
N3—H3A	0.8900	C16—H16	0.9300
N3—H3B	0.8900	C15—C14	1.385 (7)
N4—C11	1.419 (5)	C15—H15	0.9300
N4—C9	1.467 (5)	C26—H26A	0.9700
N4—C8	1.475 (5)	C26—H26B	0.9700
N5—C27	1.410 (5)	C12—C13	1.380 (6)
N5—C25	1.454 (5)	C12—H12	0.9300
N5—C24	1.468 (5)	C8—H8A	0.9700
N6—C23	1.482 (5)	C8—H8B	0.9700
N6—C26	1.495 (5)	C32—C31	1.371 (6)
N6—H6A	0.8900	C32—H32	0.9300
N6—H6B	0.8900	C24—H24A	0.9700
C18—C19	1.436 (6)	C24—H24B	0.9700
C4—C3	1.359 (6)	C29—C30	1.383 (7)
C2—C3	1.440 (6)	C29—H29	0.9300
C20—C19	1.362 (6)	C31—C30	1.383 (7)
C11—C16	1.385 (6)	C31—H31	0.9300
C11—C12	1.400 (6)	C13—C14	1.374 (7)
C9—C10	1.505 (6)	C13—H13	0.9300

C9—H9A	0.9700	C22—H22A	0.9600
C9—H9B	0.9700	C22—H22B	0.9600
C27—C28	1.398 (6)	C22—H22C	0.9600
C27—C32	1.399 (6)	C14—H14	0.9300
C3—C5	1.505 (6)	C30—H30	0.9300
C25—C26	1.514 (6)	C6—H6C	0.9600
C25—H25A	0.9700	C6—H6D	0.9600
C25—H25B	0.9700	C6—H6E	0.9600
C1—N1—C2	125.1 (3)	H21A—C21—H21B	107.8
C1—N1—H1	117.5	C29—C28—C27	121.0 (5)
C2—N1—H1	117.5	C29—C28—H28	119.5
C17—N7—C18	125.6 (3)	C27—C28—H28	119.5
C17—N7—H7	117.2	N3—C10—C9	110.7 (3)
C18—N7—H7	117.2	N3—C10—H10A	109.5
C1—N2—C4	117.3 (3)	C9—C10—H10A	109.5
C20—N8—C17	117.0 (3)	N3—C10—H10B	109.5
C7—N3—C10	112.2 (3)	C9—C10—H10B	109.5
C7—N3—H3A	109.2	H10A—C10—H10B	108.1
C10—N3—H3A	109.2	C3—C5—C6	113.3 (4)
C7—N3—H3B	109.2	C3—C5—H5A	108.9
C10—N3—H3B	109.2	C6—C5—H5A	108.9
H3A—N3—H3B	107.9	C3—C5—H5B	108.9
C11—N4—C9	117.6 (3)	C6—C5—H5B	108.9
C11—N4—C8	115.6 (3)	H5A—C5—H5B	107.7
C9—N4—C8	110.1 (3)	N3—C7—C8	110.3 (3)
C27—N5—C25	117.8 (3)	N3—C7—H7A	109.6
C27—N5—C24	116.5 (3)	C8—C7—H7A	109.6
C25—N5—C24	110.9 (3)	N3—C7—H7B	109.6
C23—N6—C26	112.2 (3)	C8—C7—H7B	109.6
C23—N6—H6A	109.2	H7A—C7—H7B	108.1
C26—N6—H6A	109.2	C11—C16—C15	120.8 (4)
C23—N6—H6B	109.2	C11—C16—H16	119.6
C26—N6—H6B	109.2	C15—C16—H16	119.6
H6A—N6—H6B	107.9	C14—C15—C16	120.3 (4)
O1—C1—N2	121.5 (4)	C14—C15—H15	119.8
O1—C1—N1	120.3 (4)	C16—C15—H15	119.8
N2—C1—N1	118.2 (4)	N6—C26—C25	109.6 (3)
O3—C18—N7	119.1 (4)	N6—C26—H26A	109.8
O3—C18—C19	125.1 (4)	C25—C26—H26A	109.8
N7—C18—C19	115.8 (3)	N6—C26—H26B	109.8
O4—C17—N8	121.8 (4)	C25—C26—H26B	109.8
O4—C17—N7	120.4 (4)	H26A—C26—H26B	108.2
N8—C17—N7	117.8 (4)	C13—C12—C11	120.0 (4)
N2—C4—C3	128.3 (4)	C13—C12—H12	120.0
N2—C4—C11	112.1 (3)	C11—C12—H12	120.0
C3—C4—C11	119.5 (3)	N4—C8—C7	110.1 (3)
O2—C2—N1	119.5 (4)	N4—C8—H8A	109.6

O2—C2—C3	124.5 (4)	C7—C8—H8A	109.6
N1—C2—C3	116.1 (3)	N4—C8—H8B	109.6
N8—C20—C19	129.3 (4)	C7—C8—H8B	109.6
N8—C20—C12	111.7 (3)	H8A—C8—H8B	108.2
C19—C20—C12	119.0 (3)	C31—C32—C27	121.1 (4)
C16—C11—C12	118.5 (4)	C31—C32—H32	119.5
C16—C11—N4	122.4 (4)	C27—C32—H32	119.5
C12—C11—N4	118.9 (4)	N5—C24—C23	110.1 (3)
N4—C9—C10	109.9 (4)	N5—C24—H24A	109.6
N4—C9—H9A	109.7	C23—C24—H24A	109.6
C10—C9—H9A	109.7	N5—C24—H24B	109.6
N4—C9—H9B	109.7	C23—C24—H24B	109.6
C10—C9—H9B	109.7	H24A—C24—H24B	108.1
H9A—C9—H9B	108.2	C30—C29—C28	120.1 (5)
C28—C27—C32	117.7 (4)	C30—C29—H29	120.0
C28—C27—N5	123.4 (4)	C28—C29—H29	120.0
C32—C27—N5	118.8 (4)	C32—C31—C30	120.7 (5)
C4—C3—C2	115.0 (4)	C32—C31—H31	119.6
C4—C3—C5	124.6 (4)	C30—C31—H31	119.6
C2—C3—C5	120.4 (3)	C14—C13—C12	121.3 (4)
N5—C25—C26	109.5 (4)	C14—C13—H13	119.3
N5—C25—H25A	109.8	C12—C13—H13	119.3
C26—C25—H25A	109.8	C21—C22—H22A	109.5
N5—C25—H25B	109.8	C21—C22—H22B	109.5
C26—C25—H25B	109.8	H22A—C22—H22B	109.5
H25A—C25—H25B	108.2	C21—C22—H22C	109.5
C20—C19—C18	114.5 (4)	H22A—C22—H22C	109.5
C20—C19—C21	124.4 (4)	H22B—C22—H22C	109.5
C18—C19—C21	121.1 (4)	C13—C14—C15	119.0 (4)
N6—C23—C24	110.4 (4)	C13—C14—H14	120.5
N6—C23—H23A	109.6	C15—C14—H14	120.5
C24—C23—H23A	109.6	C29—C30—C31	119.4 (4)
N6—C23—H23B	109.6	C29—C30—H30	120.3
C24—C23—H23B	109.6	C31—C30—H30	120.3
H23A—C23—H23B	108.1	C5—C6—H6C	109.5
C19—C21—C22	113.2 (4)	C5—C6—H6D	109.5
C19—C21—H21A	108.9	H6C—C6—H6D	109.5
C22—C21—H21A	108.9	C5—C6—H6E	109.5
C19—C21—H21B	108.9	H6C—C6—H6E	109.5
C22—C21—H21B	108.9	H6D—C6—H6E	109.5
C4—N2—C1—O1	179.0 (4)	N8—C20—C19—C21	-179.6 (4)
C4—N2—C1—N1	-0.6 (6)	C12—C20—C19—C21	2.8 (6)
C2—N1—C1—O1	-178.9 (4)	O3—C18—C19—C20	178.5 (4)
C2—N1—C1—N2	0.7 (6)	N7—C18—C19—C20	-1.4 (6)
C17—N7—C18—O3	-179.1 (4)	O3—C18—C19—C21	-0.9 (7)
C17—N7—C18—C19	0.9 (6)	N7—C18—C19—C21	179.2 (4)
C20—N8—C17—O4	179.6 (4)	C26—N6—C23—C24	-53.9 (5)

C20—N8—C17—N7	-0.8 (6)	C20—C19—C21—C22	76.2 (6)
C18—N7—C17—O4	179.9 (4)	C18—C19—C21—C22	-104.4 (5)
C18—N7—C17—N8	0.3 (6)	C32—C27—C28—C29	-1.8 (6)
C1—N2—C4—C3	1.1 (7)	N5—C27—C28—C29	173.5 (4)
C1—N2—C4—C11	-177.7 (3)	C7—N3—C10—C9	-54.1 (4)
C1—N1—C2—O2	179.0 (4)	N4—C9—C10—N3	56.9 (4)
C1—N1—C2—C3	-1.0 (6)	C4—C3—C5—C6	84.3 (5)
C17—N8—C20—C19	0.1 (7)	C2—C3—C5—C6	-92.5 (5)
C17—N8—C20—C12	177.9 (3)	C10—N3—C7—C8	53.7 (4)
C9—N4—C11—C16	8.8 (6)	C12—C11—C16—C15	1.2 (6)
C8—N4—C11—C16	-124.1 (4)	N4—C11—C16—C15	-174.7 (4)
C9—N4—C11—C12	-167.1 (4)	C11—C16—C15—C14	-0.7 (7)
C8—N4—C11—C12	60.0 (5)	C23—N6—C26—C25	54.9 (5)
C11—N4—C9—C10	164.1 (3)	N5—C25—C26—N6	-58.0 (4)
C8—N4—C9—C10	-60.6 (4)	C16—C11—C12—C13	-0.4 (6)
C25—N5—C27—C28	-13.7 (6)	N4—C11—C12—C13	175.6 (4)
C24—N5—C27—C28	121.7 (4)	C11—N4—C8—C7	-163.1 (4)
C25—N5—C27—C32	161.5 (4)	C9—N4—C8—C7	60.6 (4)
C24—N5—C27—C32	-63.1 (5)	N3—C7—C8—N4	-56.5 (4)
N2—C4—C3—C2	-1.4 (7)	C28—C27—C32—C31	1.5 (6)
C11—C4—C3—C2	177.3 (3)	N5—C27—C32—C31	-174.0 (4)
N2—C4—C3—C5	-178.3 (4)	C27—N5—C24—C23	161.1 (4)
C11—C4—C3—C5	0.4 (6)	C25—N5—C24—C23	-60.5 (5)
O2—C2—C3—C4	-178.8 (4)	N6—C23—C24—N5	55.5 (4)
N1—C2—C3—C4	1.2 (6)	C27—C28—C29—C30	1.2 (7)
O2—C2—C3—C5	-1.7 (6)	C27—C32—C31—C30	-0.6 (7)
N1—C2—C3—C5	178.3 (4)	C11—C12—C13—C14	-0.8 (7)
C27—N5—C25—C26	-160.5 (3)	C12—C13—C14—C15	1.3 (7)
C24—N5—C25—C26	61.7 (4)	C16—C15—C14—C13	-0.5 (7)
N8—C20—C19—C18	1.1 (7)	C28—C29—C30—C31	-0.3 (7)
C12—C20—C19—C18	-176.5 (3)	C32—C31—C30—C29	0.0 (7)

## Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N1—H1...O2 <sup>i</sup>	0.86	2.00	2.859 (4)	173
N3—H3 <i>A</i> ...O1 <sup>ii</sup>	0.89	2.83	3.465 (4)	129
N6—H6 <i>A</i> ...O4 <sup>iii</sup>	0.89	1.81	2.681 (5)	165
N7—H7...O3 <sup>iv</sup>	0.86	2.02	2.873 (4)	174
N3—H3 <i>A</i> ...N2 <sup>ii</sup>	0.89	1.92	2.808 (4)	174
N6—H6 <i>B</i> ...N8 <sup>v</sup>	0.89	1.92	2.798 (5)	169
C10—H10 <i>B</i> ...O2 <sup>vi</sup>	0.97	2.46	3.355 (5)	154
C26—H26 <i>A</i> ...O3 <sup>vii</sup>	0.97	2.58	3.444 (5)	147
C16—H16...C12 <sup>viii</sup>	0.93	2.80	3.462 (4)	129

Symmetry codes: (i)  $-x+1, -y-1, -z+2$ ; (ii)  $x, y+1, z$ ; (iii)  $x, y-1, z$ ; (iv)  $-x+2, -y+1, -z+2$ ; (v)  $-x+2, y-1/2, -z+3/2$ ; (vi)  $x, -y+1/2, z-1/2$ ; (vii)  $-x+2, -y, -z+2$ ; (viii)  $-x+2, y+1/2, -z+3/2$ .

## (II) 4-Phenylpiperazin-1-ium 6-chloro-5-isopropyl-2,4-dioxypyrimidin-1-ide

## Crystal data

 $C_{10}H_{15}N_2^+ \cdot C_7H_8ClN_2O_2^-$  $M_r = 350.84$ Monoclinic,  $I2/a$  $a = 20.5012$  (3) Å $b = 7.4565$  (1) Å $c = 23.1414$  (3) Å $\beta = 90.639$  (1)° $V = 3537.34$  (8) Å<sup>3</sup> $Z = 8$  $F(000) = 1488$  $D_x = 1.318$  Mg m<sup>-3</sup>Cu  $K\alpha$  radiation,  $\lambda = 1.5418$  Å

Cell parameters from 6927 reflections

 $\theta = 3.8$ – $70.0$ ° $\mu = 2.05$  mm<sup>-1</sup> $T = 101$  K

Prism, colourless

 $0.34 \times 0.13 \times 0.09$  mm

## Data collection

Agilent Xcalibur Ruby Gemini  
diffractometerDetector resolution: 10.2673 pixels mm<sup>-1</sup> $\omega$  scansAbsorption correction: multi-scan  
(CrysAlis PRO; Agilent, 2014) $T_{\min} = 0.760$ ,  $T_{\max} = 0.828$ 

13174 measured reflections

3396 independent reflections

2926 reflections with  $I > 2\sigma(I)$  $R_{\text{int}} = 0.069$  $\theta_{\max} = 70.6$ °,  $\theta_{\min} = 3.8$ ° $h = -24 \rightarrow 25$  $k = -9 \rightarrow 7$  $l = -27 \rightarrow 28$ 

## Refinement

Refinement on  $F^2$ 

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.044$  $wR(F^2) = 0.122$  $S = 1.03$ 

3346 reflections

217 parameters

0 restraints

Hydrogen site location: inferred from  
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0585P)^2 + 3.6877P]$ where  $P = (F_o^2 + 2F_c^2)/3$  $(\Delta/\sigma)_{\max} < 0.001$  $\Delta\rho_{\max} = 0.55$  e Å<sup>-3</sup> $\Delta\rho_{\min} = -0.56$  e Å<sup>-3</sup>

## Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å<sup>2</sup>)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.43825 (2)	0.80800 (7)	0.38729 (2)	0.03420 (17)
O1	0.42855 (6)	1.32956 (18)	0.26398 (6)	0.0286 (3)
O2	0.23184 (6)	1.07264 (19)	0.29852 (6)	0.0288 (3)
N3	0.33106 (7)	1.1946 (2)	0.28125 (6)	0.0227 (3)
H3	0.3124	1.2711	0.2587	0.027*
N4	0.42771 (7)	1.0889 (2)	0.32365 (6)	0.0228 (3)
N2	0.56397 (7)	1.1049 (2)	0.31180 (6)	0.0246 (3)
H3N	0.5734	1.0238	0.2849	0.029*
H2N	0.5208	1.1087	0.3152	0.029*
C11	0.39765 (9)	1.2092 (2)	0.28906 (7)	0.0228 (4)

N1	0.60877 (7)	1.3622 (2)	0.39443 (6)	0.0230 (3)
C2	0.58171 (9)	1.1911 (2)	0.41332 (8)	0.0237 (4)
H2A	0.5352	1.2036	0.4195	0.028*
H2B	0.6020	1.1556	0.4496	0.028*
C4	0.58796 (10)	1.2834 (3)	0.29310 (8)	0.0271 (4)
H4A	0.5649	1.3201	0.2582	0.032*
H4B	0.6341	1.2760	0.2845	0.032*
C14	0.39071 (9)	0.9583 (2)	0.34646 (7)	0.0236 (4)
C12	0.29194 (9)	1.0666 (3)	0.30679 (7)	0.0232 (4)
C6	0.58544 (9)	1.4868 (3)	0.49109 (8)	0.0273 (4)
H6	0.5581	1.3906	0.4992	0.033*
C13	0.32496 (9)	0.9324 (2)	0.34113 (7)	0.0231 (4)
C3	0.59378 (9)	1.0490 (3)	0.36799 (8)	0.0245 (4)
H3A	0.6403	1.0318	0.3634	0.029*
H3B	0.5750	0.9361	0.3802	0.029*
C10	0.65511 (9)	1.6458 (3)	0.42609 (8)	0.0272 (4)
H10	0.6749	1.6571	0.3903	0.033*
C15	0.28407 (10)	0.7802 (3)	0.36508 (8)	0.0286 (4)
H15	0.2393	0.8063	0.3523	0.034*
C5	0.61534 (9)	1.4980 (2)	0.43717 (8)	0.0224 (4)
C8	0.63624 (10)	1.7627 (3)	0.52142 (9)	0.0341 (5)
H8	0.6436	1.8500	0.5494	0.041*
C1	0.57729 (9)	1.4207 (3)	0.34016 (7)	0.0249 (4)
H1A	0.5953	1.5352	0.3284	0.030*
H1B	0.5309	1.4365	0.3462	0.030*
C7	0.59623 (10)	1.6182 (3)	0.53262 (8)	0.0332 (5)
H7	0.5763	1.6088	0.5684	0.040*
C9	0.66531 (10)	1.7757 (3)	0.46790 (9)	0.0320 (5)
H9	0.6921	1.8731	0.4599	0.038*
C16	0.28114 (10)	0.7739 (3)	0.43025 (8)	0.0310 (4)
H16A	0.2708	0.8909	0.4448	0.047*
H16B	0.2481	0.6902	0.4418	0.047*
H16C	0.3227	0.7365	0.4456	0.047*
C17	0.30027 (12)	0.6004 (3)	0.33815 (9)	0.0394 (5)
H17A	0.3015	0.6125	0.2969	0.059*
H17B	0.3421	0.5605	0.3522	0.059*
H17C	0.2675	0.5143	0.3483	0.059*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C11	0.0307 (3)	0.0306 (3)	0.0413 (3)	0.00185 (19)	0.0001 (2)	0.0141 (2)
O1	0.0225 (7)	0.0278 (7)	0.0355 (7)	-0.0043 (6)	0.0020 (5)	0.0107 (6)
O2	0.0210 (6)	0.0341 (8)	0.0313 (7)	-0.0042 (6)	0.0017 (5)	0.0053 (6)
N3	0.0210 (7)	0.0233 (8)	0.0238 (7)	-0.0013 (6)	0.0012 (6)	0.0032 (6)
N4	0.0233 (7)	0.0211 (8)	0.0239 (7)	0.0003 (6)	0.0022 (6)	0.0025 (6)
N2	0.0227 (7)	0.0256 (8)	0.0255 (7)	0.0009 (7)	0.0023 (6)	-0.0055 (6)
C11	0.0238 (9)	0.0216 (9)	0.0231 (8)	-0.0010 (7)	0.0033 (7)	-0.0021 (7)

N1	0.0261 (8)	0.0203 (8)	0.0225 (7)	0.0005 (6)	0.0007 (6)	0.0002 (6)
C2	0.0254 (9)	0.0221 (9)	0.0235 (8)	0.0009 (7)	0.0020 (7)	0.0010 (7)
C4	0.0302 (10)	0.0284 (10)	0.0226 (9)	-0.0025 (8)	0.0023 (7)	-0.0003 (8)
C14	0.0295 (9)	0.0207 (9)	0.0206 (8)	0.0023 (8)	0.0024 (7)	-0.0001 (7)
C12	0.0255 (9)	0.0244 (9)	0.0198 (8)	-0.0036 (8)	0.0043 (7)	-0.0030 (7)
C6	0.0300 (10)	0.0248 (10)	0.0273 (9)	-0.0002 (8)	0.0003 (8)	-0.0004 (8)
C13	0.0287 (9)	0.0209 (9)	0.0198 (8)	-0.0031 (8)	0.0042 (7)	-0.0022 (7)
C3	0.0231 (9)	0.0221 (9)	0.0284 (9)	0.0017 (7)	0.0017 (7)	-0.0005 (7)
C10	0.0241 (9)	0.0254 (10)	0.0322 (10)	0.0019 (8)	0.0017 (7)	-0.0005 (8)
C15	0.0317 (10)	0.0262 (10)	0.0279 (9)	-0.0059 (8)	0.0039 (8)	0.0007 (8)
C5	0.0201 (8)	0.0211 (9)	0.0260 (9)	0.0053 (7)	-0.0028 (7)	-0.0015 (7)
C8	0.0376 (11)	0.0290 (11)	0.0354 (10)	0.0050 (9)	-0.0084 (9)	-0.0106 (9)
C1	0.0284 (9)	0.0236 (9)	0.0227 (8)	0.0002 (8)	0.0006 (7)	0.0016 (7)
C7	0.0385 (11)	0.0350 (11)	0.0260 (9)	0.0066 (9)	0.0012 (8)	-0.0053 (8)
C9	0.0273 (10)	0.0241 (10)	0.0445 (11)	-0.0005 (8)	-0.0056 (8)	-0.0034 (9)
C16	0.0318 (10)	0.0306 (11)	0.0309 (10)	-0.0049 (9)	0.0092 (8)	0.0003 (8)
C17	0.0492 (13)	0.0299 (11)	0.0396 (11)	-0.0130 (10)	0.0139 (10)	-0.0048 (9)

*Geometric parameters (Å, °)*

C11—C14	1.7544 (18)	C6—C7	1.389 (3)
O1—C11	1.246 (2)	C6—C5	1.399 (3)
O2—C12	1.246 (2)	C6—H6	0.9300
N3—C11	1.379 (2)	C13—C15	1.519 (3)
N3—C12	1.383 (2)	C3—H3A	0.9700
N3—H3	0.8600	C3—H3B	0.9700
N4—N4	0.000 (5)	C10—C9	1.383 (3)
N4—C14	1.346 (2)	C10—C5	1.397 (3)
N4—C11	1.347 (2)	C10—H10	0.9300
N2—C4	1.485 (2)	C15—C16	1.511 (3)
N2—C3	1.490 (2)	C15—C17	1.517 (3)
N2—H3N	0.8900	C15—H15	0.9800
N2—H2N	0.8900	C8—C7	1.380 (3)
C11—N4	1.347 (2)	C8—C9	1.384 (3)
N1—C5	1.421 (2)	C8—H8	0.9300
N1—C2	1.460 (2)	C1—H1A	0.9700
N1—C1	1.472 (2)	C1—H1B	0.9700
C2—C3	1.513 (2)	C7—H7	0.9300
C2—H2A	0.9700	C9—H9	0.9300
C2—H2B	0.9700	C16—H16A	0.9600
C4—C1	1.512 (3)	C16—H16B	0.9600
C4—H4A	0.9700	C16—H16C	0.9600
C4—H4B	0.9700	C17—H17A	0.9600
C14—N4	1.346 (2)	C17—H17B	0.9600
C14—C13	1.366 (3)	C17—H17C	0.9600
C12—C13	1.442 (3)		
C11—N3—C12	125.17 (16)	C12—C13—C15	117.51 (16)

C11—N3—H3	117.4	N2—C3—C2	109.92 (15)
C12—N3—H3	117.4	N2—C3—H3A	109.7
N4—N4—C14	0 (10)	C2—C3—H3A	109.7
N4—N4—C11	0 (10)	N2—C3—H3B	109.7
C14—N4—C11	117.31 (15)	C2—C3—H3B	109.7
C4—N2—C3	111.77 (14)	H3A—C3—H3B	108.2
C4—N2—H3N	109.3	C9—C10—C5	120.54 (18)
C3—N2—H3N	109.3	C9—C10—H10	119.7
C4—N2—H2N	109.3	C5—C10—H10	119.7
C3—N2—H2N	109.3	C16—C15—C17	113.20 (18)
H3N—N2—H2N	107.9	C16—C15—C13	114.57 (16)
O1—C11—N4	121.67 (16)	C17—C15—C13	112.72 (16)
O1—C11—N4	121.67 (16)	C16—C15—H15	105.1
N4—C11—N4	0.00 (14)	C17—C15—H15	105.1
O1—C11—N3	120.23 (17)	C13—C15—H15	105.1
N4—C11—N3	118.10 (16)	C10—C5—C6	118.26 (17)
N4—C11—N3	118.10 (16)	C10—C5—N1	119.06 (16)
C5—N1—C2	116.59 (14)	C6—C5—N1	122.64 (17)
C5—N1—C1	114.83 (15)	C7—C8—C9	119.04 (19)
C2—N1—C1	110.50 (14)	C7—C8—H8	120.5
N1—C2—C3	109.78 (14)	C9—C8—H8	120.5
N1—C2—H2A	109.7	N1—C1—C4	110.37 (15)
C3—C2—H2A	109.7	N1—C1—H1A	109.6
N1—C2—H2B	109.7	C4—C1—H1A	109.6
C3—C2—H2B	109.7	N1—C1—H1B	109.6
H2A—C2—H2B	108.2	C4—C1—H1B	109.6
N2—C4—C1	110.25 (15)	H1A—C1—H1B	108.1
N2—C4—H4A	109.6	C8—C7—C6	120.75 (19)
C1—C4—H4A	109.6	C8—C7—H7	119.6
N2—C4—H4B	109.6	C6—C7—H7	119.6
C1—C4—H4B	109.6	C10—C9—C8	120.9 (2)
H4A—C4—H4B	108.1	C10—C9—H9	119.5
N4—C14—N4	0.00 (17)	C8—C9—H9	119.5
N4—C14—C13	128.81 (17)	C15—C16—H16A	109.5
N4—C14—C13	128.81 (17)	C15—C16—H16B	109.5
N4—C14—C11	111.18 (13)	H16A—C16—H16B	109.5
N4—C14—C11	111.18 (13)	C15—C16—H16C	109.5
C13—C14—C11	120.01 (14)	H16A—C16—H16C	109.5
O2—C12—N3	119.11 (17)	H16B—C16—H16C	109.5
O2—C12—C13	124.56 (17)	C15—C17—H17A	109.5
N3—C12—C13	116.32 (16)	C15—C17—H17B	109.5
C7—C6—C5	120.48 (19)	H17A—C17—H17B	109.5
C7—C6—H6	119.8	C15—C17—H17C	109.5
C5—C6—H6	119.8	H17A—C17—H17C	109.5
C14—C13—C12	114.11 (16)	H17B—C17—H17C	109.5
C14—C13—C15	128.36 (17)		
N4—N4—C11—O1	0.0 (5)	N3—C12—C13—C14	4.0 (2)



C14—N4—C11—O1	-176.64 (16)	O2—C12—C13—C15	4.8 (3)
C14—N4—C11—N4	0 (100)	N3—C12—C13—C15	-174.41 (15)
N4—N4—C11—N3	0.0 (4)	C4—N2—C3—C2	55.49 (19)
C14—N4—C11—N3	2.5 (2)	N1—C2—C3—N2	-58.11 (19)
C12—N3—C11—O1	179.96 (16)	C14—C13—C15—C16	65.7 (3)
C12—N3—C11—N4	0.8 (3)	C12—C13—C15—C16	-116.20 (19)
C12—N3—C11—N4	0.8 (3)	C14—C13—C15—C17	-65.7 (3)
C5—N1—C2—C3	-165.61 (15)	C12—C13—C15—C17	112.4 (2)
C1—N1—C2—C3	60.88 (19)	C9—C10—C5—C6	1.1 (3)
C3—N2—C4—C1	-54.5 (2)	C9—C10—C5—N1	-176.58 (17)
C11—N4—C14—N4	0 (100)	C7—C6—C5—C10	-1.1 (3)
N4—N4—C14—C13	0.0 (3)	C7—C6—C5—N1	176.42 (17)
C11—N4—C14—C13	-2.5 (3)	C2—N1—C5—C10	164.11 (16)
N4—N4—C14—C11	0.0 (3)	C1—N1—C5—C10	-64.4 (2)
C11—N4—C14—C11	177.09 (13)	C2—N1—C5—C6	-13.4 (2)
C11—N3—C12—O2	176.56 (16)	C1—N1—C5—C6	118.11 (19)
C11—N3—C12—C13	-4.2 (3)	C5—N1—C1—C4	165.63 (15)
N4—C14—C13—C12	-0.9 (3)	C2—N1—C1—C4	-59.98 (19)
N4—C14—C13—C12	-0.9 (3)	N2—C4—C1—N1	56.0 (2)
C11—C14—C13—C12	179.55 (12)	C9—C8—C7—C6	0.4 (3)
N4—C14—C13—C15	177.27 (17)	C5—C6—C7—C8	0.4 (3)
N4—C14—C13—C15	177.27 (17)	C5—C10—C9—C8	-0.3 (3)
C11—C14—C13—C15	-2.3 (3)	C7—C8—C9—C10	-0.5 (3)
O2—C12—C13—C14	-176.82 (17)		

Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N2—H2N $\cdots$ N4	0.89	1.93	2.813 (2)	174
N2—H3N $\cdots$ O1 <sup>i</sup>	0.89	1.84	2.705 (2)	164
N3—H3 $\cdots$ O2 <sup>ii</sup>	0.86	1.98	2.834 (2)	174
C3—H3A $\cdots$ O2 <sup>iii</sup>	0.97	2.54	3.394 (2)	147

Symmetry codes: (i)  $-x+1, y-1/2, -z+1/2$ ; (ii)  $-x+1/2, -y+5/2, -z+1/2$ ; (iii)  $x+1/2, -y+2, z$ .