



**Universidade de São Paulo**

**Biblioteca Digital da Produção Intelectual - BDPI**

---

Departamento de Física e Ciências Materiais - IFSC/FCM

Artigos e Materiais de Revistas Científicas - IFSC/FCM

---

2014-11

# Stochastic processes with ZN symmetry and complex Virasoro representations: the partition functions

---

Journal of Physics A, Bristol : Institute of Physics - IOP, v. 47, n. 46, p. 462001-1-462001-7, Nov. 2014

<http://www.producao.usp.br/handle/BDPI/50364>

*Downloaded from: Biblioteca Digital da Produção Intelectual - BDPI, Universidade de São Paulo*

Stochastic processes with  $Z_N$  symmetry and complex Virasoro representations. The partition functions

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2014 J. Phys. A: Math. Theor. 47 462001

(<http://iopscience.iop.org/1751-8121/47/46/462001>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 143.107.180.191

This content was downloaded on 09/12/2014 at 20:21

Please note that [terms and conditions apply](#).

## Fast Track Communication

# Stochastic processes with $Z_N$ symmetry and complex Virasoro representations. The partition functions

Francisco C Alcaraz<sup>1</sup>, Pavel Pyatov<sup>2</sup> and Vladimir Rittenberg<sup>3</sup>

<sup>1</sup> Universidade de São Paulo, Instituto de Física de São Carlos, Caixa Postal 369, 13560-590 São Carlos, São Paulo, Brazil

<sup>2</sup> National Research University Higher School of Economics, Laboratory of Mathematical Physics, 20 Myasnitskaya street, Moscow 101000, Russia & Bogoliubov Laboratory of Theoretical Physics, Joint Institute for Nuclear Research, 141980 Dubna, Moscow Region, Russia

<sup>3</sup> Physikalisches Institut, Universität Bonn, Nussallee 12, D-53115 Bonn, Germany

E-mail: [alcaraz@ifsc.usp.br](mailto:alcaraz@ifsc.usp.br), [pyatov@theor.jinr.ru](mailto:pyatov@theor.jinr.ru) and [vladimir@th.physik.uni-bonn.de](mailto:vladimir@th.physik.uni-bonn.de)

Received 11 July 2014, revised 16 September 2014

Accepted for publication 10 October 2014

Published 30 October 2014

## Abstract


In a previous letter (Alcaraz F C *et al* 2014 *J. Phys. A: Math. Theor.* **47** 212003) we have presented numerical evidence that a Hamiltonian expressed in terms of the generators of the periodic Temperley–Lieb algebra has, in the finite-size scaling limit, a spectrum given by representations of the Virasoro algebra with complex highest weights. This Hamiltonian defines a stochastic process with a  $Z_N$  symmetry. We give here analytical expressions for the partition functions for this system which confirm the numerics. For  $N$  even, the Hamiltonian has a symmetry which makes the spectrum doubly degenerate leading to two independent stochastic processes. The existence of a complex spectrum leads to an oscillating approach to the stationary state. This phenomenon is illustrated by an example.

Keywords: stochastic process, Virasoro algebra, Temperley–Lieb algebras

PACS numbers: 03.65.Bz, 03.67.-a, 05.20.-y, 05.30.-d

(Some figures may appear in colour only in the online journal)

Considering  $Z_N$  symmetric representations of the periodic Temperley–Lieb algebra  $PTL_L(x)$  [2–5], we have defined a Hamiltonian as a linear combination of the generators of this



**Figure 1.** The six link patterns configurations for  $L = 4$  sites on a cylinder and two circles without sites (noncontractible loops). The open arches and circles meet behind the cylinder. The corresponding spin presentations of the same link patterns are given on the right.

algebra. Taking  $x = 1$ , this Hamiltonian gives the time evolution of a one-dimensional stochastic process. Looking at the finite-size scaling spectra of this Hamiltonian, we have obtained numerical evidence for the appearance of Virasoro representations with complex highest weights. Moreover, the real part of the complex highest weights is smaller than the real highest weights and hence dominate the large time behavior of the systems. This observation was a big surprise and was the main content of a previous letter [1]. In the present one, we give an analytic derivation of this result and present the partition function for each sector of the model. We also present an application of our results. For  $N$  even we show that there is a symmetry in the model which makes the spectrum for any lattice size to be doubly degenerate indicating the presence of a zero fermionic mode. This letter is basically a continuation of the previous one [1], we did nevertheless our best to make it self-consistent.

The  $PTL_L(x)$  algebra has  $L$  generators  $e_k$ ,  $k = 1, 2, \dots, L$  satisfying the relations

$$e_k^2 = x e_k, \quad e_k e_{k \pm 1} e_k = e_k, \quad [e_k, e_\ell] = 0, \quad |k - \ell| > 1, \quad (1)$$

with  $e_{k+L} = e_k$ . We take  $L$  even only. We consider two quotients of the algebra

$$(AB)^N A = A, \quad (2)$$

where


$$A = \prod_{j=1}^{L/2} e_{2j}, \quad B = \prod_{j=0}^{L/2-1} e_{2j+1}, \quad (3)$$

and

$$ABA = \alpha^2 A \quad (4)$$

with  $\alpha = e^{i2\pi r/N}$ ,  $r = 0, 1, \dots, N - 1$ . One can see that the quotient (4) is a solution of equation (2) which defines the first quotient. This observation will be crucial in obtaining the partition functions mentioned above.

In order to get the  $Z_N$  symmetric representations of (1) and (2), we consider  $N$  copies of a one-dimensional periodic system with  $L$  sites. Each copy consists of  $\binom{L}{L/2}$  configurations of link patterns on a cylinder and  $n$  noncontractible loops ( $n = 0, 1, \dots, N - 1$ ) on the same



**Figure 2.** The action of the  $e_2$  generator acting the bond between the sites 2 and 3 which are the end of an arch of the size of the system  $L = 4$ . A new circle is created on the cylinder and one moves from the copy  $n = 2$  to the copy  $n = 3$ .

cylinder. This is the vector space in which the generators of the  $PTL_L(x)$  algebra act. It has the dimension  $N \times \binom{L}{L/2}$ . In figure 1 we show the six configurations for  $L = 4$  and  $n = 2$ .

An alternative way to label the states in the vector space is to use the spin representation in which the slopes in the arches are used  $+(-)$  for the beginning (ending) of an arch. The number of non-contractible loops is indicated by a supplementary label. This notation is also given in figure 1.

The action of the generators  $e_k$  on the link patterns for a given copy  $n$  is the same as the one used for the usual (non-periodic) Temperley–Lieb algebra [6] with one exception. If the generator acts on the bond connecting the beginning and the end of an arch having the size of the system  $L$ , one obtains a configuration of the copy  $n + 1$  (see figure 2).

In order to obtain the  $Z_N$  representation of the  $PTL_L(x)$  algebra with the quotient (2) one identifies the copy  $n = N$  with the copy  $n = 0$ .

In what follows, we take the parameter  $x$  in the  $PTL_L(x)$  algebra equal to one. With this choice the Hamiltonian

$$H = \sum_{k=1}^L (1 - e_k) \tag{5}$$

gives the time evolution of a stochastic process. We want to stress that the properties of the spectra which are going to be discussed below, stay valid for any value of  $x$ .

Notice that for  $N$  even the  $Z_N$  symmetric representation decomposes into a pair of identical  $\frac{N}{2} \times \binom{L}{L/2}$ -dimensional irreps<sup>4</sup>. To see this, consider two linear transformations on the  $Z_N$  space.

The transformation  $X_1$  acts diagonally in a following way. We start considering configurations with no arches hidden in the back of the cylinder (the first and fifth link patterns in figure 1). These configurations get a factor of  $(-1)^n$ . The configurations translated with one lattice unit get a factor  $(-1)^{(n+1)}$  (the second and sixth configurations in figure 1). The next translated configurations get again the factor  $(-1)^n$  and so on and so forth.

Transformation  $X_2$  permutes copies as follows

$$|\dots\rangle^{(2k)} \leftrightarrow |\dots\rangle^{(2k+1)}, \quad k = 0, 1, \dots, \frac{N}{2} - 1.$$

<sup>4</sup> For  $N$  odd the  $Z_N$  representation is irreducible.

The two transformations constitute the algebra

$$(X_1)^2 = (X_2)^2 = \text{id}, \quad X_1X_2 + X_2X_1 = 0. \quad (6)$$

This algebra has only two-dimensional equivalent irreducible representations. Since  $X_1$  and  $X_2$  commute with the action of the  $PTL_L(x)$  on the  $Z_N$  symmetric link patterns, it follows that for  $N$  even, the spectrum of  $H$  is doubly degenerate. To illustrate this observation, let us take  $L = 4$  and  $N = 2$ . The Hamiltonian splits into two stochastic Hamiltonians having each six states. The first one having the states  $|++--\rangle^{(0)}$ ,  $|--++\rangle^{(0)}$ ,  $|+-+ -\rangle^{(0)}$ ,  $| - ++ -\rangle^{(1)}$ ,  $|+ - - +\rangle^{(1)}$  and  $| - + - +\rangle^{(1)}$ , the second one having the six states in which the copies 0 and 1 are permuted.

We are interested in the spectra of  $H$  in the finite-size scaling limit. Let us keep in mind that in a stochastic process, the energies coincide with the energy gaps since the ground-state energy is zero for any system size. Since  $H$  is invariant under translations ( $e_k \mapsto e_{k+1(\text{mod } L)}$ ) and the cyclic rotations  $Z_N (|\dots\rangle^{(n)} \mapsto |\dots\rangle^{(n+1(\text{mod } N))})$ , one has  $N \times L$  sectors labeled by  $p = 0, \pm 1, \pm 2, \dots$  corresponding to the momenta  $P = 2\pi p/L$  and by  $r = 0, 1, \dots, N - 1$ , labeling the irreps of  $Z_N$ . If  $E_p^r(q)$ ,  $q = 1, 2, \dots$ , are the energy levels in the sector  $(p, r)$ , the scaling dimensions  $x_p^r(q)$  are given by  $\lim_{L \rightarrow \infty} (E_p^r(q)L) = 2\pi v_s x_p^r(q)$ , with the sound velocity  $v_s = 3\sqrt{3}/2$ .

In [1] we have diagonalized numerically  $H$  for  $N = 3$ . We went up to  $L = 30$  and looked at the lowest excitations. In the  $r = 0$  sector we confirmed the expected value  $x_0^0(1) = 0.25$ . The surprise came when we looked at the  $r = 1$  sector where we found

$$x_0^1(1) = 0.039\,05 + 0.087\,53i, \quad x_0^1(2) = 0.149\,08 - 0.118\,06i, \quad (7)$$

i.e. complex values. For  $N$  even we found  $E_p^r(q) = E_{p \pm N/2}^r(q)$  which is a consequence of the symmetry (6).

We present now our new results. In order to obtain the partition function in each sector  $r$ , we use the fact that the  $Z_N$  representation of the algebra with the quotient (2) can be decomposed into  $N$  representations of the quotient (4) [3]. The representations of the quotient (4), in the link patterns vector space, are obtained by considering a single copy but changing the action of the generators when they act on a bond connecting the beginning and the end of an arch of the system size  $L$  like in figure 2. Instead of adding a non-contractible loop, one multiplies the state in the right hand side of the figure by a fugacity  $\alpha = \exp(2\pi i r/N)$ . It was shown [2] that the quotient (4) admits also a representation in the standard spin 1/2 basis (not to be confused with the one used in figure 1) and the Hamiltonian (5) can be written in this basis. By performing a similarity transformation, the Hamiltonian is the XXZ quantum chain with a twist

$$e_k = \sigma_k^+ \sigma_{k+1}^- + \sigma_k^- \sigma_{k+1}^+ + \frac{1}{4}(1 - \sigma_k^z \sigma_{k+1}^z) + i\frac{\sqrt{3}}{4}(\sigma_{k+1}^z - \sigma_k^z), \quad k = 1, 2, \dots, L - 1,$$


$$e_L = e^{i2\pi\phi} \sigma_L^+ \sigma_1^- + e^{-i2\pi\phi} \sigma_L^- \sigma_1^+ + \frac{1}{4}(1 - \sigma_L^z \sigma_1^z) + i\frac{\sqrt{3}}{4}(\sigma_1^z - \sigma_L^z). \quad (8)$$

The twist  $\phi$  is related to the parameter  $\alpha$  by the relation

$$\alpha = 2 \cos(\pi\phi). \quad (9)$$

The vector space of the  $\binom{L}{L/2}$  link patterns configurations corresponds to the  $S^z = \sum_{k=1}^L \sigma_k^z = 0$  sector of the spin vector space.

The Hamiltonian (5), (8) is integrable using the Bethe Ansatz and the scaling dimensions (highest weights of Virasoro representations) are known [7]. They are given by the Gaussian



**Figure 3.** The density of ‘peaks’  $d^{(n)}(z)$  in the copy  $n$  as a function of  $z = 2\pi v_3 t/L$  for  $N = 3$ , and various lattice sizes  $L = 40, 80, 160, 500, 1000, 2000$  and  $4000$ .  $d_0$ , the value of the density in the stationary state, is subtracted.

model. In the  $S^z = 0$  sector they are

$$x = \frac{3}{4}(s + \phi)^2 - \frac{1}{12} + m + m', \quad p = m - m', \quad (10)$$

where  $s, m, m' = 0, \pm 1, \pm 2, \dots$

From (9) we see that for  $N$  even  $\alpha = e^{i2\pi r/N}$  and  $\alpha = e^{i2\pi(r+N/2)/N} = -e^{i2\pi r/N}$  give the same value for the twist  $\phi$  and therefore the spectrum of the Hamiltonian (5) is doubly degenerate, in agreement with our previous observation. Moreover, for the sectors  $r$  not equal to 0 or  $N/2$  the values of  $\phi$  obtained from (9) are complex, henceforth the scaling dimensions (critical exponents) (10) are complex too.

As a check, taking  $N = 3$  and  $r = 1$ , from (9) one gets

$$\phi = -0.426\ 642 - 0.137\ 279i$$


from which we get using (10) with  $s = 0$  and 1

$$x_0^1(1) = 0.039\ 904\ 99 + 0.087\ 853\ 992i; \quad x_0^1(2) = 0.149\ 0874 - 0.118\ 081\ 36i \quad (11)$$

in excellent agreement with the values (7).

We have to stress that although the spectrum of the  $Z_N$  symmetric Hamiltonian splits into  $N$  sectors, the stochastic process doesn't. The condition of positivity of the wave function describing the probability distribution function, mixes the sectors.

The existence of complex scaling dimensions has consequences on the time behavior of various correlators showing oscillatory phenomena, more so since their real part is smaller than real scaling dimensions of the  $r = 0$  sector. To illustrate the phenomenon, we looked at the density of ‘peaks’  $d^{(n)}(t, L)$  in different copies. Those are  $+-$  pairs which in the Dyck paths picture of the link patterns [8] correspond to peaks in the paths. This local observable is measured easily in Monte Carlo simulations. The time dependence of the ‘peaks’ in various sectors is determined by the initial conditions. For large values of  $t$  and large lattice sizes we expect  $d^{(n)}(t, L)$  to be a function of  $t/L$



**Figure 4.** A fit to  $d^{(0)}(z) - d_0$  as a function of  $z$ , using the parameterization (13) and (14). Monte Carlo data for lattice sizes 40, 80, 2000 and 4000 were used. The fit is indicated by dashed line.

$$d^{(n)}(t, L) = d_0 + \sum_k A_k \cos \left[ \text{Im} \left( x_0^{(n)}(k) \right) z \right] e^{-\text{Re} \left( x_0^{(n)}(k) \right) z}, \quad z = \frac{2\pi v_s t}{L}, \quad (12)$$

where  $d_0$  is the density of ‘peaks’ in the stationary state which is the same for each copy  $n$  and the  $A_k$ ’s are dependent on the initial conditions. We have computed  $d^{(n)}(t, L) - d_0$  using Monte Carlo simulations in the case  $N = 3$  for different lattice sizes. The initial state was the configuration  $|+, -, +, -, \dots, +, -\rangle^{(0)}$  in the copy  $n = 0$ . The results are shown in figure 3. One can see that, as expected, the densities are dependent on  $z$  only.

Encouraged by this observation, we did a fit to the  $n = 0$  data (see figure 4) using the parameterization

$$d^{(0)}(z) - d_0 = e^{-az} \frac{\cos b(z - z_0)}{\cos bz_0} \quad (13)$$

and obtain

$$a = 0.1379, \quad b = 0.1107, \quad z_0 = 0.060. \quad (14)$$

These values are compatible with  $x_0^{\downarrow}(2)$  given in (11). To our knowledge, it is for the first time that expressions like (13) appear in a conformal invariant theory.

Before closing this letter, we would like to notice that in a seminal paper Saleur and Sornette [9] have suggested the possible existence of complex critical exponents in non unitary conformal field theories. We have shown that they indeed exist.

### Acknowledgements

This work was supported in part by the joint DFG and RFBR grants no.RI 317/16-1 and no.12-02-5133-NNIO, by FAPESP and CNPq (Brazilian Agencies), and a grant of the Heisenberg-Landau program. PP was also supported by the RFBR grant 14-01-00474 and by the Higher School of Economics Academic Fund grant 14-09-0175. FCA thanks the Bethe Center of the Bonn University for partial financial support.



## References

- [1] Alcaraz F C, Ram A and Rittenberg V 2014 *J. Phys. A: Math. Theor.* **47** 212003
- [2] Levy D 1991 *Phys. Rev. Lett.* **67** 1971
- [3] Martin P P and Saleur H 1993 *Commun. Math. Phys.* **158** 155  
Martin P P and Saleur H 1994 *Lett. Math. Phys.* **30** 189
- [4] Graham J J and Lehrer G I 1998 *L'Enseignement Math.* **44** 173  
Graham J J and Lehrer G I 2002 *Compositio Math.* **133** 173  
Graham J J and Lehrer G I 2003 *Ann. Sci. Ecole Norm. Sup.* **36** 479
- [5] Morin-Duchesne A and Saint-Aubin Y 2013 *J. Phys. A: Math. Theor.* **46** 285207
- [6] Martin P 1990 *Potts Models and Related Problems in Statistical Mechanics* (Singapore: World Scientific)
- [7] Alcaraz F C, Barber M N and Batchelor M T 1989 *Ann. Phys.* **182** 280
- [8] Alcaraz F C and Rittenberg V 2013 *J. Stat. Mech.* P09010
- [9] Saleur H and Sornette D 1996 *J. Phys. I* **6** 327