

Soil Carbon Sequestration and Biochar as Negative Emission Technologies

Pete Smith¹

¹ Institute of Biological and Environmental Sciences, Scottish Food Security Alliance-Crops & ClimateXChange, University of Aberdeen, 23 St Machar Drive, Aberdeen, AB24 3UU, UK

***Corresponding author:** Prof Pete Smith, Tel: +44 (0)1224 272702, Fax: +44 (0)1224 272703, E-mail: pete.smith@abdn.ac.uk

Running head: Soil C and biochar as NETs

Keywords: soil, carbon, biochar, negative emission technology, sequestration

Paper type: Original research

Abstract

Despite 20 years of effort to curb emissions, greenhouse gas (GHG) emissions grew faster during the 2000s than in the 1990s, which presents a major challenge for meeting the international goal of limiting warming to less than 2°C relative to the preindustrial era. Most recent scenarios from integrated assessment models require large-scale deployment of negative emissions technologies (NETs) to reach the 2°C target. A recent analysis of NETs including direct air capture, enhanced weathering, bioenergy with carbon capture and storage and afforestation/deforestation, showed that all NETs have significant limits to implementation, including economic cost, energy requirements, land use, and water use. In this paper I assess the potential for negative emissions from soil carbon sequestration and biochar addition to land, and also the potential global impacts on land use, water, nutrients, albedo, energy and cost.

Results indicate that soil carbon sequestration and biochar have useful negative emission potential (each 0.7 GtCeq./yr), and that they potentially have lower land impact, water use, nutrient impact, albedo impact and energy requirement and cost, so have fewer disadvantages than many NETs. Limitations of soil carbon sequestration as a NET centre around issues of sink saturation and reversibility. Biochar could be implemented in combination with bioenergy with carbon capture and storage.

Current integrated assessment models do not represent soil carbon sequestration or biochar. Given the negative emissions potential of SCS and biochar and their potential advantages compared to other NETs, efforts should be made to include these options within IAMs, so that their potential can be explored further in comparison with other NETs for climate stabilisation.

Introduction

Greenhouse gas (GHG) emissions grew faster during the 2000s than in the 1990s (Le Quéré *et al.*, 2013), despite 20 years of effort to curb emissions. This continuing increase in emissions will present a major challenge for meeting the international goal of limiting warming to less than 2°C relative to the preindustrial era, particularly if stringent climate policies are not introduced rapidly (Peters *et al.*, 2013; Edenhofer *et al.*, 2014; Tavoni *et al.*, 2015). In order to avoid warming of more than 2°C with a >50% chance, most recent scenarios from integrated assessment models (IAMs) include the large-scale deployment of negative emissions technologies (NETs), i.e. technologies that result in the net removal of CO₂ / GHGs from the atmosphere (Edenhofer *et al.*, 2014; Fuss *et al.*, 2014; Krey *et al.*, 2014; Edmonds *et al.*, 2013; van Vuuren *et al.*, 2013; Rogelj *et al.*, 2013; Clarke *et al.*, 2014; Riahi *et al.*, 2015). Since society must decide which mitigation pathways are desirable to tackle climate change, information on the potential risks and opportunities afforded by all NETs is necessary.

Smith *et al.* (2015) recently reviewed and analysed the biophysical and economic limits to NET implementation for a number of NETs: (1) Bioenergy (BE; Creutzig *et al.*, 2015) with carbon capture and storage (CCS; together referred to as BECCS; Obersteiner *et al.*, 2001), (2) direct air capture of CO₂ from ambient air by engineered chemical reactions (DAC; Keith, 2009; Socolow *et al.*, 2011), (3) enhanced weathering of minerals (EW; Schuiling & Krijgsman, 2006; Köhler *et al.*, 2010; Hartmann & Kempe, 2009; Kelemen & Matter, 2008) where natural weathering to remove CO₂ from the atmosphere is accelerated, and the products stored in soils, or buried in land/deep ocean and (4) afforestation and reforestation (AR) to fix atmospheric carbon in biomass and soils (Arora & Montenegro, 2011; Canadell & Raupach, 2008; Jackson *et al.*, 2008). For reasons of tractability, the analysis of Smith *et al.* (2015) did not consider (5) manipulation of uptake of carbon by the ocean either biologically (i.e. by fertilizing nutrient limited areas; Sarmiento *et al.*, 2005; Joos *et al.*, 1991) or chemically (i.e. by enhancing alkalinity; Kheshgi, 1995), or other land based options, such as (6) changed agricultural practices (which include activities such as less invasive tillage with residue management, organic amendment, improved rotations / deeper rooting cultivars, optimized stocking density, fire management, optimised nutrient management and restoration of degraded lands; Smith *et al.*, 2008; Smith, 2012; Powlson *et al.*, 2014), or (7) converting biomass to recalcitrant biochar, for use as a soil amendment (Woolf *et al.*, 2010). IAMs have so far focused primarily on BECCS (van Vuuren *et al.*, 2013; Azar *et al.*, 2010; Kriegler *et al.*, 2013) and AR (Strengers *et al.*, 2008; Wise *et al.*, 2009; Reilly *et al.*, 2012; Humpenöder *et al.*, 2014). Although soil carbon is starting to be considered by some IAMs, it has so far only been considered as a by-product of other land management actions, such as BECCS and AR. In this study, we assess the NET potential for, and biophysical and economic limitations of, the two land-based NETs not considered in Smith *et al.* (2015), i.e. soil carbon sequestration (SCS) and soil amendment with biochar. This is timely given the proposal of the French Government to attempt to increase global soil carbon stocks by 0.4% as a climate mitigation measure. Table 1 provides a summary of the abbreviations used.

Figure 1 depicts the main flows of carbon among atmospheric, land, ocean and geological reservoirs for fossil fuel combustion (Fig. 1a), BE (Fig. 1b), CCS (Fig. 1c), and the altered carbon flows for BECCS (Fig. 1d), for soil carbon sequestration (Fig. 1e), for biochar addition to soil (Fig. 1f) and for biochar addition to soil as part of BECCS (Fig. 1g).




Figure 1. The main flows of carbon among atmospheric, land, ocean and geological reservoirs for fossil fuel combustion (a), BE (b), CCS (c), and the altered carbon flows for BECCS (d), for SCS (e), for biochar addition to soil (f) and for biochar addition to soil as part of BECCS (g). Adapted from Smith *et al.* (2015).

Table 1. Abbreviations for negative emissions technologies used in this paper.

Abbreviation	Full name of negative emission technology
NET	Negative Emission Technology
DAC	Direct Air Capture
EW	Enhanced Weathering
AR	Afforestation / Reforestation
BE	Bioenergy
CCS	Carbon Capture and Storage
BECCS	Bioenergy with Carbon Capture and Storage
SCS	Soil Carbon Storage
Biochar	Biochar addition to soil

Using data from the latest available literature, I first assess the impact on land use (area required) and GHG emissions (total GHG balance), water (water use per unit of negative emissions), nutrients (nitrogen: N, phosphorus: P and potassium: K associated with sequestered carbon), biophysical climate impacts (represented by surface albedo), energy produced or demanded (per unit of negative emissions) and cost (cost per unit of negative emissions), of SCS and biochar, all per unit of negative emission (i.e. per-t-Ceq.). I then use global estimates of potential implementation of these two NETs (GtC/yr), to estimate global impacts on each of these biophysical and cost parameters, using similar methods to those used recently by Smith *et al.* (2015).

Materials & Methods

Sources of data used to estimate impacts of SCS and biochar on a per-t-Ceq. removal basis are summarised in Table 2. Papers were selected by Web of Knowledge search for the impacts considered in this paper.

Table 2. Sources of data used to estimate impacts of SCS and biochar on a per-t-Ceq. removal basis

Impact	Data type	Source
Land use and GHG impact	Land use intensity for SCS	Smith <i>et al.</i> (2008)
	Increased productivity from SCS	Lal (2004)
	N ₂ O emissions from SCS	IPCC (2006)
	CH ₄ and N ₂ O emissions from biochar	Zhang <i>et al.</i> (2010); Scheer <i>et al.</i> (2011); Castaldi <i>et al.</i> (2011); Woolf <i>et al.</i> (2010); Meyer <i>et al.</i> (2011)
Water use	Qualitative data on improved soil water holding capacity for SCS and biochar	Lal (2004); Lehmann & Joseph (2009); Woolf <i>et al.</i> (2010)

Nutrient impacts	Nutrient content: nitrogen (N), Phosphorus (P) and Potassium (K) for SCS	Lal (2004)
	Nutrient content: nitrogen (N), Phosphorus (P) and Potassium (K) for biochar	Roberts <i>et al.</i> (2015)
Albedo impacts	Qualitative impact of SCS management practices on albedo	Luyssaert <i>et al.</i> (2014); Daughtry <i>et al.</i> (2006); Pacheco & McNairn (2010)
	Quantitative impact of biochar application on albedo	Meyer <i>et al.</i> (2012); Genesio <i>et al.</i> (2012)
Energy	Range of energy requirements for SCS management	Smith <i>et al.</i> (1998); Smith <i>et al.</i> (2008)
	Range of energy generation potentials for biochar	Roberts <i>et al.</i> (2015); Shackley <i>et al.</i> (2011); Meyer <i>et al.</i> (2011); Woolf <i>et al.</i> (2014)
Cost	Costs of SCS management actions	Smith <i>et al.</i> (2008); McKinsey & Co. (2009)
	Costs of biochar production / addition	Shackley <i>et al.</i> (2011); Meyer <i>et al.</i> (2011); Dickinson <i>et al.</i> (2014)

Per-t-C-eq. impact values were then scaled to the global level by multiplying per-t-C-eq. impact values by global implementation levels from Smith *et al.* (2008) for SCS, and from Woolf *et al.* (2010) for biochar, similar to the approach used by Smith *et al.* (2015) for other NETs.

Results

Estimating impacts of SCS and biochar on a per-t-Ceq. removal basis

Land use and GHG impacts

SCS rates vary considerably by soil type and climate region (Smith, 2012), but a global meta-analysis derived mean values for croplands and grasslands ranging from 0.03 (the lowest mean cropland SCS rate) to around 1 tC/ha/yr (for restoration of degraded land; Smith *et al.*, 2008). Inverting these values gives a land requirement per-t-Ceq. of negative emissions, i.e. a range of 1-33 ha/tC. Given that biochar is more recalcitrant than soil organic matter (Lehmann *et al.*, 2015), and that application rates to soil can be high (e.g. 30-60 t/ha: Genesio *et al.*, 2012; Zhang *et al.*, 2010) negative emissions per ha can be much higher, giving land requirements for biochar of <1 ha/tC. However, since SCS and biochar addition can be applied to all managed land without changing its current use, there is no competition for land associated with these NETs; unlike AR and BECCS, which require land use change and

compete for land for other uses (Smith *et al.*, 2010). Net removal of CO₂ from the atmosphere could be further increased by increasing plant productivity. Lal (2004) reports that an addition of 1 tC/ha on degraded cropland soils could increase crop yield by 0.5 kg/ha (cowpeas) to 40 kg/ha (wheat). If this additional yield is removed and consumed as food or feed, however, it would not contribute to net CO₂ removal from the atmosphere.

In addition to the impacts on soil carbon storage, SCS and biochar can impact other GHGs. While increased soil organic matter under SCS is considered to have small / negligible impacts on soil methane (CH₄) emissions (Smith *et al.*, 2008), increased soil C storage results in more organic nitrogen in the soil, which could be mineralised to become a substrate for nitrous oxide (N₂O) production, though the effect is difficult to quantify (IPCC, 2006). Biochar addition, on the other hand, has quantifiable impacts on non-CO₂ GHG emissions, with CH₄ emissions increasing significantly and N₂O emissions decreasing substantially when biochar is added to paddy rice soils (Zhang *et al.*, 2010). Other studies in pasture systems show the opposite effect (increased N₂O; decreased CH₄; Scheer *et al.*, 2011) and others no effect (Castaldi *et al.*, 2011). The prevention of CH₄ and N₂O emissions from biomass decay when biomass is used to produce biochar is also expected to contribute to the emission reduction delivered by biochar (Woolf *et al.*, 2010). Given the uncertainty in the relationships between both soil organic matter content and biochar addition and CH₄ and N₂O emissions, the impact on non-CO₂ GHGs for both SCS and biochar is assumed here to be zero.

Water use

Water is not used in quantity in the production of biochar (with the exception of hydrothermal carbonization; Meyer *et al.*, 2011). In terms of impacts on soil water when applied, per-t-Ceq. water impacts of increased soil organic matter storage and biochar additions to soil are difficult to quantify, though it is widely accepted that both improve soil water holding capacity (Lal, 2004; Lehmann & Joseph, 2009; Woolf *et al.*, 2010), thereby increasing retention of water in the soil / plant system. Unlike other NETs, then, SCS and biochar are likely to have a net beneficial impact on water footprint (and no negative impact), so the per-t-Ceq. impact on water use is <0. Additional water used for irrigation to grow the biochar feedstock is not included; if the biomass feedstocks needed irrigation, this would have a water footprint.

Nutrient impacts

Soil organic matter from SCS provides additional nutrients for plant uptake, since the stoichiometry of the organic matter means that for every t C/ha of soil organic matter added, nutrients, i.e. N, P and K, would increase by 80 kgN/ha, 20 kgP/ha, and 15 kgK/ha (calculated from values in Lal, 2004). Though these nutrients need to be added to maintain stable soil organic matter, it is assumed that these can be obtained through a combination of organic matter addition and nitrogen fixation. Biochar is credited with reducing nutrient losses from soils (Steiner *et al.*, 2008; Lehmann *et al.*, 2003). Though biochars are very variable, depending on feedstock and pyrolysis temperature (Lehmann & Joseph, 2009), using the average of ranges of C, N, P and K content given for lignocellulosic biochars in

Roberts *et al.* (2015), biochar containing 1 tC/ha would contain ~30 kgN/ha, ~10 kgP/ha, and ~70 kgK/ha. Nutrient benefits are, therefore, quantifiable for SCS and biochar, and in both cases are positive. Additional fertilizers to grow the biochar feedstock are not included; if the biomass feedstocks required more fertilizer than the previous land use, this would have a negative impact on nutrients.

Albedo impacts

Though soil organic carbon content is unlikely to impact directly upon albedo to a significant extent, some practices that are used for SCS (e.g. residue management) can impact upon albedo (Luyssaert *et al.*, 2014), to the extent that remote sensing can be used to detect residues on fields (Daughtry *et al.*, 2006; Pacheco & McNairn, 2010). It is conceivable that light coloured residues (e.g. cereal straw) left upon the surface of an otherwise bare soil could increase albedo, thereby providing an additional climate benefit to soil C storage. The magnitude of this impact, however, cannot currently be quantified due to uncertainty over the mix of management practices used for SCS, and the albedo impacts of each practice. It is safe to assume however, that there will be no negative impacts on albedo from SCS, as there could be from AR (Betts *et al.*, 2007).

Because biochar material tends to be dark and can be applied in large quantities, it can darken the soil surface. Biochar application at 30-60 t/ha to soil decreased surface albedo over the crop season by up to 40% relative to controls, which, in turn increased soil temperature (Genesio *et al.*, 2012). Control albedo measurements in this experiment were around 0.2 (early growing season) to 0.3 (late growing season), so a 40% reduction represents absolute albedo decreases of ~0.08-0.12 (Genesio *et al.*, 2012). These albedo changes are consistent with the mean annual albedo reduction of 0.05 calculated for application of 30-32 t ha⁻¹ biochar to a test field in Germany by Meyer *et al.* (2012).

Energy

Most practices that promote SCS are close to current practice (e.g. improved rotations, residue management etc.) and would not incur a significant energy cost. Indeed, some practices (such as reduced / zero tillage) may save energy by reducing the energy input to farm operations, such as diesel use in tillage (Frye, 1984; Kern & Johnson, 1993; Smith *et al.*, 1998). Some practices, however, may incur an energy cost (e.g. additional energy for pumping irrigation water), but given that SCS can be achieved in most cases with energy inputs that do not differ significantly from baseline practices, SCS is assumed to be energy neutral.

Assuming energy contents for lignocellulosic biochars of 16.4-35.3 MJ/kg from Roberts *et al.* (2015), and allowing for 10% and 20% process energy cost and energy loss in pyrolysis plants, respectively (from Shackley *et al.*, 2011), then expressing the range as per-t-C, we arrive at an energy generation potential of ~20-50 GJ/tC of biochar.

Cost

The cost of cropland and grazing land sequestration given by McKinsey & Co. (2009), which were recalculated from Smith *et al.* (2008), range from about -165 \$/tCeq. (for tillage and residue management) to around 40 \$/tCeq. (for degraded land restoration and improved agronomy – values converted from €/tCO₂-eq., all \$ values are US\$ for decade 2000-2010).

Estimates for costs of production to application of biochar (assuming UK conditions) are about -830 to around 1200 \$/tCeq. (converted from GB£/tCO₂-eq. values in Shackley *et al.*, 2011). The mean of this range is 185 \$/tCeq. A tighter cost range is derived from values in Meyer *et al.* (2011), using costs of 51 to 386 \$/t biochar (for yard waste and retort charcoal, respectively), giving cost ranges of 54-406 \$/tCeq. if 95% C content from slow pyrolysis is assumed, or 100-757 \$/tCeq. if 51% C content from torrefaction is assumed. The mean of these ranges are ~230 and 430 \$/tCeq., respectively. There may be economic benefits from biochar application that offset some of these costs (Dickinson *et al.*, 2014), but the benefits differ by region.

All per-t-C impacts of SCS and biochar on land use, water, nutrients, albedo, energy and cost are shown in Figure 2.




Figure 2. Impacts of SCS and biochar on land use (a), nutrients (b), albedo (c), energy (d) and cost (e), expressed on per-t-C basis. Additional water use not shown as it is zero for SCS and biochar.

Global estimates of impacts of SCS and biochar

Global implementation levels for SCS range from 0.4 to 0.7 GtCeq./yr at carbon prices ranging from ~70-370 \$/tC-eq. (equivalent to 20-100 \$/tCO₂-eq.; Smith *et al.*, 2008), with a technical potential of 1.3 GtCeq./yr. Woolf *et al.* (2010) report sustainable potential for avoided emissions for biochar, after accounting for competition for non-waste biomass, to be ~1 GtCeq./yr, with a maximum technical potential of 1.8 GtCeq./yr. Of these avoided emissions, about 30% derives from fossil fuel displacement, and about 70% is from biochar storage in soil – so the negative emission component of biochar is 0.7 GtCeq./yr, with a maximum technical potential of 1.3 GtCeq./yr. The global impacts on land use, water, nutrients, albedo, energy and cost for these levels of implementation are presented below.

Land use

For implementation levels of SCS at C prices of up to ~370 \$/tC-eq., 0.7 GtCeq./yr of SCS would require 700 (min) and 23100 (max) Mha. For implementation at maximum technical potential, the land area would be 1300 (min) to 42900 (max) Mha. Biochar, implementation at 0.7 (constrained) and 1.3 (maximum theoretical) GtCeq./yr of negative emissions would have a land footprint for spreading of biochar (assuming application rates of 50 t/ha; Genesio *et al.*, 2012) of 14 and 26 Mha, respectively.

For SCS and biochar, however, this land can remain in its current land use, and both SCS and biochar can be practiced without competing for land. The requirement for feedstock to produce biochar, however, does have a land footprint. Of the biochar potential of ~0.7 GtCeq./yr, 0.3 GtCeq./yr is derived from dedicated biomass crops (Woolf *et al.*, 2010), which is proposed to occur on 50% of abandoned, degraded cropland that is not used for other purposes (Woolf *et al.*, 2010). This 0.3 GtCeq./yr of negative emissions from biochar would require 40-260 Mha, assuming biomass yields of lignocellulosic energy crops (e.g. miscanthus: 5.8-8.6 t Ceq./ha/yr; Smith *et al.*, 2015, and that 20% (fast pyrolysis) to 88% (torrefaction) of the biomass C will be converted to biochar C (Meyer *et al.*, 2011). If yields were lower on this degraded land, the area required to deliver these negative emissions from biochar would increase proportionally.

Water use

Water use by SCS and biochar are assumed to be negligible, so global impact on water of both technologies is estimated to be ~zero.

Nutrient impact

For implementation levels of SCS at 0.7 GtCeq./yr of negative emissions, SCS would add an additional 56 Mt N/yr, 14 Mt P/yr and 10.5 Mt K/yr to the soil. Equivalent values for 1.3 GtCeq./yr of negative emissions from SCS are 104, 26, 19.5 Mt/yr of N, P and K, respectively. Biochar implemented at 0.7 GtCeq./yr of negative emissions would add 21 Mt

N/yr, 7 Mt P/yr and 49 Mt K/yr to the soil. Equivalent values for 1.3 GtCeq./yr of negative emissions from biochar are 31, 13 and 91 Mt/yr of N, P and K, respectively.

Albedo

Global impacts on albedo of SCS are zero, and for areas on which biochar is spread (14 Mha for implementation at 0.7 GtCeq./yr), albedo could be reduced by 0.08 to 0.12. To contextualise this change in albedo, the mean annual albedo reduction of 0.05 calculated for application of 30-32 t ha⁻¹ biochar to a test field in Germany by Meyer *et al.* (2012) was estimated to reduce the overall climate mitigation benefit of biochar application by 13-22% Meyer *et al.* (2012).

Energy

Global energy costs/benefits from SCS are zero, whereas biochar implemented at 0.7 GtCeq./yr of negative emissions could produce 14-35 EJ/yr energy, with maximum theoretical deployment at 1.3 GtCeq./yr of negative emissions producing up to 65 EJ/yr energy.

Cost

About 20% of the mitigation from SCS is realised at negative cost (-165-0 \$/tCeq.) and about 80% realised is between 0-40 \$/tCeq. (calculated from values in McKinsey & Co., 2009; global marginal abatement cost curve for the agriculture, which was recalculated from values in Smith *et al.*, 2008). Implementation of SCS at 0.7 GtCeq./yr of negative emissions (i.e. at C price up to 370 \$/tC-eq.) would save 7.7 B\$, comprising 16.9 B\$ of savings, and 9.2 B\$ of positive costs (values from Smith *et al.*, 2008). A marginal abatement curve is not available for biochar so total costs/savings cannot be calculated, but the global range for implementation at 0.7 GtCeq./yr of negative emissions would range from -581 B\$ (if all costs were at the low end of cost estimate range) to 1560 B\$ (if all costs were at the high end of cost estimate range). Assuming the cost to be the mean of the range of per-t-Ceq. costs (185 \$/tCeq.) given in Shackley *et al.* (2011), costs of implementation for biochar would be 130 B\$.

Discussion

SCS and biochar have lower negative emission potential (0.7 GtCeq./yr) than some other NETs, with BECCS able to deliver 3.3 GtCeq./yr by 2100 (mean from IPCC AR5 database; Smith *et al.*, 2015) and DAC also able to deliver this level of negative emissions, though the potential is comparable to that of AR (1.1 GtCeq./yr by 2100) and is greater than that for EW (0.2 GtCeq./yr by 2100; Smith *et al.*, 2015). Considering soil-based NETs is timely, given the proposal of the French Government to attempt to increase global soil carbon stocks by 0.4% as a climate mitigation measure, and their ability to improve soil quality and productivity

(Lal, 2004).

Additional land requirements are low for SCS or biochar application, as it is possible to implement SCS and spread biochar without competition for land, though land required to grow biomass feedstock for biochar could be 40-260 Mha of land. This is greater than the land requirement for DAC (negligible excluding land for renewables to provide energy), and EW (2 Mha), but ranges from an order of magnitude smaller to similar, to the land required to implement 3.3 and 1.1 GtCeq./yr of negative emissions through BECCS (380-700 Mha) and AR (320 Mha), respectively (Smith *et al.*, 2015).

While the full GHG balance of biochar application to soils is very feedstock- and technology-specific, and uncertain (between -144 and 17 kgC/tC in biomass feedstock, assuming C content of biomass to be 50% of dry matter; calculated from values in Meyer *et al.*, 2011), the magnitude and uncertainty of GHG reductions/emissions is not large enough to prevent use of biochar as a NET, though more comprehensive studies are required to assess the full climate impact (e.g. by including albedo impacts, see below, or addition of black carbon (soot) to the atmosphere; Meyer *et al.*, 2011).

Water requirements for SCS and biochar are zero, which means that they are the only NETs unlikely to have a water footprint. EW have a low water use (0.3 km³/yr), and DAC has moderate to high water use (10-300 km³/yr), whereas AR (370 km³/yr) and BECCS (720 km³/yr) each has a very large water footprint when implemented at 1.1 and 3.3 GtCeq./yr, respectively (Smith *et al.*, 2015).

Unlike DAC and EW, nutrient retention from SCS and biochar are significant, with nutrient storage measured in Mt N, P or K/yr. In comparison, AR stores several orders of magnitude less nutrients, measured in kt N/yr. The impact of BE/BECCS on ecosystem nutrient storage is variable and depends on the biomass removal regime and the vegetation that the BE crop replaces (Smith *et al.*, 2015). Although nutrient storage contributes to the (supporting) ecosystem service of nutrient cycling (UK National Ecosystem Assessment, 2011), and nutrient storage in soils helps to prevent water and air pollution (e.g. reducing ammonia emissions, and nitrate and phosphorus leaching to water courses; Smith *et al.*, 2013), this nutrient immobilization could be considered a cost, if additional fertilizer is required to provide plant available nutrients for crop growth. In the long term, however, the additional nutrients in the organic matter provide a net ecosystem benefit (Smith *et al.*, 2013).

SCS has no influence on albedo, but biochar can darken the soil surface due to the addition of substantial quantities of dark material to the soil (Genesio *et al.*, 2012), which has been shown to increase soil temperature measurably. Increases relative to un-amended soils are of the order of 0.08-0.12, and could cover an area of 14 Mha for 0.7 GtCeq./yr of negative emissions. DAC and EW have no albedo effect whilst the impact of BE/BECCS is context specific (depending on the BE crop, the vegetation replaced and the latitude), though AR is known to decrease albedo significantly when conifers replace other vegetation at high latitudes (Calvin *et al.*, 2014; Smith *et al.*, 2015).

The energy impact SCS is negligible, but biochar can produce energy during its production

by pyrolysis. Implementation at 0.7 GtCeq./yr of negative emissions would deliver 14-35 EJ/yr, with maximum theoretical implementation at 1.3 GtCeq./yr producing 65 EJ/yr. In this respect, both SCS and biochar are favoured over DAC and EW, each of which has a significant energy requirement, and similar to AR which is also neutral with respect to energy (Smith *et al.*, 2015). However, it delivers around 5-12 times less energy than BE/BECCS, but can be combined with BE/BECCS (Woolf *et al.*, 2010), so that BE/BECCS delivers energy and CO₂ capture (for BECCS), and biochar is produced in the process. There is, of course, a trade-off between energy yield and biochar production, with minimum energy penalties from biochar production of 21 to 33 GJ/tC when producing liquid and gaseous biofuels, respectively (Woolf *et al.*, 2014).


SCS could be delivered at estimated costs between -165 and 40 \$/tCeq. (McKinsey & Co., 2009), and it is estimated that much of the negative emissions could be delivered at negative cost (-16.9 B\$/yr), and the rest at low (9.2 B\$/yr) cost, with an overall saving of 7.7 B\$/yr. Assuming the cost to be the mean of the range of per-t-Ceq. costs for biochar (185 \$/tCeq.), costs of implementation for biochar at 0.7 GtCeq./yr would be 130 B\$/yr. The cost saving from SCS is the only NET which appears to have the potential to be net negative in terms of cost, whereas the cost of biochar is comparable to the investment needs to BE/BECCS of 123-138 B\$/yr in 2050 (Smith *et al.*, 2015). Costs for DAC and EW are much higher, and the cost of AR is low (Smith *et al.*, 2015).

A summary of the global impacts of SCS and biochar, and comparison with the other NETs reviewed in Smith *et al.* (2015) is given in table 3.

Table 3. A comparison of the global impacts of SCS and biochar, with other NETs reviewed in Smith *et al.* (2015). BECCS cost is investment need for electricity and biofuels by 2050; see Smith *et al.* (2015) for further details.

NET	Realistic (max) global C removal (GtCeq./yr)	Additional land requirement (max) (Mha)	Additional water requirement (km ³ /yr)	Mean (max) nutrient impact (Mt N, P, K/yr)	Albedo impact (unitless)	Energy requirement (max) (EJ/yr)	Estimated cost (B\$)	Reference
BECCS	3.3	380-700	720	Variable	Variable	-170	138 / 123	Smith <i>et al.</i> (2015)
DAC	3.3	Very low (unless solar PV used for energy)	10-300	None	None	156	>> BECCS	Smith <i>et al.</i> (2015)
EW	0.2 (1.0)	2 (10)	0.3 (1.5)	None	None	46	>BECCS	Smith <i>et al.</i> (2015)
AR	1.1 (3.3)	320 (970)	370 (1040)	2.2 (16.8)	Negative; or reduced GHG benefit where not negative	Very low	<<BECCS	Smith <i>et al.</i> (2015)
SCS	0.7 (1.3)	0	0	N:56, P:14, K:10.5 (N:104, P:26, K:19.5)	0	0	-7.7	This study
Biochar	0.7 (1.3)	40-260	0	N:21, P:7, K:49 (N:31, P:13, K:91)	0.08-0.12	-14 to -35 (-65)	130	This study

1 A summary graphic, comparing the impacts of SCS and biochar on land use, water, nutrients,
 2 albedo, energy and cost, and the comparison with other NETs, is presented in Figure 3.
 3



4
 5 **Figure 3.** Comparison of the global impacts of SCS and biochar and other NETs on land use,
 6 water, nutrients, albedo, energy and cost. NETs to which SCS and biochar are compared are
 7 BECCS, DAC, EW and AR as assessed by Smith *et al.* (2015). Water requirement shown as
 8 water drops with quantities in km³/yr; all other units are indicated on the figure. Adapted
 9 from Smith *et al.* (2015).

10 SCS and biochar therefore provide negative emissions with fewer potential disadvantages
 11 than many other NETs. Though the negative emissions potential is lower than for DAC and
 12 BECCS, it is not insignificant, and is comparable to the potential for AR and greater than for
 13 EW (Smith *et al.*, 2015). The potential for biochar could be enhanced by combining with
 14 BECCS, so that biochar is produced as a co-product of energy generation from biomass
 15 (Woolf *et al.*, 2010), and the system can be modified to favour carbon allocation either to
 16 CO₂ for CCS, or biochar for use as a soil amendment (Woolf *et al.*, 2014). Given that the
 17 same feedstock is used for biochar production and BE/BECCS, the negative emission
 18 potentials for the two options are not additive.

19
 20 SCS and biochar negative emission potentials are assessed here independently, and are not
 21 additive, since some SCS options rely on retention of crop / forest residues and additions of
 22 organic matter to the soil (Smith *et al.*, 2008; Smith, 2012), which means that these

23 amendments would not be available as feedstocks for biochar production. On the other hand,
24 if these residues are used for biochar production, they may cause a decline in soil organic
25 carbon content (Woolf *et al.*, 2010).

26

27 A drawback of SCS, which also affects AR but not other NETs, is that of sink saturation. We
28 express SCS negative emission potential here as a yearly value, but the potential is time
29 limited. SCS potential is large at the outset, but decreases as soils approach a new, higher
30 equilibrium value (Smith, 2012), such that the potential decreases to zero when the new
31 equilibrium is reached. This sink saturation occurs after 10-100 years, depending on the SCS
32 option, soil type and climate zone (slower in colder regions), with IPCC using a default
33 saturation time of 20 years (IPCC, 1997; IPCC, 2006). Since sinks derived from SCS are also
34 reversible (Smith, 2012), practices need to be maintained, even when the sink is saturated so
35 any yearly costs will persist even after the emission potential has reduced to zero at sink
36 saturation. Sink saturation also means that SCS implemented in 2020 will no longer be
37 effective as a NET after 2040 (assuming 20 years for sink saturation). The importance of this
38 for NETs, is that NETs are most frequently required in the second half of this century (Fuss *et al.*,
39 2014; Smith *et al.*, 2015), so SCS, like AR, may no longer be available after 2050, or will
40 be less effective, if they are implemented for mitigation relatively soon.

41

42 The same issues associated in sink saturation apply partly to biochar, though the issue is less
43 pronounced as biochar is more recalcitrant, and equilibrium (if it occurs) would be expected
44 to take much longer, so that biochar should still be effective as a NET in the second half of
45 this century even if implemented relatively soon. Some authors have suggested that biochar
46 may induce a priming effect, such that organic matter decomposed more rapidly when
47 biochar is add, but evidence for this effect is inconclusive (Zimmerman *et al.*, 2011).

48

49 This analysis has shown that SCS and biochar have potential for delivering negative
50 emissions, and have some advantages over other NETs with respect to other biophysical,
51 energy and cost impacts. However, neither SCS nor biochar production are represented in the
52 current generation of IAMs. Given the negative emissions potential of SCS and biochar and
53 their potential advantages compared to other NETs, efforts should be made to include these
54 options within IAMs, so that their potential can be explored further in comparison with other
55 NETs.

56

57

58 **Acknowledgements**

59 Thanks to Steve J. Davis who created the original figures from which the figures presented
60 here were adapted. This output contributes to the Belmont Forum/FACCE-JPI funded DEVIL
61 project (NE/M021327/1), the Global Carbon Project and the MaGNET program
62 (<http://www.cger.nies.go.jp/gcp/magnet.html>), and also contributes to the EU FP7 SmartSoil
63 project (project number: 289694).

64

65 **References**

- 66 Arora, VK. & Montenegro, A. (2011) Small temperature benefits provided by realistic
67 afforestation efforts. *Nature Geoscience* **4**, 514-518.
- 68 Azar, C. *et al.* (2010) The feasibility of low CO₂ concentration targets and the role of bio-
69 energy carbon-capture and storage. *Climatic Change* **100**, 195-202 (2010).
- 70 Betts, RA., Falloon, PD., Goldewijk, K.K., Ramankutty, N. (2007) Biogeophysical effects of
71 land use on climate: Model simulations of radiative forcing and large-scale temperature
72 change. *Agricultural and Forest Meteorology* **142**, 216-233.
- 73 Calvin, K. *et al.* (2014) Trade-offs of different land and bioenergy policies on the path to
74 achieving climate targets. *Climatic Change* **123**, 691-704.
- 75 Canadell, JG., Raupach, MR. (2008) Managing forests for climate change mitigation. *Science*
76 **320**, 1456-1457.
- 77 Castaldi, S., Riondino, M., Baronti, S., Esposito, FR., Marzaioli, R., Rutigliano, FA., Vaccari,
78 FP., Miglietta, F. (2011) Impact of biochar application to a Mediterranean wheat crop on soil
79 microbial activity and greenhouse gas fluxes. *Chemosphere* **85**, 1464-1471.
- 80 Clarke, L. *et al.* (2014) Assessing transformation pathways. In: *Climate Change 2014:*
81 *Mitigation of Climate Change. Contribution of Working Group III to the Fifth Assessment*
82 *Report of the Intergovernmental Panel on Climate Change* (ed. Edenhofer, O. *et al.*).
83 Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- 84 Creutzig F. *et al.* (2015) Bioenergy and climate change mitigation: an assessment. *Global*
85 *Change Biology Bioenergy* **7**, 916-944.
- 86 Daughtry, CST., Doraiswamy, PC., Hunt, ER., Stern, AJ., McMurtrey, JE., Prueger, JH.
87 (2006) Remote sensing of crop residue cover and soil tillage intensity. *Soil & Tillage*
88 *Research* **91**, 101–108.
- 89 Dickinson, D., Balduccio, L., Buysse, J., Ronsse, F., van Huylenbroeck, G., Prins. W. (2014)
90 Cost-benefit analysis of using biochar to improve cereals agriculture. *Global Change Biology*
91 *Bioenergy* **7**, 850–864.
- 92 Edenhofer, O. *et al.* (Eds) (2014) *Climate Change 2014: Mitigation of Climate Change.*
93 *Contribution of Working Group III to the Fifth Assessment Report of the Intergovernmental*
94 *Panel on Climate Change.* Cambridge University Press, Cambridge, United Kingdom and
95 New York, NY, USA.
- 96 Edmonds, J. *et al.* (2013) Can radiative forcing be limited to 2.6 Wm⁻² without negative
97 emissions from bioenergy and CO₂ capture and storage? *Climatic Change* **118**, 29-43.
- 98 Frye, WW. (1984) Energy requirement in no tillage. In: *No Tillage Agriculture. Principles*
99 *and Practices* (eds. Phillips, RE., Phillips, SH.), pp. 127-151. Van Nostrand Reinhold, New
100 York.
- 101 Fuss, S., Canadell, JG., Peters, GP. *et al.* (2014) Betting on negative emissions. *Nature*
102 *Climate Change* **4**, 850–853.

103 Genesio, L., Miglietta, F., Lugato, E., Baronti, S., Pieri, M., Vaccari, FP. (2012) Surface
 104 albedo following biochar application in durum wheat. *Environmental Research Letters*
 105 (2012): 1-8. *IOP Science*. <http://iopscience.iop.org/1748-9326/7/1/014025>.

106 Hartmann, J., Kempe, S. (2008) What is the maximum potential for CO₂ sequestration by
 107 “stimulated” weathering on the global scale? *Naturwissenschaften* **95**, 1159-1164.

108 Humpenöder, F. *et al.* (2014) Investigating afforestation and bioenergy CCS as climate
 109 change mitigation strategies. *Environmental Research Letters* **9**, 064029 (13pp).

110 IPCC (1997) *Revised 1996 IPCC guidelines for national greenhouse gas inventories*
 111 *workbook, vol. 2*. Cambridge, UK: Cambridge University Press.

112 IPCC (2006) *Revised 2006 IPCC guidelines for national greenhouse gas inventories, vol. 3*.
 113 Cambridge, UK: Cambridge University Press.

114 Jackson, R.B. *et al.* (2008) Protecting climate with forests. *Environmental Research Letters* **3**,
 115 044006.

116 Joos, F., Sarmiento, JL., Siegenthaler, U. (1991) Estimates of the effect of Southern Ocean
 117 iron fertilization on atmospheric CO₂ concentrations. *Nature* **349**, 772-775.

118 Keith, D. (2009) Why capture CO₂ from the atmosphere. *Science* **325**, 1654-1655.

119 Kelemen, P. B. & Matter, J. M. (2008) In situ carbonation of peridotite for CO₂ storage.
 120 *Proceedings of the National Academy of Sciences USA* **105**, 17295-17300.

121 Kern, JS., Johnson, SE. (1993) Conservation tillage impacts on national soil and atmospheric
 122 carbon levels. *Soil Science Society of America Journal* **57**, 200-210.

123 Kheshgi, HS. (1995) Sequestering atmospheric carbon dioxide by increasing ocean alkalinity.
 124 *Energy* **20**, 915-922.

125 Köhler, P., Hartmann, J., Wolf-Gladrow, DA. (2010) Geoengineering potential of artificially
 126 enhanced silicate weathering of olivine. *Proceedings of the National Academy of Sciences*
 127 *USA* **107**, 20228-20233.

128 Krey, V., Luderer, G., Clarke, L., Kriegler, E. (2014) Getting from here to there – energy
 129 technology transformation pathways in the EMF27 scenarios. *Climatic Change* **123**, 369-382.

130 Kriegler, E. *et al.* (2013) What does the 2°C target imply for a global climate agreement in
 131 2020? The LIMITS study on Durban Platform scenarios. *Climate Change Economics* **04**,
 132 1340008. doi: 10.1142/S2010007813400083.

133 Lal, R. (2004) Soil carbon sequestration impacts on global climate change and food security.
 134 *Science* **304**, 1623-1627.

135 Le Quéré, C. *et al.* (2013) The global carbon budget 1959-2011. *Earth System Science Data*
 136 **5**, 165-185.

137 Lehmann, J. & Joseph, S. (2009) *Biochar for Environmental Management: Science and*
 138 *Technology*. Earthscan Books Ltd., London, UK.

- 139 Lehmann, J. *et al.* (2003) Nutrient availability and leaching in an archaeological Anthrosol
140 and a Ferralsol of the Central Amazon basin: fertilizer, manure and charcoal amendments .
141 *Plant & Soil* **249**, 343-357.
- 142 Lehmann J., Czimczik, C., Laird, D., Sohi, S. (2015) Stability of biochar in soil. In: *Biochar*
143 *for Environmental Management: Science, Technology and Implementation* (eds. Lehmann, J.,
144 Joseph, S.), pp. 235-282, Taylor and Francis, London, UK.
- 145 Luysaert, S., Jammet, M., Stoy, PC. *et al.* (2014) Land management and land-cover change
146 have impacts of similar magnitude on surface temperature. *Nature Climate Change* **4**, 389–
147 393.
- 148 McKinsey & Co. (2009) *Pathways to a Low-Carbon Economy. Version 2 of the Global*
149 *Greenhouse Gas Abatement Cost Curve*. McKinsey & Co. Available at:
150 [http://www.mckinsey.com/en/Client_Service/Sustainability/Latest_thinking/Pathways_to_a_l](http://www.mckinsey.com/en/Client_Service/Sustainability/Latest_thinking/Pathways_to_a_low_carbon_economy.aspx)
151 [ow_carbon_economy.aspx](http://www.mckinsey.com/en/Client_Service/Sustainability/Latest_thinking/Pathways_to_a_low_carbon_economy.aspx) (accessed 05 September 2015).
- 152 Meyer, S., Glaser, B., Quicker, P. (2011) Technical, economical, and climate-related aspects
153 of biochar production technologies: a literature review. *Environmental Science &*
154 *Technology*, **45**, 9473–9483.
- 155 Meyer, S., Bright, RM., Fischer, D., Schulz, H., Glaser, B. (2012) Albedo impact on the
156 suitability of biochar systems to mitigate global warming. *Environmental Science &*
157 *Technology*, **46**, 12726-12734.
- 158 Obersteiner, M. *et al.* (2001) Managing climate risk. *Science* **294**, 786-787.
- 159 Pacheco, A, McNairn, H. (2010) Evaluating multispectral remote sensing and spectral
160 unmixing analysis for crop residue mapping. *Remote Sensing of the Environment* **114**, 2219–
161 2228.
- 162 Peters, GP. *et al.* (2013) The challenge to keep global warming below 2°C. *Nature Climate*
163 *Change* **3**, 4-6.
- 164 Powelson, DS., Stirling, CM., Jat, ML., Gerard, BG., Palm, CA., Sanchez, PA., Cassman, KG.
165 (2014) Limited potential of no-till agriculture for climate change mitigation. *Nature Climate*
166 *Change* **4**, 678–683.
- 167 Rau, GH., Knauss, KG., Langer, WH., Caldeira K. (2007) Reducing energy-related CO₂
168 emissions using accelerated weathering of limestone. *Energy* **32**,1471-1477.
- 169 Reilly, J. *et al.* (2012) Using land to mitigate climate change: hitting the target, recognizing
170 the trade-offs. *Environmental Science & Technology* **46**, 5672-5679.
- 171 Riahi, K. *et al.* (2015) Locked into Copenhagen pledges - Implications of short-term emission
172 targets for the cost and feasibility of long-term climate goals. *Technological Forecasting and*
173 *Social Change Part A* **90**, 8-23.

174 Roberts, DA., Paul, NA., Dworjanyn, SA., Bird, MI., de Nys, R. (2015) Biochar from
175 commercially cultivated seaweed for soil amelioration. *Scientific Reports* **5**, 9665. doi:
176 10.1038/srep09665.

177 Rogelj, J., McCollum, DL., Reisinger, A., Meinshausen, M., Riahi, K. (2013) Probabilistic
178 cost estimates for climate change mitigation. *Nature* **493**, 79-83.

179 Sarmiento, J.L., Gruber, N., Brzezinski, M.A., Dunne, J.P. (2004) High-latitude controls of
180 thermocline nutrients and low latitude biological productivity. *Nature* **427**, 56-60.

181 Scheer, C., Grace, PR., Rowlings, DW., Kimber, S., Zwieten, LV. (2011) Effect of biochar
182 amendment on the soil-atmosphere exchange of greenhouse gases from an intensive
183 subtropical pasture in northern New South Wales, Australia. *Plant & Soil* **345**, 47-58.

184 Schuiling, RD., Krijgsman P. (2006) Enhanced weathering: an effective and cheap tool to
185 sequester CO₂. *Climatic Change* **74**, 349–354.

186 Shackley, S., Hammond, J., Gaunt, J., Ibarrola, R. (2011) The feasibility and costs of biochar
187 deployment in the UK. *Carbon Management* **2**, 335–356.

188 Smith, P. (2012) Soils and climate change. *Current Opinion in Environmental Sustainability*
189 **4**, 539–544.

190 Smith, P., Ashmore, M., Black, H., Burgess, P.J., Evans, C., Quine, T., Thomson, A.M.,
191 Hicks, K. & Orr, H. 2013. The role of ecosystems and their management in regulating
192 climate, and soil, water and air quality. *Journal of Applied Ecology* **50**, 812–829.

193 Smith, P., Davis, S.J., Creutzig, F. *et al.* (2015) Biophysical and economic limits to negative
194 CO₂ emissions. *Nature Climate Change* (in press).

195 Smith, P., Gregory, P., van Vuuren, D. *et al.* (2010) Competition for land. *Philosophical*
196 *Transactions of the Royal Society, B*. **365**, 2941-2957.

197 Smith, P., Martinio, D., Cai, Z., *et al.* (2008) Greenhouse gas mitigation in agriculture.
198 *Philosophical Transactions of the Royal Society B*. **363**, 789-813.

199 Smith, P., Powlson, DS., Glendining, MJ., Smith, JU. (1998) Preliminary estimates of the
200 potential for carbon mitigation in European soils through no-till farming. *Global Change*
201 *Biology* **4**, 679-685.

202 Socolow, R. *et al.* (2011) *Direct air capture of CO₂ with chemicals: A technology assessment*
203 *for the APS Panel on Public Affairs*. American Physical Society. 92pp.

204 Steiner, C., Glaser, B., Teixeira, WG., Lehmann, J., Blum, WEH., Zech, W. (2008) Nitrogen
205 retention and plant uptake on a highly weathered central Amazonian Ferralsol amended with
206 compost and charcoal. *Journal of Plant Nutrition & Soil Science* **171**, 893 – 899.

207 Strengers, BJ., Minnen, JGV., Eickhout B. (2008) The role of carbon plantations in
208 mitigating climate change: potentials and costs. *Climatic Change* **88**, 343-366.

209 Tavoni, M. *et al.* (2015) Post-2020 climate agreements in the major economies assessed in
210 the light of global models. *Nature Climate Change* **5**, 119-126.

- 211 UK National Ecosystem Assessment (2011) *The UK National Ecosystem Assessment*
212 *Technical Report*. UNEP-WCMC, Cambridge.
- 213 van Vuuren, D. P. *et al.* (2013) The role of negative CO₂ emissions for reaching 2 °C -
214 insights from integrated assessment modelling. *Climatic Change* **118**, 15-27.
- 215 Wise, M. *et al.* (2009) Implications of limiting CO₂ concentrations for land use and energy.
216 *Science* **324**, 1183–1186.
- 217 Woolf, D., Amonette, JE., Street-Perrott. A., Lehmann, J., Joseph, S. (2010) Sustainable
218 biochar to mitigate global climate change. *Nature Communications* **1**, Article 56,
219 doi:10.1038/ncomms1053 (2010).
- 220 Woolf, D., Lehmann, J., Fisher, EM., Angenent, LT. (2014) Biofuels from pyrolysis in
221 perspective: trade-offs between energy yields and soil-carbon additions. *Environmental*
222 *Science & Technology*, **48**, 6492-6499.
- 223 Zhang, A., Cui, LQ., Pan, G., Li, LQ., Hussain, Q., Zhang, XH., Zheng, JW., Crowley, D.
224 (2010) Effect of biochar amendment on yield and methane and nitrous oxide emissions from
225 a rice paddy from Tai Lake Plain, China. *Agriculture, Ecosystems & Environment* **139**, 469-
226 475.
- 227 Zimmerman, A., Gao, B., Ahn, M.-Y. (2011) Positive and negative carbon mineralization
228 priming effects among a variety of biochar-amended soils. *Soil Biology and Biochemistry* **43**,
229 1169–1179.