

ANALYTIC FAMILIES OF QUANTUM HYPERBOLIC INVARIANTS

STÉPHANE BASEILHAC¹, RICCARDO BENEDETTI²¹ Institut de Mathématiques et de Modélisation, Université Montpellier 2, Case Courrier 51, 34095 Montpellier Cedex 5, France (sbaseilh@math.univ-montp2.fr)² Dipartimento di Matematica, Università di Pisa, Largo Bruno Pontecorvo 5, 56127 Pisa, Italy (benedett@dm.unipi.it)

ABSTRACT. We organize the quantum hyperbolic invariants (QHI) of 3-manifolds into sequences of rational functions indexed by the odd integers $N \geq 3$ and defined on moduli spaces of geometric structures refining the character varieties. In the case of one-cusped hyperbolic 3-manifolds M we generalize the QHI and get rational functions $\mathcal{H}_N^{h_f, h_c, k_c}$ depending on a finite set of cohomological data (h_f, h_c, k_c) called *weights*. These functions are regular on a determined Abelian covering of degree N^2 of a Zariski open subset, canonically associated to M , of the geometric component of the variety of augmented $PSL(2, \mathbb{C})$ -characters of M . New combinatorial ingredients are a weak version of branchings which exists on every triangulation, and state sums over weakly branched triangulations, including a sign correction which eventually fixes the sign ambiguity of the QHI. We describe in detail the invariants of three cusped manifolds, and present the results of numerical computations showing that the functions $\mathcal{H}_N^{h_f, h_c, k_c}$ depend on the weights as $N \rightarrow +\infty$, and recover the volume for some specific choices of the weights.

CONTENTS

1. Introduction	2
1.1. QHI of cusped manifold patterns	3
1.2. QHI of other patterns	4
1.3. State sums over weakly branched triangulations	5
1.4. Plan of the paper	5
2. Structured triangulations and gluing varieties	6
3. Analytic configurations	13
3.1. Local analytic configurations	13
3.2. Globalization	15
4. Cusped manifolds	17
4.1. The augmented $PSL(2, \mathbb{C})$ -character variety and A -polynomial	17
4.2. Rich components of $G(T)$	18
4.3. Refined analytic configurations for one-cusped manifolds	19
4.4. Relation with Chern-Simons theory	28
5. QHFT partition functions	32
6. \mathcal{N} -graph calculus	32
6.1. \mathcal{N} -graphs representing the same weakly branched triangulation	33
6.2. Changing the weak branching	33
6.3. Transits	38
7. State sum invariance over weakly branched triangulations	40
7.1. Formal conversion of tensor networks	40
7.2. The relations between $\mathcal{R}_N(\Delta, b, w, f, c)$ and $\mathcal{R}_N(\Delta, b_\beta, w', f', c')$	42
7.3. Proof of Proposition 7.1.	43
8. Resolution of the sign anomaly	43
8.1. Behaviour of the sign anomaly under QH transits	43
8.2. Behaviour of the sign anomaly under a change of weak branching	46

9. Examples	46
9.1. Notations	47
9.2. The figure-eight knot complement	47
9.3. The complement of the knot 5_2	49
9.4. The sister of the figure-eight knot complement	52
References	53

1. INTRODUCTION

In the series of papers [3, 4, 5] we defined a family of complex valued quantum invariants $\mathcal{H}_N(\mathcal{P})$ of “patterns” \mathcal{P} of geometric nature, called *quantum hyperbolic invariants* (QHI) and indexed by the odd integers $N \geq 3$. Roughly, a pattern consists of a compact oriented 3-manifold equipped with a representation of the fundamental group into $PSL(2, \mathbb{C})$, plus some compatible cohomological datas (see the next subsections for a complete definition). The QHI generalize the Kashaev invariants of links in S^3 [6], subject to the celebrated Volume Conjecture. They define a $(2 + 1)$ -dimensional quantum field theory, and, when applied to mapping cylinders of surface diffeomorphisms, they coincide with the invariants derived from the local version of finite dimensional quantum Teichmüller theory (see [1], [2] and [5]).

In [3, 4, 5], the invariance of the QHI was proved only up to sign and multiplication by N -th roots of unity. The eventual existence and the meaning of such a phase anomaly, as well as the determination of the asymptotical behaviour of $\mathcal{H}_N(\mathcal{P})$ as $N \rightarrow +\infty$, are two main open issues of the theory. In order to tackle them, it seemed necessary to develop a functional approach that would clarify the intrinsic nature of the various combinatorial and geometric ingredients involved in the definition of the QHI. To this aim, we achieve in this paper the following goals:

QHI of cusped manifolds: We extend the QHI of any one-cusped hyperbolic 3-manifold M to invariants defined on the geometric component $X_0(M)$ of the variety of augmented $PSL(2, \mathbb{C})$ -characters of M . In [4, 5] the QHI of cusped hyperbolic manifolds were defined only at the hyperbolic holonomy.

State sums over weakly branched triangulations: In order to achieve the previous goal it is necessary to introduce state sums over triangulations that do not support any branching, replaced by relaxed structures called *weak branchings*. These states sums give rise to QHI generalizing all the previously defined ones.

Analytic families of QHI: We recast all the QHI into sequences $\{\mathcal{A}_N(Y)\}$ of families of complex analytic spaces and maps, indexed by the odd integers $N \geq 3$. Each family $\mathcal{A}_N(Y)$ is associated to a *topological support* Y , and provides concrete models of geometric structures over Y called *patterns*. Patterns over cusped manifolds have an intrinsic meaning in terms of the $PSL(2, \mathbb{C})$ -version of the A -polynomial, and natural relationships with Chern-Simons theory.

Fixing the sign ambiguity: We include a sign correction in the state sum formulas of the QHI, which removes their sign ambiguity under a mild assumption on the “bulk c -weight” (see below). The sign correction depends on the combinatorics of the weak branching and is a by-product of [9]. It becomes trivial when dealing (when possible) with branched triangulations, so that the QHI of [3, 4, 5] are eventually defined up to multiplication by N -th roots of unity.

In a sequel to this paper (in collaboration with C. Frohman), we will develop our approach concerning the asymptotical behaviour of the QHI. Basically, given a topological support Y and a sequence $\{\mathcal{P}_N\}$ of patterns over Y , we study the limit of the family $\{\mathcal{A}_N(Y)\}$ as $N \rightarrow +\infty$, instead of $\{\mathcal{H}_N(\mathcal{P}_N)\}$ for a single sequence $\{\mathcal{P}_N\}$. Then, assuming that $\mathcal{H}_\infty(\{\mathcal{P}_N\}) := \limsup_{N \rightarrow \infty} (\log |\mathcal{H}_N(\mathcal{P}_N)|/N)$ is finite (this is the case for all natural sequences $\{\mathcal{P}_N\}$), we consider the following problems:

- (1) *Determine the nature (regularity) of $\mathcal{H}_\infty(\{\mathcal{P}_N\})$ as a function of the patterns over Y .*
- (2) *Describe the asymptotical behaviour of $\{\mathcal{H}_N(\mathcal{P}_N)\}$ as $N \rightarrow \infty$ in terms of classical geometric invariants of Y : Chern-Simons invariants, torsions, twisted cohomology, etc.*

The Kashaev-Murakami-Murakami Volume Conjecture is a particular case of (2), for constant sequences of patterns associated to links in S^3 (see Theorem 1.2 below).

In the rest of this Introduction we describe with more details the content of the paper.

1.1. QHI of cusped manifold patterns. In this paper we call *cusped manifold* an oriented, connected, non-compact complete hyperbolic 3-manifold of finite volume with *one* cusp. Hence a cusped manifold M is diffeomorphic to the interior of a compact 3-manifold denoted by V , with one torus boundary component.

A *pattern* $\mathcal{P} = (Y_{\mathcal{P}}, \rho, (h, k))$ over M consists of a *topological support* $Y_{\mathcal{P}}$ together with additional geometric structures determined by the couple $(\rho, (h, k))$. The topological support takes the form $Y_{\mathcal{P}} = (V, (h_c, k_c))$, where $(h_c, k_c) \in H^1(V; \mathbb{Z}/2\mathbb{Z}) \times H^1(\partial V; \mathbb{Z})$ is a so-called *c-weight*, defined by a “bulk c-weight” h_c and a “boundary c-weight” k_c satisfying

$$(1) \quad r(k_c) = i^*(h_c)$$

where $r : H^1(\partial V; \mathbb{Z}) \rightarrow H^1(\partial V; \mathbb{Z}/2\mathbb{Z})$ is the reduction mod(2), and the map $i^* : H^1(V; \mathbb{Z}/2\mathbb{Z}) \rightarrow H^1(\partial V; \mathbb{Z}/2\mathbb{Z})$ is induced by the inclusion map $i : \partial V \rightarrow V$. The pattern \mathcal{P} is obtained by completing $Y_{\mathcal{P}}$ with a couple $(\rho, (h_f, k_f))$ where ρ is a *PSL(2, \mathbb{C})-character* of V , i.e. a conjugacy class of representations of $\pi_1(V)$ in $PSL(2, \mathbb{C})$, and $(h_f, k_f) \in H^1(V; \mathbb{Z}/2\mathbb{Z}) \times H^1(\partial V; \mathbb{C})$ is a so-called *f-weight* (relative to ρ), defined by a “bulk f-weight” h_f and a “boundary f-weight” k_f satisfying the following constraint. Up to conjugacy the restriction of ρ to the torus ∂V is valued in the group of complex affine transformations of the plane; the linear part of this restriction defines a class in $H^1(\partial V; \mathbb{C}^*)$. Let $d \in H^1(\partial V; \mathbb{C}/2i\pi\mathbb{Z})$ be the log of this class, with imaginary part in $]-\pi, \pi]$. One requires that for all $a \in H_1(\partial V; \mathbb{Z})$,

$$(2) \quad k_f(a) = d(a) \bmod(i\pi); \quad (k_f(a) - d(a))/i\pi = i^*(h_f)(a) \bmod(2).$$

Collecting the bulk and boundary weights we will often write \mathcal{P} as $(V, \rho, (h, k))$, where

$$(h, k) = ((h_f, h_c), (k_f, k_c)).$$

Notation. For every $n \in \mathbb{N}$, we write “ $a \equiv_n b$ ” to mean that a and b are equal up to multiplication by a power of $\exp(2i\pi/n)$. If n is odd, then $a \equiv_{2n} b$ if and only if $a \equiv_n \pm b$. We denote by μ_n the group of n -th roots of unity, acting on \mathbb{C} by multiplication.

In [4, 5], for every cusped manifold M and odd $N \geq 3$ we defined quantum hyperbolic invariants

$$(3) \quad \mathcal{H}_N(M) := \mathcal{H}_N(V, \rho_{\text{hyp}}, (0, 0)) \in \mathbb{C}/\mu_{2N}$$

that is, for the pattern $\mathcal{P} = (V, \rho_{\text{hyp}}, (0, 0))$ where ρ_{hyp} is the hyperbolic holonomy of M and all weights vanish. The following theorem summarizes our new results for patterns over M (all terms are defined in Section 4).

Theorem 1.1. *Let M be an arbitrary cusped manifold, $X(M)$ the variety of augmented $PSL(2, \mathbb{C})$ -characters of M , and $X_0(M) \subset X(M)$ the irreducible component of ρ_{hyp} . There is a canonical non empty Zariski open subset $\Omega(M)$ of $X_0(M)$ containing ρ_{hyp} such that:*

(1) *For every odd integer $N \geq 3$ and every pattern $(V, \rho, (h, k))$ such that $\rho \in \Omega(M)$, there is a quantum hyperbolic invariant $\mathcal{H}_N(V, \rho, (h, k)) \in \mathbb{C}/\mu_{2N}$ satisfying $\mathcal{H}_N(V, \rho_{\text{hyp}}, (0, 0)) = \mathcal{H}_N(M)$.*

(2) *(Analytic Families) Fix a topological support $(V, (h_c, k_c))$ and $h_f \in H^1(V; \mathbb{Z}/2\mathbb{Z})$. For every odd integer $N \geq 3$, the invariants $\mathcal{H}_N(V, \rho, (h, k))$ define a regular rational function $\mathcal{H}_N^{h_f, h_c, k_c} : \tilde{\Omega}(M)_N \rightarrow \mathbb{C}/\mu_{2N}$ on a determined $(\mathbb{Z}/N\mathbb{Z})^2$ -covering space $\tilde{\Omega}(M)_N$ of $\Omega(M)$.*

(3) *(Resolution of the sign ambiguity) If $N \equiv 1 \pmod{4}$, or $N \equiv 3 \pmod{4}$ and $h_c = 0$, then the statements (1) and (2) above hold true by replacing μ_{2N} with μ_N .*

Comments:

a) $X(M)$ is a complex algebraic variety and ρ_{hyp} a regular point of $X(M)$. Hence there is a unique irreducible component $X_0(M)$ of $X(M)$ containing ρ_{hyp} . As M has only one cusp, $X_0(M)$ is an algebraic curve and $\Omega(M)$ the complement of a finite set of points.

b) Theorem 1.1 (2) is a rough qualitative formulation of the N -th analytic family $\mathcal{A}_N(Y)$ associated to the topological support $Y = (V, (h_c, k_c))$ of M .

c) The arguments of Theorem 1.1 (1)-(2) apply verbatim to prove the invariance of simplicial formulas defining the $PSL(2, \mathbb{C})$ -Chern-Simons section, realized as an analytic equivariant function $\eta^* \mathcal{H}_1^{\mathcal{F}_{EP}, h_f} : \tilde{\Omega}(M)_\infty \rightarrow \mathbb{C}$ on a \mathbb{Z}^2 -covering $\tilde{\Omega}(M)_\infty$ of $\Omega(M)$ (see Section 4.4).

1.2. QHI of other patterns. In order to explain the nature of the QHI in Theorem 1.1 (1), and why this result is not straightforward, it is useful to recall a few general facts from [3, 4, 5].

QHFT partition functions. The QHI of QHFT patterns have been defined in [5]. The topological supports of QHFT patterns have the form $(V, L, (h_c, k_c))$ where:

- V is a compact oriented connected 3-manifold with (possibly empty) boundary ∂V made by torus components; if $\partial V = \emptyset$ we will use the notation W instead of V .
- L is a *non-empty* link in the interior of V ;
- the bulk & boundary c -weights $(h_c, k_c) \in H^1(V; \mathbb{Z}/2\mathbb{Z}) \times H^1(\partial V; \mathbb{Z})$ satisfy (1).

The QHFT patterns are obtained by completing $(V, L, (h_c, k_c))$ with a couple $(\rho, (h_f, k_f))$, where ρ is any $PSL(2, \mathbb{C})$ -character of V and (h_f, k_f) are bulk and boundary f -weights satisfying (2) with respect to ρ .

When $V = W$ is a closed 3-manifold, (k_c, k_f) disappears so that $h = (h_c, h_f) \in H^1(W; \mathbb{Z}/2\mathbb{Z})^2$. In [3, 4] we defined the QHI $\mathcal{H}_N(W, L, \rho, h)$ in that situation, for every character ρ and weight h . A specialization is $H_N(S^3, L) := \mathcal{H}_N(S^3, L, \rho_{\text{triv}}, 0)$, where ρ_{triv} is the trivial character of S^3 . In [6] we obtained the following result, which establishes a connection with Jones invariants.

Theorem 1.2. *For every link L in S^3 and every odd integer $N \geq 3$ we have*

$$H_N(S^3, L) \equiv_N \langle L \rangle_N = J_N(L)(e^{2i\pi/N})$$

where $\langle L \rangle_N$ is the link invariant defined by the enhanced Yang-Baxter operator extending the Kašaevev R -matrix, and $J_N(L)(q) \in \mathbb{Z}[q^{\pm 1}]$ is the colored Jones polynomial, normalized so that $J_N(K_U)(q) = 1$ on the unknot K_U .

Remark 1.3. The second equality is due to [32]. In [3, 4, 5] we quoted occasionally the first one as a motivating fact. Later we realized that we were unable to derive a complete proof from the existing literature (in particular [24, 25]), so we provided an independent one in [6], under the ambiguity \equiv_{2N} . The above statement with \equiv_N follows from the state sum sign correction introduced in the present paper (see also Remarks 2.11 and 5.5).

When $\partial V \neq \emptyset$ the QHFT partition functions are more sophisticated (see [5]). In particular the link L contains an essential simple closed curve on each boundary component which actually encodes a Dehn filling instruction. Anyway, also in this case the invariants $\mathcal{H}_N(V, L, \rho, (h, k))$ are defined for arbitrary characters and weights.

Relation with the QHI of cusped manifolds. The patterns over cusped manifolds and the QHFT patterns are complementary in the sense that the link L is *empty* in the former. In [4, 5] the proof of invariance of $\mathcal{H}_N(M) = \mathcal{H}_N(V, \rho_{\text{hyp}}, (0, 0))$ differs to many extents from the one for the QHFT partition functions. It uses the “volume rigidity” for cusped manifolds (see eg. [20]), Thurston’s hyperbolic Dehn filling theorem, a construction of certain auxiliary invariants $\mathcal{H}_N(V, \rho_{\text{hyp}}, (0, 0), a)$ that depend a priori on an additional datum “ a ”, and finally a surgery formula. Set $H_N(W, L, \rho) := \mathcal{H}_N(W, L, \rho, 0)$, where W is closed (see our notations above). By combining all these results we proved:

Theorem 1.4. ([5], Section 6.2) *Let W_n be a sequence of closed hyperbolic Dehn fillings of M whose holonomies ρ_n (considered as $PSL(2, \mathbb{C})$ -characters on M) converge to ρ_{hyp} in $X(M)$. Denote by L_n the geodesic core of the solid torus that fills V to produce W_n . For every odd $N \geq 3$ and every additional datum “ a ” we have*

$$(4) \quad \lim_{n \rightarrow \infty} H_N(W_n, L_n, \rho_n) \equiv_{2N} \mathcal{H}_N(V, \rho_{\text{hyp}}, (0, 0), a).$$

Hence “ a ” is eventually immaterial, and the limit defines $\mathcal{H}_N(M) \in \mathbb{C}/\mu_{2N}$.

The normalization $(h, k) = (0, 0)$ on the right side of (4) is a by-product of the proof. Under some additional assumptions on M (for instance if M is “very gentle” according to [4, 5]; then “ a ” = \emptyset),

we could avoid the delicate surgery argument and define the invariants $\mathcal{H}_N(V, \rho_{\text{hyp}}, h, k)$ for arbitrary weights relative to the hyperbolic holonomy.

1.3. State sums over weakly branched triangulations. One complication with the construction of the QHI of cusped manifolds in [4, 5] depended on a technical difficulty that we overcome in the present paper, and that we are going to illustrate.

For every topological support $Y_{\mathcal{P}}$, denote by \hat{V} the compact space obtained by filling each boundary component of ∂V with the cone over it. If $\partial V = \emptyset$ then $\hat{V} = W$; \hat{V} has a finite set of non-manifold points, the vertices of the filling cones. For every pattern \mathcal{P} supported by $Y_{\mathcal{P}}$, $\mathcal{H}_N(\mathcal{P})$ is computed by state sums over certain “decorated” triangulations T of \hat{V} , depending on the choice of a point z_{ρ} in the associated gluing variety $G(T)$ such that z_{ρ} represents the character ρ (see Section 2), and on a suitable encoding of the weights (h, k) . Moreover, T is equipped with a *branching*; equivalently, T carries a structure of Δ -*complex* in the sense of [22].

In the case of QHFT partition functions this is not so demanding: T can be a “quasi-regular” triangulation (every edge has distinct endpoints), a branching b can be induced for example by a total ordering of the vertices, and z_{ρ} can be realized by means of a so-called “idealization” of $PSL(2, \mathbb{C})$ -valued 1-cocycles on (T, b) .

On another hand, for a cusped manifold M we use *ideal* triangulations T of \hat{V} such that the gluing variety $G(T)$ contains a point z_h representing the hyperbolic holonomy ρ_{hyp} , and having coordinates with non-negative imaginary part (we say that z_h is *non-negative*). Such triangulations exist for every M , for instance any maximal subdivision of the canonical Epstein-Penner cell decomposition has this property. However we do not know if every cusped manifold has such a triangulation T admitting a branching b . For instance, the canonical Epstein-Penner decomposition of the “figure-8-knot sister” (m003 in Snappea’s census) is made of two regular hyperbolic ideal tetrahedra and does not carry any branching. The “very gentle” manifolds M mentioned after Theorem 1.4 admit *by definition* a branched triangulation (T, b) with a non-negative point z_h in $G(T)$.

In order to get Theorem 1.1 (1) we relax branchings to *weak branchings* which exist on every triangulation, and this leads us to include *2-face tensors* in the state sum formulas. In this setup, as well as to cover arbitrary characters of M in $\Omega(M)$, the proof of the state sum invariance requires additional arguments with respect to [4, 5].

1.4. Plan of the paper. Let V be as in Section 1.2 and \hat{V} as in Section 1.3.

In Section 2 we recall a few general facts about triangulations endowed with *pre-branchings*, *weak branchings*, or *branchings*, and the associated *gluing varieties* of M .

In Section 3 we construct the *analytic configuration* $\mathcal{A}_N(T, \tilde{b}, c)$ for every odd integer $N \geq 3$, weakly branched triangulation (T, \tilde{b}) of \hat{V} , and *rough charge* c on T (suitably specialized “global charges” will eventually encode the c -weights below). In particular $\mathcal{A}_N(T, \tilde{b}, c)$ contains an infinite Abelian covering of the gluing variety, $p_{\infty} : G(T, \tilde{b})_{\infty} \rightarrow G(T)$, and an analytic function $\mathcal{H}_N(T, \tilde{b}, c) : G(T, \tilde{b})_{\infty} \rightarrow \mathbb{C}$. In the case of QHFT patterns it is described qualitatively by:

Proposition 1.5. *For every topological support $(V, L, (h_c, k_c))$ there is a weakly branched triangulation (T, \tilde{b}) of \hat{V} and a global charge c on T such that:*

- (1) c encodes the c -weight (h_c, k_c) .
- (2) For every QHFT pattern $(V, L, \rho, (h, k))$ with topological support $(V, L, (h_c, k_c))$, there is a point $u \in G(T, \tilde{b})_{\infty}$ such that $p_{\infty}(u)$ represents the character ρ and for every odd $N \geq 3$ the scalar $\mathcal{H}_N(V, L, \rho, (h, k)) \equiv_{2N} \mathcal{H}_N(T, \tilde{b}, c)(u)$ does not depend on the choice of (T, \tilde{b}, c) and u .

In the case of patterns over a cusped manifold M , we have:

Proposition 1.6. *For every topological support $Y = (V, (h_c, k_c))$ over M , there is a determined Zariski open subset $\Omega(M)$ of $X_0(M)$ containing the hyperbolic holonomy ρ_{hyp} , and there is a weakly branched ideal triangulation (T, \tilde{b}) of \hat{V} , and a global charge c on T such that:*

- (1) c encodes the c -weight (h_c, k_c) .
- (2) The gluing variety $G(T)$ contains a non-negative point z_h representing ρ_{hyp} .

- (3) *There is an irreducible component Z of $G(T)$, a Zariski open subset Ω_Z of Z containing z_h , and a homeomorphism $\rho : \Omega_Z \rightarrow \Omega(M)$ extending a regular rational isomorphism between Zariski open subsets, such that $\rho(z)$ is the holonomy of V represented by z .*
- (4) *For every pattern $(V, \rho, (h, k))$ over M with topological support Y and for every $\rho \in \Omega(M)$, there is a point $u \in Z_\infty := p_\infty^{-1}(Z)$ such that $\rho = \rho(p_\infty(u))$, and for every odd $N \geq 3$ the scalar $\mathcal{H}_N(V, \rho, (h, k)) \equiv_{2N} \mathcal{H}_N(T, \tilde{b}, c)(u)$ does not depend on the choice of (T, \tilde{b}, c) and u .*

We will use concrete models of the finite coverings and regular rational maps in Theorem 1.1 (2) by considering a suitable factorization of $\mathcal{H}_N(T, \tilde{b}, c) : Z_\infty \rightarrow \mathbb{C}$.

In Section 4 we develop the content of Proposition 1.6 and prove Theorem 1.1 (1)-(2) and the analogous result when $N = 1$ (see the comment (c) and Corollary 4.17).

In Section 5 we indicate briefly how to deal with QHFT partition functions.

In Section 6 we collect a few facts about a diagrammatic calculus for weakly branched triangulations. This calculus is used in Section 7, which contains the invariance proof of the state sums defined over such triangulations.

In Section 8 we prove Theorem 1.1 (3) and the analogous result for QHFT partition functions. As a by product we show that global compensations of the local sign ambiguities imply that the QHI $\mathcal{H}_N(\mathcal{P})$ defined by means of branched triangulations have no sign ambiguity. This applies to QHFT partition functions and to very gentle cusped manifolds.

In Section 9 we describe the quantum hyperbolic invariants of three cusped manifolds: the figure-eight knot complement, its “sister”, and the complement of the knot 5_2 . We present the results of numerical computations showing that the functions $\mathcal{H}_N^{h_f, h_c, k_c}$ depend on the weights as $N \rightarrow +\infty$, and recover the volume for some specific choices of the weights.

Notations. In general we denote by N the odd integers ≥ 3 , including the case $N = 1$ when specified, and we put $m := (N - 1)/2$ and $\zeta := \exp(2\pi i/N)$. The set $\mathcal{I}_N = \{0, \dots, N - 1\}$ is identified to the group $\mathbb{Z}/N\mathbb{Z}$ and $[n]_N \in \mathcal{I}_N$ denotes the residue modulo N of $n \in \mathbb{N}$. We let $\delta_N(n) := 1$ (resp. 0) if $[n]_N = 0$ (resp. $[n]_N \neq 0$).

Acknowledgments. The first author’s work was supported by the ANR projects *Géométrie et Topologie Quantiques* (ANR-08-JCJC-0114-01) and “Extensions des théories de Teichmüller-Thurston” (ANR-09-BLAN-0116-01). Many thanks are due to Columbia University, Fukuoka University, the MSC at Tsinghua University and the University of Utah at Moab where parts of this work was presented. The second author’s work is supported by the italian FIRB project *Geometry and topology of low-dimensional manifolds*. The numerical computations presented in Section 9 were obtained by using the Maple software.

2. STRUCTURED TRIANGULATIONS AND GLUING VARIETIES

Triangulations. Let V be as in Section 1.2, that is, a compact oriented connected 3-manifold with (possibly empty) boundary ∂V made by toric components. Denote by \hat{V} the space obtained by taking the cone over each boundary component of ∂V . A triangulation T of \hat{V} is a collection of oriented tetrahedra $\Delta_1, \dots, \Delta_s$ together with a complete system \sim of pairings of their 2-faces via orientation reversing affine isomorphisms, such that the oriented quotient space

$$T := \coprod_{i=1}^s \Delta_i / \sim$$

is homeomorphic to \hat{V} , preserving the orientations. We will distinguish between the 2-faces, edges and vertices of the disjoint union $\coprod_{i=1}^s \Delta_i$, and the ones of T after the 2-face pairings. In particular we denote by $E(\{\Delta_i\})$ and $E(T)$ the set of edges of $\coprod_{i=1}^s \Delta_i$ and T respectively, and we write $E \rightarrow e$ to mean that an edge $E \in E(\{\Delta_i\})$ is identified to $e \in E(T)$ under the 2-face pairings. The 2-faces of each tetrahedron Δ_i have the boundary orientation defined by the rule: “*first the outgoing normal*”. When $\partial V \neq \emptyset$, the non-manifold points of \hat{V} are necessarily vertices of T . A triangulation T of \hat{V} is called *ideal* if the set of vertices of T coincides with the set of non-manifold points of \hat{V} .

Gluing varieties. Let T be as above. For every tetrahedron Δ_j choose a vertex v^j . Order the edges of the opposite 2-face F^j so that the induced cyclic ordering is the *opposite* of the boundary orientation. Denote these edges by E_0^j, E_1^j, E_2^j . Give a label $u_r^j \in \mathbb{C}_*$ to E_r^j and the opposite edge, where $\mathbb{C}_* := \mathbb{C} \setminus \{0, 1\}$ and $r \in \{0, 1, 2\}$. The gluing variety of T is the algebraic subset of \mathbb{C}_*^{3s} with coordinates $(u_0^1, u_1^1, u_2^1, \dots, u_0^s, u_1^s, u_2^s)$ and defining equations (r is mod(3)):

$$\forall j \in \{1, \dots, s\}, r \in \{0, 1, 2\}, u_{r+1}^j(1 - u_r^j) = 1$$

$$\forall e \in E(T), \prod_{E \rightarrow e} u(E) = 1.$$

By the first set of s ‘‘tetrahedral relations’’, we see that the gluing variety is the graph of an explicit regular rational map defined on an algebraic subset of \mathbb{C}_*^s defined by the second set of ‘‘edge relations’’. The auxiliary choices of ordered edges E_r^j being immaterial, these algebraic varieties are canonically isomorphic. We denote them by $G(T)$.

Every point $u \in G(T)$ represents a $PSL(2, \mathbb{C})$ -character $\rho(u)$ of V . Its components u_0^j, u_1^j, u_2^j can be interpreted as the cross-ratio parameters of an isometry class of oriented hyperbolic ideal tetrahedra associated to Δ_j , that we denote by (Δ_j, u^j) . Their imaginary parts have a same sign $\epsilon^j \in \{-1, 0, 1\}$ (by convention $\epsilon^j = 0$ if the imaginary parts are zero). By a classical result of Schläfli, the algebraic volume of (Δ_j, u^j) is given by

$$(5) \quad \text{Vol}_{\text{alg}}(\Delta_j, u^j) := \epsilon(u^j) \text{Vol}(\Delta_j, u^j) = D_2(u_0^j)$$

where Vol is the geometric positive volume and D_2 the Bloch-Wigner dilogarithm. When the components u_r^j are real, (Δ_j, u^j) is degenerate and both sides of (5) vanish. By summing the algebraic volumes of the (Δ_j, u^j) s we get a *volume function*

$$(6) \quad \text{Vol} : G(T) \rightarrow \mathbb{R}.$$

If $G(T)$ is non empty, for every point $u \in G(T)$, $\text{Vol}(u)$ coincides with the (intrinsically defined) volume $\text{Vol}(\rho(u))$ of the character $\rho(u)$. In general $G(T)$ might be empty, but \hat{V} has always triangulations T such that $G(T)$ is non trivial. A first general result concerns its dimension.

Theorem 2.1. ([35], [33]; see also [8]) *Assume that V has one torus boundary component. Let T be an ideal triangulation of \hat{V} . If the gluing variety $G(T)$ is non empty, then it is a complex algebraic set of dimension ≥ 1 .*

This result depends on the combinatorial properties of T . If the interior of V is a cusped manifold M , then it has the canonical Epstein-Penner (EP) cell decomposition by embedded convex hyperbolic ideal polyhedra (see for instance [8]). Then we have:

Proposition 2.2. *The maximal subdivisions of the EP cell decomposition of M define a finite set $\mathcal{T}_{EP}(M)$ of ideal triangulations of \hat{V} , such that for every $T \in \mathcal{T}_{EP}(M)$ the gluing variety $G(T)$ contains a non-negative point $u_h \in G(T)$ such that $\rho(u_h) = \rho_{\text{hyp}}$, and $\text{Vol}(u_h) = \text{Vol}(M)$.*

Every non-degenerate hyperbolic ideal tetrahedron of (T, u_h) has strictly positive (geometric) volume, but in general one cannot avoid some degenerate tetrahedra.

For other manifolds the nature of $G(T)$ is not so well-known. For instance consider the case of a closed 3-manifold W . A triangulation T of W is called *quasi-regular* if every edge of T has distinct vertices in T . It is clear that every triangulation has a quasi-regular subdivision. Take one, and fix a total ordering of the vertices. For every edge e with endpoints v and v' , orient e from v to v' if $v < v'$. Every simplicial $PSL(2, \mathbb{C})$ -valued 1-cocycle z on T , defined by using this edge orientation, represents a character $\rho(z)$ of W . Then, to any sufficiently generic cocycle z one can associate a point $u \in G(T)$ such that $\rho(z) = \rho(u)$, and we have ([3]):

Proposition 2.3. *Let T be a quasi-regular triangulation of W . For every character ρ of W there is a point $u \in G(T)$ such that $\rho = \rho(u)$.*

Clearly, the point u is far to be unique. For instance, if $W = S^3$ every point of $G(T)$ represents the trivial character. A similar, slightly more elaborated result holds for all other topological supports of QHFT patterns; it uses triangulations of \hat{V} obtained from quasi-regular relative triangulations $(T, \partial T)$ of $(V, \partial V)$ by adding a cone over each component of ∂T .

The method used to prove Proposition 2.3 is reminiscent of Thurston's spinning construction, and is strictly related to it when W is hyperbolic. In that case, the following result, which is proved by using the spinning construction, agrees with Proposition 2.3 in the case of quasi regular triangulations.

Proposition 2.4. ([30]) *Let T be a triangulation of a closed oriented hyperbolic 3-manifold W such that no edge is a null-homotopic loop in W . Then there exists $u \in G(T)$ such that $\text{Vol}(u) = \text{Vol}(W)$, and moreover its holonomy $\rho(u)$ is the hyperbolic holonomy.*

Variations on branched triangulations. Define a *pre-branched tetrahedron* (Δ, σ) as an oriented tetrahedron Δ with a choice σ of co-orientations of the 2-faces, such that two co-orientations are ingoing and two are outgoing. As every 2-face has the boundary orientation, by duality σ can be interpreted as a system of 2-face orientations.




FIGURE 1. Pre-branched tetrahedron.

Figure 1 shows a pre-branched tetrahedron (Δ, σ) embedded in \mathbb{R}^3 , with coordinates (x_1, x_2, x_3) such that the plane of the picture is $\{x_3 = 0\}$. We put on Δ the orientation induced from \mathbb{R}^3 , and assume that the two 2-faces above (resp. below) the plane are those with outgoing (resp. ingoing) co-orientations. This specifies two *diagonal edges* and four *square edges*. Every square edge is oriented as the common boundary edge of two 2-faces with opposite co-orientations. So the square edges form an oriented quadrilateral. Using the orientation of Δ , one can also distinguish among the square edges two pairs of opposite edges, called *A-edges* and *B-edges*. The orientation of the diagonal edges is not determined.

An oriented tetrahedron Δ becomes a 3-simplex by ordering its vertices. This is equivalent to a system b of orientations of the edges, called (local) *branching*, such that the vertex v_j has j incoming edges ($j \in \{0, \dots, 3\}$). The 2-faces of (Δ, b) are ordered as the opposite vertices, and b induces a branching b_F on each 2-face F . The branchings b and b_F define orientations on Δ and F respectively, the *b-* and *b_F-orientations*. The *b-* and *b_F-orientations* may coincide or not with the orientation of Δ . We encode this by a sign, $*_b \in \{-1, +1\}$. The boundary orientation and the b_F -orientation agree on two 2-faces. Hence b defines a pre-branching σ_b . On another hand, given a pre-branching σ on Δ there are exactly four branchings such that $\sigma_b = \sigma$. They can be obtained by choosing an *A* (resp. *B*) edge, reversing its orientation, and completing the resulting square edge orientations to define a branching b (this can be done in a single way; see Figure 2). Note that $*_b = 1$ (resp. $*_b = -1$) if and only if we have chosen an *A* (resp. *B*) square edge, and the square edge is $[v_0, v_3]$. The diagonal edges are $[v_0, v_2]$ and $[v_1, v_3]$.

One can extend these notions to triangulations T of \hat{V} :

- A *pre-branched triangulation* (T, σ) is formed by pre-branched tetrahedra (Δ_j, σ_j) such that the 2-face co-orientations match under the 2-face pairings.
- A *weakly-branched triangulation* (T, \tilde{b}) is formed by branched tetrahedra (Δ_j, b_j) such that the induced pre-branched tetrahedra (Δ_j, σ_{b_j}) form a pre-branched triangulation (T, σ) .
- A *branched triangulation* (T, b) is formed by branched tetrahedra (Δ_j, b_j) such that the branchings (ie. the edge orientations) match under the 2-face pairings.




FIGURE 2. Branched tetrahedra inducing the same pre-branched tetrahedron.

Remark 2.5. Branched triangulations of \hat{V} and Δ -complexes over \hat{V} [22] are equivalent notions. In particular the simplicial 3-chain $\sum_j *b_j(\Delta_j, b_j)$ represents the fundamental class in $H_3(\hat{V}; \mathbb{Z})$.

Let T be a triangulation of \hat{V} . Denote by \bar{V} the compact 3-manifold with boundary obtained by removing a small open 3-ball around every vertex of T which is a manifold point. Clearly, $V = \bar{V}$ if and only if T is an ideal triangulation. A pre-branched triangulation (T, σ) of \hat{V} can be described in a very concrete way in terms of the standard spine P of \bar{V} dual to T :

Lemma 2.6. *There is a 1-to-1 duality correspondence between the sets of pre-branchings of T and those of P , where a prebranching of P is defined as an orientation of the singular locus $\text{Sing}(P)$ such that every vertex has two outgoing and two ingoing edges.*

The proof is evident, as every edge of $\text{Sing}(P)$ is dual to a 2-face of T . The notion of (weakly) branched triangulation (T, \bar{b}) has a natural dual counterpart as (weakly) branched spine (P, \bar{b}) .

The branched boundary of a pre-branched triangulation. The triangulation T of \hat{V} yields a decomposition of \bar{V} into truncated tetrahedra. Their triangular 2-faces form a triangulation ∂T of $\partial \bar{V}$. If (T, σ) is a pre-branched triangulation, then ∂T has a branching $b_{\partial \sigma}$ defined on the 2-faces of ∂T as in Figure 3.




FIGURE 3. Branched boundary of pre-branched triangulations.

The corners of every 2-simplex of $(\partial T, b_{\partial\sigma})$ have a label in $\{A, B, \emptyset\}$, and a sign ± 1 obtained by comparing the boundary and the $b_{\partial\sigma}$ -orientations. Such a $\{A, B, \emptyset\}$ -labelling may be defined more generally for any branched triangulation of a closed oriented surface.

Lemma 2.7. *Each vertex of a branched triangulation of a closed oriented surface has an even number of adjacent corners with the empty label \emptyset . Hence every edge e of a pre-branched triangulation (T, σ) of \hat{V} has an even number of diagonal edges $E \in E(\{\Delta_i\})$ such that $E \rightarrow e$.*

Proof. For each 2-face the edges adjacent to the corner with label \emptyset have opposite b -orientations, while they agree elsewhere. The existence of an orientation on any transverse loop implies the first claim. The second follows by considering the triangulation ∂T . \square

Remark 2.8. Give the labels $1 \in \mathbb{Z}/2\mathbb{Z}$ to the diagonal edges and $0 \in \mathbb{Z}/2\mathbb{Z}$ to the square edges of every pre-branched tetrahedron (Δ, σ) of (T, σ) . Then Lemma 2.7 and Lemma 2.10 below imply that \mathbb{Z}_2 -valued *taut* structures on T always exist (a notion borrowed from F. Luo's work).

Networks and \mathcal{N} -graphs. Consider an oriented graph $(\text{Sing}(P), \bar{\sigma})$ as in Lemma 2.6. Put around every vertex v the dual branched tetrahedron (Δ_v, b_v) so that its 2-faces intersect transversely the edges of $\text{Sing}(P)$. Thus each edge connects a 2-face F^i of the ‘‘initial’’ branched tetrahedron (Δ_{v_i}, b_{v_i}) with a 2-face F^f of the ‘‘final’’ one (Δ_{v_f}, b_{v_f}) , identified in T . The gluing map $\phi_e : F^i \rightarrow F^f$ is determined by a color $r(e) \in \mathbb{Z}/3\mathbb{Z}$ defined as follows. Set $J_3 = \{0, 1, 2\}$. Denote by $S(J_3)$ the symmetric group on J_3 , by $A(J_3)$ the subgroup of even permutations, and by u_j^i and u_j^f the vertices of F^i and F^f ($j \in J_3$). The map ϕ_e is determined by the permutation $\tau_e \in A(J_3)$ such that $\phi_e(u_j^i) = u_{\tau_e(j)}^f$. Then we put

$$r(e) := \alpha^{-1}(\tau_e) \in \mathbb{Z}/3\mathbb{Z}$$

where $\alpha : \mathbb{Z}/3\mathbb{Z} \rightarrow A(J_3)$ is the isomorphism given by $\alpha(j) = (012)^j$. We define $\mathcal{N}(T, \tilde{b})$ as the oriented graph $(\text{Sing}(P), \bar{\sigma})$ endowed with the $\mathbb{Z}/3\mathbb{Z}$ -edge colors $r(e)$, and with the correspondence, for every vertex v , between the 2-faces of (Δ_v, b_v) and the germs of edges adjacent to v . Clearly we have:

Lemma 2.9. *A weakly branched triangulation (T, \tilde{b}) is branched if and only if all the $\mathbb{Z}/3\mathbb{Z}$ -edge colors $r(e)$ are equal to 0.*




Lemma 2.10. *Any triangulation T of \hat{V} admits pre-branchings and hence compatible weak-branchings.*

We stress that this is no longer true for genuine branchings (see an example in Figure 7). If T is quasi-regular, then every total ordering of the vertices induces a branching of T .

It is useful to represent $\mathcal{N}(T, \tilde{b})$ by a \mathcal{N} -graph Γ , defined as follows. Basically Γ is a planar immersion of $(\text{Sing}(P), \bar{\sigma})$ with normal crossings. It has two kinds of vertices: *essential crossings*, represented by a solid dot, corresponding to the vertices of P , and connected by the oriented edges of Γ , and non essential (‘‘immaterial’’) crossings due to the immersion. An essential crossing encodes a branched tetrahedron (Δ_v, b_v) as shown in Figure 4; the labels on the arc endpoints correspond to the b_v -ordering of the dual 2-faces of (Δ_v, b_v) . A full decoding is provided by Figure 5 in the case $*_b = 1$, showing a branched tetrahedron, the dual ‘‘butterfly’’ with co-oriented (hence oriented) wings, and its conversion into a portion of a oriented branched surface. An arc of Γ corresponding to an edge e of $(\text{Sing}(P), \bar{\sigma})$ inherits the $\mathbb{Z}/3\mathbb{Z}$ -color $r(e)$. If $r(e) = 0$ we omit it.


Given Γ , an easy decoding procedure (extending the one of Figure 5) produces an embedding in \mathbb{R}^3 of a closed regular neighbourhood $N(P)$ of $\text{Sing}(P)$ in the standard spine P . We believe that it is enough to show this in the case of the simplest cusped manifolds: M_0 , the figure-eight-knot complement in S^3 , and its ‘‘sister’’ M_1 , which is the complement of a knot in the lens space $L_{5,1}$ (see [31]). In both cases the Epstein-Penner decomposition is an ideal triangulation made by two regular hyperbolic ideal tetrahedra. Denote by T_0 and T_1 the corresponding triangulations of \hat{V}_0 and \hat{V}_1 . Figure 6 (resp. Figure 7) shows an \mathcal{N} -graph and its decoding for a branching b of T_0 (resp. weak branching \tilde{b} of T_1). It is not hard to verify that T_1 *does not carry any branching*.

Edge decorations. We will use several labellings of the edges $E \in E(\{\Delta_i\})$ of a weakly branched triangulation, called *decorations*, such that for every tetrahedron opposite edges have the same label.

FIGURE 4. \mathcal{N} -graph crossings for $*_b = \pm 1$.FIGURE 5. Decoding of an \mathcal{N} -graph crossing ($*_b = 1$).FIGURE 6. A \mathcal{N} -graph of (T_0, b) and its decoding.

An example is the decoration of the square edges with A or B according to the pre-branching, and of the diagonal edges with \emptyset . Every decoration d is determined on each tetrahedron Δ_j by the triple $(d_0, d_1, d_2) := (d(E_0^j), d(E_1^j), d(E_2^j))$. In terms of decoded \mathcal{N} -graphs, they are placed as in Figure 8, where we understand that the over/under crossing arcs are labelled by d_0 , and we show also A, B, \emptyset .

Remark 2.11. In the case of genuine branchings, \mathcal{N} -graphs were used in [6] under the name of *normal o-graphs*, with the opposite convention for the sign $*_b$ (in accordance with the usual crossing

FIGURE 7. A \mathcal{N} -graph of (T_1, \tilde{b}) and its decoding.FIGURE 8. Edge decorations on \mathcal{N} -graphs.

signs of link diagrams). This choice gives the equality $H_N(S^3, L) \cong_N J_N(\bar{L})$, where \bar{L} is the mirror image of the link L . The present convention yields the statement of Theorem 1.2.

The model $G(T, \tilde{b})$ of the gluing variety. If (T, \tilde{b}) is a weakly branched triangulation, we can use the weak branching \tilde{b} to fix the auxiliary choices used in the definition of the gluing variety $G(T)$. On every branched tetrahedron (Δ_j, b_j) let $v^j = v_3^j$, and order the edges of F_3^j as $E_0^j = [v_0^j, v_1^j]$, $E_1^j = [v_1^j, v_2^j]$, and $E_2^j = [v_0^j, v_2^j]$. If $*_{b_j} = +1$, this ordering is compatible with the cyclic ordering induced by the opposite of the boundary orientation of F_3^j . If $*_{b_j} = -1$ this is no longer true, and since the coordinates u_r^j , $r \in \{0, 1, 2\}$, of $G(T)$ are the cross-ratio moduli of an isometry class of oriented hyperbolic tetrahedron associated to Δ_j , u_r^j should be replaced by $(u_r^j)^{-1}$ in order to compensate the choice of opposite orientation. Hence we define a new system of coordinates $(w_r^j) \in \mathbb{C}_*^{3s}$ of $G(T)$ by labelling E_r^j with w_r^j , setting

$$(7) \quad (w_0^j, w_1^j, w_2^j) = \begin{cases} (u_0^j, u_1^j, u_2^j) & \text{if } *_{b_j} = +1 \\ (1/u_2^j, 1/u_1^j, 1/u_0^j) & \text{if } *_{b_j} = -1. \end{cases}$$

This gives us the model of $G(T)$ that we are going to use. We denote it by $G(T, \tilde{b})$. The defining equations of $G(T, \tilde{b})$ are the edge relations, for all edges e of T , given by

$$(8) \quad \prod_{E \rightarrow e} w(E)^{*E} = 1$$

where if E is an edge of (Δ_j, b_j) we have $w(E) = w_i^j$ if and only if E is E_i^j or the opposite edge, and $*_E := *_{b_j}$. The volume function on $G(T, \tilde{b})$ takes now the form $\text{Vol}(w) = \sum_j *_{b_j} D_2(w_0^j)$. Clearly $\text{Vol}(u) = \text{Vol}(w(u))$.

Example 2.12. It is easy to recover the edge equations of $G(T, \tilde{b})$ from a decoded \mathcal{N} -graph representing it: at each essential crossing one places the cross-ratio variables (w_0, w_1, w_2) like the decorations (d_0, d_1, d_2) in Figure 8, and take the products of cross-ratio variables along the boundary lines of $N(P)$. For example, consider the cusped manifold M_1 and (T_1, \tilde{b}) . Assign (w_0, w_1, w_2) to the top

crossing of Figure 7, and (W_0, W_1, W_2) to the bottom one. Note that $*_b = +1$ for both of them. Then we get the equations

$$(9) \quad w_0 w_1^2 W_0 W_1^2 = 1, \quad w_0 w_2^2 W_0 W_2^2 = 1.$$

Using the relation $w_{j+1} = 1/(1 - w_j)$ and the similar one for W_j , they reduce to the unique quadratic equation

$$w_1(w_1 - 1)W_1(W_1 - 1) = 1.$$

The parameter space of “positive solutions” is the half plane $\text{Im } w_1 > 0$ with the ray $0.5 + si$ removed, where $\sqrt{15}/2 \leq s < +\infty$ [?]. The complete hyperbolic structure is realized at $w_1 = W_1 = \exp(i\pi/3)$ (two regular ideal tetrahedra). Similarly, using (T_0, b) one recovers Thurston’s celebrated treatment of the figure-eight knot complement M_0 .

3. ANALYTIC CONFIGURATIONS

3.1. Local analytic configurations. Take an oriented 3-simplex (Δ, b) . As in the previous section the 2-face F_3 is opposite to the vertex v_3 , and the edges of F_3 are ordered as

$$E_0 = [v_0, v_1], \quad E_1 = [v_1, v_2], \quad E_2 = [v_0, v_2].$$

3.1.1. Quantum hyperbolic 3-simplices. Recall that the edge decorations of (Δ, b) are equal on opposite edges, and hence specified by triples $d = (d_0, d_1, d_2)$ where $d_r = d(E_r)$ ($r \in \{0, 1, 2\}$). A *quantum hyperbolic 3-simplex* is a tuple (Δ, b, w, f, c) where:

- $w = (w_0, w_1, w_2)$ is the system of cross ratios (7).
- $f = (f_0, f_1, f_2) \in \mathbb{Z}^3$ is such that $l_k = \log(w_k) + i f_k \pi$, $k \in \{0, 1, 2\}$, satisfies $l_0 + l_1 + l_2 = 0$.
- $c = (c_0, c_1, c_2) \in \mathbb{Z}^3$ satisfies $c_0 + c_1 + c_2 = 1$.

Here, \log is the branch of logarithm with imaginary part in $(-\pi, \pi]$. For every $k \in \{0, 1, 2\}$ set

$$(10) \quad l_{k,N,*_b,c} = \frac{1}{N}(\log(w_k) + i\pi(N+1)(f_k - *_b c_k)).$$

We call f a *flattening*, c a *charge*, l_k a *classical log-branch*, and $l_{k,N,*_b,c}$ a *quantum log-branch*. If $\text{Im}(w) > 0$ (resp. $\text{Im}(w) < 0$), then c is a charge if and only if $-c$ (resp. c) is a flattening.

We organize the edge decorations w, f, c as follows. Denote by $p_\infty : \mathcal{W}_\infty \rightarrow \mathbb{C}_*$ the maximal abelian covering map. We realize \mathcal{W}_∞ as the quotient space $\mathcal{W}_\infty = (\mathcal{D} \times \mathbb{Z}^2) / \sim$, where \mathcal{D} is the result of gluing two copies of $(-\infty, 0)$ and $(1, +\infty)$ to the boundary of $\mathbb{C} \setminus ((-\infty, 0) \cup (1, +\infty))$, and \sim is the equivalence relation

$$(11) \quad \begin{aligned} (x + i0; p, q) &\sim (x - i0; p + 2, q) && \text{if } x \in (-\infty, 0) \\ (x + i0; p, q) &\sim (x - i0; p, q + 2) && \text{if } x \in (1, +\infty). \end{aligned}$$

Setting $l(y; m) := \log(y) + im\pi$, $m \in \mathbb{Z}$, the bijective map

$$(12) \quad \left(x; \frac{l(x; p) - \log(x)}{i\pi}, \frac{l((1-x)^{-1}; q) - \log((1-x)^{-1})}{i\pi}\right) \rightarrow (l(x; p), l((1-x)^{-1}; q))$$

identifies \mathcal{W}_∞ with the Riemann surface of the maps

$$(13) \quad \phi_{\epsilon, \epsilon'} : x \rightarrow (\log(x) + i\epsilon\pi, \log((1-x)^{-1}) + i\epsilon'\pi) \quad , \quad \epsilon, \epsilon' \in \{0, 1\}.$$

For every sign $* = \pm 1$ and couple $c = (c_0, c_1) \in \mathbb{Z}^2$ define the analytic map $l_{N,*_b,c} : \mathcal{W}_\infty \rightarrow (\mathbb{C}^*)^2$ by

$$l_{N,*_b,c}([x; p, q]) = \left(\frac{1}{N}(\log(x) + i\pi(N+1)(p - *_b c_0)), \frac{1}{N}(\log((1-x)^{-1}) + i\pi(N+1)(q - *_b c_1))\right).$$

Then:

- \mathbb{C}_* (resp. \mathcal{W}_∞) with coordinate w_0 (resp. $[w_0; f_0, f_1]$) is a parametrization of the set of edge decorations w (resp. (w, f)) of (Δ, b) .
- Setting $c_2 = 1 - c_0 - c_1$, the components of $l_{N,*_b,c}([w_0; f_0, f_1])$ determine the three quantum log-branches $l_{k,N,*_b,c}$ of (Δ, b, w, f, c) .

- Put $\text{EXP}(z_1, \dots, z_{2s}) = (\exp(z_1), \dots, \exp(z_{2s}))$, $s \geq 1$. The map $\text{EXP} \circ l_{N,*,c} : \mathcal{W}_\infty \rightarrow (\mathbb{C}^*)^2$ provides a distinguished system of N th-roots of the cross ratios w_k :

$$(14) \quad w'_k := \exp(l_{k,N,*,c}([w_0; f_0, f_1])).$$

The proof of the following Lemma is straightforward (see [4] and [5], Remark 2.1 and 2.18).

Lemma 3.1. (1) The N th-roots (w'_0, w'_1, w'_2) depend only on the residues mod (N) of the flattening f and the charge c , and verify the relation

$$(15) \quad w'_0 w'_1 w'_2 = -\zeta^{*b m}$$

where as usual $N := 2m + 1$ and $\zeta := \exp(2i\pi/N)$.

(2) For any charge c and N -th roots u_k of the cross ratios w_k verifying the relation (15), there is a flattening f such that $u_k = w'_k$ as in (14).

(3) The image \mathcal{W}_N of \mathcal{W}_∞ via the map $\text{EXP} \circ l_{N,*,c}$ is the curve in $(\mathbb{C}^*)^2$ with defining equation

$$u_0^N + (u_1^{-1})^N = 1.$$

There is a natural N^2 -to-1 rational regular map $p_N : \mathcal{W}_N \rightarrow \mathbb{C}_*$, $p_N(u_0, u_1) := u_0^N$.

3.1.2. *Tetrahedral tensors.* Given a branched tetrahedron (Δ, b) and an operator $A \in \text{End}(\mathbb{C}^N \otimes \mathbb{C}^N)$, associate a copy V_j of \mathbb{C}^N to the 2-face F_j , and consider A as a map

$$(16) \quad A = \begin{cases} (A_{k,l}^{i,j}) : V_3 \otimes V_1 \rightarrow V_2 \otimes V_0 & \text{if } *b = +1 \\ (A_{i,j}^{k,l}) : V_2 \otimes V_0 \rightarrow V_3 \otimes V_1 & \text{if } *b = -1. \end{cases}$$

Note that the pair of indices (i, j) (resp. (k, l)) corresponds to the space $V_2 \otimes V_0$ (resp. $V_3 \otimes V_1$), and to the outgoing (resp. ingoing) 2-face co-orientations induced by b if $*b = 1$, and the converse if $*b = -1$. The entries $A_{k,l}^{i,j}$ and $A_{i,j}^{k,l}$ are taken in the standard basis of $\mathbb{C}^N \otimes \mathbb{C}^N$. Graphically this is encoded in Figure 9 (compare with Figure 4).




FIGURE 9. Position of tensor indices.

Using (16) we define the tetrahedral tensors $\mathcal{R}_N(\Delta, b, w, f, c)$ of a quantum hyperbolic 3-simplex by

$$\mathcal{R}_1(\Delta, b, w, f) := \mathcal{R}_{(1,*,c)}([w_0; f_0, f_1]) \quad \text{if } N = 1$$

and

$$\mathcal{R}_N(\Delta, b, w, f, c) := \mathcal{R}_{(N,*,c)} \circ \text{EXP} \circ l_{N,*,c}([w_0; f_0, f_1]) \quad \text{if } N \geq 3 \text{ is odd}$$

where $\mathcal{R}_{(1,*)} : \mathcal{W}_\infty \rightarrow \text{GL}(\mathbb{C} \otimes \mathbb{C}) \cong \mathbb{C}^*$ and $\mathcal{R}_{(N,*,c)} : \mathcal{W}_N \rightarrow \text{GL}(\mathbb{C}^N \otimes \mathbb{C}^N)$ are defined as follows. The map $\mathcal{R}_{(1,*)}$ is analytic. It is given by

$$(17) \quad \mathcal{R}_{(1,*)}([x; p, q]) := \exp\left(*\frac{2}{i\pi} \left(L(x) + \frac{i\pi}{2} (p \log(1-x) + q \log(x)) \right)\right)$$

where L is the Rogers dilogarithm

$$L(x) = -\frac{\pi^2}{6} - \frac{1}{2} \int_0^x \frac{\log(t)}{1-t} + \frac{\log(1-t)}{t} dt.$$

The maps $\mathcal{R}_{(N,*,c)}$ are regular rational maps called *matrix dilogarithms*. They verify non commutative versions of the five term relations satisfied by L. For every $x \in \mathbb{C}^*$, put $x^{1/N} := \exp(\log(x)/N)$, extended to $0^{1/N} := 0$ by continuity. Set

$$[x] := N^{-1} \frac{1-x^N}{1-x}, \quad g(x) := \prod_{j=1}^{N-1} (1-x\zeta^{-j})^{j/N}, \quad h(x) := g(x)/g(1).$$

Let $\mathcal{F}_N = \{(u', v') \in \mathbb{C}^2 \mid (u')^N + (v')^N = 1\}$, and $\omega : \mathcal{F}_N \times \mathbb{Z}/N\mathbb{Z} \rightarrow \mathbb{C}$ the regular rational map given by $\omega(u', v'|0) := 1$ and $\omega(u', v'|n) := \prod_{j=1}^{[n]_N} v'(1-u'\zeta^j)^{-1}$. Define $\mathcal{L}_N : \mathcal{F}_N \rightarrow \text{Aut}(\mathbb{C}^N \otimes \mathbb{C}^N)$ by (recall that $m = (N-1)/2$)

$$(18) \quad \mathcal{L}_N(u', v')_{k,l}^{i,j} = h(u') \zeta^{kj+(m+1)k^2} \omega(u', v'|i-k) \delta_N(i+j-l).$$

We have

$$(\mathcal{L}_N(u', v')^{-1})_{i,j}^{k,l} = \frac{[u']}{h(u')} \zeta^{-kj-(m+1)k^2} \frac{\delta_N(i+j-l)}{\omega(u'/\zeta, v'|i-k)}.$$

We put

$$(19) \quad \begin{aligned} \mathcal{R}_{(N,+1,c)}(u_0, u_1)_{k,l}^{i,j} &:= (u_0^{-c_1} u_1^{c_0})^{\frac{N-1}{2}} \mathcal{L}_N(u_0, u_1^{-1})_{k,l}^{i,j} \\ \mathcal{R}_{(N,-1,c)}(u_0, u_1)_{k,l}^{i,j} &:= (u_0^{-c_1} u_1^{c_0})^{\frac{N-1}{2}} (\mathcal{L}_N(u_0, u_1^{-1})^{-1})_{i,j}^{k,l}. \end{aligned}$$

Remark 3.2. The map $\mathcal{R}_{(1,*)}$ does not depend on the charge c and $\mathcal{R}_{(N,*,c)}$ depends on c only by the factor $(u_0^{-c_1} u_1^{c_0})^{(N-1)/2}$, but \mathcal{R}_N depends on c also via the map $l_{N,*,c}$.

Definition 3.3. *The local analytic configurations over (Δ, b, c) are the family of spaces and maps indexed by the odd integers $N \geq 3$, given by*

$$\mathcal{A}_N(\Delta, b, c) := \{\mathcal{W}_\infty, \mathcal{W}_N, p_\infty, p_N, l_{N,*,c}, \mathcal{R}_{(1,*)}, \mathcal{R}_{(N,*,c)}\}.$$

3.2. Globalization. In addition to the tetrahedral tensors R_N , when dealing with arbitrary triangulations we need also *face tensors* \mathcal{Q}_N , defined as follows. For every odd $N \geq 3$, consider the symmetric $N \times N$ matrices S and T with entries

$$S_j^i = N^{-1/2} \zeta^{ij}, \quad T_j^i = \zeta^{i^2(m+1)} \delta_N(i+j).$$

We show in Lemma 7.3 that S and T generate an N -dimensional projective representation of $SL(2, \mathbb{Z})$. We put $\mathcal{Q}_N := S \cdot T^{-1}$ (see Section 7.1 for the notation “.”). This $N \times N$ matrix has entries

$$(\mathcal{Q}_N)_j^i = \sum_{k=0}^{N-1} S_j^k (T^{-1})_k^i = N^{-\frac{1}{2}} \zeta^{-ij-(m+1)j^2},$$

and is projectively of order 3, namely $\mathcal{Q}_N^3 = \phi_N^{-1} \text{I}_N$ with $\phi_N = \left(\frac{m+1}{N}\right)$ if $N \equiv 1 \pmod{4}$, and $\phi_N = \left(\frac{m+1}{N}\right) i$ if $N \equiv 3 \pmod{4}$.

Let (T, \tilde{b}) be a weakly branched triangulation of \hat{V} made of s branched tetrahedra (Δ_j, b_j) . For every collection of charges c^j on the Δ_j s and every point $[w; f] := ([w_0^1; f_0^1, f_1^1], \dots, [w_0^s; f_0^s, f_1^s]) \in \mathcal{W}_\infty^s$ we call c a *rough global charge* and (T, \tilde{b}, w, f, c) a *rough QH triangulation*. Working dually, consider the oriented graph \mathcal{N} associated to (T, \tilde{b}) as before Lemma 2.9. For every odd $N \geq 1$ define a *QH tensor network* over \mathcal{N} as follows:

- associate $R_N(\Delta_j, b_j, w^j, f^j, c^j)$ to the structured vertex v_j of \mathcal{N} dual to (Δ_j, b_j) ;
- if $N \geq 3$, associate $\mathcal{Q}_N^{r(e)}$ to each edge e of \mathcal{N} with $\mathbb{Z}/3\mathbb{Z}$ -color $r(e)$, so that the domain and target of the linear map associated to $\mathcal{Q}_N^{r(e)}$ correspond respectively to the initial and final endpoints of e ; if $N = 1$, replace $\mathcal{Q}_N^{r(e)}$ by the scalar $1 \in \mathbb{C}^*$.

Next we define a normalization factor $a_N(T, \tilde{b})$. Formally put $a_1(T, \tilde{b}) = 1$. Denote by $q(T, \tilde{b})$ the number of edges e of $\text{Sing}(P)$ such that $r(e) = 2 \in \mathbb{Z}/3\mathbb{Z}$. Fix an auxiliary orientation on every edge

e of T . Let $n_+(e)$ (resp. $n_-(e)$) be the number of abstract *diagonal* edges $E \rightarrow e$ such that the \tilde{b} -orientation of E agrees (resp. does not agree) with the orientation of e . By Lemma 2.7, $n_+(e) - n_-(e)$ is an even integer. Set $\epsilon_N := (-1)^{\frac{N-1}{2}}$. For every odd $N \geq 3$ define

$$(20) \quad a_N(T, \tilde{b}) := N^{-v} \phi_N^{-q(T, \tilde{b})} c_N(T, \tilde{b}), \quad c_N(T, \tilde{b}) := \epsilon_N^{v+l-\frac{1}{2} \sum_e (n_+(e) - n_-(e))}$$

where ϕ_N is as above, and l, v are respectively the number of edges and the number of vertices of T which are manifold points.

Remark 3.4. Clearly $c_N(T, \tilde{b})$ does not depend on the choice of the auxiliary edge orientations, and $c_N(T, \tilde{b}) = 1$ if $N - 1 \equiv 0 \pmod{4}$. If \tilde{b} is a branching b , using the edge b -orientation we see that $l - \frac{1}{2} \sum_e (n_+(e) - n_-(e)) = l - t = \chi(\tilde{V}) = \chi(P) = -v$, where χ denotes the Euler characteristic and t the number of tetrahedra of T . Hence in this case $c_N(T, b) = 1$, and $a_N(T, b) = N^{-v}$.

A N -state of \mathcal{N} is an assignment of a label in $\{0, \dots, N-1\}$ to each edge; every N -state σ determines a matrix element, denoted by $R_N(\Delta_j, b_j, w^j, f^j, c^j)_\sigma$ or $\mathcal{Q}_{N, \sigma}^{r(e)}$, of each tensor of the QH tensor network.

Definition 3.5. The *QH state sum function* $\mathcal{H}_N(T, \tilde{b}, c) : \mathcal{W}_\infty^s \rightarrow \mathbb{C}$ is the total contraction of the QH tensor network over \mathcal{N} , normalized by $a_N(T, \tilde{b})$:

$$(21) \quad \mathcal{H}_N(T, \tilde{b}, c)([w; f]) := a_N(T, \tilde{b}) \sum_\sigma \prod_j R_N(\Delta_j, b_j, w^j, f^j, c^j)_\sigma \prod_e \mathcal{Q}_{N, \sigma}^{r(e)}.$$

Sometimes we denote $\mathcal{H}_N(T, \tilde{b}, c)([w; f])$ by $\mathcal{H}_N(\mathcal{T})$, where $\mathcal{T} := (T, \tilde{b}, w, f, c)$, and when the context is clear, \mathcal{H}_N by $\mathcal{H}_N(T, \tilde{b}, c)$. Note that \mathcal{H}_1 is a product of analytic scalar functions and does not depend on c . When $N \geq 3$, $\mathcal{H}_N(T, \tilde{b}, c) = \mathcal{H}'_N(T, \tilde{b}, c) \circ (\text{EXP} \circ l_{N, *, c})$ where $\mathcal{H}'_N(T, \tilde{b}, c)$ is a rational regular function defined on the algebraic variety \mathcal{W}_N^s .

In order to make (21) actual invariant state sums we need to restrict $[w; f] \in \mathcal{W}_\infty^s$ to a suitable subspace, that we denote $G_0(T, \tilde{b})_\infty$ below (later we will do similarly for charges, see Definition 4.7 and 5.1). Consider the model $G(T, \tilde{b})$ of the gluing variety, and the products of the maps of Section 3.1 over the Δ_j s (keeping, with slight abuse, the same notations):

$$p_\infty : \mathcal{W}_\infty^s \rightarrow \mathbb{C}_*^s, \quad p_N : \mathcal{W}_N^s \rightarrow \mathbb{C}_*^s \\ l_{N, *, c} : \mathcal{W}_\infty^s \rightarrow (\mathbb{C}^*)^{2s}, \quad \text{EXP} \circ l_{N, *, c} : \mathcal{W}_\infty^s \rightarrow \mathcal{W}_N^s \subset (\mathbb{C}^*)^{2s}.$$

Set

$$G(T, \tilde{b})_\infty := p_\infty^{-1}(G(T, \tilde{b})), \quad G(T, \tilde{b}, c)_N := \text{EXP} \circ l_{N, *, c}(G(T, \tilde{b})_\infty).$$

The restrictions of p_∞ and p_N to these spaces are covering maps of $G(T, \tilde{b})$, that make a commutative diagram with $\text{EXP} \circ l_{N, *, c} : G(T, \tilde{b})_\infty$ is a closed analytic subset of \mathcal{W}_∞^s , $G(T, \tilde{b}, c)_N$ an algebraic subset of \mathcal{W}_N^s , and $\text{Sing}(G(T, \tilde{b})_*) = p_*^{-1}(\text{Sing}(G(T, \tilde{b})))$.

Define the *total cross-ratio* of an edge e of T by (see the notations at the end of Section 2)

$$W(e) := \prod_{E \rightarrow e} w(E)^{*E}$$

so that $[w; f] \in G(T, \tilde{b})_\infty$ if and only if $W(e) = 1$ for all e . Similarly, define the *total log-branch*, *total N -th root modulus*, and *total charge* at e by

$$(22) \quad L(e) := \sum_{E \rightarrow e} *E l(E), \quad W'_N(e) := \prod_{E \rightarrow e} w'(E)^{*E}, \quad C(e) := \sum_{E \rightarrow e} c(E)$$

where $l(E) = \log(w_k^j) + i f_k^j \pi$, $w(E) = w_k^j$ and $c(E) = c_k^j$ if and only if E is an edge of (Δ_j, b_j, w^j) , E is either E_k^j or the opposite edge, and $*E = *_{b_j}$. Fix an auxiliary ordering of the, say n , edges of (T, \tilde{b}) . We have analytic maps

$$t_L : \mathcal{W}_\infty^s \rightarrow \mathbb{C}^n, \quad t_L([w; f]) := (L(e_1), \dots, L(e_n))$$

$$t_{N, W', c} : \mathcal{W}_\infty^s \rightarrow \mathbb{C}^n, \quad t_{N, W', c}([w; f]) := (W'_N(e_1), \dots, W'_N(e_n))$$

the dependence of $L(e_k)$ (resp. $W'_N(e_k)$) on $[w; f]$ (resp. $[w; f]$ and c) being understood. The following Lemma is evident.

Lemma 3.6. *The restriction of t_L to $G(T, \tilde{b})_\infty$ takes values in $(2i\pi\mathbb{Z})^n$ and the restriction of $t_{N,W',c}$ takes values in the set μ_N of N th-roots of unity. Both maps are discrete-valued, hence constant on the closure of each connected component of the set of non-singular points of $G(T, \tilde{b})_\infty$. Moreover, the restriction of $t_{N,W',c}$ factorizes as*

$$t_{N,W',c} = t'_{N,W} \circ (\text{EXP} \circ l_{N,*b,c})$$

where $t'_{N,W}$ is a rational regular map defined on $G(T, \tilde{b}, c)_N$ and constant on irreducible components.

Finally define the analytic subset

$$G_0(T, \tilde{b})_\infty = t_L^{-1}(0, \dots, 0) \subset G(T, \tilde{b})_\infty$$

and, for every N , the algebraic subvariety

$$G_0(T, \tilde{b}, c)_N = \text{EXP} \circ l_{N,*b,c}(G_0(T, \tilde{b})_\infty) \subset G(T, \tilde{b}, c)_N.$$

As usual, keep the same notation for the restriction of any map already defined.

Definition 3.7. The N -th analytic configuration over (T, \tilde{b}, c) is the family of spaces and maps

$$\mathcal{A}_N(T, \tilde{b}, c) := \{G_0(T, \tilde{b})_\infty, G_0(T, \tilde{b}, c)_N, p_\infty, p_N, l_{N,*b,c}, \mathcal{H}_1, \mathcal{H}_N\}.$$

We stress that every map $t_{N,W',c}$ is constant on $G_0(T, \tilde{b})_\infty$, and its value depends on the fixed rough charge c , precisely on the total edge charges $(C(e_1), \dots, C(e_n))$.

4. CUSPED MANIFOLDS

Let M be a cusped manifold, and V be as in Section 1.1, that is, a compact oriented connected 3-manifold with one torus boundary component such that M is diffeomorphic to the interior of V . We use the notation \hat{V} as in Section 2.

4.1. The augmented $PSL(2, \mathbb{C})$ -character variety and A -polynomial. Fix a geometric basis (l, m) of the fundamental group $\pi_1(\partial V) \cong \mathbb{Z} \times \mathbb{Z}$. It is given by a couple of oriented simple closed curves on ∂V which meet at one point, transversely and positively. If M is the complement of a hyperbolic knot K in S^3 we can take a canonical longitude l and a meridian m of K .

There is a conjugacy class $[\sigma]$ of representations of $\pi_1(\partial V)$ in $PSL(2, \mathbb{C})$, each one with images isomorphic to $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$, acting on the Riemann sphere $\mathbb{CP}^1 = \partial\mathbb{H}^3$ without a common fixed point. The class $[\sigma]$ is obtained as follows: pick two geodesic lines in \mathbb{H}^3 , say γ_l and γ_m , that meet at one point forming a right angle; then define σ by $\sigma(x) = r_x$, where $x \in \{l, m\}$ and r_x is the rotation by π around γ_x .

By definition, the $PSL(2, \mathbb{C})$ -character variety of M is the algebro-geometric quotient

$$X'(M) := \text{Hom}'(\pi_1(M), PSL(2, \mathbb{C})) // PSL(2, \mathbb{C})$$

where $PSL(2, \mathbb{C})$ acts by conjugation, and

$$\text{Hom}'(\pi_1(M), PSL(2, \mathbb{C})) \subset \text{Hom}(\pi_1(M), PSL(2, \mathbb{C}))$$

is the subset of representations that do not restrict to an element of $[\sigma]$ on $\pi_1(\partial V)$. Given a peripheral subgroup $\pi_1(\partial V)$ of $\pi_1(M) \cong \pi_1(V)$, denote by

$$R(M) \subset \text{Hom}(\pi_1(V), PSL(2, \mathbb{C})) \times \mathbb{CP}^1$$

the set of couples (r, z) such that z is fixed by $r(\pi_1(\partial V))$. In particular, couples such that $[r]$ restricts to $[\sigma]$ on ∂V are excluded. The *augmented $PSL(2, \mathbb{C})$ -character variety* of M is the algebro-geometric quotient

$$X(M) := R(M) // PSL(2, \mathbb{C})$$

where $PSL(2, \mathbb{C})$ acts by conjugation on $R(M)$ and by Möbius transformations on \mathbb{CP}^1 . Hence, any *augmented character* of M , $\rho = [(\rho, z)] \in X(M)$, is a character of representations $r : \pi_1(M) \rightarrow PSL(2, \mathbb{C})$ together with a choice of fixed points of the peripheral subgroups, invariant by conjugation. The hyperbolic holonomy of M defines a point $\rho_{\text{hyp}} \in X(M)$. In a similar way we can define the $PSL(2, \mathbb{C})$ -character variety $X'(\partial V)$ of ∂V and its augmented version $X(\partial V)$. The inclusion $i : \partial V \rightarrow V$ induces regular maps $i^* : X'(M) \rightarrow X'(\partial V)$ and $i^* : X(M) \rightarrow X(\partial V)$.

Theorem 4.1. (1) ([17, 11]; see [37]) Both $X'(M)$ and $X(M)$ are complex algebraic affine varieties, and the natural projection $q : X(M) \rightarrow X'(M)$ is a regular map.

(2) ([23]) ρ_{hyp} (resp. $q(\rho_{\text{hyp}})$) is a regular point of $X(M)$ (resp. $X'(M)$). Hence it belongs to a unique irreducible component $X_0(M)$ of $X(M)$ (resp. $X'_0(M)$ of $X'(M)$), which is a complex algebraic curve.

(3) The restriction of q to $X_0(M)$, $q_0 : X_0(M) \rightarrow X'_0(M)$, is generically $2 : 1$.

(4) ([18]) The restricted map $i^* : X'_0(M) \rightarrow X'_0(\partial V)$ (resp. $i^* : X_0(M) \rightarrow X_0(\partial V)$) is generically $1 : 1$. Hence it is a birational isomorphism onto its image.

For any $\rho \in X(\partial V)$ which is non trivial, let $\bar{\rho}$ be a representative of ρ such that $\bar{\rho}(\pi_1(\partial V))$ fixes $z = \infty \in \mathbb{CP}^1$. For any non zero class $\gamma \in \pi_1(\partial V)$, $\bar{\rho}(\gamma)$ acts on \mathbb{C} as $w \mapsto \gamma_\rho w + b$, where $\gamma_\rho \in \mathbb{C}^*$ and $b \in \mathbb{C}$. In general γ_ρ is a squared eigenvalue of $\bar{\rho}(\gamma) \in PSL(2, \mathbb{C})$; if $\bar{\rho}(\gamma)$ is loxodromic, then $\gamma_\rho \neq 1$, and the two reciprocally inverse eigenvalues are distinguished by the augmentation, which selects an endpoint, whence an orientation, of γ . Consider the so called *holonomy map*

$$(23) \quad \begin{array}{ccc} \text{hol}_\gamma : X(\partial V) & \longrightarrow & \mathbb{C}^* \\ \rho & \longmapsto & \gamma_\rho. \end{array}$$

By using the above cusp basis (l, m) we get an algebraic isomorphism

$$\text{hol}_m \times \text{hol}_l : X(\partial V) \rightarrow \mathbb{C}^* \times \mathbb{C}^*.$$

Define the rational map

$$\mathfrak{h} : X(M) \rightarrow \mathbb{C}^* \times \mathbb{C}^*, \quad \mathfrak{h} = (\text{hol}_m \times \text{hol}_l) \circ i^*.$$

Following [14], [16], and Dunfield's appendix in [10], consider the plane curve $A(M)$ defined as the closure of the 1-dimensional part of the image $\mathfrak{h}(X(M))$. The (suitably normalized) polynomial generating the ideal of $A(M)$ is by definition the *PSL(2, \mathbb{C}) A-polynomial* of M . We denote by $A_0(M)$, and call *geometric component of $A(M)$* , the closure of $\mathfrak{h}(X_0(M))$. It follows from Theorem 4.1 (4) that:

Corollary 4.2. *The restricted map $\mathfrak{h} : X_0(M) \rightarrow A_0(M)$ is a birational isomorphism.*

4.2. Rich components of $G(T)$. Let T be any ideal triangulation of \hat{V} such that the gluing variety $G(T)$ is non empty. There is a natural regular map (see eg. [16] or the appendix of [10])

$$(24) \quad \rho : G(T) \rightarrow X(M).$$

As $G(T)$ is a complex algebraic curve, $\mathfrak{h}(\rho(G(T)))$ is a union of irreducible components of the plane curve $A(M)$. Assume that $G(T)$ contains a *non-negative* point z_h such that $\rho(z_h) = \rho_{\text{hyp}}$. Recall that “non-negative” means that for every tetrahedron Δ_j of T the cross-ratios z_h^j have non-negative imaginary parts. The point z_h is not necessarily a regular point of the gluing variety, hence in general it could be contained in several irreducible components of $G(T)$.

Definition 4.3. An irreducible component of $G(T)$ is *rich* if it contains z_h and an infinite sequence of points z_n that converge to z_h (in the “strong” topology of $G(T)$ as an analytic space), and correspond to compact closed hyperbolic Dehn fillings of M that converge geometrically to M .

Proposition 4.4. *For every gluing variety $G(T)$ containing a non-negative point z_h with $\rho(z_h) = \rho_{\text{hyp}}$, the set of rich components of $G(T)$ is non empty and finite.*

Proof. This follows from the proof of the hyperbolic Dehn filling Theorem given in [36], and the fact that the algebraic curve $G(T)$ has a finite number of irreducible components. \square

Remark 4.5. In [35], the hyperbolic Dehn filling Theorem was proved assuming that M (allowing several cusps) has an ideal triangulation T such that $G(T)$ contains a *strictly positive* point z_h representing the hyperbolic holonomy. This proof uses also that z_h is a regular point of $G(T)$. By elaborating on the analysis of [35], the paper [15] has provided a proof of this regularity result, extended to every strictly positive point of $G(T)$. In [8], the proof of [35] was presented with some mild modifications avoiding the assumption that z_h is regular. Unfortunately, it did not give yet a complete proof of the hyperbolic Dehn filling Theorem based on gluing varieties, since it is not known if every

M has strictly positive geometric ideal triangulations. The paper [36] filled this gap; it uses any gluing variety $G(T)$ with a non-negative and possibly singular point z_h , which always exists; the proof uses also the arguments of [8]. In [29] it is proved that strictly positive geometric ideal triangulations exist “virtually”, that is, every cusped M has a finite covering \tilde{M} having such an ideal triangulation. On the other hand, passing to \tilde{M} increases the number of cusps.

Proposition 4.6. *Let Z be any rich component of $G(T)$. Then the restricted maps $\rho : Z \rightarrow X_0(M)$ and $\mathfrak{h} \circ \rho : Z \rightarrow A_0(M)$ are generically 1 : 1 and hence are both birational isomorphisms. More precisely, there are maximal non empty Zariski open subsets Ω_Z^0 and Ω_Z of Z , with $\Omega_Z^0 \subset \Omega_Z$, such that:*

- (1) $\rho(\Omega_Z)$ and $\rho(\Omega_Z^0)$ are Zariski open subsets of $X_0(M)$ (resp. $\mathfrak{h} \circ \rho(\Omega_Z)$ and $\mathfrak{h} \circ \rho(\Omega_Z^0)$ are Zariski open subsets of $A_0(M)$).
- (2) The restriction of ρ and $\mathfrak{h} \circ \rho$ to Ω_Z is 1 : 1 onto its image, and $z_h \in \Omega_Z$. The restriction of ρ and $\mathfrak{h} \circ \rho$ to Ω_Z^0 is a regular rational isomorphism onto its image.

Proof. This goes step by step as the proof of Theorem 3.1 in [18]. The key ingredients are the Gromov-Thurston “volume rigidity” for closed hyperbolic manifolds, the fact that the volume of representations yields a well defined function on the normalization of the smooth projective model of $A_0(M)$, and the existence on $X_0(M)$ of infinite sequences of points corresponding to the holonomies of compact closed hyperbolic Dehn fillings of M that converge geometrically to M . In the present situation this last fact is ensured by the definition of a rich component, and the volume of representations lifts to the function (6) on $G(T)$. \square

If T has a weak branching \tilde{b} the above discussion can be rephrased in terms of the gluing variety $G(T, \tilde{b})$ of Section 2. Recall that a system of cross ratios w^j on the branched tetrahedra (Δ_j, b_j) of (T, \tilde{b}) is non negative if $z^j = (w^j)^{*j}$ is non negative.

4.3. Refined analytic configurations for one-cusped manifolds. Let (T, \tilde{b}) be a weakly branched ideal triangulation of \hat{V} .

Definition 4.7. A *global charge* c on T is a rough global charge satisfying the following additional global constraint: *For every edge e of T , the total edge charge $C(e)$ is constant and equal to 2.*

For every global charge c the variety $G_0(T, b, c)_N$ has at every edge e of T the defining equation

$$(25) \quad W'(e) = \zeta^{-1}$$

where $\zeta := \exp(2\pi i/N)$. Note that $\zeta^{-1} = \zeta^{2m}$. We have:

Proposition 4.8. (1) *Every global charge c on (T, \tilde{b}) determines a c -weight (h_c, k_c) of V .*

(2) *For every c -weight (h_c, k_c) of V and every ideal triangulation (T, \tilde{b}) of \hat{V} , there exists a global charge c on (T, \tilde{b}) with c -weight equal to (h_c, k_c) .*

Proposition 4.9. *Assume that the gluing variety $G(T, \tilde{b})$ is non empty, and let $\rho : G(T, \tilde{b}) \rightarrow X(M)$ be as in (24).*

(1) *Every point $[w; f] \in G_0(T, \tilde{b})_\infty$ determines an f -weight (h_f, k_f) of $(V, \rho(w))$.*

(2) *For every point $\rho \in \rho(G(T, \tilde{b}))$ and every f -weight (h_f, k_f) of (V, ρ) , there exists a point $[w; f]$ of $G_0(T, \tilde{b})_\infty$ with f -weight equal to (h_f, k_f) .*

Proposition 4.8 and 4.9 are reformulations of results of Neumann [33, 34]. We recall below the constructions underlying the statements (1), which claim the existence of the pairs $(\gamma(c), \gamma'_2(c))$ and $(L([w; f]), \gamma'_2(f))$ in Definition 4.10. The statements (2) are much more subtle. For the sake of completeness, we recall the proofs at the end of the section (part of the material will be used in Section 4.4).

Denote by T_0 the cell complex obtained from T by removing an open cone neighborhood of its vertex. Hence T_0 is the result of gluing tetrahedra with truncated vertices, and the polyhedron underlying T_0 is homeomorphic to V . Denote by ∂T_0 the induced triangulation of ∂V . Represent any non zero class in $H_1(\partial V; \mathbb{Z})$ by *normal loops*, that is, a disjoint union of oriented essential simple

closed curves in ∂V , transverse to the edges of ∂T_0 and such that no curve enters and exits a 2-simplex by a same edge. The intersection of a normal loop, say C , with a 2-simplex F consists of a disjoint union of arcs, each of which turns around a vertex of F ; if F is a cusp section of the tetrahedron Δ of T , for every vertex v of F we denote by E_v the edge of Δ containing v , and by $*_v$ the branching sign of Δ . We write $C \rightarrow E_v$ to mean that some subarcs of C turn around v . We count them algebraically, by using the orientation of C : if there are s_+ (resp. s_-) such subarcs whose orientation is compatible with (resp. opposite to) the orientation of ∂V as viewed from v , then we set $\text{ind}(C, v) := s_+ - s_-$. For every point $[w; f] \in G_0(T, \tilde{b})_\infty$ and every global charge c on (T, \tilde{b}) , one defines cohomology classes $L([w; f]) \in H^1(\partial V; \mathbb{C})$ and $\gamma(c) \in H^1(\partial V; \mathbb{Z})$ by (C is a normal loop on ∂V):

$$(26) \quad \begin{aligned} L([w; f])([C]) &:= \sum_{C \rightarrow E_v} *_v \text{ind}(C, v) l(E_v) \\ &= \sum_{C \rightarrow E_v} *_v \text{ind}(C, v) (\log(w(E_v)) + \pi\sqrt{-1}f(E_v)) \end{aligned}$$

$$(27) \quad \gamma(c)([C]) := \sum_{C \rightarrow E_v} \text{ind}(C, v) c(E_v).$$

Additional classes $\gamma'_2(c), \gamma'_2(f) \in H^1(V; \mathbb{Z}/2\mathbb{Z})$ are defined similarly, by using normal loops in T and taking the sum mod(2) of the charges or flattenings, respectively, of the edges we face along the loops. The reduction mod(2) of $\gamma(c)$ coincides with the image of $\gamma'_2(c)$ under the map induced on cohomology by the inclusion $i : \partial V \rightarrow V$. Hence $r(\gamma(c)) = i^*(\gamma'_2(c))$. Also, denoting by $d_w \in H^1(\partial V; \mathbb{C}/2i\pi\mathbb{Z})$ the log of the linear part of the restriction of $\rho(w)$ to $\pi_1(\partial V)$, for all $a \in H_1(\partial V; \mathbb{Z})$ we have

$$(28) \quad L([w; f])(a) = d_w(a) \bmod(i\pi), \quad (L([w; f])(a) - d_w(a))/i\pi = i^*(\gamma'_2(f))(a) \bmod(2).$$

This proves that the following definition agrees with the defining constraints of weights (see (1) and (2) in the Introduction).

Definition 4.10. The f -weight (k_f, h_f) of a point $[w; f] \in G_0(T, \tilde{b})_\infty$, and the c -weight (k_c, h_c) of a global charge c on (T, \tilde{b}) , are defined respectively as the pairs of cohomology classes

$$(L([w; f]), \gamma'_2(f)) \in H^1(\partial V; \mathbb{C}) \times H^1(V; \mathbb{Z}/2\mathbb{Z})$$

and

$$(\gamma(c), \gamma'_2(c)) \in H^1(\partial V; \mathbb{Z}) \times H^1(V; \mathbb{Z}/2\mathbb{Z}).$$

Example 4.11. (*Example 2.12 continued*). Every point w of the gluing variety $G(T_1, \tilde{b})$ gives rise to a class $\exp(d_w) \in H^1(\partial V; \mathbb{C}^*)$. For a suitable choice of geometric basis (λ, μ) of $H_1(\partial V; \mathbb{Z})$ we have

$$\exp(d_w)(\lambda) = w_0 w_1 W_0^{-1} W_1^{-1}, \quad \exp(d_w)(\mu) = w_1^{-2} W_2^2.$$

Denote by f_j and $l_j := \log(w_j) + i\pi f_j$ (resp. F_j and $L_j := \log(W_j) + i\pi F_j$) the flattening and log-branch variables at the top (resp. bottom) crossing of Figure 7. The equations for the log-branch variables are (see (9))

$$(29) \quad l_0 + l_1 + l_2 = L_0 + L_1 + L_2 = 0, \quad l_0 + 2l_1 + L_0 + 2L_1 = 0, \quad l_0 + 2l_2 + L_0 + 2L_2 = 0.$$

For simplicity let us deal with the positive points w of the (unique) rich component of $G(T_1, \tilde{b})$ which contains the hyperbolic solution, $w_1 = W_1 = \exp(i\pi/3)$. For every such a positive point the equations for the flattening variables are

$$(30) \quad f_0 + f_1 + f_2 = F_0 + F_1 + F_2 = -1, \quad f_0 + 2f_1 + F_0 + 2F_1 = -2, \quad f_0 + 2f_2 + F_0 + 2F_2 = -2.$$

Indeed, under positivity of w the pairs of first two equations in the systems (30) and (29) are clearly equivalent one to each other. Although a bit subtler, this is true also for the remaining pairs of equations, because the consistency relations (9) imply that the sum of the arguments of the cross ratios around edges is *exactly* equal to 2π (see [8], Lemma E.6.1).

The equations for the charge variables are formally obtained from those for the flattening ones by replacing in (30) each f_j with $-c_j$ and each F_j with $-C_j$. Solving the systems we get

$$\begin{aligned} f_0 &= -(2f_1 + F_1 - F_2 + 1), & F_0 &= -(F_1 + F_2 + 1), & f_2 &= f_1 + F_1 - F_2 \\ c_0 &= -(2c_1 + C_1 - C_2 - 1), & C_0 &= -(C_1 + C_2 - 1), & c_2 &= c_1 + C_1 - C_2 \\ l_0 &= -(2l_1 + L_1 - L_2), & L_0 &= -(L_1 + L_2), & l_2 &= l_1 + L_1 - L_2. \end{aligned}$$

Let us turn to the boundary c -weight $k_c \in H^1(\partial V; \mathbb{Z})$ and, for every positive w as above, the boundary f -weight $k_f = k_f(w) \in H^1(\partial V; \mathbb{C})$. They are given by

$$\begin{aligned} k_c(\lambda) &= c_0 + c_1 - C_0 - C_1, & k_c(\mu) &= 2C_2 - 2c_1 \\ k_f(\lambda) &= l_0 + l_1 - L_0 - L_1, & k_f(\mu) &= 2L_2 - 2l_1. \end{aligned}$$

At the complete solution we have

$$k_f(\lambda)/i\pi = f_0 + f_1 - F_0 - F_1, \quad k_f(\mu)/i\pi = 2F_2 - 2f_1.$$

Hence $k_f/i\pi \in H^1(\partial V; \mathbb{Z})$, and $k_f(\mu)/i\pi = 0 \pmod{2}$. Moreover

$$\begin{aligned} f_0 &= k_f(\lambda)/i\pi - k_f(\mu)/2i\pi - 2f_1 - 1, & f_2 &= f_1 + k_f(\mu)/2i\pi - k_f(\lambda)/i\pi \\ F_0 &= k_f(\lambda)/i\pi - 3k_f(\mu)/2i\pi - 2f_1 - 1, & F_1 &= k_f(\mu)/i\pi - k_f(\lambda)/i\pi + f_1 \\ F_2 &= f_1 + k_f(\mu)/2i\pi. \end{aligned}$$

Hence for any given boundary f -weight k_f , there is a one-parameter family of flattenings that realize k_f , depending on the free parameter f_1 . Similarly for the charges we have

$$c_0 = k_c(\lambda) - k_c(\mu)/2 - 2c_1 + 1, \quad c_2 = c_1 + k_c(\mu)/2 - k_c(\lambda)$$

$$C_0 = k_c(\lambda) - 3k_c(\mu)/2 - 2c_1 + 1, \quad C_1 = k_c(\mu) - k_c(\lambda) + c_1, \quad C_2 = c_1 + k_c(\mu)/2$$

where the free parameter is now c_1 . By taking $f_1 = c_1 = 0$, the relevant pairs of parameters that enter the formulas of the tensors \mathcal{R}_N are

$$(f_0, f_1) = (k_f(\lambda)/i\pi - 1 - k_f(\mu)/2i\pi, 0), \quad (c_0, c_1) = (k_c(\lambda) + 1 - k_c(\mu)/2, 0)$$

$$(F_0, F_1) = (-1 + k_f(\lambda)/i\pi - 3k_f(\mu)/2i\pi, k_f(\mu)/i\pi - k_f(\lambda)/i\pi)$$

$$(C_0, C_1) = (1 + k_c(\lambda) - 3k_c(\mu)/2, k_c(\mu) - k_c(\lambda))$$

stresssing in this way a pure dependence on the weights. Finally, the state sums are

$$\mathcal{H}_N(T_1, \tilde{b}, w, W, f, F, c, C) = \sum_{i,j,l,k,I,J=0}^{N-1} \mathcal{R}_N(1, w, f, c)_{k,l}^{i,j} \mathcal{R}_N(1, W, F, C)_{j,i}^{I,J} (\mathcal{Q}^2)_I^l (\mathcal{Q})_J^k$$

where we quoted that $*_b = 1$ for both tetrahedra. \square

Let now Z be a rich component of $G(T, \tilde{b})$. Denote $Z_\infty := p_\infty^{-1}(Z)$ (an analytic subspace of $G_0(T, \tilde{b})_\infty$), and let $Z_{\infty,0}$ be the union of connected components of Z_∞ made of points $[w; f]$ such that

$$(31) \quad i^*(\gamma'_2(f)) = 0 \in H^1(\partial V; \mathbb{Z}/2\mathbb{Z}).$$

Recall the basis (l, m) of $\pi_1(\partial V)$. Because of (28), for every point $[w; f] \in Z_{\infty,0}$ we have

$$(32) \quad (e^{L([w;f])}(l), e^{L([w;f])}(m)) = \mathfrak{h} \circ \rho(w),$$

and for every global charge c on (T, \tilde{b}) , the formula

$$\begin{aligned}
L_{N,c}([w; f])(C) &:= \sum_{C \rightarrow E_v} *_v \text{ind}(C, v) l_{N, *_v, c}(E_v) \\
&= \sum_{C \rightarrow E_v} \frac{*_v}{N} \text{ind}(C, v) (\log(w(E_v)) + \pi\sqrt{-1}(N+1)(f(E_v) - *_v c(E_v))) \\
(33) \quad &= \frac{1}{N} (L([w; f])([C]) - \pi\sqrt{-1}(N+1)\gamma(c)([C])) \\
&\quad + \underbrace{\pi\sqrt{-1} \sum_{C \rightarrow E_v} *_v \text{ind}(C, v) f(E_v)}_{\in 2\mathbb{Z}}
\end{aligned}$$

yields a cohomology class $L_{N,c}([w; f]) \in H^1(\partial V; \mathbb{C}/2\pi i\mathbb{Z})$ such that

$$(34) \quad ((e^{L_{N,c}([w; f])(l)})^N, (e^{L_{N,c}([w; f])(m)})^N) = \mathfrak{h} \circ \rho(w).$$

If we allow $[w; f]$ to live in $Z_\infty \setminus Z_{\infty,0}$, the sum in (33) can be odd, and the equalities (32) & (34) hold true only up to signs in each component. More precisely, for all $[w; f] \in Z_\infty$ the sum in (33) is an integer expression of the residue class $\gamma'_2(f)([C]) \in \mathbb{Z}/2\mathbb{Z}$, so that we have a well-defined cohomology class

$$(35) \quad L_{N,c}([w; f]) \in H^1(\partial V; \mathbb{C}/2\pi i\mathbb{Z})$$

such that

$$L_{N,c}([w; f]) = \frac{1}{N} (L([w; f]) - \pi i(N+1)\gamma(c) + \pi\sqrt{-1} i^*(\gamma'_2(f))) \pmod{2\pi i\mathbb{Z}}$$

where, as usual, $i^* : H^1(V; \mathbb{Z}/2\mathbb{Z}) \rightarrow H^1(\partial V; \mathbb{Z}/2\mathbb{Z})$ is induced by the inclusion map $i : \partial V \rightarrow V$. The spaces $Z_{\infty,0}$ and $A_0(M)$ are related as follows (see Proposition 4.6). Let

$$A_0(M)_\infty := \pi_\infty^{-1}(A_0(M)), \quad A_0(M)_N := \text{EXP} \circ l_{N, k_c}(A_0(M)_\infty)$$

where

$$(36) \quad \pi_\infty : \tilde{\mathbb{C}}^2 \rightarrow (\mathbb{C}^*)^2, \quad ([\lambda; p], [\lambda; q]) \mapsto (\lambda, \mu)$$

is the universal covering map, and for every integral class $k_c \in H^1(\partial V; \mathbb{Z})$, define $l_{N, k_c} : \tilde{\mathbb{C}}^2 \rightarrow \mathbb{C}^2$ by

$$(37) \quad l_{N, k_c}([\lambda; p], [\mu; q]) := \left(\frac{1}{N} (\log(\lambda) + 2\pi\sqrt{-1}p - \pi\sqrt{-1}(N+1)k_c(l)), \right. \\ \left. \frac{1}{N} (\log(\mu) + 2\pi\sqrt{-1}q - \pi\sqrt{-1}(N+1)k_c(m)) \right).$$

The restricted map $\pi_\infty : A_0(M)_\infty \rightarrow A_0(M)$ and the map $\pi_N : A_0(M)_N \rightarrow A_0(M)$, $(u, v) \mapsto (u^N, v^N)$, define respectively a $\mathbb{Z} \times \mathbb{Z}$ - and a $\mathbb{Z}/N \times \mathbb{Z}/N$ -covering of $A_0(M)$.

Definition 4.12. The *classical* and *quantum log-holonomies* are the maps defined on $Z_{\infty,0}$ by the components of the classes $L([w; f]) \in H^1(\partial V; \mathbb{C})$ and $\exp \circ L_{N,c}([w; f]) \in H^1(\partial V; \mathbb{C}^*)$, $[w; f] \in Z_{\infty,0}$, in the basis (l, m) of $\pi_1(\partial V)$:

$$\begin{aligned}
\text{loghol}_\infty : \quad Z_{\infty,0} &\longrightarrow A_0(M)_\infty \\
[w; f] &\longmapsto (L([w; f])(l), L([w; f])(m)). \\
\text{loghol}_{N,c} : \quad Z_{\infty,0} &\longrightarrow A_0(M)_N \\
[w; f] &\longmapsto (e^{L_{N,c}([w; f])(l)}, e^{L_{N,c}([w; f])(m)}).
\end{aligned}$$

Note that (32) & (34) imply that loghol_∞ (resp. $\text{loghol}_{N,c}$) maps into $A_0(M)_\infty$ (resp. $A_0(M)_N$). The identification (12) of \mathcal{W}_∞^s as a product of Riemann surfaces, and the fact that $Z_\infty \subset G_0(T, \tilde{b})_\infty$ is an analytic subspace, imply that both loghol_∞ and $\text{loghol}_{N,c}$ are analytic maps.

Remark 4.13. If one removes the assumption (31), the maps loghol_∞ and $\text{loghol}_{N,c}$ extend to the whole of Z_∞ . Then loghol_∞ maps Z_∞ to the union of four connected components $A_0(M)_\infty^{\varepsilon,\varepsilon'} := (\pi_\infty^{\varepsilon,\varepsilon'})^{-1}(A_0(M))$, where $\pi_\infty^{\varepsilon,\varepsilon'} : \tilde{\mathbb{C}}^2 \rightarrow (\mathbb{C}^*)^2$, $([\lambda;p], [\mu;q]) \mapsto (\varepsilon\lambda, \varepsilon'\mu)$, $\varepsilon, \varepsilon' \in \{-1, +1\}$. The target of $\text{loghol}_{N,c}$ becomes the union of the spaces $\text{EXP} \circ l_{N,k_c}(A_0(M)_\infty^{\varepsilon,\varepsilon'})$. The component $A_0(M)_\infty = A_0(M)_\infty^{0,0}$ is a natural one: we explain in Section 4.4 its relation with the Chern-Simons line bundle over ∂V .

We deduce immediately from the formulas (26) and (37):

Lemma 4.14. *For every N and every global charge c on (T, \tilde{b}) we have a commutative diagram:*

$$\begin{array}{ccccc}
A_0(M) & \xleftarrow{\pi_N} & A_0(M)_N & & \\
\swarrow \mathfrak{h} \circ \rho & & \uparrow \text{loghol}_{N,c} & & \\
Z & \xleftarrow{p_\infty} & Z_{\infty,0} & & \\
\swarrow \mathfrak{h} \circ \rho & & \downarrow e^{l_{N,k_c}} & & \\
A_0(M) & \xleftarrow{\pi_\infty} & A_0(M)_\infty & & \\
& & \swarrow \text{loghol}_\infty & &
\end{array}$$

where l_{N,k_c} is defined by (37), using the charge boundary weight k_c , and we denote $\text{EXP} \circ l_{N,k_c}$ by $e^{l_{N,k_c}}$.

Let now \mathcal{F} be any finite family of rich components, possibly contained in different gluing varieties associated to different weakly branched triangulations of \hat{V} . Recall Proposition 4.6. Define the following non empty Zariski open subsets of $X_0(M)$:

$$\Omega_{\mathcal{F}} = \bigcap_{Z \in \mathcal{F}} \rho(\Omega_Z), \quad \Omega_{\mathcal{F}}^0 = \bigcap_{Z \in \mathcal{F}} \rho(\Omega_Z^0).$$

If $Z \in \mathcal{F}$ is a rich component of $G(T, \tilde{b})$, define

$$\Omega_{\mathcal{F}}(Z) := \rho|_Z^{-1}(\Omega_{\mathcal{F}}), \quad \Omega_{\mathcal{F}}^0(Z) := \rho|_Z^{-1}(\Omega_{\mathcal{F}}^0)$$

and

$$\Omega_{\mathcal{F}}(Z)_{\infty,0} := p_\infty^{-1}(\Omega_{\mathcal{F}}(Z)) \cap Z_{\infty,0}, \quad \Omega_{\mathcal{F}}(Z)_N := \text{EXP} \circ l_{N,*b,c}(\Omega_{\mathcal{F}}(Z)_{\infty,0}).$$

We have:

- $\Omega_{\mathcal{F}}^0 \subset \Omega_{\mathcal{F}}$.
- The restriction of ρ to $\Omega_{\mathcal{F}}(Z)$ is a rational regular map which is 1 : 1 onto $\Omega_{\mathcal{F}}$. Moreover, the non negative point $w_h^Z \in Z$ such that $\rho(w_h^Z) = \rho_{\text{hyp}}$ belongs to $\Omega_{\mathcal{F}}(Z)$.
- For every component $Z \in \mathcal{F}$, the restriction of ρ to $\Omega_{\mathcal{F}}^0(Z)$ is a rational regular isomorphism onto $\Omega_{\mathcal{F}}^0$.

The following Lemma is an immediate consequence of Proposition 4.9.

Lemma 4.15. *Let \mathcal{F} be any finite family of rich components of gluing varieties associated to weakly branched ideal triangulations of \hat{V} . For every component Z in \mathcal{F} , every point $\rho \in \Omega_{\mathcal{F}}$, and every f -weight (k_f, h_f) of (V, ρ) such that $i^*(h_f) = 0$, there is a point $[w; f]_Z \in \Omega_{\mathcal{F}}(Z)_{\infty,0}$ such that $(L([w; f]_Z), \gamma_2'(f)) = (k_f, h_f)$.*

Put

$$A_{0,\mathcal{F}}(M) := \mathfrak{h}(\Omega_{\mathcal{F}})$$

and

$$A_{0,\mathcal{F}}(M)_\infty := \pi_\infty^{-1}(A_{0,\mathcal{F}}(M)), \quad A_{0,\mathcal{F}}(M)_N := \text{EXP} \circ l_{N,k_c}(A_{0,\mathcal{F}}(M)_\infty).$$

Replacing $G(T, \tilde{b})$ with $\Omega_{\mathcal{F}}(Z) \subset Z \subset G(T, \tilde{b})$, and $G_0(T, \tilde{b})_\infty$ with $\Omega_{\mathcal{F}}(Z)_{\infty,0} \subset Z_{\infty,0} \subset G_0(T, \tilde{b})_\infty$, and taking the restrictions of each of the spaces and maps that form the sequence of analytic configurations $\{\mathcal{A}_N(T, \tilde{b}, c)\}$, one obtains a sequence of sub-configurations

$$\mathcal{A}_N(c, Z, \mathcal{F}) := \{\Omega_{\mathcal{F}}(Z)_{\infty,0}, \Omega_{\mathcal{F}}(Z)_N, p_\infty, p_N, l_{N,*b,c}, \mathcal{H}_1, \mathcal{H}_N\}.$$

By Lemma 4.14 it fits into a commutative diagram:

$$\begin{array}{ccccc}
& & & & \Omega_{\mathcal{F}}(Z)_N \\
& & & & \uparrow \mathcal{H}'_N \\
A_{0,\mathcal{F}}(M) & \xleftarrow{\pi_N} & A_{0,\mathcal{F}}(M)_N & \xleftarrow{\text{loghol}_{N,c}} & \mathbb{C} \\
& \searrow \mathfrak{h} \circ \rho & \downarrow \lambda & \nearrow e^{l_{N,*b,c}} & \\
& & \Omega_{\mathcal{F}}(Z) & \xleftarrow{p_\infty} & \Omega_{\mathcal{F}}(Z)_{\infty,0} \\
& \searrow \mathfrak{h} \circ \rho & \downarrow e^{l_{N,k_c}} & \nearrow \text{loghol}_\infty & \\
A_{0,\mathcal{F}}(M) & \xleftarrow{\pi_\infty} & A_{0,\mathcal{F}}(M)_\infty & \xleftarrow{\text{loghol}_\infty} & \mathbb{C} \\
& & & & \nearrow \mathcal{H}_1
\end{array}$$

The following two results show that if we fix the bulk weight h_f together with the c -weight (h_c, k_c) , then \mathcal{H}_1 and \mathcal{H}_N factor through loghol_∞ and $\text{loghol}_{N,c}$, and hence induce maps on $A_{0,\mathcal{F}}(M)_\infty$ and $A_{0,\mathcal{F}}(M)_N$, respectively.

Theorem 4.16. *Let \mathcal{F} be any finite family of rich components Z of gluing varieties $G(T_Z, \tilde{b}_Z)$ associated to weakly branched ideal triangulations (T_Z, \tilde{b}_Z) of \hat{V} . Let $\rho \in \Omega_{\mathcal{F}}$, the f -weight (h_f, k_f) and $[w; f]_Z \in \Omega_{\mathcal{F}}(Z)_{\infty,0}$ be as in Lemma 4.15, and (h_c, k_c) be any c -weight realized on the triangulation T_Z by a global charge c_Z . Then:*

(1) *For every odd $N \geq 3$ the scalar $\mathcal{H}'_N(\mathcal{P}) := \mathcal{H}_N(T_Z, \tilde{b}_Z, c_Z)([w; f]_Z)$ is an invariant of the pattern $\mathcal{P} = (V, \rho, h_f, k_f, h_c, k_c)$ up to multiplication by $2N$ -th roots of unity, and depends on $k_f \pmod{\pi i N}$ only.*

(2) *The scalar $\mathcal{H}_1(\mathcal{P}) := \mathcal{H}_1(T_Z, \tilde{b}_Z)([w; f]_Z)$ is an invariant of $\mathcal{P} = (V, \rho, h_f, k_f)$ up to multiplication by 6th roots of unity, and is defined exactly if \tilde{b}_Z is a branching.*

Proof. We have to show that, up to the phase ambiguities, $\mathcal{H}'_N(\mathcal{P})$ and $\mathcal{H}_1(\mathcal{P})$ do not depend on the encoding of \mathcal{P} , ie. the choice of (T_Z, \tilde{b}_Z, c_Z) , Z and $[w; f]_Z$. For that we use the QH transits of [3, 4, 5], that we adapt to weakly branched triangulations in Section 7, in the case $N \geq 3$; for $N = 1$ this adaptation follows easily from Section 4.2-4.3 of [4] and the argument of Proposition 8.2. As a result, given two realizations of the pattern \mathcal{P} , say:

- $[w; f]_Z \in \Omega_{\mathcal{F}}(Z)_{\infty,0}$, included into some configuration $\mathcal{A}_N(c_Z, Z, \mathcal{F})$, and
- $[w'; f']_{Z'} \in \Omega_{\mathcal{F}}(Z')_{\infty,0}$, included into another configuration $\mathcal{A}_N(c_{Z'}, Z', \mathcal{F})$,

there exists a finite sequence of QH transits relating $[w; f]_Z$ to $[w''; f'']_{Z''} \in \Omega_{\mathcal{F}}(Z'')_{\infty,0}$ included into a configuration $\mathcal{A}_N(c_{Z''}, Z'', \mathcal{F})$ differing from $\mathcal{A}_N(c_{Z'}, Z', \mathcal{F})$ at most by the global charge, and such that

$$\mathcal{H}_N(T_Z, \tilde{b}_Z, c_Z)([w; f]_Z) \equiv_{2N} \mathcal{H}_N(T_{Z''}, \tilde{b}_{Z''}, c_{Z''})([w''; f'']_{Z''}).$$

It follows from the definition of $\Omega_{\mathcal{F}} \subset X_0(M)$ that $w'' = w'$. A further transit argument for flattenings and charges shows finally that

$$\mathcal{H}_N(T_{Z''}, \tilde{b}_{Z''}, c_{Z''})([w'; f'']_{Z''}) \equiv_{2N} \mathcal{H}_N(T_{Z'}, \tilde{b}_{Z'}, c_{Z'})([w'; f']_{Z'}).$$

Since the QH state sums (21) depend only on the reduction $\text{mod}(N)$ of flattenings, the dependence of \mathcal{H}'_N , $N \geq 3$, on $k_f \pmod{\pi i N}$ follows immediately. \square

The above constructions depend on the choice of \mathcal{F} . A canonical choice is the family \mathcal{F}_{EP} of rich components Z contained in any weakly branched EP-triangulation of \hat{V} (see Proposition 2.2). By definition, the analytic configuration $\mathcal{A}_N(Y)$ of $Y = (V, (h_c, k_c))$ is the family of analytic sub-configurations $\mathcal{A}_N(c, Z, \mathcal{F}_{\text{EP}})$ for varying charges and rich components $Z \in \mathcal{F}_{\text{EP}}$.

Corollary 4.17. *The invariants $\mathcal{H}'_1{}^{\mathcal{F}_{\text{EP}}}(\mathcal{P})$ and $\mathcal{H}_N{}^{\mathcal{F}_{\text{EP}}}(\mathcal{P})$ of patterns \mathcal{P} over M , such that $\mathcal{P} = (V, \rho, h_f, k_f, h_c, k_c)$ with $\rho \in \Omega_{\mathcal{F}_{\text{EP}}}$, fixed c -weight (h_c, k_c) , and fixed bulk f -weight $h_f \in H^1(V; \mathbb{Z}/2\mathbb{Z})$ such that $i^*(h_f) = 0$, define respectively an analytic function*

$$\mathcal{H}'_1{}^{\mathcal{F}_{\text{EP}}, h_f} : A_{0, \mathcal{F}_{\text{EP}}}(M)_\infty \rightarrow \mathbb{C}$$

given by (cosets of) edges), and the *signed* antisymmetric bilinear form $\langle \cdot, \cdot \rangle$ on J given by

$$(38) \quad \langle \cdot, \cdot \rangle = \bigoplus_j *_j \langle \cdot, \cdot \rangle_j$$

where $\langle \cdot, \cdot \rangle_j$ is the standard antisymmetric bilinear form on J_j , defined in the basis $\{e_0^j, e_1^j\}_j$ by

$$(39) \quad \langle e_0^j, e_1^j \rangle_j = 1 \quad , \quad \langle e_0^j, e_1^m \rangle = 0 \text{ for } m \neq j.$$

Hence $\{e_0^j, e_1^j\}$ is a symplectic basis of $\langle \cdot, \cdot \rangle$ restricted to J_j when Δ_j has positive branching orientation $*_j = +1$, and is an anti-symplectic basis of J_j otherwise. Equivalently, $\{e_0^j, e_1^j\}$ is symplectic if the edges $e_0^j, (e_1^j)^j, e_2^j$ have positive cyclic ordering as viewed from their common endpoint v_0 in Δ_j (in [33] the branchings are not needed and not used; one fixes positively cyclically ordered edges $e_0^j, (e_1^j)^j, e_2^j$ independently for all Δ_j , so that the signs $*_j$ occurring here are systematically replaced by $+1$). Define linear maps $\beta : C_1 \rightarrow E(T)$ and $\beta^* : E(T) \rightarrow C_1$ by

$$\beta(e) = \sum_{E \rightarrow e} E$$

and

$$\beta^*(e_k^j) = *_j \left(p(e_{k+1}^j) - p(e_{k+2}^j) + p((e'_{k+1})^j) - p((e'_{k+2})^j) \right)$$

and the same for $\beta^*((e'_k)^j)$, where indices are regarded mod(3), and $p : E(T) \rightarrow C_1$ is the identification map, assigning to an abstract edge its coset in T . They induce maps (we keep the same notations)

$$\beta : C_1 \rightarrow J \quad , \quad \beta^* : J \rightarrow C_1.$$

Clearly, for any edge e of T we have

$$(40) \quad (\beta^*(e_k^j), e) = \langle e_k^j, \beta(e) \rangle.$$

Also, β^* splits as $\beta^* : J \xrightarrow{\beta_1} \bar{J} \xrightarrow{\beta_2} C_1$, where $\beta_1(e_k^j) := *_j(e_{k+1}^j - e_{k+2}^j)$ and $\beta_2(e_k^j) := p(e_k^j) + p((e'_k)^j)$. Similarly to (40), β_1 and the natural projection $q : \bar{J} \rightarrow J$ are adjoint maps: for all $x \in \bar{J}$ and $a \in J$ we have

$$(41) \quad (\beta_1(a), x) = \langle a, q(x) \rangle.$$

Denote by C_0 the \mathbb{Z} -module freely generated by the vertices of T , and by $\alpha^* : C_1 \rightarrow C_0$ the map which assigns to an edge the sum of its vertices. One has $\text{Im}(\beta^*) \subset \text{Ker}(\alpha^*)$, and interpreting the elements of $\text{Ker}(\alpha^*)$ as unoriented 1-cycles in T , there is an isomorphism

$$(42) \quad \text{Ker}(\alpha^*) / \text{Im}(\beta^*) \cong H_1(T; \mathbb{Z}/2\mathbb{Z}).$$

It is not hard to see that $\text{Im}(\beta) \subset \text{Ker}(\beta^*)$. Put $H := \text{Ker}(\beta^*) / \text{Im}(\beta)$. The 2-form $\langle \cdot, \cdot \rangle$ on J descends to H . Since $\text{Ker}(\beta^*) = \text{Im}(\beta)^\perp$, $\langle \cdot, \cdot \rangle$ is non degenerate on $H / \text{Tors}(H)$. In fact, it is shown in [33] that there is an exact sequence

$$(43) \quad 0 \longrightarrow H \xrightarrow{(\gamma', \gamma'_2)} H^1(\partial V; \mathbb{Z}) \oplus H^1(V; \mathbb{Z}/2\mathbb{Z}) \xrightarrow{r-i^*} H^1(\partial V; \mathbb{Z}/2\mathbb{Z}) \longrightarrow 0.$$

Moreover, denoting by \cdot the intersection product on $H_1(\partial V; \mathbb{Z})$, and by $\gamma = PD \circ \gamma'$ the map γ' followed by the Poincaré duality isomorphism, for all $a, b \in H$ we have

$$(44) \quad \gamma(a) \cdot \gamma(b) = 2\langle a, b \rangle.$$

The map γ is defined as follows. Recall the notations introduced before Definition 4.10. For any 2-face F of ∂T_0 which is a boundary section of the truncated tetrahedron of T_0 corresponding to Δ_j , let us write $F \rightarrow \Delta_j$, and denote by $s_k^{j,F}$ the edge of F which is opposite to the vertex of F that belongs to the edge e_k^j or $(e'_k)^j$, with the positive orientation as viewed from that vertex. Then, the linear map

$$(45) \quad \begin{aligned} \bar{\gamma} : \bar{J} &\longrightarrow C_1(\partial T_0) \\ e_k^j &\longmapsto \sum_{F \rightarrow \Delta_j} s_k^{j,F}. \end{aligned}$$

descends to

$$(46) \quad \gamma : H \rightarrow H_1(\partial V; \mathbb{Z}).$$

Conversely, let $q : \bar{J} \rightarrow J$ be the natural projection. Denote by $\partial T'_0$ the cellulation of ∂V dual to ∂T_0 . Represent classes in $H_1(\partial V; \mathbb{Z})$ by simplicial loops in $\partial T'_0$. For all $F \rightarrow \Delta_j$, denote by $a_k^{j,F}$ the simplicial arc in $\partial T'_0 \cap F$ which faces the vertex of F that belongs to the edge e_k^j or $(e'_k)^j$, with the positive orientation as viewed from that vertex. Define a linear map

$$\begin{aligned} \bar{\delta} : C_1(\partial T'_0) &\longrightarrow \bar{J} \\ a_k^{j,F} &\longmapsto e_k^j. \end{aligned}$$

Then, $q\bar{\delta}$ descends to a map $\delta : H_1(\partial V; \mathbb{Z}) \rightarrow H$. Moreover, for all $x \in H_1(\partial V; \mathbb{Z})$ we have

$$(47) \quad \gamma \circ \delta(x) = 2x.$$

The identity (44) follows from (47), together with the simple fact that for any simplicial loop C in $\partial T'_0$, we have

$$(48) \quad C \cdot \bar{\gamma}(e_k^j) = *_j \langle \bar{\delta}(C), e_k^j \rangle_j,$$

where the sign $*_j$ comes from the orientation induced by the basis $\{e_0^j, e_1^j\}$ of J_j , as discussed after (39). Hence γ and δ are adjoint maps with respect to the form $\langle \cdot, \cdot \rangle$ on J and the intersection form on $H_1(\partial V; \mathbb{Z})$: for all $x \in H_1(\partial V; \mathbb{Z})$ and $a \in H$ we have

$$(49) \quad \gamma(a) \cdot x = \langle a, \delta(x) \rangle.$$

Taking coefficients in $\mathbb{Z}/2\mathbb{Z}$ and simplicial loops in T'_0 in normal position with respect to T_0 , one defines similarly to δ a map $\delta_2 : H_1(V; \mathbb{Z}/2\mathbb{Z}) \rightarrow H \otimes (\mathbb{Z}/2\mathbb{Z})$. The map γ'_2 in the above exact sequence is defined for all $a \in H \otimes (\mathbb{Z}/2\mathbb{Z})$ and $c \in H_1(V; \mathbb{Z}/2\mathbb{Z})$ by

$$\gamma'_2(a)(c) = \langle a, \delta_2(c) \rangle.$$

With this material at hands, one can show that the maps γ' and γ'_2 produce the boundary and the bulk weights of V . This is done for the charge weights in Section 6 of [33], and for the flattening weights in Section 9 of [34], with the further constraint that the weights are taken equal to 0. For the sake of completeness, let us explain the case of the flattening weights, the adaptation to the charge weights being not difficult.

Consider the weakly branched ideal triangulation (T, \tilde{b}) of $(V, \rho, (h, k))$. By assumption, there is a point $w \in G(T, \tilde{b})$ having ρ as holonomy. Take for each 3-simplex (Δ_j, b_j) the (local) flattening of the form $(\underline{l}_0^j, \underline{l}_1^j, \underline{l}_2^j) := (\log(w_0^j), \log(w_1^j), \log(w_2^j) + \varepsilon i\pi)$, $\varepsilon \in \{-1, +1\}$, and set $\underline{l} := (\underline{l}_0^j, \underline{l}_1^j, \underline{l}_2^j)_j$. Consider the vector

$$(50) \quad v_{\underline{l}} := \sum_{j=1}^s \underline{l}_1^j e_0^j - \underline{l}_0^j e_1^j \in J \otimes \mathbb{C}.$$

We have

$$\beta_1(v_{\underline{l}}) = \sum_{j=1}^s *_j (\underline{l}_0^j e_0^j + \underline{l}_1^j e_1^j - (\underline{l}_0^j + \underline{l}_1^j) e_2^j) = \sum_{j=1}^s *_j (\underline{l}_0^j e_0^j + \underline{l}_1^j e_1^j + \underline{l}_2^j e_2^j) \in \bar{J} \otimes \mathbb{C}.$$

So $\beta_1(v_{\underline{l}})$ represents the map assigning the value $*_j \underline{l}_k^j$ to the abstract edges e_k^j and $(e'_k)^j$ of T . Recall that for every $\bar{a} \in \bar{J}$, $\beta_2(\bar{a})$ computes at every edge $e \in C_1$ the sum of the coefficients of \bar{a} attached to the abstract edges having coset e . Then, $\beta^* = \beta_2 \circ \beta_1$, the gluing equation (8), and the argument of Lemma 2.7 imply that $\beta^*(v_{\underline{l}}) \in 2\pi i C_1$ (note that in the Lemma the label \emptyset is attached to edges of the type e_2^j and $(e'_2)^j$). By (42) we deduce that $\beta^*(v_{\underline{l}})/\pi i$ represents the 0 class in $H_1(T; \mathbb{Z}/2\mathbb{Z})$, and so there exists $\underline{f} \in J$ such that $\beta^*(\underline{f}) = \beta^*(v_{\underline{l}})/\pi i$. Put $v_l := v_{\underline{l}} - \pi i \underline{f}$, and write its coefficients as

$$v_l = \sum_{j=1}^s \underline{l}_1^j e_0^j - \underline{l}_0^j e_1^j = \sum_{j=1}^s (\log(w_1^j) + \pi i f_1^j) e_0^j - (\log(w_0^j) + \pi i f_0^j) e_1^j.$$

Since $\beta^*(v_l) = 0$, the collection $l := (l_0^j, l_1^j, l_2^j)_j$ satisfies the edge relations $L(e) = 1$ (see (22)). The tetrahedral relations are satisfied because any element of $\text{Im}(\beta_1)$ has coefficient sum equal to 0 in

each module \bar{J}_j . So l defines a flattening of w . Consider the class $[v_l] \in H \otimes \mathbb{C}$. One can see that $\gamma'([v_l]) = L([w; f])$ as follows. Consider the map

$$\begin{aligned} \bar{\gamma}'' : \bar{J} \otimes \mathbb{C} &\longrightarrow C^1(\partial T'_0; \mathbb{C}) \\ x &\longmapsto (C \mapsto (x, \bar{\delta}(C))). \end{aligned}$$

Denote by $A \in J \otimes \mathbb{C}$ a representative of $a \in H \otimes \mathbb{C}$. Then $\bar{\gamma}''$ induces a map

$$\begin{aligned} \gamma'' : H \otimes \mathbb{C} &\longrightarrow H^1(\partial V; \mathbb{C}) \\ a &\longmapsto \bar{\gamma}''(\beta_1(A)). \end{aligned}$$

Using successively (41) and (49), we get

$$(51) \quad \gamma''(a)([C]) = (\beta_1(A), \bar{\delta}(C)) = \langle a, q\bar{\delta}([C]) \rangle = \gamma(a) \cdot [C].$$

Hence $\gamma'' = \gamma'$. Clearly, $\gamma''([v_l]) = \bar{\gamma}''(\beta_1(v_l))$ is given by (26). This proves $\gamma'([v_l]) = L([w; f])$. The collection $f := (f_0^j, f_1^j, f_2^j)_j$ satisfies also the edge relations mod(2), hence it defines a class $\gamma'_2(f) \in H^1(V; \mathbb{Z}/2\mathbb{Z})$ as explained after (26). By construction we have the compatibility relations (28), so the pair $(L([w; f]), \gamma'_2(f))$ is a flattening weight of (V, ρ) . Finally, take any other flattening weight (k'_f, h'_f) of (V, ρ) . Comparing (2) and (28) we see that $k'_f - L([w; f]) \in H^1(V; \pi i \mathbb{Z})$ and $(k'_f - L([w; f]))/\pi i = i^*(h'_f - \gamma'_2(f)) \text{ mod}(2)$. By the exact sequence (43), there exists $a \in H$ such that $\gamma'(a) = (k'_f - L([w; f]))/\pi i$ and $\gamma'_2(a) = h'_f - \gamma'_2(f)$. Then the vector $\beta_1(v_l + \pi i a) \in \bar{J} \otimes \mathbb{C}$ represents a flattening of w with weight (h'_f, k'_f) . \square

4.4. Relation with Chern-Simons theory. Recall that V is the compact 3-manifold obtained by completing the cusp of M with a torus. Chern-Simons theory with gauge group $PSL(2, \mathbb{C})$ associates to ∂V the *Chern-Simons bundle* $\mathcal{L}_{\partial V} \rightarrow X(\partial V)$, which is a \mathbb{C}^* -bundle with canonical connection 1-form and canonical inner product, and to V the parallel *Chern-Simons section* $s_V : X(V) \rightarrow i^* \mathcal{L}_{\partial V}$, where $i : X(V) \rightarrow X(\partial V)$ is the restriction map ([20, 27]).

We are going to show that when the bulk f -weight h_f is 0 the bottom row of the diagram of Theorem 4.17 encodes the restriction of the pair $(i^* \mathcal{L}_{\partial V}, s_V)$ to $\Omega_{\mathcal{FEP}} \subset X_0(M)$, identified as a component of $X(V)$. For arbitrary h_f one gets a twisted version of $(i^* \mathcal{L}_{\partial V}, s_V)$. The proof relies on a certain number of results that we need to recall. Denote by $\text{CS}(M') \in \mathbb{R}/\mathbb{Z}$ the Chern-Simons invariant of the Levi-Civita connection of a closed riemannian 3-manifold M' , and by $\text{CS}(M)$ Meyerhoff's extension to cusped hyperbolic manifolds M . By definition, $\text{CS}(M)$ is the limit of $\text{CS}(M'_n)$ for any sequence (M'_n) of closed hyperbolic Dehn fillings of M converging to M in Dehn surgery space. In [34], Corollary 14.6 & Theorem 14.7, Neumann proved (using the notations of the present paper):

Theorem 4.19. *Let Y be M or a closed hyperbolic Dehn filling M' of M . Denote by ρ_Y the hyperbolic holonomy of Y , identified as a character of $V \subset M'$ in the case $Y = M'$. If $Y = M$, put $k_Y = 0$, and if $Y = M'$, let k_Y be any boundary f -weight of V relative to ρ_Y such that k_Y vanishes on the meridian of the added solid torus. Then*

$$\mathcal{H}_1(V, \rho_Y, (0, k_Y)) = \exp \left(\frac{2}{\pi} \text{Vol}(Y) + 2\pi i \text{CS}(Y) \right).$$

The expression on the right hand side is related to the Chern-Simons section s_V as follows. To any compact closed oriented 3-manifold Y , $PSL(2, \mathbb{C})$ -Chern-Simons theory associates a *function* $s_Y : X'(Y) \rightarrow \mathbb{C}^*$ defined on the variety $X'(Y)$ of $PSL(2, \mathbb{C})$ -characters of Y . If Y is hyperbolic with holonomy ρ_Y , a classical result of Yoshida [40] gives

$$(52) \quad s_Y(\rho_Y) = \exp \left(\frac{2}{\pi} \text{Vol}(Y) + 2\pi i \text{CS}(Y) \right).$$

Assume that $Y = V \cup_{\varphi} (D^2 \times S^1)$ is a closed hyperbolic Dehn filling of M whose holonomy ρ_Y lies in a sufficiently small neighborhood D of ρ_{hyp} in $X(V) = X(M)$ (here we identify $\rho(Y)$ with an augmented character of V). Then Kirk-Klassen ([27], pages 554–556) showed that

$$(53) \quad s_Y(\rho_Y) = \langle s_V(\rho_Y), s_{D^2 \times S^1}(\rho_Y) \rangle$$

where $\langle \cdot, \cdot \rangle$ is the canonical inner product of $i^* \mathcal{L}_{\partial V}$, and on the right hand side ρ_Y denotes also the augmented character induced on the glued solid torus $D^2 \times S^1$. Explicitly, if we fix the gauge on $i^* \mathcal{L}_{\partial V}$ by taking as coordinates on D the ones corresponding under the map \mathfrak{h} to the *standard* logarithms $\log(\lambda)$ and $\log(\mu)$, which are equal to 0 at ρ_{hyp} , we have

$$(54) \quad s_V(\rho_{hyp}) = \exp\left(\frac{2}{\pi} \text{Vol}(M) + 2\pi i \text{CS}(M)\right).$$

Then, if the Dehn filling instruction φ maps the meridian $\partial(D^2 \times *)$ to $l^q m^p$ and the longitude $* \times S^1$ to $l^s m^r$ (so $ps - qr = 1$), the formula (53) splits as

$$(55) \quad s_Y(\rho_Y) = s_V(\rho_{hyp}) \times \exp\left(-\frac{1}{2\pi i} \int_{\rho_{hyp}}^{\rho_Y} (\log(\lambda) d \log(\mu) - \log(\mu) d \log(\lambda))\right) \exp(-(s \log(\lambda) + r \log(\mu))).$$

The first exponential gives the variation of s_V between ρ_{hyp} and ρ_Y (s_V being a parallel section of $i^* \mathcal{L}_{\partial V}$, this exponential is the holonomy of the connection 1-form of $i^* \mathcal{L}_{\partial V}$ between the two points); the product with $s_V(\rho_{hyp})$ is thus the value of $s_V(\rho_Y)$ at the chosen gauge. The second exponential is the value of $s_{D^2 \times S^1}(\rho_Y)$ in the gauge corresponding to the Dehn filling: $\log(\text{hol}_{\partial(D^2 \times *)}) = q \log(\lambda) + p \log(\mu) = 2\pi i$, $\log(\text{hol}_{* \times S^1}) = s \log(\lambda) + r \log(\mu)$. Its argument is $-(\text{length} + i \text{ rotation angle})$ of the geodesic core of the surgery torus. The arguments of both exponential can be collected in a single integral by changing the coordinates $\log(\lambda)$, $\log(\mu)$ on D to $\log(\lambda) + 2\pi i r$, $\log(\mu) - 2\pi i s$, that is, by working on a different leaf of the Riemann surface of these maps. Then the gauge $\log(\text{hol}_{\partial(D^2 \times *)})$ vanishes, like k_Y in Theorem 4.19.

We are going to show that $(i^* \mathcal{L}_{\partial V}, s_V)|_{\Omega_{\mathcal{F}_{EP}}}$ is determined by $\mathcal{H}_1^{\mathcal{F}_{EP}, 0} : A_{0, \mathcal{F}_{EP}}(M)_\infty \rightarrow \mathbb{C}$. Because of Theorem 4.19 and (54), it is enough to identify the variation of $\mathcal{H}_1^{\mathcal{F}_{EP}, 0}$ with the one of s_V , given by (55). This is the content of the following result.

Proposition 4.20. *The function $\mathcal{H}_1^{\mathcal{F}_{EP}, 0}$ descends to a parallel section \mathcal{S}^0 of a flat trivial \mathbb{C}^* -bundle $\mathcal{L}(M) \rightarrow A_{0, \mathcal{F}_{EP}}(M)$ with canonical connection 1-form and inner product, such that $\mathfrak{h}^*(\mathcal{L}(M), \mathcal{S}^0)$ is isomorphic to $(i^* \mathcal{L}_{\partial V}, s_V)|_{\Omega_{\mathcal{F}_{EP}}}$.*

Proof. We define $\mathcal{L}(M)$ from the variation of the invariant $\mathcal{H}_1^{\mathcal{F}_{EP}}$ of Theorem 4.16, like the Chern-Simons line bundle is defined from the Chern-Simons action in [20, 27]. Hence $\mathcal{L}(M)$ is the quotient of $A_{0, \mathcal{F}_{EP}}(M)_\infty \times \mathbb{C}^*$ under the action of \mathbb{Z}^2 given by

$$(56) \quad \forall a, b \in \mathbb{Z}, (a, b) \cdot ([\lambda; p], [\mu; q], z) := ([\lambda; p + a], [\mu; q + b], z e^{b \log(\lambda) - a \log(\mu)}).$$

We have a \mathbb{C}^* -bundle projection

$$\pi'_\infty : \begin{array}{ccc} \mathcal{L}(M) & \longrightarrow & A_{0, \mathcal{F}_{EP}}(M) \\ [[\lambda; p], [\mu; q], z] & \longmapsto & (\lambda, \mu). \end{array}$$

Denote by $\mathcal{L}(M)^{-1}$ the inverse \mathbb{C}^* -bundle, which is the same as $\mathcal{L}_{\mathcal{F}_{EP}}(-M)$, where $-M$ denotes M with reversed orientation. An inner product is defined by the bundle map

$$\langle \cdot, \cdot \rangle : \begin{array}{ccc} \mathcal{L}(M) \times \mathcal{L}(M)^{-1} & \longrightarrow & \mathbb{C}^* \\ (([\lambda; p], [\mu; q], z_1), ([\lambda; p], [\mu; q], z_2)) & \longmapsto & z_1 z_2 \end{array}$$

where \mathbb{C}^* is regarded as the bundle over a point. Also, a flat analytic connection 1-form is defined by the restriction to $A_{0, \mathcal{F}_{EP}}(M)_\infty$ of the 1-form on $\tilde{\mathbb{C}} \times \tilde{\mathbb{C}}$ given by (we put $l_a(\lambda) := \log(\lambda) + 2\pi i a$)

$$(57) \quad \eta := -\frac{1}{2\pi i} (l_p(\lambda) dl_q(\mu) - l_q(\mu) dl_p(\lambda)).$$

By Lemma 4.15, given any even class $\theta \in H^1(\partial V; 2\mathbb{Z})$ and any point $[w; f]_Z \in \Omega_{\mathcal{F}_{EP}}(Z)_{\infty, 0}$, the flattening f can be modified to a flattening f' such that $[w; f']_Z \in \Omega_{\mathcal{F}_{EP}}(Z)_{\infty, 0}$ and the corresponding boundary weights satisfy

$$L([w; f']_Z) - L([w; f]_Z) = \pi i \theta.$$

As in the proof of Proposition 4.8 (2) and 4.9 (2), denote by v_l (resp. $v_{l'}$) the vector in $H \otimes \mathbb{C}$ associated to the classical log-branch at $[w; f]_Z$ (resp. $[w; f']_Z$), and similarly denote by $v_d \in H \otimes (\mathbb{C}/2\pi i\mathbb{Z})$ the vector associated to $(\log(w_0^j), \log(w_1^j), \log(w_2^j))_j$. Recall the class $d_w \in H^1(\partial V; \mathbb{C}/2\pi i\mathbb{Z})$ in (28). Then $d_w = \gamma'(v_d)$ and

$$\pi i \theta = \gamma'(v_{l'} - v_l) = \gamma' \left(\pi i \sum_{j=1}^s ((f')_1^j - f_1^j) e_0^j - ((f')_0^j - f_0^j) e_1^j \right).$$

Hence, denoting $\mathcal{H}_1^{\mathcal{F}_{\text{EP}}} := \mathcal{H}_1^{\mathcal{F}_{\text{EP}}}(T_Z, \tilde{b}_Z)$ we have

$$\begin{aligned} \mathcal{H}_1^{\mathcal{F}_{\text{EP}}}([w; f']_Z) \mathcal{H}_1^{\mathcal{F}_{\text{EP}}}([w; f]_Z)^{-1} &= \\ &= \exp \left(\sum_{j=1}^s *_{j} \left(((f')_1^j - f_1^j) \log(w_0^j) - ((f')_0^j - f_0^j) \log(w_1^j) \right) \right) \\ &= \exp \left(-\frac{1}{\pi i} \langle v_{l'} - v_l, v_d \rangle \right) \\ &= \exp \left(-\frac{1}{2\pi i} \langle \langle \gamma'(v_{l'} - v_l), \gamma'(v_d) \rangle \rangle \right) \\ (58) \quad &= \exp \left(\frac{1}{2} (\theta(m) \log(\text{hol}_l(\rho(w))) - \theta(l) \log(\text{hol}_m(\rho(w)))) \right) \end{aligned}$$

where the map hol_C , C a curve in ∂V , is defined in (23), and $\langle \langle \cdot, \cdot \rangle \rangle$ denotes the cup product followed by the evaluation on the fundamental class. The first equality follows from (17) and (21), the second is by the definition (38) of the symplectic pairing on $H \otimes \mathbb{C}$, the third follows from (44) and Poincaré duality, and the last one uses that the symplectic basis of $H_1(\partial V; \mathbb{Z})$ is (l, m) . By setting $\theta(m) := 2b$ and $\theta(l) := 2a$, we see from (56) and (58) that the function $\mathcal{H}_1^{\mathcal{F}_{\text{EP}}, h_f}$ (with arbitrary bulk weight h_f for the moment) descends from $A_{0, \mathcal{F}_{\text{EP}}}(M)_{\infty}$ to a section of $\mathcal{L}(M)$. Denote it

$$S^{h_f} : A_{0, \mathcal{F}_{\text{EP}}}(M) \rightarrow \mathcal{L}(M).$$

We show that this section is parallel with respect to the connection η by a computation similar to (58). As remarked in [33], Lemma 10.2, for any \mathbb{Q} -vector space E , the skew symmetric bilinear form $\langle \cdot, \cdot \rangle$ on J induces a symmetric bilinear map

$$\begin{aligned} B : (H \otimes E) \otimes (H \otimes E) &\longrightarrow E \wedge E \\ (a \otimes v) \otimes (b \otimes w) &\longmapsto \langle a, b \rangle v \wedge w. \end{aligned}$$

Consider the map \cdot_E on $H_1(\partial V; E) = H_1(\partial V; \mathbb{Z}) \otimes E$ induced by the intersection product on $H_1(\partial V; \mathbb{Z})$:

$$\begin{aligned} \cdot_E : H_1(\partial V; E) \otimes H_1(\partial V; E) &\longrightarrow E \wedge E \\ (x \otimes v) \otimes (y \otimes w) &\longmapsto (x \cdot y) v \wedge w. \end{aligned}$$

Denote again by γ the map $\gamma \otimes \text{id} : H \otimes E \rightarrow H_1(\partial V; E)$. By (44) we have a commutative diagram

$$\begin{array}{ccc} (H \otimes E) \otimes (H \otimes E) & \xrightarrow{B} & E \wedge E \\ \gamma \otimes \gamma \downarrow & & \downarrow 2\times \\ H_1(\partial V; E) \otimes H_1(\partial V; E) & \xrightarrow{\cdot_E} & E \wedge E \end{array}$$

In particular, let $E = C^{\infty}(\mathcal{W}_{\infty})$ and denote by $l_0^j, l_1^j \in E$ the classical log-branch functions at the edges e_0^j, e_1^j of Δ_j . Put $a = \sum_{j=1}^s e_0^j \otimes l_1^j - e_1^j \otimes l_0^j \in H \otimes E$. For any point $[w; f]_Z \in \Omega_{\mathcal{F}_{\text{EP}}}(Z)_{\infty, 0}$ we have $\gamma(a)([w; f]_Z) = l \otimes L([w; f]_Z)(m) - m \otimes L([w; f]_Z)(l)$; dually the class $\gamma'(a) \in H^1(\partial V; E)$, $\gamma'(a) = PD \circ \gamma(a)$, is given by $\gamma'(a)([w; f]_Z) = L([w; f]_Z) \in H^1(\partial V; \mathbb{C})$. Moreover, in $E \wedge E$ we have

$$2 \sum_{j=1}^s *_{j} (l_0^j \wedge l_1^j) = B(a, a) = \frac{1}{2} \gamma(a) \cdot_E \gamma(a) = L(\cdot)(l) \wedge L(\cdot)(m).$$

Applying the homomorphism $E \wedge E \rightarrow \Omega^1(\mathcal{W}_\infty)$, $f \wedge g \mapsto fdg - gdf$, we deduce

$$2 \sum_{j=1}^s *_{j} (l_0^j dl_1^j - l_1^j dl_0^j) = L(\cdot)(l)dL(\cdot)(m) - L(\cdot)(m)dL(\cdot)(l).$$

Hence, for any two points $[w; f]_Z, [w'; f']_Z$ connected by a continuous path $(\gamma^1, \dots, \gamma^s)$ in $\Omega_{\mathcal{F}_{\text{EP}}}(Z)_{\infty,0}$, we have

$$(59) \quad \mathcal{H}_1^{\mathcal{F}_{\text{EP}}}([w'; f']_Z) \mathcal{H}_1^{\mathcal{F}_{\text{EP}}}([w; f]_Z)^{-1} = \exp \left(2 \sum_{j=1}^s *_{j} \int_{\gamma^j} \eta \right) = \exp \left(\int_{\gamma} \eta \right)$$

where $\gamma \subset A_{0, \mathcal{F}_{\text{EP}}}(M)_{\infty}$ is the image of $(\gamma^1, \dots, \gamma^s)$ under the map loghol_{∞} . Hence \mathcal{S}^{h_f} is a parallel section of $\mathcal{L}(M)$. Multiplying coordinates by $4\pi i$ identifies $\mathfrak{h}^* \mathcal{L}(M)$ with the restriction of $i^* \mathcal{L}_{\partial V}$ to $\Omega_{\mathcal{F}_{\text{EP}}}$ (see [27], end of Section 3 and pages 555-556). Finally, as described before the statement, from (55) and (59) we get $\mathfrak{h}^* \mathcal{S}^0 = (s_V)|_{\Omega_{\mathcal{F}_{\text{EP}}}}$. \square

Remark 4.21. 1) In [27] the Chern-Simons bundle is defined over the variety of characters $X'(\partial V)$ instead of the variety of augmented characters $X(\partial V)$; the coordinates on $X'(\partial V)$ are $1/2\pi i$ times the logarithms of the eigenvalues at a pair of meridian and longitude curve on ∂V . A fundamental domain for the action of $\mathbb{Z} \times \mathbb{Z}$ on coordinates is $([0; 1/2] \times \mathbb{R}) \times ([0; 1] \times \mathbb{R})$. The rescaling factor $4\pi i$ at the end of the proof replaces these coordinates by the logarithms of the squared eigenvalues, so that a fundamental domain for the action of $\mathbb{Z} \times \mathbb{Z}$ on $A_{0, \mathcal{F}_{\text{EP}}}(M)_{\infty}$ is $([0; 2i\pi] \times \mathbb{R})^2$.

2) The bundle $\mathcal{L}(M)$ is the restriction to $A_{0, \mathcal{F}_{\text{EP}}}(M)$ of a bundle $\mathcal{L}(T^2)$ over $X(T^2) = X(\partial V)$ defined by the action (56). Of course $\mathcal{L}(T^2)$ is isomorphic to $\mathcal{L}_{\partial V}$. Its connection 1-form (57) has curvature $F_{\eta} = (-1/i\pi)d \log(\lambda) \wedge d \log(\mu)$, its Euler class is represented by $(-1/2(i\pi)^2)d \log(\lambda) \wedge d \log(\mu)$, and its Euler number is -2 .

Remarks 4.22. *Problems and perspectives :*

a) The \mathbb{Z}^2 -equivariant analytic function $\mathcal{H}_1^{\mathcal{F}_{\text{EP}},0}$ on $A_{0, \mathcal{F}_{\text{EP}}}(M)_{\infty}$ is the classical analog of the regular rational functions $\mathcal{H}_N^{\mathcal{F}_{\text{EP}},0, h_c, k_c}$ on $A_{0, \mathcal{F}_{\text{EP}}}(M)_N$. One expects that a refinement of the \mathbb{Z}^2 -action appears in the quantum world, involving the quantum torus algebra generated by \hat{l}, \hat{m} such that $\hat{l}\hat{m} = \zeta \hat{m}\hat{l}$, where ζ is a primitive N -th root of 1 and (l, m) a geometric basis of $H_1(\partial V; \mathbb{Z})$:

Problem 4.23. *Lift the functions $\mathcal{H}_N^{\mathcal{F}_{\text{EP}}, h_f, h_c, k_c} : A_{0, \mathcal{F}_{\text{EP}}}(M)_N \rightarrow \mathbb{C}/\mu_{2N}$ to \mathbb{C} -valued functions defined on some covering space of $A_{0, \mathcal{F}_{\text{EP}}}(M)_N$.*

b) By Theorem 1.1 (2), the functions $\mathcal{H}_N^{\mathcal{F}_{\text{EP}}, h_f, h_c, k_c}$ yield \mathbb{C}/μ_{2N} -valued rational functions on a covering space $\tilde{X}_0(M)_N$ of $X_0(M)$. On another hand, the field $\mathbb{C}(\tilde{X}_0(M)_N)$ is a finite extension of $\mathbb{C}(X_0(M))$, and $\mathbb{C}(X_0(M))$ is generated by the functions of augmented characters associated to hol_l and hol_m by pull-back via the map $\mathfrak{h} : X_0(M) \rightarrow A_0(M)$.

Problem 4.24. *Describe $\mathcal{H}_N^{\mathcal{F}_{\text{EP}}, h_f, h_c, k_c}$ in terms of functions of augmented characters, or as a meromorphic function on the smooth projective model of $\tilde{X}_0(M)_N$.*

For instance, a property that is obvious from the state sum formulas is that the poles of $\mathcal{H}_N^{\mathcal{F}_{\text{EP}}, h_f, h_c, k_c}$ cover ideal points of $X_0(M)$.

By a theorem of Bullock [12], the ring of $SL(2, \mathbb{C})$ -characters $\mathbb{C}[X(M)]$ is isomorphic to $K_{-1}(M)/\sqrt{0}$, where $K_{-1}(M)$ is the Kauffman bracket skein module of M specialized at $A = -1$, with its natural algebra structure, and $\sqrt{0}$ its nilradical.

Problem 4.25. *Find skein theoretic constructions of $\mathcal{H}_N^{\mathcal{F}_{\text{EP}}, h_f, h_c, k_c}$.*

5. QHFT PARTITION FUNCTIONS

For simplicity we consider only the QHFT patterns with topological support (W, L, h_c) , where W is a closed oriented connected 3-manifold. In that case only the bulk c -weight h_c occurs. In order to build the analytic configurations of (W, L, h_c) we consider *quasi-regular* triangulations T of W (see Proposition 2.3), and we require that L is realized by a *Hamiltonian* subcomplex H of the 1-skeleton of T . Such a pair (T, H) is called a *distinguished* triangulation of (W, L) .

Definition 5.1. A *global charge* c on (T, H) is a rough global charge on T satisfying the following additional global constraint on the total edge charges: $C(e) = 0$ if e is an edge of H , and $C(e) = 2$ otherwise.

Note that any global charge on (T, H) encodes H (i.e. the link L). Let us turn to the combinatorial encoding of (W, L, h_c) . Similarly to Proposition 4.8 and 4.9 we have (see [3, 4, 5]):

Proposition 5.2. (1) Every pair (W, L) has quasi-regular distinguished triangulations (T, H) .

(2) Every global charge c on (T, H) determines a bulk c -weight h_c of W .

(3) For every bulk c -weight h_c of W and every distinguished triangulation (T, H) of (W, L) there is a global charge c on (T, H) with bulk c -weight equal to h_c .

Let us fix a quasi-regular distinguished triangulation (T, H) of (W, L) , a global charge c on (T, H) , and a branching b on T (it exists).

Proposition 5.3. (1) Every point $[w; f] \in G_0(T, b)_\infty$ determines a character $\rho = \rho(w)$ and a bulk f -weight h_f , so that (W, L, ρ, h_f, h_c) is a pattern with topological support (W, L, h_c) .

(2) For every pattern (W, L, ρ, h_f, h_c) there is a point $[w; f] \in G_0(T, b)_\infty$ with holonomy ρ and bulk f -weight h_f .

The construction of the QHI of (W, L, h_c) is achieved by the following Proposition; it is improved in Section 8 with respect to the QHI sign anomaly. The proof is similar to that of Theorem 4.16. For every odd $N \geq 3$ consider the analytic configurations $\mathcal{A}_N(T, b, c)$. The defining equations of the algebraic variety $G_0(T, b, c)_N$ are:

- $W'(e) = \zeta^{-1}$, for every edge e of T not contained in H ;
- $W'(e) = 1$, for every edge e of T contained in H .

Proposition 5.4. For every pattern $\mathcal{P} = (W, L, \rho, h_f, h_c)$, every distinguished branched quasi-regular triangulation (T, b, H) of (W, L) , every global charge c on (T, H) with bulk c -weight equal to h_c , and every point $[w; f] \in G_0(T, b)_\infty$ with holonomy ρ and bulk f -weight h_f , the scalars

- $\mathcal{H}_1(\mathcal{P}) := \mathcal{H}_1(T, b)([w; f])$
- $\mathcal{H}_N(\mathcal{P}) := {}_{\equiv 2N} \mathcal{H}_N(T, b, c)([w; f])$

do not depend on the choice of (T, b, c) and $[w; f]$, and hence define invariants of \mathcal{P} .


By definition, the analytic configuration $\mathcal{A}_N(W, L, h_c)$ is the family of analytic configurations $\mathcal{A}_N(T, b, c)$ over the triples (T, b, c) as in Proposition 5.4.

Remark 5.5. The state sum normalization factor $a_N(T, \tilde{b})$ includes a term N^{-v} . Other choices are possible. In [6] we used instead N^{2-v} in order to get Theorem 1.2.

For general QHFT patterns the constructions are more elaborated; in particular, one requires that H is Hamiltonian in the 1-skeleton of T from which the non-manifold vertices have been removed, and in the normalization factor N^{-v} , v denotes the number of manifold (i.e. internal) vertices. Anyway, the results have basically the same flavour.

6. \mathcal{N} -GRAPH CALCULUS

As every QH state sum is the total contraction of a tensor network supported by a \mathcal{N} -graph, one needs to describe the main features of a “calculus” based on such diagrams.

FIGURE 10. Some Reidemeister moves of type III for \mathcal{N} -graphs

6.1. \mathcal{N} -graphs representing the same weakly branched triangulation. Using the \mathcal{N} -graph decoding of Section 2 it is easy to see that two \mathcal{N} -graphs Γ and Γ' encode the same weakly branched triangulation (T, \tilde{b}) if and only if they are equivalent under the relation generated by:

- Plane isotopy.
- Switching the over/under arcs at an accidental crossing.
- The \mathcal{N} -graph versions of oriented Reidemeister moves, called *R-moves* (i.e. formally the usual ones if only accidental crossings are involved, or the Reidemeister move of type III shown in Figure 10 (the orientation being understood) when a dotted crossing is involved).

6.2. Changing the weak branching. Given a \mathcal{N} -graph Γ representing a weakly branched triangulation (T, \tilde{b}) , and an arbitrary change of weak branching $(T, \tilde{b}) \rightarrow (T, \tilde{b}')$, we want to describe a systematic way to modify Γ to a \mathcal{N} -graph Γ' representing (T, \tilde{b}') . We begin with the local changes of branching on a given tetrahedron.

6.2.1. The S_4 action $(\Delta, b) \rightarrow (\Delta, b_\beta)$. Consider a branched tetrahedron (Δ, b) with b -ordered vertices v_0, v_1, v_2, v_3 and 2-facets F_0, F_1, F_2, F_3 , as usual. Let $S(J_4)$ be the symmetric group on the set $J_4 = \{0, 1, 2, 3\}$. There is a natural 1–1 correspondence between the elements $\beta \in S(J_4)$ and the branchings b_β on Δ such that $b = b_{\text{Id}}$. The branching b induces a branching $b(j)$ on each facet F_j , that is a labelling of the vertices of F_j by elements of $J_3 = \{0, 1, 2\}$. For every $\beta \in S(J_4)$, also the branching b_β induces a branching $b_\beta(j)$ on F_j (note that these facets are still ordered with respect to b , not b_β), and the transition from $b(j)$ to $b_\beta(j)$ is encoded by a permutation $\epsilon_\beta(j) \in S(J_3)$. Clearly we have:

Lemma 6.1. (1) *The branching signs $*_{b_\beta}$ and $*_b$ coincide if and only if $\beta \in A(J_4)$.*

(2) *The 2-facet F_j has the same b - and b_β -transverse co-orientations if and only if $\epsilon_\beta(j) \in A(J_3)$.*

Let $C(b)$ denote a dotted \mathcal{N} -graph crossing that encodes (Δ, b) as in Figure 4. We call $C(b)$ a $2-2$ \mathcal{N} -tangle; we consider it as a planar tangle properly embedded in a 2-disk D , with endpoints on ∂D labelled by J_4 . One can assume that the crossing is central to D , and that the arcs joining it to the endpoints are contained in two oriented diameters of D . We want to describe an algorithm that produces, for every $\beta \in S(J_4)$, a new decorated $2-2$ \mathcal{N} -tangle $C(b_\beta)$ in D , with the same unoriented arc-germs as $C(b)$ at ∂D . Hence the arcs of $C(b)$ and $C(b_\beta)$ will be in natural 1–1 correspondence and oriented by b and b_β respectively. The tangle $C(b_\beta)$ will verify the following properties:

- $C(b_\beta)$ contains one dotted crossing. According to the last Lemma, $*_{b_\beta} = *_b$ if and only if $\beta \in A(J_4)$. Hence, given β , we know the sign $*_{b_\beta}$.
- $C(b_\beta)$ can replace $C(b)$ in every \mathcal{N} -graph Γ' obtained after a change of weak branching $\tilde{b} \rightarrow \tilde{b}'$ acting as $b \rightarrow b_\beta$ on the corresponding branched tetrahedron (Δ, b) of (T, \tilde{b}) .
- Each of the four arc-germs of $C(b_\beta)$ at ∂D is labelled by an element of $S(J_3)$, which will eventually contribute to the edge coloring of Γ' .

An auxiliary construction. Consider the two *basic 2–2 tangles* in a 2-disk D' , made by two *oriented* simple arcs a, a' that either are disjoint or intersect transversely at one point, without any over/under information at the crossing point. A J_4 -labelling of the 4 free endpoints of $a \cup a'$ is *admissible* if it respects the following conditions:

(a) The endpoints labelled by 1, 3 (resp. 0, 2) belong to different arcs and are both either the initial or the final endpoint of the corresponding arc.

(b) 3, 2 (resp. 1, 0) label the endpoints of the same arc.

These conditions are satisfied by the labels of the tangles of Figure 4; recall that in this figure, $*_b = 1$ (resp. $*_b = -1$) if and only if 1, 3 label initial (resp. final) endpoints. By the same rule we give every basic tangle an admissible J_4 -labelling, denoted by \mathcal{B} , and a sign $*_{\mathcal{B}} = \pm 1$. Note also that in Figure 4, the arc with endpoints labelled by 0, 1 passes over the other arc. Now we convert \mathcal{B} into a 2–2 \mathcal{N} -tangle as follows:




FIGURE 11. Basic to \mathcal{N} -tangles.




FIGURE 12. More basic to \mathcal{N} -tangles.

(c) If $a \cap a' = \emptyset$, perform an oriented Reidemeister move of type II of the arc with labels 0, 1 over the other arc, thus creating two crossings. Only one of them can be made into a dotted crossing so that its sign $*_b$ agrees with $*_{\mathcal{B}}$ (see some examples in Figure 11).

(d) If a and a' cross at one point, let A be the arc with 0, 1 labels, and I a small open interval in the interior of the arc with 2, 3 labels, such that I contains the crossing point. Let A pass over I , put a dot at the crossing point, and turn I so that the crossing sign and $*_{\mathcal{B}}$ agree. Finally complete $I \cup A$ to an \mathcal{N} -tangle with the same endpoints as $a \cap a'$, by introducing 0 or 2 accidental crossings on opposite sides of A with respect to $I \cap A$ (see some examples in Figure 12). \square

The algorithm $C(b) \rightarrow C(b_{\beta})$:

- (1) Remove the interior of a smaller concentric sub-disk D' of D . Each endpoint of $C(b)$, say x_j , $j \in J_4$, is connected by an oriented sub-arc to a point $x'_{\beta(j)}$ on $\partial D'$.
- (2) There is only one basic 2–2 tangle \mathcal{B} in D' such that the b_{β} -labelling of the endpoints $x'_{\beta(j)}$ is admissible and $*_{\mathcal{B}} = *_{b_{\beta}}$. As in the auxiliary construction, convert \mathcal{B} into an 2–2 \mathcal{N} -tangle in D' ; glue it to the four sub-arcs $[x_j, x'_{\beta(j)}]$, and denote by $C'(b_{\beta})$ the resulting 2–2 \mathcal{N} -tangle in D .
- (3) The 2–2 \mathcal{N} -tangle $C(b_{\beta})$ is obtained from $C'(b_{\beta})$ by giving a label $\hat{\epsilon}_{\beta}(j) \in S(J_3)$ to each sub-arc $[x_j, x'_{\beta(j)}]$ as follows. Recall the permutation $\epsilon_{\beta}(j) \in S(J_3)$ defined before Lemma 6.1; the b - and b_{β} -orientations of $[x_j, x'_{\beta(j)}]$ coincide one to each other iff $\epsilon_{\beta}(j) \in A(J_3)$. Then set:
 - $\hat{\epsilon}_{\beta}(j) = \epsilon_{\beta}(j)$, if the b_{β} -orientation of $[x_j, x'_{\beta(j)}]$ coincides with the orientation pointing towards the center of D ;
 - $\hat{\epsilon}_{\beta}(j) = \epsilon_{\beta}(j)^{-1}$ otherwise. \square

6.2.2. *Iterations.* Assume that $\beta = \alpha \cdot \gamma$, where by definition $\alpha \cdot \gamma := \gamma \circ \alpha$. Then we can produce $C(b_{\beta})$ by iterating two times the preceding algorithm. We get a sequence of nested disks

$$D'' \subset D' \subset D,$$

a subdivision of the segment $[x_j, x''_{\alpha \cdot \gamma(j)}]$ into sub-arcs

$$[x_j, x''_{\alpha \cdot \gamma(j)}] = [x_j, x'_{\alpha(j)}] \cup [x'_{\alpha(j)}, x''_{\alpha \cdot \gamma(j)}],$$

and a factorization of the permutation associated to $[x_j, x''_{\alpha \cdot \gamma(j)}]$:

$$\epsilon_{\beta}(j) = \epsilon_{\alpha}(j) \cdot \epsilon_{\gamma}(\alpha(j)).$$

The “adjacent” transpositions (01), (12), (23) form a standard set of generators of S_4 . Let $\beta = \tau_1 \cdots \tau_k$ where each τ_j is such a generator. Then we can produce again $C(b_{\beta})$ by iterating k times the algorithm. We get a sequence of nested disks

$$D_k \subset \dots \subset D_1 \subset D,$$

a subdivision of the arc $[x_j, x_{\beta(j)}^{(k)}]$ of $C(b) \setminus \text{Int}(D_k)$ into sub-arcs

$$[x_j, x_{\tau_1(j)}^{(1)}] \cup [\dots] \cup [x_{\tau_1 \cdots \tau_{k-1}(j)}^{(k-1)}, x_{\beta(j)}^{(k)}],$$

and a factorization of the permutation $\epsilon_{\beta}(j) \in S_3$ associated to the arc $[x_j, x_{\beta(j)}^{(k)}]$:

$$\epsilon_{\beta}(j) = \epsilon_{\tau_1}(j) \cdots \epsilon_{\tau_k}(\tau_1 \cdots \tau_{k-1}(j)) =: f_1 \cdots f_k.$$

Lemma 6.2. *Every transposition $f_i \in S_3$ belongs to the set $\{\text{Id}, (01), (12)\}$.*

Proof. It is enough to check the case $\beta = \tau_j$ (see Figure 13). \square

For future application, it is convenient to associate a symbol to each sub-arc of $C(b_{\beta})$, that indicates its label $f \in \{\text{Id}, (01), (12)\}$ and depends on its b -orientation:

- If the b -orientation is outgoing (i.e. opposite to the one pointing towards the center of D), then let this symbol be $< \text{Id} <$, or $< f >$ (a “source”) if f is a transposition;
- if the b -orientation is ingoing, then let this symbol be $> \text{Id} >$, or $> f <$ (a “pit”) if f is a transposition.

Under concatenation of sub-arcs, adjacent symbols are separated by \ll or \gg . Here are examples:

$$\cdots \langle f \gg \rangle g \langle \cdots, \cdots \rangle f \ll \langle g \rangle \cdots, \cdots \rangle id \gg \rangle id \rangle \cdots$$

$$\cdots \langle id \ll \langle id \langle \cdots, \cdots \rangle id \gg \rangle f \langle \cdots, \cdots \rangle id \ll \langle f \rangle \cdots, \text{ etc.}$$

Letting $? \in \{\ll, \gg\}$, in the case of $\epsilon_\beta(j) = f_1 \cdots f_k$ we get:

- $\langle f_1? \cdots ? f_k \langle \text{ or } \rangle f_1? \cdots ? f_k \rangle$ if b and b_β induce on $[x_j, x_{\beta(j)}^{(k)}]$ the same orientation;
- $\langle f_1? \cdots ? f_k \rangle \text{ or } \rangle f_1? \cdots ? f_k \langle$ otherwise.

We can also make simplifications of the form:

$$?s \gg \rangle f \ll \langle f \gg \rangle t? = ?s \gg \rangle t?$$

and so on, so that the concatenation of symbols of sub-arcs eventually reduces to $\rangle \text{Id} \langle$, $\langle \text{Id} \langle$, or alternating pits and sources carrying the transpositions (01) and (12). In Figure 13 we show graphically the result when $*_b = +1$ and β varies in $\{(01), (12), (23)\}$; we dropped the notation Id for the symbol $\rangle \text{Id} \langle$. The case $*_b = -1$ is similar.




FIGURE 13. $C(b_{ij})$, $*_b = +1$.

6.2.3. Oriented C -moves. We call $C(b) \rightarrow C(b_\beta)$ a C -move. If $C(b)$ and $C(b_\beta)$ define the same pre-branching then we say that it is an *oriented C -move*. Set


$$\sigma = (0123) = (23) \cdot (12) \cdot (01), \quad \tau = (02) \cdot (13).$$

σ generates a subgroup \mathfrak{G} of $S(J_4)$ isomorphic to $(\mathbb{Z}/4\mathbb{Z}, +)$ via the map $\sigma \mapsto 1$.

Lemma 6.3. *For both $*_b = \pm 1$, $\beta \in S_4$ induces an oriented C -move $C(b) \rightarrow C(b_\beta)$ iff $\beta \in \mathfrak{G}$.*

Proof. Looking at Figure 2 anti-clockwise from the top left, and denoting by b the first branching (so that $*_b = +1$), the others are successively b_τ , $b_{\sigma \cdot \tau}$, $b_{\tau \cdot \sigma \cdot \tau}$, $b_{\sigma^{-1} \cdot \tau \cdot \sigma \cdot \tau} = b$. Hence $\beta \in \mathfrak{G}$. One can do similarly when $*_b = -1$. \square

Figure 14 shows the oriented C -moves $C(b) \rightarrow C(b_{\sigma^{-1}})$ for $*_b = +1$ and $C(b) \rightarrow C(b_\sigma)$ for $*_b = -1$.

FIGURE 14. Generating oriented C -moves.

6.2.4. *Globalization.* Let Γ be an \mathcal{N} -graph encoding (T, \tilde{b}) , and \tilde{b}' another weak branching on T . We want to produce an \mathcal{N} -graph Γ' encoding (T, \tilde{b}') . For every abstract branched tetrahedron (Δ_j, b_j) of (T, \tilde{b}) there is a permutation $\beta_j \in S(J_4)$ such that the restriction of the transit $\tilde{b} \rightarrow \tilde{b}'$ to Δ_j is $b_j \rightarrow (b_j)_{\beta_j}$. Applying at every crossing of Γ the algorithm $C(b_j) \rightarrow C((b_j)_{\beta_j})$, we get a diagram Γ' . Forgetting the orientations, there is a natural 1 – 1 correspondence between the edges of Γ and Γ' . Let e be an edge of Γ with color $r(e)$. Denote by \bar{e} the corresponding edge of Γ' . According to the last step of the algorithm, \bar{e} is labelled by “initial” and “final” permutations $\hat{e}^i(\bar{e}), \hat{e}^f(\bar{e}) \in S(J_3)$. There is an alternative:

- Both $\hat{e}^i(\bar{e})$ and $\hat{e}^f(\bar{e})$ belong to $A(J_3)$ (in such a case e and \bar{e} have compatible orientations); then the color $r(\bar{e})$ is defined by

$$\hat{e}^i(\bar{e}) \cdot (012)^{r(e)} \cdot \hat{e}^f(\bar{e}) = (012)^{r(\bar{e})}.$$

- Both $\hat{e}^i(\bar{e})$ and $\hat{e}^f(\bar{e})$ do not belong to $A(J_3)$ (in such a case e and \bar{e} have opposite orientations); then the color $r(\bar{e})$ is defined by

$$\hat{e}^i(\bar{e}) \cdot (012)^{-r(e)} \cdot \hat{e}^f(\bar{e}) = (012)^{r(\bar{e})}.$$

The oriented C -moves have the nice feature to be local and independent one to each other. On the contrary, a family of permutations $\beta_j \in S(J_4)$ as above must satisfy non trivial global constraints to induce a change of weak branching $\tilde{b} \rightarrow \tilde{b}'$. Fortunately, the following local/global calculus on \mathcal{N} -graphs (introduced in [9]) covers the general case.

Lemma 6.4. *Any change of \mathcal{N} -graphs $\Gamma \rightarrow \Gamma'$ corresponding to a change of weak branching $(T, \tilde{b}) \rightarrow (T, \tilde{b}')$ is a composition of a finite sequence of moves $\Gamma_j \rightarrow \Gamma_{j+1}$ of the following types:*

- (1) *The moves of Subsection 6.1;*
- (2) *The oriented C -moves;*
- (3) *(A non-local “circuit” move). Let γ be a simple circuit of Γ_j , ie. a circuit with one component. Assume that each edge of γ is the upper strand at each crossing through which γ passes. Then Γ_{j+1} is obtained by acting on each of these crossings with the transposition $(23) \in S_4$. Equivalently, Γ_{j+1} is obtained by reversing the orientation of γ and keeping the $\mathbb{Z}/3\mathbb{Z}$ -color $r(e)$ unchanged at every edge e of $\text{Sing}(P)$.*

Proof. The equivalence between the two descriptions of the non-local “circuit” move is clear. Let $\Gamma \rightarrow \Gamma'$ be as in the statement. The set \mathcal{S} of edges of $\text{Sing}(P)$ where the pre-branchings associated to \tilde{b} and \tilde{b}' disagree is the union of non-overlapping simple circuits $\gamma_1, \dots, \gamma_r$ oriented by \tilde{b} . Using

oriented C -moves one can modify Γ so that γ_1 eventually passes over at each crossing where it goes through. Then change its orientation. Doing similarly with $\gamma_2, \dots, \gamma_r$, one ends up with $\mathcal{S} = \emptyset$. \square

6.3. Transits. It is well-known that two “naked” triangulations T and T' of \hat{V} can be connected by a finite sequence of 3-dimensional Pachner’s moves, also called MP and bubble moves. The MP moves, also called “2 \leftrightarrow 3” moves, are illustrated in Figure 15 in terms of triangulations and dual spines. Branched versions of the bubble moves are shown in Figure 16 in terms of \mathcal{N} -graphs.




FIGURE 15. The MP move.

Definition 6.5. Let (T, σ) and (T', σ') be pre-branched triangulations of \hat{V} . A move $T \rightarrow T'$ (either bubble or MP, positive or negative) lifts to a *pre-branching transit* $(T, \sigma) \rightarrow (T', \sigma')$ if and only if σ and σ' coincide on every common 2-face of T and T' . We say that $T \rightarrow T'$ lifts to a \tilde{b} -transit $(T, \tilde{b}) \rightarrow (T', \tilde{b}')$ of weakly-branched triangulations if it induces a pre-branching transit $(T, \sigma_{\tilde{b}}) \rightarrow (T', \sigma_{\tilde{b}'})$. If (T, b) and (T', b') are branched triangulations, then $(T, b) \rightarrow (T', b')$ is a *b-transit* if b and b' coincide on every common edge of T and T' .

Figure 16 shows the different bubble b -transits in terms of normal \mathcal{N} -graphs (recall Remark 2.11), and the decorations A, B of the dual square edges according to Figure 8. Figure 17 and 18 show an example of MP b -transit in terms of branched triangulations and normal \mathcal{N} -graphs; note that $*_b = +1$ for all 3-simplices (resp. dotted crossings). It is called the *Schaeffer b-transit*, and plays a distinguished role in the study of matrix dilogarithms (see Section 8).

The following Lemma will be useful in applications to the QH state sums. Denote by \cong_{pb} the equivalence relation on the set of weakly branched triangulations, defined by \tilde{b} -transits $(T, \tilde{b}) \rightarrow (T', \tilde{b}')$.

Lemma 6.6. *The equivalence relation \cong_{pb} is generated by the following two transformations:*

- *Change the weak-branching $(T, \tilde{b}) \rightarrow (T', \tilde{b}')$ by moves preserving the pre-branching (ie. dually do oriented C -moves, see Lemma 6.4).*
- *Perform \tilde{b} -transits $(T, \tilde{b}) \rightarrow (T', \tilde{b}')$ which look like b -transits at the sub-patterns of tetrahedra involved in the move.*

Proof. The proof for bubble moves is easy and basically illustrated in Figure 16 in terms of \mathcal{N} -graphs. Consider a positive MP \tilde{b} -transit $(T, \tilde{b}) \rightarrow (T', \tilde{b}')$, and the two tetrahedra of T modified by the transit. At their common 2-face F one of the following possibilities is realized:

- (1) there are exactly two square edges which are both *not monochromatic* (i.e. each one inherits different A, B labels from the pre-branchings of the two tetrahedra);




FIGURE 16. Bubble b -transits.




FIGURE 17. Schaeffer's b -transits.





FIGURE 18. Normal \mathcal{N} -graph Schaeffer's b -transit.

(2) there is exactly one square edge which is monochromatic (either A or B), and the other two edges are monochromatic (either B or A coupled with the empty label).

Moreover, this information at F determines completely the co-orientations of the three 2-faces produced by the transit, and hence determines the pre-branching transit $(T, \sigma_{\bar{b}}) \rightarrow (T', \sigma_{\bar{b}'})$. Then, it is enough to realize the above possibilities by means of b -transits (changing the weak branchings before and/or after the transit, if necessary, by moves preserving the pre-branching). Figure 19 shows it in terms of (decoded) normal \mathcal{N} -graphs. Clearly this realization is not unique. \square

FIGURE 19. Branched realization of MP pb -transits.

7. STATE SUM INVARIANCE OVER WEAKLY BRANCHED TRIANGULATIONS

Let (T, \tilde{b}, w, f, c) be a weakly branched QH triangulation of a pattern \mathcal{P} ; w and f verify the defining equations of the spaces in Definition 3.7, and c those of Definition 4.7 (resp. 5.1) if \mathcal{P} is a pattern over a cusped manifold (resp. a QHFT pattern).

We want to prove that $\mathcal{H}_N(\mathcal{P}) := \pm \mathcal{H}_N(T, \tilde{b}, w, f, c)$ is an invariant of \mathcal{P} . In the case when (T, b) is a branched triangulation we achieved this result in [3, 4, 5] by introducing the notion of *QH transit* $(T, b, w, f, c) \rightarrow (T', b', w', f', c')$. A QH transit is such that: $(T, b) \rightarrow (T', b')$ is a b -transit, the tuples (w, f, c) and (w', f', c') coincide on the “common” tetrahedra of T and T' , and for every common edge e we have (see Section 3.2 for the notations, where $W_T(e)$ is $W(e)$, and so on)

$$(60) \quad W_T(e) = W_{T'}(e) , \quad L_T(e) = L_{T'}(e) , \quad C_T(e) = C_{T'}(e).$$

QH triangulations related by QH transits encode a same pattern \mathcal{P} . Moreover:

- $\mathcal{H}_N(T, b, w, f, c)$ is invariant under any QH transit.
- Any two branched QH triangulations of \mathcal{P} are related by a finite sequence of QH transits and branching changes.
- $\mathcal{H}_N(T, b, w, f, c)$ is invariant under any change of the branching b .

Clearly these three facts prove the invariance of $\mathcal{H}_N(\mathcal{P})$. We follow the same strategy when \tilde{b} is an arbitrary weak branching. Since the QH transits are local transformations, we define QH transits $(T, \tilde{b}, w, f, c) \rightarrow (T', \tilde{b}', w', f', c')$ by requiring that the \tilde{b} -transit $(T, \tilde{b}) \rightarrow (T', \tilde{b}')$ restricts to a genuine b -transit on the tetrahedra modified by the move. Then, trivially $\mathcal{H}_N(T, \tilde{b}, w, f, c)$ is again invariant under QH transits, and using Lemma 6.6 and the results of [3, 4, 5] it is easy to see that any two weakly branched QH triangulations of \mathcal{P} are related by a finite sequence of QH transits and changes of weak branching. So it remains to show:

Proposition 7.1. *For any change of weak branching $\tilde{b} \rightarrow \tilde{b}'$ we have*

$$\mathcal{H}_N(T, \tilde{b}, w, f, c) \equiv_N \pm \mathcal{H}_N(T, \tilde{b}', w, f, c).$$

In order to develop the proof we need to fix some conventions regarding the contraction of tensors, and to describe the relations between tetrahedral tensors related by a change of branching. This is the content of the two following subsections.

7.1. Formal conversion of tensor networks. Recall the notation $V = \mathbb{C}^N$. We use the canonical isomorphism $V \rightarrow V^*$ sending the standard basis to the dual basis. Let $A \in \text{End}(\mathbb{C}^N \otimes \mathbb{C}^N)$ be

associated to a $2 - 2$ \mathcal{N} -tangle as in Figure 9, according to the conventions of Section 3.1. Let $\beta \in S(J_4)$. Assuming that $*_b = 1$, under the natural isomorphisms

$$(61) \quad \text{Hom}(V_3 \otimes V_1, V_2 \otimes V_0) \cong V_2 \otimes V_0 \otimes (V_3 \otimes V_1)^* \cong V_2 \otimes V_0 \otimes V_1^* \otimes V_3^*$$

we see that A belongs to $V_2 \otimes V_0 \otimes V_1^* \otimes V_3^*$, where as usual V_j is the copy of \mathbb{C}^N associated to the j -th face of (Δ, b) . Consider $W_2 \otimes W_0 \otimes W_1^* \otimes W_3^*$ if $*_{b_\beta} = 1$, and $W_3 \otimes W_1 \otimes W_0^* \otimes W_2^*$ if $*_{b_\beta} = -1$, where W_j is the copy of \mathbb{C}^N associated to the j -th-face according to b_β . By using $\text{Id} : V_j \rightarrow W_{\beta(j)}$ or the canonical isomorphisms $V_j \rightarrow W_{\beta(j)}^*$, we get further canonical isomorphisms

$$V_2 \otimes V_0 \otimes V_1^* \otimes V_3^* \cong W_2 \otimes W_0 \otimes W_1^* \otimes W_3^* \quad \text{and} \quad V_2 \otimes V_0 \otimes V_1^* \otimes V_3^* \cong W_1 \otimes W_3 \otimes W_0^* \otimes W_2^*.$$

Denote by A_β the operator supported by (Δ, b_β) , defined as the image of A via such an isomorphism. We call A_β the *formal β -conversion* of A . There are explicit identities between the matrix elements of A and A_β . Following our conventions for the index positions, we get for example

$$A_{s,t}^{p,q} = (A_{(13)(02)})_{t,s}^{q,p} = (A_{(23)})_{s,q}^{p,t} = (A_{(02)})_{q,p}^{s,t} = (A_{(01)})_{p,t}^{s,q} = (A_{(12)})_{t,q}^{s,p}.$$

Let \mathcal{A} be a QH tensor network supported by a weakly branched triangulation (T, \tilde{b}) . For any change of weak branching $\tilde{b} \rightarrow \tilde{b}'$, the formal conversions of the tetrahedral tensors of \mathcal{A} match to produce a new tensor network $\mathcal{A}_{\tilde{b}'}$, supported by (T, \tilde{b}') . The state sums associated to \mathcal{A} and $\mathcal{A}_{\tilde{b}'}$, being total contractions, it is tautologically evident that they take the same values.




FIGURE 20. Formal conversion of a square matrix.




FIGURE 21. Relations for M -conversion.

The formal conversion of tensors can be widely applied. Here is a second example that we use below. Let $M = (M_j^i)$ be any $N \times N$ matrix. We interpret M as an endomorphism of $V = \mathbb{C}^N$. There are two formal conversions of M , M_u and M_d , which are bilinear forms on V^* and on V respectively, defined by:

$$(62) \quad (M_u)^{i,j} = M_j^i = (M_d)_{i,j}.$$

Note that we have also $(M_u)^{i,j} = (M^t)_i^j = (M_d)_{i,j}$, where M^t is the transpose matrix of M , that is, the adjoint endomorphism. Figure 20 shows a graphical encoding of these identities. If M is symmetric, the position of the indices i, j is immaterial. Such formal conversions satisfy a few relations shown in Figure 21, where A, B and C denote $N \times N$ matrices and we assume that $AC = CA$. Note that

in every case the index k is traced out, and the top/right picture represents the matrix element of a composite endomorphism

$$A \cdot B := B \circ A.$$

7.2. The relations between $\mathcal{R}_N(\Delta, b, w, f, c)$ and $\mathcal{R}_N(\Delta, b_\beta, w', f', c')$. Consider a change of branching $b \rightarrow b_\beta$, $\beta \in S(J_4)$. It is enough to treat the case where β is one of the standard generators $(01), (12), (23) \in S(J_4)$. Consider the algorithm $C(b) \rightarrow C(b_\beta)$ of Section 6.2. Associate a tensor network to $C(b_\beta)$ as follows:

- Associate $\mathcal{R}_N(\Delta, b_\beta, w', f', c')$ to the dotted crossing of $C(b_\beta)$.
- Using the matrices S and T of Section 3.2, replace $> (01) <$ with $> T <$, $> (12) <$ with $> S <$, $< (01) >$ with $< T^{-1} >$, $< (12) >$ with $< S^{-1} >$, and $> \text{Id} >$ with $> I >$.

Contracting the tensor network we get a tensor $\hat{\mathcal{R}}_N(T, b_\beta, w', f', c')$ which has the same type as $\mathcal{R}_N(\Delta, b, w, f, c)$, since both have the same oriented arc-germs at ∂D . The next Proposition rephrases the content of Lemma 3.3 of [3] and Corollary 5.6 of [4]. Matrix indices are raised or lowered using the standard inner product of \mathbb{C}^N (see Section 7.1).

Proposition 7.2. *If $\beta = (01), (12), (23)$, then*

$$\mathcal{R}_N(T, b, w, f, c) \equiv_N \pm \hat{\mathcal{R}}_N(T, b_\beta, w', f', c').$$

More precisely, exchanging the roles of b and b_β and setting $\epsilon_N = (-1)^{\frac{N-1}{2}}$, we have:

$$\mathcal{R}_N(\Delta, b_{(01)}, w', f', c')_{k,j}^{i,l} \equiv_N \epsilon_N^{c_0} T_{k,k'} \mathcal{R}_N(\Delta, b, w, f, c)_{i',j}^{k',l} (T^{-1})^{i',i}$$

$$\mathcal{R}_N(\Delta, b_{(12)}, w', f', c')_{j,l}^{i,k} \equiv_N \epsilon_N^{c_1} T_{l,l'} \mathcal{R}_N(\Delta, b, w, f, c)_{i',j}^{k,l'} (S^{-1})^{i',i}$$

$$\mathcal{R}_N(\Delta, b_{(23)}, w', f', c')_{i,l}^{k,j} \equiv_N \epsilon_N^{c_0} S_{l,l'} \mathcal{R}_N(\Delta, b, w, f, c)_{i,j'}^{k,l'} (S^{-1})^{j',j}.$$

The main properties of S and T are described in the following Lemma.

Lemma 7.3. *We have:*

- (1) $S^4 = I_N$.
- (2) $(TS)^3 = \phi_N S^2$ where $\phi_N \in \{\pm 1, \pm i\}$ is given by

$$\phi_N = \begin{cases} \left(\frac{m+1}{N} \right) & \text{if } N \equiv 1 \pmod{4} \\ \left(\frac{m+1}{N} \right) i & \text{if } N \equiv 3 \pmod{4}. \end{cases}$$

- (3) $(S^{-1}T)^3 = \phi_N I_N$.

Proof. (1) is immediate. We have $((TS)^2)_i^j = N^{-1} \zeta^{-(m+1)i^2 - ij} \sum_{k=0}^{N-1} \zeta^{(m+1)(k-i-j)^2}$. For all non vanishing coprime integers a, b with $b > 0$ set

$$G(a, b) = \sum_{x \pmod{b}} e^{2\pi\sqrt{-1}ax^2/b}.$$

The sum in $((TS)^2)_i^j$ has this form for $a = m+1$, $b = N$. By [28, page 86-87] we have

$$G(a, b) = \left(\frac{a}{b} \right) G(1, b), \quad b \text{ odd}$$

where $G(1, b) = \sqrt{b}$ if $b \equiv 1 \pmod{4}$, and $G(1, b) = i\sqrt{b}$ if $b \equiv 3 \pmod{4}$. Hence $((TS)^2)_i^j = \phi_N N^{-1/2} \zeta^{-(m+1)i^2 - ij}$, and (2) follows easily from this. As $TS^{-1} = S(S^{-1}T)S^{-1}$, it is enough to prove the last statement for TS^{-1} . This follows: $(TS)^4 = (TS)(TS)^3 = \phi_N TS^3 = \phi_N TS^{-1}$; $(\phi_N TS^{-1})^3 = ((TS)^4)^3 = \phi_N^4 I_N$; finally $(TS^{-1})^3 = \phi_N I_N$. \square

7.3. Proof of Proposition 7.1. By Lemma 6.4 we can assume that $(T, \tilde{b}) \rightarrow (T, \tilde{b}')$ corresponds to an oriented C -move or a circuit move on the associated \mathcal{N} -graphs. A circuit move preserves the $\mathbb{Z}/3\mathbb{Z}$ -color $r(e)$ of every edge e , and produces one extremal pit and one extremal source associated to the matrices S and S^{-1} at the endpoints of each edge of the circuit. Hence $\mathcal{H}_N(T, \tilde{b}, w, f, c)$ is preserved, as it is the contraction of a tensor network. Consider the generating oriented C -move associated to $\sigma = (0123) \in S(J_4)$, and the tensor $\hat{\mathcal{R}}_N(T, b_\sigma, w', f', c')$ resulting from the formal conversion of the sequence

$$C(b) \longrightarrow C(b_{(23)}) \longrightarrow C(b_{(23)(12)}) \longrightarrow C(b_{(23)(12)(01)}) = C(b_\sigma)$$

as before Proposition 7.2. From Figure 14 we see that an edge e' of $C(b_\sigma)$ carries the matrix $\mathcal{Q}_N = S \cdot T^{-1}$ if $r(e') = 1 + r(e) \in \mathbb{Z}/3\mathbb{Z}$ (e being the arc of $C(b)$ associated to e'), and that e' carries $\mathcal{Q}_N^{-1} = T \cdot S^{-1}$ if $r(e') = -1 + r(e) \in \mathbb{Z}/3\mathbb{Z}$. Otherwise it carries the identity matrix. By the relation $\mathcal{Q}_N^3 = \phi_N^{-1} \mathbf{I}_N$ proved in Lemma 7.3, the sum in $\mathcal{H}_N(T, \tilde{b}, w, f, c)$ is changed by a factor $\pm i$ when $r(e') = 1 + r(e)$ and $r(e) = 2$, or when $r(e') = -1 + r(e)$ and $r(e) = 0$. The term $\phi_N^{-q(T, \tilde{b})}$ of $a_N(T, \tilde{b})$ compensates exactly such factors. \square

8. RESOLUTION OF THE SIGN ANOMALY

In this Section we prove Theorem 1.1 (3) and the analogous result for QHFT partition functions:

Proposition 8.1. *Let \mathcal{P} be a QHFT pattern or a pattern over a cusped manifold. If $N - 1 \equiv 0 \pmod{4}$, or $N - 1 \equiv 2 \pmod{4}$ and the bulk c -weight of \mathcal{P} vanishes, then $\mathcal{H}_N(\mathcal{P})$ is defined up to multiplication by N -th roots of unity.*

Proof. We combine several results described below in Sections 8.1 and 8.2. By Corollary 8.6, for any odd $N \geq 3$ the invariance of $\mathcal{H}_N(\mathcal{T})$ under QH transits holds true up to multiplication by N -th roots of unity. If $N - 1 \equiv 0 \pmod{4}$, we have $\epsilon_N = 1$, so $\mathcal{H}_N(\mathcal{T})$ has no sign anomaly with respect to any change of weak branching (Proposition 7.2). If $N - 1 \equiv 0 \pmod{4}$ and $h_c = 0$, the same is true by Lemma 8.7 and 8.8. Then the conclusion follows as explained before Proposition 7.1 \square

Let \mathcal{P} be a pattern for which $\mathcal{H}_N(\mathcal{P})$ can be defined by using branched triangulations, as in [3, 4, 5]. Then $a_N(T, b) = N^{-v}$ or 1, so we can wonder if the sign anomalies occurring in Proposition 7.2 disappear because of global compensations. This is eventually true.

Proposition 8.2. *For any pattern \mathcal{P} as above, $\mathcal{H}_N(\mathcal{P})$ is defined up to multiplication by a N -th root of unity, with no assumption on the weights of \mathcal{P} .*

Proof. By a result of Costantino [13], any change of branching can be achieved by means of a finite sequence of branching transits. The conclusion follows from Corollary 8.6. \square

Remark 8.3. One can ask if a weakly branched version of Costantino's result holds true: given weakly branched triangulations (T, \tilde{b}) and (T, \tilde{b}') of \hat{V} that differ only by the weak branchings, are they related by a sequence of oriented C -moves and "local" b -transits? The answer is negative. In fact, we know that the state sums \mathcal{H}_N have no sign anomaly with respect to QH transits and oriented C -moves, but there are examples of weakly branched triangulations (T, \tilde{b}) admitting circuit moves along circuits γ such that $h_c(\gamma) \neq 0 \in \mathbb{Z}/2\mathbb{Z}$.

8.1. Behaviour of the sign anomaly under QH transits. Recall the formula (19). The matrix dilogarithms \mathcal{R}_N satisfy Proposition 7.2, whereas the *basic* dilogarithms \mathcal{L}_N satisfy the relations ([4], Proposition 5.3):

$$(63) \quad \begin{aligned} \mathcal{L}_N(\Delta, b_{(01)}, w', f', c')_{k,j}^{i,l} &\equiv_N (w'_0)^{\frac{1-N}{2}} T_{k,k'} \mathcal{L}_N(\Delta, b, w, f, c)_{i',j'}^{k',l} (T^{-1})^{i',i} \\ \mathcal{L}_N(\Delta, b_{(12)}, w', f', c')_{j,l}^{i,k} &\equiv_N (w'_1)^{\frac{N-1}{2}} T_{l,l'} \mathcal{L}_N(\Delta, b, w, f, c)_{i',j'}^{k,l'} (S^{-1})^{i',i} \\ \mathcal{L}_N(\Delta, b_{(23)}, w', f', c')_{i,l}^{k,j} &\equiv_N (w'_0)^{\frac{1-N}{2}} S_{l,l'} \mathcal{L}_N(\Delta, b, w, f, c)_{i',j'}^{k,l'} (S^{-1})^{j',j}. \end{aligned}$$

Define the transits of N -th root cross ratio moduli like in (60), by replacing $W(e)$ with the total N -th root modulus $W'(e)$ at all edges e of T and T' . Clearly any QH transit induces a transit of N -th root cross ratio moduli.

Theorem 8.4. ([4], Theorem 5.2) *The basic matrix dilogarithm identity supported by the Schaeffer $2 \leftrightarrow 3$ transit of N -th root cross ratio moduli holds true up to an N -th root of unity anomaly.*

The proof goes as follows. The basic matrix dilogarithm identity supported by the Schaeffer $2 \leftrightarrow 3$ QH transit is equivalent to the following equality in $\text{Aut}(\mathbb{C}^N \otimes \mathbb{C}^N \otimes \mathbb{C}^N)$:

$$(64) \quad \Psi_{23}^1(V)\Psi_{12}^3(U) = \Psi_{12}^4(U)\Psi_{13}^2(-UV)\Psi_{23}^0(V)$$

where U and V are explicit $N^2 \times N^2$ matrices satisfying $U^N = -Id \otimes Id$, $V^N = -Id \otimes Id$, and $VU = \zeta UV$, ζ being a primitive N -th root of 1, and for a tensor A satisfying $A^N = -Id$ (on the appropriate space: \mathbb{C}^N , $\mathbb{C}^N \otimes \mathbb{C}^N$, etc.) one sets

$$(65) \quad \Psi^i(A) = \sum_{l=0}^{N-1} A^l \prod_{s=1}^l \frac{((w_0^i)')^{-1}((w_1^i)')^{-1}}{1 - ((w_0^i)')^{-1}\zeta^{-s}}.$$

In (64), $\Psi_{23}^1(V)$ means $\Psi^1(V)$ acting on the second and third tensorands, i.e. $Id \otimes \Psi^1(V)$, etc. One can check that Ψ^i is uniquely determined up to multiplication by scalars by the functional relation

$$(66) \quad \Psi^i(\zeta^{-1}A) = \Psi^i(A) \left(\frac{1 - ((w_0^i)')^{-1}((w_1^i)')^{-1}A}{((w_0^i)')^{-1}} \right) = \Psi^i(A) ((w_0^i)' - ((w_1^i)')^{-1}A).$$

The identity (64) lifts to the algebra generated by U and V , so we have to show

$$(67) \quad \Psi^1(V)\Psi^3(U) = \Psi^4(U)\Psi^2(-UV)\Psi^0(V)$$

when $U^N = -1$, $V^N = -1$, and $VU = \zeta UV$. To do it, one uses crucially the relations between N -th root cross-ratio moduli implied by the Schaeffer QH transit. One shows that UV commutes with $P(-UV) := \Psi^4(U)^{-1}\Psi^1(V)\Psi^3(U)\Psi^0(V)^{-1}$; hence $P(-UV)$ is a function of UV . Moreover, it satisfies

$$(68) \quad P(-\zeta^{-1}UV) = P(-UV) ((w_0')^2 - ((w_1')^2)^{-1} (-UV)).$$

Comparing with (66) we see that $P(-UV)$ and $\Psi^2(-UV)$ are equal up to a multiplicative constant $C := C((w_0')^2, (w_1')^2)$. We introduce an N -th root of the reciprocal of $\det(\Psi^i(A))$ in each entry of $\Psi^i(A)$ so that $\det(\Psi^i(A)) = 1$ (this is the term $h((w_0^i)') = g((w_0^i)')/g(1)$ in (18)). Then $\det(P(-UV)) = 1 = C^N \det(\Psi^2(-UV))$ and $C^N = 1$. As a result the tensor $\mathcal{L}_N(w_0', w_1')$ can be expressed in terms of $\Psi(A)$ for a suitable A (see [4], formulas (32) and (33)). \square

The basic matrix dilogarithm identities hold true up to an N -th root of unity anomaly only for some $2 \leftrightarrow 3$ branching transits. From (63) one deduces the following table of the anomalies produced by the symmetries of the Schaeffer $2 \leftrightarrow 3$ transit (it is enough to consider those induced by the transpositions (01), \dots , (34) on the vertex ordering used in Figure 17). The transits of N -th root moduli imply that the anomalies are equal for the symmetries induced by (01) and (34), but differ for (12) and (23) because of the occurrence of the opposite exponents $(1-N)/2$, $(N-1)/2$ for the N -th root moduli $(w_0^k)'$, $(w_1^k)'$ respectively. For instance $(w_1^0)'(w_0^4)' = (w_1^1)'$, but we see $((w_1^0)')^{\frac{N-1}{2}}((w_0^4)')^{\frac{1-N}{2}} \neq ((w_1^1)')^{\frac{N-1}{2}}$.

	Δ^1	Δ^3	Δ^0	Δ^2	Δ^4
(01)	1	$((w_0^3)')^{\frac{1-N}{2}}$	1	$((w_0^2)')^{\frac{1-N}{2}}$	$((w_0^4)')^{\frac{1-N}{2}}$
(12)	1	$((w_1^3)')^{\frac{N-1}{2}}$	$((w_0^0)')^{\frac{1-N}{2}}$	1	$((w_1^4)')^{\frac{N-1}{2}}$
(23)	$((w_1^1)')^{\frac{N-1}{2}}$	1	$((w_1^0)')^{\frac{N-1}{2}}$	1	$((w_0^4)')^{\frac{1-N}{2}}$
(34)	$((w_0^1)')^{\frac{1-N}{2}}$	1	$((w_0^0)')^{\frac{1-N}{2}}$	$((w_0^2)')^{\frac{1-N}{2}}$	1

Replacing the tensors \mathcal{L}_N with \mathcal{R}_N we have:

Proposition 8.5. *The matrix dilogarithm identities hold true up to an N -th root of unity anomaly for all QH transits.*

Proof. Since we have the identity (64), in the case of the Schaeffer QH transit identity it is enough to compare the symmetrization factors at both sides. They are $(\epsilon_N := (-1)^{\frac{N-1}{2}})$:

$$\epsilon_N^{f_0^1 c_1^1 + f_1^1 c_0^1 + f_0^3 c_1^3 + f_1^3 c_0^3} \exp\left(\frac{N-1}{2N} (-c_{10}^1 + c_{01}^1 - c_{10}^3 + c_{01}^3)\right)$$

and

$$\epsilon_N^{f_0^0 c_1^0 + f_1^0 c_0^0 + f_0^2 c_1^2 + f_1^2 c_0^2 + f_0^4 c_1^4 + f_1^4 c_0^4} \exp\left(\frac{N-1}{2N} (-c_{10}^0 + c_{01}^0 - c_{10}^2 + c_{01}^2 - c_{10}^4 + c_{01}^4)\right).$$

A computation shows that the transits of log-branches and charges imply that the two exponentials are equal, and with the flattening transit mod(2) that the signs are equal. To prove the claim for all others $2 \leftrightarrow 3$ QH transits we compare the anomalies resulting from symmetries on the Schaeffer QH transit. Since all QH transits are branched, the actions by the matrices S, T due to Proposition 7.2 cancel out along interior faces. The following table shows the sign anomalies. Again, the transit of charges implies that they are equal at both sides.

	Δ^1	Δ^3	Δ^0	Δ^2	Δ^4
(01)	1	ϵ_N^3	1	ϵ_N^2	ϵ_N^4
(12)	1	ϵ_N^3	ϵ_N^0	1	ϵ_N^4
(23)	ϵ_N^1	1	ϵ_N^0	1	ϵ_N^4
(34)	ϵ_N^1	1	ϵ_N^0	ϵ_N^2	1

As usual, the identities for the bubble moves are formal consequence of those for the $2 \leftrightarrow 3$ moves. This concludes the proof. \square

Corollary 8.6. *Let $(T, \tilde{b}, w, f, c) \rightarrow (T', \tilde{b}', w', f', c')$ be any QH transit of weakly branched QH triangulations of a pattern \mathcal{P} . Then*

$$\mathcal{H}_N(T, \tilde{b}, w, f, c) \equiv_N \mathcal{H}_N(T', \tilde{b}', w', f', c').$$

Proof. If the transit is supported by a MP move it is immediate that $a_N(T, \tilde{b}) = a_N(T', \tilde{b}')$, hence the conclusion follows from Corollary 8.5. If it is supported by a positive bubble move, spelling the contribution of each term we realize that if

$$a_N(T, \tilde{b}) = N^{-v} \phi_N^{-q(T, \tilde{b})} c_N(T, \tilde{b}), \quad c_N(T, \tilde{b}) = \epsilon_N^{v+l-\frac{1}{2}\sum_e(n_+(e)-n_-(e))}$$

then

$$a_N(T', \tilde{b}') = N^{-(v+1)} \phi_N^{-q(T', \tilde{b}')} c_N(T', \tilde{b}'), \quad c_N(T', \tilde{b}') = \epsilon_N^{(v+1)+(l+3)-(2+\frac{1}{2}\sum_e(n_+(e)-n_-(e))}$$

and $\phi_N^{-q(T', \tilde{b}')} = \phi_N^{-q(T, \tilde{b})}$. Hence $a_N(T', \tilde{b}') = N^{-1} a_N(T, \tilde{b})$, as it must be in order to ensure the invariance under the bubble move, which increases by 1 the number of internal vertices. \square

8.2. Behaviour of the sign anomaly under a change of weak branching. A change of weak branching $\tilde{b} \rightarrow \tilde{b}'$ modifies $a_N(T, \tilde{b})$ by its sub-factors $\phi_N^{-q(T, \tilde{b})}$ and

$$c_N^*(T, \tilde{b}) := \epsilon_N^{l - \frac{1}{2} \sum_e (n_+(e) - n_-(e))}.$$

By the proof of Proposition 7.1, $\phi_N^{-q(T, \tilde{b})}$ and the unnormalized QH state sum have reciprocal variations. So it remains to analyze the behaviour of c_N^* with respect to oriented C -moves and the circuit move of Lemma 6.4.

It is relevant here to clarify the nature of c_N^* . By extending Chapter 7 of [7], the paper [9] provides a combinatorial realization of spin structures on 3-manifolds, based on weakly branched triangulations instead of branched ones. In particular, one associates to every weakly branched triangulation (T, \tilde{b}) a framing $\nu_{\tilde{b}}$ of the manifold defined along $\text{Sing}(P)$, and computes a cellular cochain $\alpha_{\tilde{b}} \in C^2(P; \mathbb{Z}/2\mathbb{Z})$ representing the *obstruction* to extend $\nu_{\tilde{b}}$ over the whole of the spine P . Consider the cellular 2-chain $R(P) = \sum_R R \in C_2(P; \mathbb{Z}/2\mathbb{Z})$, where R varies among the 2-regions of P . It is a fact that

$$l - \frac{1}{2} \sum_e (n_+(e) - n_-(e)) = \alpha_{\tilde{b}}(R(P)) \pmod{2}.$$

How $\alpha_{\tilde{b}}$, whence $\alpha_{\tilde{b}}(R(P))$, varies with the weak branching is part of the theory developed in [9]. As a consequence we get:

Lemma 8.7. *For any change of weak branching $\tilde{b} \rightarrow \tilde{b}'$ preserving the induced pre-branching we have*

$$\mathcal{H}_N(T, \tilde{b}, w, f, c) \equiv_N \mathcal{H}_N(T, \tilde{b}', w, f, c).$$

Proof. It is enough to show the result for one generating oriented C -move $\sigma = (0123)$. By local inspection one checks that $\alpha_{\tilde{b}'}(R(P)) = \alpha_{\tilde{b}}(R(P)) + 1$, so $c_N^*(T, \tilde{b}') = \epsilon_N c_N^*(T, \tilde{b})$. On the other hand, using Proposition 7.2, the factorization $(0123) = (23) \cdot (12) \cdot (01)$, and the relation $c_0 + c_1 + c_2 = 1$, we see that σ changes the unnormalized QH state sum by $\epsilon_N^{c_1} \epsilon_N^{c_2} \epsilon_N^{c_0} = \epsilon_N$. Hence the two sign variations are equal. \square

Lemma 8.8. *For any change of weak branching $\tilde{b} \rightarrow \tilde{b}'$ induced by one circuit move along a circuit γ we have*

$$\mathcal{H}_N(T, \tilde{b}, w, f, c) \equiv_N \epsilon_N^{h_c(\gamma)} \mathcal{H}_N(T, \tilde{b}', w, f, c)$$

where h_c is the bulk c -weight of \mathcal{P} . In particular, $\mathcal{H}_N(\mathcal{P})$ has no sign anomaly if $h_c = 0$.

Proof. Let us realize the circuit move in terms of \mathcal{N} -graphs. By the third identity of Proposition 7.2 the move changes the state sum by one factor $\epsilon_N^{c_0}$ for each crossing of γ . It is easy to check that their product computes $h_c(\gamma)$. On the other hand $\alpha_{\tilde{b}'}(R(P)) = \alpha_{\tilde{b}}(R(P))$, so that $c_N^*(T, \tilde{b}') = c_N^*(T, \tilde{b})$. \square

9. EXAMPLES

In this section we describe the quantum hyperbolic invariants of the figure-eight knot complement, its “sister”, and the complement of the knot 5_2 .

We give also some samples of numerical computations obtained by using Maple, which are part of our current analytical and numerical exploration of the QHI. At first we quote that they corroborate the invariance of the QH state sums with respect to the choice of flattenings or charges having the same weights. In the case of the sister manifold, they confirm also the invariance of the QH state sums supported by weakly branched triangulations. Moreover, the numerical computations that we present below are proving:

- The actual dependence of the invariants on the weights (and not only on the characters ρ).
- This dependence persists as $N \rightarrow \infty$.
- For some weights the invariants seem to grow exponentially with N , and yield instances of the volume conjecture, and for some other weights they do not.

Other interesting phenomena happen, that deserve to be understood. For instance, in the case of the complements of the knots 4_1 and 5_2 , it appears that the asymptotical behaviour of the absolute values of the invariants do not depend on the values on the meridian of the charge weights or the flattening weights.

9.1. Notations. Let (Δ, b, w, f, c) be a quantum 3-simplex. Put $d_k := f_k - *_j c_k$, $k \in \{0, 1, 2\}$. We call d_k an *edge color*. In order to simplify the formulas we denote the N -th roots of the cross-ratios w_k by bold letters \mathbf{w}_k (rather than w'_k , as we did in (14)), and call them *q-shape parameters*. So

$$\mathbf{w}_k := \exp\left(\frac{1}{N}(\log(w_k) + \pi i(N+1)d_k)\right).$$

For any cusped manifold M and any QH triangulation \mathcal{T} of M , we have the tetrahedral and edge relations (see (15), (22) and (25))

$$\sum_{k=0,1,2} (\log(w_k) + \pi i d_k) = - *_b \pi i, \quad \sum_{E \rightarrow e} *_E (\log(w(E)) + \pi i d(E)) = -2\pi i$$

and

$$\prod_{k=0,1,2} \mathbf{w}_k = e^{-*_b \frac{\pi i}{N}}, \quad \prod_{E \rightarrow e} \mathbf{w}(E)^{*_E} = e^{-\frac{2\pi i}{N}}$$

where, as usual, $d(E) = d_k$ and $\mathbf{w}(E) = \mathbf{w}_k$ if E is the edge E_k or the opposite edge of a quantum 3-simplex (Δ, b, w, f, c) of \mathcal{T} , and $*_E := *_b$. We denote by $G_N(T, \tilde{b})$ the variety thus defined by the q-shape parameters, and call it the *QH gluing variety* (NB: it coincides with the variety $G(T, \tilde{b}, c)_N$ of Section 3.2; to simplify notations, here we drop the reference to the charge c).

As usual, denote by V a compact 3-manifold with one torus boundary component such that M is diffeomorphic to the interior of V . We compute the invariant $\mathcal{H}_N(V, \rho, k_f, k_c, h_f, h_c)$ by means of the function $\mathcal{H}'_N(T, \tilde{b}, c)$ on $G_N(T, \tilde{b})$ introduced after Definition 3.5. As $\mathcal{H}'_N(T, \tilde{b}, c)$ is formulated in terms of q-shape parameters and charges, we express equivalently $\mathcal{H}_N(V, \rho, k_f, k_c, h_f, h_c)$ as $\mathcal{H}_N(V, \rho, \kappa, k_c, h, h_c)$, using the classes κ and h obtained from the flattening weights k_f and h_f by replacing the flattenings f_k by the edge colors d_k in the respective formulas (see Definition 4.10, and (26) for κ). Clearly $\kappa = k_f - \pi i k_c$ and $h = h_c + h_f$. We call κ and h the *reduced boundary and bulk weights*. By (1) and (28) one has the compatibility relations

$$(69) \quad \kappa(a) = d_w(a) \pmod{i\pi\mathbb{Z}}, \quad (\kappa(a) - d_w(a))/i\pi = i^*(h)(a) \pmod{2\mathbb{Z}}$$

for all $a \in H_1(\partial V; \mathbb{Z})$.

We consider triangulations with ordered tetrahedra. We denote by $u, v, w \dots$ (resp. $\mathbf{u}, \mathbf{v}, \mathbf{w} \dots$, resp. $a, b, c \dots$, resp. $a', b', c' \dots$) the triples of cross-ratios (resp. q-shape parameters, resp. edge colors, resp. charges) of the tetrahedra, in the same order.

9.2. The figure-eight knot complement. The Epstein-Penner decomposition of M is an ideal triangulation T having a branching b and two tetrahedra, with opposite b -orientations. We showed the dual spine in Figure 6. We denote by Δ^0 the tetrahedron with positive b -orientation. In [5] we described the flattenings at positive points of the gluing variety $G(T, b)$, as well as the charges and the QH state sums carried by (T, b) .

9.2.1. The QH gluing variety. The gluing variety $G(T, b)$ is the irreducible plane curve with coordinates (u_0, v_0) and defining equation $u_1 u_2^2 v_0^{-2} v_1^{-1} = 1$, which may be written as $u_0^2 v_0^2 = (1 - u_0)(1 - v_0)$ (as usual $u_{i+1} = 1/(1 - u_i)$ and similarly for the v_i 's). The point $(e^{\frac{\pi i}{3}}, e^{-\frac{\pi i}{3}})$ realizes the complete hyperbolic structure. Let l and m be the canonical longitude and the meridian of the figure-eight knot, and let $(u_0, v_0) \in G(T, b)$. The dilation factors of the holonomy $\rho := \rho(u_0, v_0)$ are given by

$$\text{hol}_l(\rho) = u_0^2 u_2^{-2} = \frac{u_0^4}{(1 - u_0)^2}, \quad \text{hol}_m(\rho) = u_2 v_2 = \frac{(u_0 - 1)(v_0 - 1)}{u_0 v_0}.$$

The tetrahedral and edge relations between q-shape parameters are

$$\begin{aligned} \mathbf{u}_0 \mathbf{u}_1 \mathbf{u}_2 &= e^{-\frac{\pi i}{N}}, \quad \mathbf{v}_0 \mathbf{v}_1 \mathbf{v}_2 = e^{+\frac{\pi i}{N}} \\ \mathbf{u}_1 \mathbf{u}_2^2 \mathbf{v}_0^{-2} \mathbf{v}_1^{-1} &= e^{-\frac{2\pi i}{N}}, \quad \mathbf{u}_1 \mathbf{u}_0^2 \mathbf{v}_2^{-2} \mathbf{v}_1^{-1} = e^{-\frac{2\pi i}{N}}. \end{aligned}$$

Together with the identities $(\mathbf{u}_{i+1})^N = 1/(1 - (\mathbf{u}_i)^N)$ and $(\mathbf{v}_{i+1})^N = 1/(1 - (\mathbf{v}_i)^N)$, these relations define $G_N(T, b)$. Using the tetrahedral relations it is easy to see that the two edge relations are equivalent. Recall the reduced weights h and κ introduced in Section 9.1. By (69) we have

$$(70) \quad \mathbf{u}_0^2 \mathbf{u}_2^{-2} = e^{\frac{\kappa(l)}{N} + \pi i h(l)}, \quad \mathbf{u}_2 \mathbf{v}_2 = e^{\frac{\kappa(m)}{N} + \pi i h(m)}.$$

As M is a knot complement, the bulk weight h is determined by κ , and similarly h_c is determined by k_c . Moreover, $h(l) = 0$, so h is non zero if and only if $h(m) = a_2 + b_2 = 1 \pmod{2}$.

Over the point $(e^{\frac{\pi i}{3}}, e^{-\frac{\pi i}{3}})$ we have $\kappa(m) \in \pi i \mathbb{Z}$ and $\kappa(l) \in 2\pi i \mathbb{Z}$, and the tetrahedral relations, the edge relations, and the compatibility relations with κ imply the following expressions of the edge colors a_k and b_k in terms of a_0 and κ : $a_2 = -2 - a_0 - a_1$, $b_2 = 2 - b_0 - b_1$, and

$$(71) \quad a_1 = -2a_0 + \frac{\kappa(l)}{2\pi i} - 2, \quad b_0 = \frac{\kappa(m)}{\pi i} - a_0, \quad b_1 = 2a_0 - \frac{2\kappa(m)}{\pi i} - \frac{\kappa(l)}{2\pi i} + 2.$$

For points $(\mathbf{u}, \mathbf{v}) \in G_N(T, b)$ lying over a sufficiently small neighborhood of the point $(e^{\frac{\pi i}{3}}, e^{-\frac{\pi i}{3}}) \in G(T, b)$, the same expressions are valid if one replaces $\kappa(\cdot)/\pi i$ with $(\kappa(\cdot) - \log(\text{hol}(\rho)))/\pi i$.

Similarly the charges a'_k and b'_k are given by

$$(72) \quad a'_1 = -2a'_0 + \frac{k_c(l)}{2} + 1, \quad b'_0 = a'_0 - k_c(m), \quad b'_1 = -2a'_0 + 2k_c(m) + \frac{k_c(l)}{2} + 1.$$

Let M' be a closed hyperbolic (p, q) -Dehn filling of M represented by $(u_0, v_0) \in G(T, b)$ in a small neighborhood of $(e^{\frac{\pi i}{3}}, e^{-\frac{\pi i}{3}})$. The holonomy ρ factors through the quotient map $\pi_1(M) \rightarrow \pi_1(M')$ to define the holonomy ρ' of M' . One has

$$(73) \quad p \log(\text{hol}_m(\rho')) + q \log(\text{hol}_l(\rho')) = 2\pi i,$$

and letting $r, s \in \mathbb{Z}$ be such that $ps - qr = 1$, the reduced weight κ satisfies $\kappa(m^p l^q) = 0$ if and only if

$$(74) \quad a_1 = r - 2 - 2a_0, \quad b_0 = -2s - a_0, \quad b_1 = 2 - r + 4s + 2a_0.$$

Similar formulas express the identity $k_c(m^p l^q) = 0$.

9.2.2. The state sum formulas. Recall the notations of Section 3.1.2. For any $(\mathbf{u}, \mathbf{v}) \in G_N(T, b)$ and any charge c on T , the QH state sum

$$(75) \quad \mathcal{H}'_N(T, b, c)(\mathbf{u}, \mathbf{v}) = (\mathbf{u}_0^{-a'_1} \mathbf{u}_1^{a'_0} \mathbf{v}_0^{-b'_1} \mathbf{v}_1^{b'_0})^{\frac{N-1}{2}} \frac{[\mathbf{v}_0] g(\mathbf{u}_0)}{g(\mathbf{v}_0)} \sum_{\alpha, \beta=0}^{N-1} \zeta^{\beta^2 - \alpha^2} \frac{\omega(\mathbf{u}_0, \mathbf{u}_1^{-1} | N - \beta)}{\omega(\mathbf{v}_0 / \zeta, \mathbf{v}_1^{-1} | N - \alpha)}$$

computes the invariant $\mathcal{H}_N(V, \rho, \kappa, k_c, h, h_c)$, and defines a (non-constant) rational function on $G_N(T, b)$. By a surgery theorem proved in [5], at points (\mathbf{u}, \mathbf{v}) where κ and k_c satisfy $\kappa(m^p l^q) = k_c(m^p l^q) = 0$ as above, $\mathcal{H}_N(V, \rho, \kappa, k_c, h, h_c)$ coincides with the N -th quantum hyperbolic invariant of the closed manifold (M', L, ρ') , where L is the surgery core.

9.2.3. Some numerical results. Denote by $\rho(A, B) := \rho(u_0(A, B), v_0(A, B))$ the holonomy of M given by the shape parameter $u_0(A, B) := e^{\frac{\pi i}{3}} + A + iB$ and the solution $v_0(A, B)$ of the gluing equation $u_0(A, B)^2 v_0(A, B)^2 - (1 - u_0(A, B))(1 - v_0(A, B)) = 0$ such that $u_0(A, B)$ has positive imaginary part and $v_0(A, B)$ has negative imaginary part. So $\rho(0, 0)$ is the hyperbolic holonomy.

Recall that h and h_c are determined by κ and k_c respectively. In the following table we present some values of $|\mathcal{H}_{15}(V, \rho(A, B), \kappa, k_c, h, h_c)|$ up to 4 digits, where (A, B) and $k_c(l)$ are as indicated on the top and the left of the corresponding column and row ($(A, B)_{(2,3)} := (0.06734378, \dots, -4.2400885\dots)$ defines the holonomy of the $(2, 3)$ -Dehn filling of M), and we put $\kappa(m) = k_c(m) = 0$ and $\kappa(l) = k_f(l) - \pi i k_c(l)$, where $k_f(l) = L([w; f])(l)$ has the collection of shape parameters $w := w(A, B)$ determined by $(u_0(A, B), v_0(A, B))$, and a (compatible) flattening f such that $L([w(0, 0); f]) = -2\pi i$ at the complete hyperbolic structure.

	(0, 0)	(1, 0)	(1, 0.5)	(1, 1)	$(A, B)_{(2,3)}$
$k_c(l) = 6$	2.5587...	2.6504...	2.0018...	1.6118...	5.0307...
$k_c(l) = 4$	58.5466...	58.3761...	47.0533...	39.9892...	95.0326...
$k_c(l) = 2$	2.1356...	2.0279...	1.9058...	1.8138...	2.7491...
$k_c(l) = 0$	77.4851...	77.5401...	77.5885...	77.6997...	76.4850...
$k_c(l) = -2$	0.1118...	0.1620...	0.1672...	0.1746...	0.0650...
$k_c(l) = -4$	77.4851...	77.5401...	95.8738...	112.7032...	47.4247...
$k_c(l) = -6$	2.1356...	2.0279...	2.7647...	3.5183...	1.0549...
$k_c(l) = -8$	58.5466...	58.3761...	89.7752...	124.5239...	21.8881...
$k_c(l) = -10$	2.5587...	2.6504...	4.2536...	6.1199...	0.7577...
$k_c(l) = -12$	33.2019...	33.1224...	63.6250...	104.4917...	7.4274...

Note that the table shows that the invariants depend on the character and the weights. Also, the first two columns show a symmetry about $k_c(l) = -2$ (that is, when $(A, B) = (0, 0)$, about $\kappa(l) = 0$), corresponding to a change of orientation of l . The dominant rows are for $k_c(l) = 0$ or -4 (that is, when $(A, B) = (0, 0)$, about $\kappa(l) = -2\pi i$ or $2\pi i$).

Now consider the behaviour of $|\mathcal{H}_N(V, \rho(A, B), \kappa, k_c, h, h_c)|$ as $N \rightarrow +\infty$. Put $G_N(\kappa(l)) := \pi \log (|\mathcal{H}_{N+2}(V, \rho(0, 0), \kappa, k_c, h, h_c)|/|\mathcal{H}_N(V, \rho(0, 0), \kappa, k_c, h, h_c)|)$, where the arguments are as above except that we fix $(A, B) = (0, 0)$ (the complete hyperbolic structure). Let $\mathcal{H}_N''(T, b, c)$ be $\mathcal{H}_N'(T, b, c)$ divided by $(\mathbf{u}_0^{-a_1'} \mathbf{u}_1^{a_0'} \mathbf{v}_0^{-b_1'} \mathbf{v}_1^{b_0'})^{\frac{N-1}{2}}$, the product of the matrix dilogarithm symmetrization factors. Note that $G_N := \pi \log (|\mathcal{H}_{N+2}''(T, b, c)(\mathbf{u}, \mathbf{v})|/|\mathcal{H}_N''(T, b, c)(\mathbf{u}, \mathbf{v})|)$ and $G_N(\kappa(l))$ are equivalent as $N \rightarrow +\infty$. The following table gives a sample of values of G_{151} , where $\kappa(l)$ takes the values indicated in the first row.

	$-6\pi i$	$-4\pi i$	$-2\pi i$	0	$2\pi i$
G_{151}	2.03069...	-0.49036...	2.02968...	0.48922...	2.02968...

Note that the dependence on the weights persists as $N \gg 1$. Further numerical computations show that $G_N(\pm 2\pi i)$ converges to $\text{Vol}(M) \approx 2.02988321\dots$ as $N \gg 1$.

9.3. The complement of the knot 5_2 . The ideal triangulation T of M with smallest complexity has three tetrahedra $\Delta^0, \Delta^1, \Delta^2$, and it has a branching b that gives each tetrahedron the negative branching orientation. This triangulation is the one provided by Snappea; we keep the same ordering of tetrahedra. The branching b is determined by the ordering it induces on the set of vertices of Δ^0 , which is obtained from that provided by Snappea by applying the permutation $(v_0, v_1, v_2, v_3) \mapsto (v_1, v_2, v_0, v_3)$.

9.3.1. *The QH gluing variety.* The edge relations of (T, b) are $u_0 u_2 v_0^2 w_1 w_2 = 1$, $u_1 u_2 v_2 v_1^2 w_0 w_1 = 1$, and $u_0 u_1 v_2 w_0 w_2 = 1$. By the tetrahedral relations they reduce to two independent relations, so the gluing variety of (T, b) is the irreducible curve in \mathbb{C}_*^3 with coordinates (u_0, v_0, w_0) and defining equations

$$u_1 v_0^{-2} w_0 = 1, \quad u_2 v_2^{-1} w_1 = 1.$$

The complete hyperbolic structure is realized by the point $(u_{hyp}, v_{hyp}, w_{hyp})$ given up to 8 digits by (the imaginary parts are negative since all tetrahedra have the negative branching orientation)

$$\begin{aligned} u_{hyp} &= 0.21507987.. - i 1.30714121... \\ v_{hyp} &= 0.33764102... - i 0.56227951... \\ w_{hyp} &= 0.33764102... - i 0.56227951... \end{aligned}$$

Since $G(T, b)$ is irreducible and $(u_{hyp}^{-1}, v_{hyp}^{-1}, w_{hyp}^{-1})$ is a positive point, it is a rich variety. Let l and m be the canonical longitude and the meridian of the knot, and let $(u_0, v_0, w_0) \in G(T, b)$. The dilation factors of the holonomy $\rho := \rho(u_0, v_0, w_0)$ are given by

$$\text{hol}_l(\rho) = u_1^2 u_2^{-3} v_2 v_0^{-2} w_1 = \frac{u_0^3 (v_0 - 1)}{(1 - u_0)^2 (u_0 - 1)^3 v_0^3 (1 - w_0)}$$

$$\text{hol}_m(\rho) = u_1 v_0^{-1} = \frac{1}{(1 - u_0) v_0}.$$

The tetrahedral and edges relations between q-shape parameters are (note that the exponents -1 on the left of the edge relations are due to the negative branching orientation of the tetrahedra)

$$\mathbf{u}_0 \mathbf{u}_1 \mathbf{u}_2 = e^{+\frac{\pi i}{N}}, \quad \mathbf{v}_0 \mathbf{v}_1 \mathbf{v}_2 = e^{+\frac{\pi i}{N}}, \quad \mathbf{w}_0 \mathbf{w}_1 \mathbf{w}_2 = e^{+\frac{\pi i}{N}}.$$

$$(\mathbf{u}_0 \mathbf{u}_2 \mathbf{v}_0^2 \mathbf{w}_1 \mathbf{w}_2)^{-1} = e^{-\frac{2\pi i}{N}}, \quad (\mathbf{u}_1 \mathbf{u}_2 \mathbf{v}_2 \mathbf{v}_1^2 \mathbf{w}_0 \mathbf{w}_1)^{-1} = e^{-\frac{2\pi i}{N}}, \quad (\mathbf{u}_0 \mathbf{u}_1 \mathbf{v}_2 \mathbf{w}_0 \mathbf{w}_2)^{-1} = e^{-\frac{2\pi i}{N}}.$$

Together with the identity $(\mathbf{u}_{i+1})^N = 1/(1 - (\mathbf{u}_i)^N)$ and the similar ones for the $(\mathbf{v}_i)^N$'s and the $(\mathbf{w}_i)^N$'s, these relations define $G_N(T, b)$. Using the tetrahedral relations it is easy to see that any of the edge relations is a consequence of the other two. By definition, the weights satisfy

$$\mathbf{u}_1^2 \mathbf{u}_2^{-3} \mathbf{v}_2 \mathbf{v}_0^{-2} \mathbf{w}_1 = e^{\frac{\kappa(l)}{N} + \pi i h(l)}, \quad \mathbf{u}_1 \mathbf{v}_0^{-1} = e^{\frac{\kappa(m)}{N} + \pi i h(m)}.$$

As M is a knot complement, the bulk weight h is determined by κ , and similarly h_c is determined by k_c . Moreover $h(l) = 0$, so h is non zero if and only if $h(m) = a_1 + b_0 = 1 \pmod{2}$.

At the complete hyperbolic structure we have $\kappa(m) \in \pi i \mathbb{Z}$ and $\kappa(l) \in 2\pi i \mathbb{Z}$, and the tetrahedral relations, the edge relations, and the compatibility relations with κ imply the following expressions of the edge colors a_k, b_k and c_k , in terms of a_0, a_1 and κ : $a_2 = 2 - a_0 - a_1$, $b_2 = 2 - b_0 - b_1$, $c_2 = 2 - c_0 - c_1$, and

$$b_0 = a_1 - \frac{\kappa(m)}{\pi i}, \quad b_1 = -4 + 2a_0 + a_1 + 2\frac{\kappa(m)}{\pi i} - \frac{\kappa(l)}{2\pi i}$$

$$c_0 = -4 + a_1 - 2\frac{\kappa(m)}{\pi i}, \quad c_1 = -a_0 - a_1 - \frac{\kappa(m)}{\pi i} + \frac{\kappa(l)}{2\pi i}.$$

If $(\mathbf{u}, \mathbf{v}, \mathbf{w}) \in G_N(T, b)$ lies over a point in a small neighborhood of $(u_{hyp}, v_{hyp}, w_{hyp}) \in G(T, b)$, then the same expressions are valid if one replaces $\kappa(\cdot)/\pi i$ with $(\kappa(\cdot) - \log(\text{hol}(\rho)))/\pi i$.

Similarly, the charges a'_k, b'_k and c'_k are given by

$$b'_0 = a'_1 - k_c(m), \quad b'_1 = -1 + 2a'_0 + a'_1 + 2k_c(m) - k_c(l)/2$$

$$c'_0 = a'_1 - 2k_c(m), \quad c'_1 = 1 - a'_0 - a'_1 - k_c(m) + k_c(l)/2.$$

9.3.2. *The state sum formulas.* For any $(\mathbf{u}, \mathbf{v}, \mathbf{w}) \in G(T, b)$ and any charge c on T we have (here we denote by $\mathcal{R}_N(\Delta, \mathbf{w})$ the tensor $\mathcal{R}_{N,*,b,c}(\mathbf{w}_0, \mathbf{w}_1)$)

$$\begin{aligned} \mathcal{H}'_N(T, b, c)(\mathbf{u}, \mathbf{v}, \mathbf{w}) &= \sum_{a, \dots, f=0}^{N-1} \mathcal{R}_N(\Delta^0, \mathbf{u})_{b,c}^{d,a} \mathcal{R}_N(\Delta^1, \mathbf{v})_{e,a}^{f,b} \mathcal{R}_N(\Delta^2, \mathbf{w})_{d,f}^{c,e} \\ &= (\mathbf{u}_0^{-a'_1} \mathbf{u}_1^{a'_0} \mathbf{v}_0^{-b'_1} \mathbf{v}_1^{b'_0} \mathbf{w}_0^{-c'_1} \mathbf{w}_1^{c'_0})^{\frac{N-1}{2}} \frac{[\mathbf{u}_0][\mathbf{v}_0][\mathbf{w}_0]g(1)^3}{g(\mathbf{u}_0)g(\mathbf{v}_0)g(\mathbf{w}_0)} \\ &\quad \times \sum_{b,c,d=0}^{N-1} \frac{\zeta^{(b+c-d)(c+d)}}{\omega(\mathbf{u}_0/\zeta, \mathbf{u}_1^{-1}|b-d)\omega(\mathbf{v}_0/\zeta, \mathbf{v}_1^{-1}|d)\omega(\mathbf{w}_0/\zeta, \mathbf{w}_1^{-1}|d-c)} \end{aligned}$$

The QH state sum $\mathcal{H}'_N(T, b, c)(\mathbf{u}, \mathbf{v}, \mathbf{w})$ computes the invariant $\mathcal{H}_N(V, \rho, \kappa, k_c, h, h_c)$ and defines a (non-constant) rational function on $G_N(T, b)$. As for the figure-eight knot complement, and using the same notations, at a point $(\mathbf{u}, \mathbf{v}, \mathbf{w})$ such that $\kappa(m^{plq}) = k_c(m^{plq}) = 0$, $\mathcal{H}_N(V, \rho, \kappa, k_c, h, h_c)$ coincides with the N -th quantum hyperbolic invariant of the triple (M', L, ρ') obtained by hyperbolic (p, q) -Dehn filling of M .

9.3.3. *Some numerical results.* Keeping the same notations as in Section 9.2.3, we present below some values of $|\mathcal{H}_{15}(V, \rho(A, B), \kappa, k_c, h, h_c)|$ up to 4 digits, and some values of G_{121} . For each of the shape parameter $u_0(A, B) := u_{hyp} + A + iB$ that we consider, with the exception of $u_0(0, 0)$, there are two distinct points $(u_0(A, B), v_0(A, B), w_0(A, B))_{\pm} \in G(T, b)$ such that both $u_0(A, B)$, $v_0(A, B)$ and $w_0(A, B)$ have negative imaginary parts. We take again $\kappa(m) = k_c(m) = 0$, but now $k_c(l) = 0$ and hence $\kappa(l) = k_f(l) = L([w; f])(l)$, where $w := w_{\pm}(A, B)$ is determined by $(u_0(A, B), v_0(A, B), w_0(A, B))_{\pm}$, and f is a (compatible) flattening such that, at the complete hyperbolic structure, $k_f^0 := L([w(0, 0); f])$ takes the value indicated on the left of each row.

	(0, 0)	(-0.5, 0.5) ₊	(0.5, 0.2) ₊	(0.7, 0.3) ₊	(0.7, 0.3) ₋
$-6\pi i$	428.2809...	437.3709...	474.4262...	447.6316...	488.3855...
$-4\pi i$	55.2674...	70.5945...	65.7471...	90.4874...	68.6357...
$-2\pi i$	470.6170...	487.7969...	518.2639...	450.2483...	532.0331...
0	71.9995...	74.9737...	76.7206...	96.6462...	77.9037...
$2\pi i$	470.6170...	494.8640...	515.6044...	416.9562...	527.6381...
$4\pi i$	55.2674...	45.3442...	62.4615...	109.4872...	65.4050...
$6\pi i$	428.2809...	456.9464...	466.5497...	356.3560...	476.1700...

	$-6\pi i$	$-4\pi i$	$-2\pi i$	0	$2\pi i$
G_{121}	2.85352...	0.20745...	2.83431...	0.15541...	2.83431...

Further numerical computations show that $G_N(\pm 2\pi i)$ converges as $N \gg 1$ to $\text{Vol}(M) \approx 2.8281220883\dots$

9.4. The sister of the figure-eight knot complement. ($M = m003$ in Snappea's census). This cusped manifold is obtained by $(5, 1)$ Dehn surgery on a component of the Whitehead link (see [39], [31]). The figure eight knot complement and its sister M have the smallest volume of any orientable, cusped hyperbolic 3-manifold. Hyperbolic Dehn surgery on M induced by $(5, 1)$ and $(5, 2)$ Dehn surgeries on the Whitehead link yields the Fomenko-Matveev-Weeks manifold, which has the smallest volume of any closed orientable hyperbolic 3-manifold (approximately 0.9427...) [21].

The Epstein-Penner decomposition of M is the ideal triangulation T with two tetrahedra considered in Example 2.12 and 4.11. We have already described a weak branching \tilde{b} on T where both tetrahedra have positive branching orientations, and computed the charges, the flattenings at positive points of the gluing variety $G(T, \tilde{b})$, and the QH state sums carried by (T, \tilde{b}) .

9.4.1. The QH gluing variety. The edge relations of (T, \tilde{b}) are $u_0 u_1^2 v_0 v_1^2 = 1$ and $u_0 u_2^2 v_0 v_2^2 = 1$. By the tetrahedral relations they are equivalent, so the gluing variety $G(T, \tilde{b})$ is the irreducible plane curve with coordinates (u_0, v_0) and defining equation $u_0 u_1^2 v_0 v_1^2 = 1$. The point $(e^{\frac{\pi i}{3}}, e^{\frac{\pi i}{3}})$ realizes the complete hyperbolic structure (thus Δ^0 and Δ^1 are regular ideal tetrahedra, as in the case of the figure-eight knot complement). There is a basis (l, m) of $\pi_1(\partial \tilde{M})$ where the dilation factors of the holonomy $\rho := \rho(u_0, v_0)$ are given by

$$(76) \quad \text{hol}_l(\rho) = u_0 u_1 v_0^{-1} v_1^{-1} = \frac{u_0(1-v_0)}{(1-u_0)v_0}, \quad \text{hol}_m(\rho) = u_1^{-2} v_2^2 = \frac{(1-u_0)^2(v_0-1)^2}{v_0^2}.$$

The tetrahedral and edges relations between q-shape parameters are

$$\begin{aligned} \mathbf{u}_0 \mathbf{u}_1 \mathbf{u}_2 &= e^{-\frac{\pi i}{N}}, \quad \mathbf{v}_0 \mathbf{v}_1 \mathbf{v}_2 = e^{-\frac{\pi i}{N}} \\ \mathbf{u}_0 \mathbf{u}_1^2 \mathbf{v}_0 \mathbf{v}_1^2 &= e^{-\frac{2\pi i}{N}}, \quad \mathbf{u}_0 \mathbf{u}_2^2 \mathbf{v}_0 \mathbf{v}_2^2 = e^{-\frac{2\pi i}{N}}. \end{aligned}$$

Together with the identity $(\mathbf{u}_{i+1})^N = 1/(1 - (\mathbf{u}_i)^N)$ and the similar one for the $(\mathbf{v}_i)^N$'s, these relations define $G_N(T, \tilde{b})$. Using the tetrahedral relations it is easy to see that the two edge relations are equivalent. The reduced weights satisfy the compatibility relations

$$\mathbf{u}_0 \mathbf{u}_1 \mathbf{v}_0^{-1} \mathbf{v}_1^{-1} = e^{\frac{k(l)}{N} + \pi i h(l)}, \quad \mathbf{u}_1^{-2} \mathbf{v}_2^2 = e^{\frac{k(m)}{N} + \pi i h(m)}.$$

Here we note that $H_1(M; \mathbb{Z}) \cong \mathbb{Z} \oplus \mathbb{Z}/5\mathbb{Z}$, and that the curve m or l generates $H_1(M; \mathbb{Z}/2\mathbb{Z}) \cong \mathbb{Z}/2\mathbb{Z}$. So the bulk weight h is determined by κ , and similarly h_c is determined by k_c . Moreover $h(m) = 0 \pmod{2}$, so h is non zero if and only if $h(l) = a_0 + a_1 + b_0 + b_1 = a_2 + b_2 = 1 \pmod{2}$.

At the complete hyperbolic structure we see that $\kappa(l) \in \pi i \mathbb{Z}$ and $\kappa(m) \in 2\pi i \mathbb{Z}$, and the tetrahedral relations, the edge relations, and the compatibility relations with κ imply the following expressions of the edge colors a_k and b_k in terms of a_1 and κ :

$$(77) \quad \begin{aligned} a_0 &= \frac{\kappa(l)}{\pi i} - \frac{\kappa(m)}{2\pi i} - 2a_1 - 2, \quad a_2 = a_1 - \frac{\kappa(l)}{\pi i} + \frac{\kappa(m)}{2\pi i} \\ b_0 &= \frac{\kappa(l)}{\pi i} - \frac{3}{2} \frac{\kappa(m)}{\pi i} - 2a_1 - 2, \quad b_1 = -\frac{\kappa(l)}{\pi i} + \frac{\kappa(m)}{\pi i} + a_1, \quad b_2 = \frac{\kappa(m)}{2\pi i} + a_1. \end{aligned}$$

For points $(\mathbf{u}, \mathbf{v}) \in G_N(T, \tilde{b})$ lying over a small neighborhood of $(e^{\frac{\pi i}{3}}, e^{\frac{\pi i}{3}}) \in G(T, \tilde{b})$, the same expressions are valid if one replaces $\kappa(\cdot)/\pi i$ with $(\kappa(\cdot) - \log(\text{hol}(\rho)))/\pi i$.

9.4.2. The state sum formulas. For any $(\mathbf{u}, \mathbf{v}) \in G_N(T, \tilde{b})$ and any charge c on T we have (again we denote by $\mathcal{R}_N(\Delta, \mathbf{w})$ the tensor $\mathcal{R}_{N,*,b,c}(\mathbf{w}_0, \mathbf{w}_1)$)

$$\begin{aligned} \mathcal{H}'_N(T, \tilde{b}, c)(\mathbf{u}, \mathbf{v}) &= \sum_{i,j,k,l,I,J=0}^{N-1} \mathcal{R}_N(\Delta^0, \mathbf{u})_{k,l}^{i,j} \mathcal{R}_N(\Delta^1, \mathbf{v})_{j,i}^{I,J} (\mathcal{Q}^2)_I^l \mathcal{Q}_J^k \\ &= (\mathbf{u}_0^{-a_1} \mathbf{u}_1^{a'_0} \mathbf{v}_0^{-b_1} \mathbf{v}_1^{b'_0})^{\frac{N-1}{2}} \frac{g(\mathbf{u}_0)g(\mathbf{v}_0)}{N\phi_N g(1)^2} \\ &\quad \times \sum_{i,j,k,I=0}^{N-1} \zeta^{(i+k)(j+I)-ik+\frac{i^2+I^2}{2}} \omega(\mathbf{u}_0, \mathbf{u}_1^{-1}|i-k)\omega(\mathbf{v}_0, \mathbf{v}_1^{-1}|I-j) \end{aligned}$$

The QH state sum $\mathcal{H}'_N(T, \tilde{b}, c)(\mathbf{u}, \mathbf{v})$ computes the invariant $\mathcal{H}_N(V, \rho, \kappa, k_c, h, h_c)$ and defines a (non-constant) rational function on $G_N(T, \tilde{b})$. As for the previous examples, at a point (\mathbf{u}, \mathbf{v}) such that $\kappa(m^{pl^q}) = k_c(m^{pl^q}) = 0$, $\mathcal{H}_N(V, \rho, \kappa, k_c, h, h_c)$ coincides with the N -th quantum hyperbolic invariant of the triple (M', L, ρ') obtained by hyperbolic (p, q) -Dehn filling of M .

9.4.3. *Some numerical results.* We keep the same notations as in Section 9.2.3 and we take again $\kappa(m) = k_c(m) = 0$, but now both $u_0(A, B)$ and $v_0(A, B)$ have positive imaginary part, and the value of $\kappa(l)$ is given by $k_c(l)$ (on the left of each row), $w(A, B)$, and a flattening f such that $L([w(0, 0); f]) = 0$. The following table presents values of $|\mathcal{H}_{15}(V, \rho(A, B), \kappa, k_c, h, h_c)|$ up to 4 digits.

	(0, 0)	(1, 0)	(-1, 0)	(0, 1)	(1, 1)
2	4.7755...	5.4995...	4.5346...	3.1943...	2.6295...
1	173.2621...	182.5736...	173.3850...	139.8353...	126.3516...
0	0.2500...	0.6231...	0.3624...	0.2367...	0.1498...
-1	173.2621...	114.3211...	173.3850...	214.7305...	236.1129...
-2	4.7755...	6.8535...	4.5346...	7.3011...	9.3949...

REFERENCES

- [1] H. Bai, *Quantum Teichmüller spaces and Kashaev's 6j-symbols*, Alg. Geom. Topol. 7 (2007) 1541–1560
- [2] H. Bai, F. Bonahon, X. Liu, *Local representations of the quantum Teichmüller space*, arXiv:math.GT/0707.2151
- [3] S. Baseilhac, R. Benedetti, *Quantum hyperbolic invariants of 3-manifolds with $PSL(2, \mathbb{C})$ -characters*, Topology 43 (6) (2004) 1373–1423
- [4] S. Baseilhac, R. Benedetti, *Classical and quantum dilogarithmic invariants of flat $PSL(2, \mathbb{C})$ -bundles over 3-manifolds*, Geom. Topol. 9 (2005) 493–570
- [5] S. Baseilhac, R. Benedetti, *Quantum hyperbolic geometry*, Alg. Geom. Topol. 7 (2007) 845–917
- [6] S. Baseilhac, R. Benedetti, *The Kashaev and quantum hyperbolic invariants of links*, J. Gökova Geom. Topol. 5 (2011) 31–85
- [7] R. Benedetti, C. Petronio, *Branched Standard Spines of 3-manifolds*, Lecture Notes in Mathematics 1653, Springer (1997)
- [8] R. Benedetti, C. Petronio, *Lectures on Hyperbolic Geometry*, Universitext, Springer Verlag (1992)
- [9] R. Benedetti, C. Petronio, *Spin structures on 3-manifolds via arbitrary triangulations*, to appear in Algebraic and Geometric Topology
- [10] D.W. Boyd, F. Rodriguez-Villegas, *Mahler's measure and the dilogarithm (II)*, with an appendix by N.M. Dunfield, arXiv:math.NT/0308041
- [11] S. Boyer, X. Zhang, *On Culler–Shalen seminorms and Dehn filling*, Ann. of Math. (2) 148 (3) (1998) 737–801
- [12] D. Bullock, *Rings of $SL_2(\mathbb{C})$ -characters and the Kauffman bracket skein module*, Comment. Math. Helv. 72 (4) (1997) 521–542
- [13] F. Costantino, *A calculus for branched spines of 3-manifolds*, Math. Zeitschrift 251 (2) (2005) 427–442
- [14] D. Cooper, M. Culler, H. Gillet, D.D. Long, P.B. Shalen, *Plane curves associated to character varieties of 3-manifolds*, Invent. Math. 118 (1994) 47–84
- [15] Y.-E. Choi, *Positively oriented ideal triangulations on hyperbolic 3-manifolds*, Topology 43 (2004) 1345–1371
- [16] A. Champanerkar, *A-polynomial and Bloch invariants of hyperbolic 3-manifolds*, PhD Thesis, Columbia University (2003)
- [17] M. Culler, P.B. Shalen, *Varieties of group representations and splittings of 3-manifolds*, Ann. of Math (2) 117 (1) (1983) 109–146
- [18] N. Dunfield, *Cyclic surgery, degree of maps of character curves, and volume rigidity for hyperbolic manifolds*, Invent. Math. 136 (1999) 623–657
- [19] S. Freed, *Classical Chern-Simons Theory, 1*, Adv. in Math. 113 (1995) 237–303
- [20] S. Francaviglia, *Hyperbolic volume of representations of fundamental groups of cusped 3-manifolds*, Internat. Math. Res. Not. 9 (2004), 425–459
- [21] D. Gabai, R. Meyerhoff, P. Milley, *Minimum volume cusped hyperbolic three-manifolds*, Journal Amer. Math. Soc. 22 (4) (2009) 1157–1215

- [22] A. Hatcher, *Algebraic Topology*, Cambridge Univ. Press (2002)
- [23] M. Kapovich, *Hyperbolic Manifolds and Discrete Groups*, Birkhäuser, Prog. Math. 183 (2010)
- [24] R.M. Kashaev, *Quantum dilogarithm as a 6j-symbol*, Mod. Phys. Lett. A Vol. 9 (40) (1994) 3757–3768
- [25] R.M. Kashaev, *A link invariant from quantum dilogarithm*, Mod. Phys. Lett. A Vol. 10 (40) (1995) 1409–1418
- [26] R. M. Kashaev, *The hyperbolic volume of knots from the quantum dilogarithm*, Lett. Math. Phys. 39 (1997) 269–275
- [27] P. Kirk, E. Klassen, *Chern-Simons invariants of 3-manifolds decomposed along tori and the circle bundle over the representation space of T^2* , Comm. Math. Phys. 153 (3) (1993) 521–557
- [28] S. Lang, *Algebraic Number Theory*, Springer (1994)
- [29] F. Luo, S. Schleimer, S. Tillmann *Geodesic ideal triangulations exist virtually*, Proc. Amer. Math. Soc. 136 (2008) 2625–2630
- [30] F. Luo, S. Tillmann, T. Yang *Thurston’s spinning construction and solutions to the hyperbolic gluing equations for closed hyperbolic 3-manifolds*, Proc. Amer. Math. Soc. 141 (1) (2013) 335–350.
- [31] S.V. Matveev, A.T. Fomenko, *Constant energy surfaces of Hamiltonian systems, enumeration of three-dimensional manifolds in increasing order of complexity, and computation of volumes of closed hyperbolic manifolds*, Russian Math. Surveys 43:1 (1988) 3–24
- [32] H. Murakami, J. Murakami, *The colored Jones Polynomials and the simplicial volume of a knot*, Acta Math. 186 (2001), 85–104
- [33] W. D. Neumann, *Combinatorics of triangulations and the Chern-Simons invariant for hyperbolic 3-manifolds*, Topology’90 (Columbus, OH, 1990), De Gruyter, Berlin (1992) 243–271
- [34] W.D. Neumann, *Extended Bloch group and the Cheeger-Chern-Simons class*, Geom. Topol. 8 (2004) 413–474
- [35] W.D. Neumann, D. Zagier, *Volumes of hyperbolic 3-manifolds*, Topology 24, 307–332 (1985)
- [36] C. Petronio, J. Porti, *Negatively oriented ideal triangulations and a proof of Thurston’s hyperbolic Dehn filling theorem*, Exp. Math. 18 (2000) 1–35
- [37] P.B. Shalen, *Representations of 3-manifold groups*, Handbook of geometric topology, North-Holland, Amsterdam (2002) 955–1044
- [38] W.P. Thurston, *The geometry and topology of 3-manifolds*, Princeton Univ. Press (1979)
- [39] J. Weeks, *Hyperbolic structures on 3-manifolds*, Ph.D. thesis, Princeton Univ. (1985)
- [40] T. Yoshida, *The η -invariant of hyperbolic 3-manifolds*, Invent. Math 81 (1985) 473–514