

VEGETATION LANDSCAPE MANAGEMENT OF "NATURAL RESERVE OF CHIARONE", MASSACIUCCOLI LAKE BASIN (TUSCANY, IT)

ANDREA BERTACCHI¹, TIZIANA LOMBARDI¹, IACOPO TOMEI¹, MARCELLO LABATE², ANDREA FONTANELLI², LUCIA PICCHI²

¹Department of Agriculture, Food and Environment (DAFE), University of Pisa, Via del Borghetto 80, 56124 Pisa, Italy;

²<http://www.oaslipummassaciuccoli.org> - Via del Porto 6, loc. Massaciuccoli 55050 Massarosa (LU)

The Massaciuccoli Lake basin is located in NW of Tuscany (Italy) (43°50'N 10°19'E) and it is integral part of the Regional Park of San Rossore, Migliarino and Massaciuccoli (SIR-pSIC-ZPS n° 25 "Lago e Padule di Massaciuccoli" (IT5120017). This endorheic lake and the surrounding marshlands, with over 2,000 hectares of surface, form the largest retrodunal wetland of Tuscany. The lake covers about 690 hectares. The origin is probably related to a lagoon lateral of the mouth of the protohistoric Auser river, transformed into a closed basin and isolated from the sea by the advance towards the interior of the dune systems. Currently the lake presents a depth that, in the areas not affected by the quarries, is between 1 and 2.5 m. The average level of the surface of the lake remains for long periods of the year below the marine one. Over the last century this ecological system has undergone profound changes both because of land drainage, both because of the industrial and agricultural development of the surrounding areas that have heavily polluted lake, whose waters are still affected by serious eutrophication and ecological degradation.

Since the '50s and '60s the human settlement in the territory of the Massaciuccoli basin has grown in all its forms, since the urban impact due to tourism, particularly in the municipalities of Viareggio and Massarosa and the intense work of mechanical reclamation for the acquisition of farmland, they have a negative impact on the ecosystem, causing a gradual deterioration of water quality. The presence of a large amount of nutrients (N, P, C), show that the lake has undergone a major change going from the clear waters and complex trophic nets are characterized by the presence of submerged macrophytes and zooplankton, to a situation of algal blooms and simpler trophic nets, resulting in the loss of biodiversity.

The characteristic climate of the area is Mediterranean humid and is characterized by average winter temperatures of 7 °C and summer average of 22 °C.

The rainfall (800 mm per year) is greater in autumn - winter, coinciding which the waters of the lake reached the highest levels

In order to preserve the great natural value of the lake in 1979, with the birth of the regional protected area, was established in the marshy area around the village of Massaciuccoli, the "Natural Reserve of Chiarone" (47 ha of mainland and 53 of lake). Since 1985 the association LIPU (Italian League for Bird Protection) has obtained the management of the reserve, and its activity is directed to purposes of environmental education, environmental restoration, monitoring and conservation of specific habitats.




Fig. 1 Geographic location of study area.

This report shows the results of several years of investigation on the main vegetational types (Tab. 1-9) (Phytosociological surveys of vegetation – using Braun-Blanquet method – and floristic detections are carried out in 2014-2015), the presence and distribution of endemic/rare/protected species, the changes on the structure of floating islands of peats and the main actions of conservation and management of these wetlands.

Tab. 1 Vegetation of *Phragmitetum australis* (Gams 1927) Shmala 1939 (Fig. 2 A) (*Frangula alnus* facies (rel 13); *Periploca graeca* facies (rel 14))

Rel. no.	2	6	8	10	12	13	14
Surface (m ²)	25	25	25	25	25	25	25
Coverage (%)	100	100	100	100	100	100	100
n° species	10	7	10	12	11	9	7

charact. sp. of *Phragmitetum australis* Phragmites australis (Cav.) Trin. ex Steud.

Sp. *Phragmitetum-Phragmitetalia*

Calystegia sepium L.

Solidago canadensis L.

Eupatorium cannabinum L.

Euphorbia palustris L.

Typha europaea L.

Stachys palustris L.

Lytrum salicaria L.

Hibiscus palustris L.

Typha angustifolia L.

Scrophularia laciniata Pall.

Mentha aquatica L.

Osmunda regalis L.

Iris pseudacorus L.

Osmunda cinnamomea L.

Periploca graeca L.

sp. *Alnion-Alno-Quercetalia*

Frangula alnus Mill.

Salicetum cinereae

Salix cinerea L.

*Tab. 2 Vegetation of *Cladietum marisci* (Allorge 1922) (Fig. 2 B)*

Zobrist 1935

Rel. no.

Surface (m²)

Coverage (%)

n° species

charact. sp. of *Cladietum marisci* Cladietum marisci (L.) Pohl

Sp. *Phragmitetum-Phragmitetalia*

Phragmites australis (Cav.) Trin. ex Steud.

Calystegia sepium L.

Solidago canadensis L.

Hibiscus palustris L.

Thelypteris palustris Schott.

Lytrum salicaria (L.)

Typha angustifolia L.

Scrophularia laciniata Pall.

Mentha aquatica L.

Osmunda regalis L.

Iris pseudacorus L.

Osmunda cinnamomea L.

Periploca graeca L.

Salix cinerea L.

*Tab. 3 Vegetation of *Cladietum marisci* (Allorge 1922) Zobrist 1935 a *Phragmitetum australis**

Jekel(Cav.) Trin. ex Steud. (Fig. 2 C)

Surface (m²)

Coverage (%)

n° species

charact. sp. of *Cladietum marisci* Cladietum marisci (L.) Pohl

Sp. *Phragmitetum-Phragmitetalia*

Phragmites australis (Cav.) Trin. ex Steud.

Calystegia sepium L.

Solidago canadensis L.

Hibiscus palustris L.

Thelypteris palustris Schott.

Lytrum salicaria (L.)

Typha angustifolia L.

Scrophularia laciniata Pall.

Mentha aquatica L.

Osmunda regalis L.

Iris pseudacorus L.

Osmunda cinnamomea L.

Periploca graeca L.

Salix cinerea L.

*Tab. 4 Vegetation of *Typhetum angustifoliae* (Allorge 1921) Pign. 1953 (Fig. 2 D)*

Rel. no.

Surface (m²)

Coverage (%)

n° species

charact. sp. of *Typhetum angustifoliae* Typha angustifolia L.

Calystegia sepium L.

Phragmites australis (Cav.) Trin. ex Steud.

Scrophularia laciniata Pall.

Lytrum salicaria (L.)

Holoschoenus vulgaris Link

Schoenoplectus lacustris Pall.

Hydrocoleus vulgaris Link

Eupatorium cannabinum L.

Euphorbia palustris L.

Typha latifolia L.

Iris pseudacorus L.

Gallium palustre L.

Lysimachia vulgaris L.

Solidago canadensis L.

*Tab. 5 Vegetation of *Eleocharitetum palustris* Schenn. 1919 (Fig. 2 E)*

Rel. no.

Surface (m²)

Coverage (%)

n° species

charact. sp. of *Eleocharitetum palustris* Eleocharis palustris (L.)

Samolus valerandi (L.)

Holoschoenus vulgaris Link

Hydrocoleus vulgaris (L.)

Gallium palustre L.

Lysimachia vulgaris L.

Solidago canadensis L.

*Tab. 7 Vegetation of *Nymphaeum albae* Vilm. 1947 (Fig. 2 G) and *Myriophyllum-Nupharum* Koch 1926 *Najas marina* faces (rel 37)*

Rel. no.

Surface (m²)

Coverage (%)

n° species

charact. sp. of *Nymphaeum albae* Nymphaea alba L.

Nuphar lutea L.

Myriophyllum verticillatum L.

Najas marina L.

*Tab. 9 Vegetation of *Sphagnum palustris* mown meadows (Fig. 2 M)*

Rel. no.

Surface (m²)

Coverage (%)

n° species

Rel. no.

Surface (m²)

Coverage (%)

n° species

Rel. no.

Surface (m²)

<p