



Syddansk Universitet

On the uniqueness of the injective III1 factor

Haagerup, Uffe

Published in:
Documenta Mathematica

Publication date:
2016

Document version
Final published version

Document license
CC BY

Citation for pulished version (APA):
Haagerup, U. (2016). On the uniqueness of the injective III1 factor. Documenta Mathematica, 21(2016), 1193-1226.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

ON THE UNIQUENESS OF THE INJECTIVE III_1 FACTOR

UFFE HAAGERUP

Received: August 2, 2016

Communicated by Joachim Cuntz

ABSTRACT. We give a new proof of a theorem due to Alain Connes, that an injective factor N of type III_1 with separable predual and with trivial bicentralizer is isomorphic to the Araki–Woods type III_1 factor R_∞ . This, combined with the author’s solution to the bicentralizer problem for injective III_1 factors provides a new proof of the theorem that up to $*$ -isomorphism, there exists a unique injective factor of type III_1 on a separable Hilbert space.

2010 Mathematics Subject Classification: 46L36

Preamble by Alain Connes

Uffe Haagerup solved the hardest problem of the classification of factors, namely the uniqueness problem for injective factors of type III_1 . The present paper, taken from his unpublished notes, presents a direct proof of this uniqueness by showing that any injective factor of type III_1 is an infinite tensor product of type I factors so that the uniqueness follows from the Araki–Woods classification. The proof is typical of Uffe’s genius, the attack is direct, and combines his amazing control of completely positive maps and his sheer analytical power, together with his solution to the bicentralizer problem. After his tragic death, Hiroshi Ando volunteered to type the manuscript¹. Some pages were missing from the notes, but eventually Cyril Houdayer and Reiji Tomatsu suggested a missing proof of Lemma 3.4 and Theorem 3.1. We heartily thank Hiroshi, Cyril and Reiji for making the manuscript available to the community. We also thank Søren Haagerup for giving permission to publish his father’s paper.

¹The manuscript is typed by Hiroshi Ando (Chiba University) in cooperation with Cyril Houdayer (Université Paris-Sud), Toshihiko Masuda (Kyushu University), Reiji Tomatsu (Hokkaido University), Yoshimichi Ueda (Kyushu University) and Wojciech Szymanski (University of Southern Denmark).

CONTENTS

1	INTRODUCTION	1194
2	PRELIMINARIES	1196
2.1	Notation	1196
2.2	Connes–Woods’ characterization of ITPFI factors	1197
2.3	Bicentralizers on type III ₁ factors	1198
2.4	Almost unitary equivalence in Hilbert N -bimodules	1199
3	COMPLETELY POSITIVE MAPS FROM $m \times m$ -MATRICES INTO AN INJECTIVE FACTOR OF TYPE III ₁	1200
4	\mathbb{Q} -STABLE STATES ON III ₁ FACTORS	1208
5	PROOF OF MAIN THEOREM	1213

1 INTRODUCTION

The problem, whether all injective factors of type III₁ on a separable Hilbert space are isomorphic, has been settled affirmatively. The proof of the uniqueness of injective III₁ factors falls in two parts, namely (see §2.3 for the definition of the bicentralizer):

THEOREM 1.1 ([Con85]). *Let M be an injective factor of type III₁ on a separable Hilbert space, such that the bicentralizer B_φ is trivial (i.e., $B_\varphi = \mathbb{C}1$) for some normal faithful state φ on M , then M is $*$ -isomorphic to the Araki–Woods factor R_∞ .*

THEOREM 1.2 ([Haa87]). *For any normal faithful state φ on an injective factor M of type III₁ on a separable Hilbert space, one has $B_\varphi = \mathbb{C}1$.*

In this paper we give an alternative proof of Theorem 1.1 above, based on the technique of our simplified proof [Haa85] of Connes’ Theorem [Con76] “injective \Rightarrow hyperfinite” in the type II₁ case². The key steps in our proof of Theorem 1.1 are listed below:

STEP 1

By use of continuous crossed products, we prove that the identity map on an injective factor N of type III₁ has an approximate factorization

$$\begin{array}{ccc}
 & R & \\
 S_\lambda \nearrow & & \searrow T_\lambda \\
 N & \xrightarrow{\text{id}_N} & N
 \end{array}$$

²Typewriter’s note: Haagerup used this technique to give a new proof of the uniqueness of injective type III _{λ} ($0 < \lambda < 1$) factor. This result has been published in [Haa89].

through the hyperfinite factor R of type II₁, such that $(S_\lambda)_{\lambda \in \Lambda}$ and $(T_\lambda)_{\lambda \in \Lambda}$ are nets of normal unital completely positive maps, and for a fixed normal faithful state φ on N (chosen prior to S_λ and T_λ), there exist normal faithful states $(\psi_\lambda)_{\lambda \in \Lambda}$ on R , such that for all $t \in \mathbb{R}$ and $\lambda \in \Lambda$,

$$\begin{aligned}\varphi \circ T_\lambda &= \psi_\lambda, & \psi_\lambda \circ S_\lambda &= \varphi, \\ \sigma_t^\varphi \circ T_\lambda &= T_\lambda \circ \sigma_t^{\psi_\lambda}, \\ \sigma_t^{\psi_\lambda} \circ S_\lambda &= S_\lambda \circ \sigma_t^\varphi,\end{aligned}$$

and $\|S_\lambda \circ T_\lambda(x) - x\|_\varphi \xrightarrow{\lambda \rightarrow \infty} 0$ for all $x \in N$, where $\|y\|_\varphi := \varphi(y^*y)^{\frac{1}{2}}$ ($y \in N$).

STEP 2

From Step 1, we deduce that a certain normal faithful state φ (\mathbb{Q} -stable state defined in §4) on an injective factor N of type III₁ has the following property: for any finite set of unitaries u_1, \dots, u_n in N and for every $\gamma, \delta > 0$, there exists a finite-dimensional subfactor F of N such that

$$\varphi = \varphi|_F \otimes \varphi|_{F^c},$$

and such that there exist unitaries v_1, \dots, v_n in F and a unital completely positive map $T: F \rightarrow N$ such that

$$\begin{aligned}\varphi \circ T &= \varphi, \\ \|\sigma_t^\varphi \circ T - T \circ \sigma_t^{\varphi|_F}\| &\leq \gamma|t|, \quad t \in \mathbb{R},\end{aligned}$$

and

$$\|T(v_k) - u_k\|_\varphi < \delta, \quad k = 1, \dots, n.$$

STEP 3

We prove that if $N, \varphi, F, u_1, \dots, u_n, v_1, \dots, v_n$ are as in Step 2, then for every σ -strong neighborhood \mathcal{V} of 0 in N , there exists a finite set of operators a_1, \dots, a_p in N such that

- (a) $\sum_{i=1}^p a_i^* a_i \in 1 + \mathcal{V}$ and $\sum_{i=1}^p a_i^* a_i \leq 1$,
- (b) $\varepsilon_{F, \varphi} \left(\sum_{i=1}^p a_i a_i^* \right) \in 1 + \mathcal{V}$ and $\varepsilon_{F, \varphi} \left(\sum_{i=1}^p a_i a_i^* \right) \leq 1$,
- (c) $\sum_{i=1}^p \|a_i \xi_\varphi - \xi_\varphi a_i\|^2 < \delta'$,
- (d) $\sum_{i=1}^p \|a_i u_k - v_k a_i\|_\varphi^2 < \delta', \quad k = 1, \dots, n$.

Here ξ_φ denotes the unique representing vector of φ in a natural cone. The above $\delta' > 0$ depends on γ and δ in Step 2, and δ' is small when γ and δ are small. Here, $\varepsilon_{F,\varphi}$ is the φ -invariant conditional expectation of N onto F . Moreover in (c), the standard Hilbert space H of N is regarded as a Hilbert N -bimodule, by putting $\eta a := Ja^*J\eta$ ($a \in N$, $\eta \in H$).

Assume now that the bicentralizer of any normal faithful state on N is trivial. Then by an averaging argument, we can exchange (b) by

$$(b') \quad \sum_{i=1}^p a_i a_i^* \in 1 + \mathcal{V} \quad \text{and} \quad \sum_{i=1}^p a_i a_i^* \leq 1.$$

STEP 4

From (a), (b'), (c) and (d) above, we derive that there exists a unitary operator $w \in N$ such that

$$\|w\xi_\varphi - \xi_\varphi w\| < \varepsilon$$

and

$$\|wu_k - v_k w\|_\varphi < \varepsilon, \quad k = 1, \dots, n,$$

where ε is small when δ' is small and \mathcal{V} is a small σ -strong neighborhood of 0 in N . The key part of Step 4 is a theorem about general Hilbert N -bimodules, which was proved in [Haa89].

STEP 5

From Step 4, we get that for every finite set of unitaries $u_1, \dots, u_n \in N$ and every $\varepsilon > 0$, there exists a finite dimensional subfactor F_1 (namely w^*Fw) of N and n unitaries v'_1, \dots, v'_n in F_1 (namely $w^*v_k w$, $k = 1, \dots, n$), such that

$$(i) \quad \|v'_k - v_k\|_\varphi < \varepsilon$$

and

$$(ii) \quad \|\varphi - \varphi|_{F_1} \otimes \varphi|_{F_1^c}\| < \varepsilon.$$

The last inequality follows from the fact, that when w almost commutes with ξ_φ , it almost commutes with φ too. The properties (i) and (ii) above show that φ satisfies the product condition of Connes–Woods [CW85] and thus N is an ITPFI factor. But it is well-known that R_∞ is the only ITPFI factor of type III₁ (cf. [AW68] and [Con73]).

2 PRELIMINARIES

2.1 NOTATION

We use M, N, \dots to denote von Neumann algebras and ξ, η, \dots to denote vectors in a Hilbert space. Let M be a von Neumann algebra. $\mathcal{U}(M)$ denotes the

unitary group of M . For a faithful normal state φ on M , we denote by Δ_φ (resp. J_φ) the modular operator (resp. modular conjugation operator) associated with φ , and the modular automorphism group of φ is denoted by σ^φ . The norm $\|x\|_\varphi = \varphi(x^*x)^{\frac{1}{2}}$ defines the strong operator topology (SOT) on the unit ball of M . The centralizer of φ is denoted by M_φ .

2.2 CONNES–WOODS’ CHARACTERIZATION OF ITPFI FACTORS

Recall that a von Neumann algebra M with separable predual is called *hyperfinite* if there exists an increasing sequence $M_1 \subset M_2 \subset \dots$ of finite-dimensional *-subalgebras such that $M = (\bigcup_{n=1}^\infty M_n)''$. A factor M is called an *Araki–Woods factor* or an *ITPFI* (infinite tensor product of factors of type I) factor, if it is isomorphic to the factor of the form

$$\bigotimes_{i \in I} (M_i, \varphi_i),$$

where I is a countable infinite set and each M_i (resp. φ_i) is a σ -finite type I factor (resp. a faithful normal state). Araki and Woods classified most ITPFI factors:

THEOREM 2.1 ([AW68]). *There exists a unique ITPFI factor with separable predual for each type I_∞, II₁, II_∞ and III_λ, λ ∈ (0, 1]. In particular, all ITPFI factors of type III₁ are isomorphic to*

$$R_\infty := \bigotimes_{n \in \mathbb{N}} (M_3(\mathbb{C}), \text{Tr}(\rho \cdot)),$$

where $\rho := \frac{1}{1+\lambda+\mu} \text{diag}(1, \lambda, \mu)$ and $0 < \lambda, \mu$ satisfies $\frac{\log \lambda}{\log \mu} \notin \mathbb{Q}$.

It is clear that an ITPFI factor with separable predual is hyperfinite. The converse is also true for factors not of type III₀, but false in general. Namely, Connes–Woods [CW85] characterized hyperfinite factors of type III₀ with separable predual which are isomorphic to ITPFI factors by the approximate transitivity of their flow of weights, while the existence of hyperfinite factors of type III₀ with separable predual which are not isomorphic to ITPFI factors had been shown in [Con72]. Let N be a von Neumann algebra, and let F be a finite dimensional subfactor of N with relative commutant $F^c := F' \cap N$ in N . Then it is elementary to check, that the map

$$\sum_{i=1}^n x_i \otimes y_i \mapsto \sum_{i=1}^n x_i y_i, \quad x_i \in F, y_i \in F^c \quad (1 \leq i \leq n)$$

is an isomorphism of $F \otimes F^c$ onto N . If ω_1 is a normal state on F and ω_2 is a normal state on F^c , we let $\omega_1 \otimes \omega_2$ denote the corresponding state on N , i.e.,

$$(\omega_1 \otimes \omega_2)(xy) = \omega_1(x)\omega_2(y), \quad x \in F, y \in F^c.$$

In our proof of

$$[N \text{ injective III}_1 \text{ and } B_\varphi = \mathbb{C}1] \Rightarrow N \cong R_\infty,$$

we shall need the following criterion for a factor to be ITPFI:

PROPOSITION 2.2 ([CW85, Lemma 7.6]). *Let N be a factor on a separable Hilbert space. Then N is ITPFI if and only if N admits a normal faithful state φ with the following property: for every finite set x_1, \dots, x_n of operators in N , for every $\varepsilon > 0$, and every strong* neighborhood \mathcal{V} of 0 in N , there exists a finite dimensional subfactor F of N , such that*

$$x_k \in F + \mathcal{V}, \quad k = 1, \dots, n$$

and

$$\|\varphi - \varphi|_F \otimes \varphi|_{F^c}\| < \varepsilon.$$

2.3 BICENTRALIZERS ON TYPE III₁ FACTORS

In this subsection, we recall Connes' bicentralizers. Let M be a σ -finite von Neumann algebra, and let φ be a normal faithful state on M . We denote by $\text{AC}(\varphi)$ the set of all norm-bounded sequences $(x_n)_{n=1}^\infty$ in M such that $\lim_{n \rightarrow \infty} \|\varphi x_n - x_n \varphi\| = 0$ holds.

DEFINITION 2.3 (Connes). The *bicentralizer* of φ is the set B_φ of all $x \in M$ such that $\lim_{n \rightarrow \infty} \|xa_n - a_n x\|_\varphi = 0$ holds for all $(a_n)_{n=1}^\infty \in \text{AC}(\varphi)$.

Since B_φ is a von Neumann subalgebra of M [Haa87, Proposition 1.3], it holds that $\lim_{n \rightarrow \infty} \|xa_n - a_n x\|_\varphi^\sharp = 0$.

It was conjectured by Connes that for all factors of type III₁ with separable predual, the bicentralizer B_φ of any normal faithful state φ on M is trivial, i.e., $B_\varphi = \mathbb{C}1$ holds. This is still an open problem. We will need the following result on type III₁ factors, known as the Connes–Størmer transitivity:

THEOREM 2.4 ([CS78]). *Let M be a type III₁ factor with separable predual. Then for every faithful normal states φ, ψ on M and $\varepsilon > 0$, there exists a unitary $u \in \mathcal{U}(M)$ such that $\|u\varphi u^* - \psi\| < \varepsilon$ holds.*

Connes showed that by the Connes–Størmer transitivity, for a type III₁ factor M with separable predual, the triviality of B_φ for one fixed faithful normal state φ on M implies the triviality of B_ψ for every faithful normal state ψ (see [Haa87, Corollary 1.5] for the proof). He also showed that the triviality of the bicentralizer is equivalent to the following property (the proof is given in [Haa87, Proposition 1.3 (2)]):

PROPOSITION 2.5 (Connes). *Let M be a von Neumann algebra with a normal faithful state φ . Then $B_\varphi = \mathbb{C}1$ holds, if and only if the following condition is satisfied: for every $a \in M$ and $\delta > 0$,*

$$\overline{\text{conv}}\{u^* a u; u \in \mathcal{U}(M), \|u\varphi - \varphi u\| \leq \delta\} \cap \mathbb{C}1 \neq \emptyset,$$

where $\overline{\text{conv}}$ is the closure of the convex hull in the σ -weak topology.

We will use the following variant of Proposition 2.5.

PROPOSITION 2.6. *Let M be a type III₁ factor with separable predual, and let φ be a normal faithful state on M whose modular automorphism group σ^φ leaves a finite-dimensional subfactor F globally invariant. Let $\varepsilon_{F,\varphi}: M \rightarrow F$ be the normal faithful φ -preserving conditional expectation. Assume that $B_\varphi = \mathbb{C}1$. Then for every $\delta > 0$ and $a \in M$, we have*

$$\varepsilon_{F,\varphi}(a) \in \overline{\text{conv}}\{u^*au; u \in \mathcal{U}(F^c), \|u\xi_\varphi - \xi_\varphi u\| \leq \delta\}. \quad (1)$$

Here, ξ_φ is the representing vector of φ in the natural cone.

Proof. The proof is essentially the same as Proposition 2.5, so we only indicate the outline. Note that by Araki-Powers-Størmer inequality, for every $u \in \mathcal{U}(M)$ one has:

$$\|\xi_\varphi - u\xi_\varphi u^*\|^2 \leq \|\varphi - u\varphi u^*\| \leq \|\xi_\varphi - u\xi_\varphi u^*\| \cdot \|\xi_\varphi + u\xi_\varphi u^*\|.$$

Therefore in the arguments below, we may replace the condition “ $\|u\xi_\varphi - \xi_\varphi u\| \leq \delta$ ” in Proposition 2.6 with the condition “ $\|u\varphi - \varphi u\| \leq \delta$ ”, as we take $\delta > 0$ to be arbitrarily small. As was pointed out in [Haa87, Remark 1.4], it follows from the proof of Proposition 2.5 that the condition $B_\varphi = \mathbb{C}1$ is equivalent to the next condition that for all $a \in M$ and $\delta > 0$,

$$\varphi(a)1 \in \bigcap_{\delta > 0} \overline{\text{conv}}\{u^*au; u \in \mathcal{U}(M), \|u\xi_\varphi - \xi_\varphi u\| < \delta\}. \quad (2)$$

Let $a \in M$. Since $M \cong F \otimes F^c$ with $\varphi = \varphi|_F \otimes \varphi|_{F^c}$, we may now apply (2) to $F^c (\cong M)$ and $\varphi|_{F^c}$ to obtain

$$\varepsilon_{F,\varphi}(a) = \text{id}_F \otimes \varphi|_{F^c}(a) \in \overline{\text{conv}}\{u^*au; u \in \mathcal{U}(F^c), \|u\xi_\varphi - \xi_\varphi u\| \leq \delta\}.$$

Note that we used the fact that $\|\varphi u - u\varphi\| = \|\psi u - u\psi\|$, where $\psi := \varphi|_{F^c}$ and $u \in \mathcal{U}(F^c)$ thanks to the existence of a normal faithful φ -preserving conditional expectation from M onto F^c . \square

2.4 ALMOST UNITARY EQUIVALENCE IN HILBERT N -BIMODULES

We recall a result about almost unitary equivalence in Hilbert bimodules established in [Haa89] which is a generalization of [Haa85, Theorem 4.2]. Let N be a von Neumann algebra, and H be a normal Hilbert N -bimodule, i.e., H is a Hilbert space on which there are defined left and right actions by elements from N :

$$(x, \xi) \mapsto x\xi, \quad (x, \xi) \mapsto \xi x, \quad x \in N, \quad \xi \in H$$

such that the above maps $N \times H \rightarrow H$ are bilinear and

$$(x\xi)y = x(\xi y), \quad x, y \in N, \quad \xi \in H.$$

Moreover, $x \mapsto L_x$, where $L_x \xi := x\xi$ ($\xi \in H$) is a normal unital $*$ -homomorphism, and $x \mapsto R_x$, where $R_x \xi := \xi x$ ($\xi \in H$) is a normal unital $*$ -antihomomorphism.

DEFINITION 2.7. Let N be a von Neumann algebra, let (N, H) be a normal Hilbert N -bimodule, and let $\delta \in \mathbb{R}_+$. Two n -tuples (ξ_1, \dots, ξ_n) and (η_1, \dots, η_n) of unit vectors in H are called δ -related, if there exists a family $(a_i)_{i \in I}$ of operators in N , such that

$$\sum_{i \in I} a_i^* a_i = \sum_{i \in I} a_i a_i^* = 1$$

and

$$\sum_{i \in I} \|a_i \xi_k - \eta_k a_i\|^2 < \delta, \quad k = 1, \dots, n.$$

We will use the following result which relates the δ -relatedness to approximate unitary equivalence in Hilbert N -bimodules:

THEOREM 2.8 ([Haa89, Theorem 2.3]). *For every $n \in \mathbb{N}$ and $\varepsilon > 0$, there exists a $\delta = \delta(n, \varepsilon) > 0$, such that for all von Neumann algebra N and δ -related n -tuples (ξ_1, \dots, ξ_n) and (η_1, \dots, η_n) of unit vectors in a normal Hilbert N -bimodule H , there exists a unitary $u \in \mathcal{U}(N)$ such that*

$$\|u \xi_k - \eta_k u\| < \varepsilon, \quad k = 1, \dots, n.$$

REMARK 2.9. As can be seen in the proof of [Haa89, Theorem 2.3], in order to show that the conclusion of Theorem 2.8 holds, it suffices to show the following: for every σ -strong neighborhood \mathcal{V} of 0 in N , there exist $a_1, \dots, a_p \in N$ such that

$$\sum_{i=1}^p \|a_i \xi_k - \eta_k a_i\|^2 < \delta, \quad k = 1, \dots, n \quad (3)$$

$$\sum_{i=1}^p a_i^* a_i \leq 1, \quad \sum_{i=1}^p a_i a_i^* \leq 1 \quad (4)$$

$$\sum_{i=1}^p a_i^* a_i \in 1 + \mathcal{V}, \quad \sum_{i=1}^p a_i a_i^* \in 1 + \mathcal{V}. \quad (5)$$

This is because we can obtain the conclusions of [Haa89, Lemma 2.5] out of (3), (4) and (5), which is enough to prove Theorem 2.8. We will use this variant in the proof of Lemma 5.6.

3 COMPLETELY POSITIVE MAPS FROM $m \times m$ -MATRICES INTO AN INJECTIVE FACTOR OF TYPE III₁

The main result of this section is:

THEOREM 3.1. *Let N be an injective factor of type III₁ with separable predual, and let φ be a faithful normal state on N . Then for every finite set u_1, \dots, u_n of unitaries in N , and every $\varepsilon, \delta > 0$, there exists $m \in \mathbb{N}$, a unital completely*

positive map $T: M_m(\mathbb{C}) \rightarrow N$, and n unitaries v_1, \dots, v_n in $M_m(\mathbb{C})$, such that $\psi = \varphi \circ T$ is a normal faithful state on $M_m(\mathbb{C})$, and

$$\begin{aligned} \|\sigma_t^\varphi \circ T - T \circ \sigma_t^\psi\| &\leq \delta|t|, \quad t \in \mathbb{R}, \\ \|T(v_k) - u_k\|_\varphi &< \varepsilon, \quad k = 1, \dots, n. \end{aligned}$$

In the following we let $M = N \rtimes_{\sigma^\varphi} \mathbb{R}$ be the crossed product of N by σ^φ with generators $\pi_{\sigma^\varphi}(x)$ ($x \in N$) and $\lambda(s)$ ($s \in \mathbb{R}$). We identify $\pi_{\sigma^\varphi}(x)$ with $x \in N$. Let a be the (unbounded) self-adjoint operator for which $\lambda(s) = \exp(isa)$ ($s \in \mathbb{R}$). For $f \in L^1(\mathbb{R})$, we define the Fourier transform \hat{f} by

$$\hat{f}(s) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-ist} f(t) dt, \quad s \in \mathbb{R}.$$

In the sequel, von Neumann algebra-valued integrals are understood to be the σ -weak sense. Let $(\theta_s^\varphi)_{s \in \mathbb{R}}$ be the dual action of σ^φ on M . By [Haa79-2], there exists a normal faithful semifinite operator-valued weight $P: M_+ \rightarrow \hat{N}_+$ (\hat{N}_+ is the extended positive part of N) given by

$$P(x) = \int_{-\infty}^{\infty} \theta_s^\varphi(x) ds, \quad x \in M_+. \quad (6)$$

Following [CT77], if we put

$$\mathfrak{m} := \text{span} \left\{ x \in M_+; \sup_{c>0} \left\| \int_{-c}^c \theta_t^\varphi(x) dt \right\| < \infty \right\},$$

then the formula (6) for $x \in \mathfrak{m}$ makes sense and $P(x) \in N$. Moreover, $\mathfrak{m} \ni x \mapsto P(x) \in N$ defines a positive linear map.

For all $x \in \mathfrak{m}$, the σ -weak integral $\int_{-c}^c \theta_t^\varphi(x) dt$ is σ -strongly convergent as $c \rightarrow \infty$. The range of P is contained in $\pi_{\sigma^\varphi}(N)$, because $\pi_{\sigma^\varphi}(N)$ is the fixed point algebra in M under the dual action.

LEMMA 3.2. *Let $t \mapsto x(t)$ be a σ -strongly* continuous function from \mathbb{R} to N such that $t \mapsto \|x(t)\|$ is in $L^1(\mathbb{R}) \cap L^\infty(\mathbb{R})$. Put*

$$x := \int_{-\infty}^{\infty} \lambda(t)x(t) dt \in M.$$

Then $x^*x \in \mathfrak{m}$, and

$$P(x^*x) = 2\pi \int_{-\infty}^{\infty} x(t)^*x(t) dt.$$

Proof. Note first, that

$$\begin{aligned} x^*x &= \iint_{\mathbb{R}^2} x(s)^* \lambda(t-s)x(t) ds dt \\ &= \iint_{\mathbb{R}^2} x(s)^* \lambda(t)x(s+t) ds dt. \end{aligned}$$

Put $f_n(s) = e^{-s^2/(4n)}$ ($s \in \mathbb{R}$), and

$$g_n(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} f_n(s)e^{-its} ds = \left(\frac{n}{\pi}\right)^{\frac{1}{2}} e^{-nt^2} \quad (t \in \mathbb{R}).$$

Using that $\theta_s^\varphi(y) = y$ ($s \in \mathbb{R}, y \in N$), $\theta_s^\varphi(\lambda(t)) = e^{-ist}\lambda(t)$ ($s, t \in \mathbb{R}$)³ and the Fubini Theorem, we have for every $\psi \in M_*$,

$$\begin{aligned} \langle \psi, \int_{-\infty}^{\infty} \theta_u^\varphi(x^*x)f_n(u) du \rangle &= \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-itu} f_n(u)\psi(x(s)^*\lambda(t)x(s+t)) ds dt du \\ &= \int_{-\infty}^{\infty} g_n(t) \left(\int_{-\infty}^{\infty} 2\pi\psi(x(s)^*\lambda(t)x(s+t)) ds \right) dt. \end{aligned}$$

Since $t \mapsto \int_{-\infty}^{\infty} \psi(x(s)^*\lambda(t)x(s+t))ds$ is in $C_0(\mathbb{R})$ and $g_n \xrightarrow{n \rightarrow \infty} \delta_0$ (weak* in $C_0(\mathbb{R})^*$), we have

$$\lim_{n \rightarrow \infty} \langle \psi, \int_{-\infty}^{\infty} \theta_u^\varphi(x^*x)f_n(u) du \rangle = \langle \psi, 2\pi \int_{-\infty}^{\infty} x(s)^*x(s) ds \rangle.$$

Since $\psi \in M_*$ is arbitrary, $\theta_u^\varphi(x^*x) \geq 0$ and $f_n \nearrow 1$ uniformly on compact sets, it follows that

$$\lim_{n \rightarrow \infty} \int_{-\infty}^{\infty} \theta_u^\varphi(x^*x)f_n(u)du = 2\pi \int_{-\infty}^{\infty} x(s)^*x(s) ds \quad (\sigma\text{-strongly}).$$

Therefore $x^*x \in \mathfrak{m}$, and $P(x^*x) = 2\pi \int_{-\infty}^{\infty} x(t)^*x(t) dt$. □

LEMMA 3.3. *Let a be the (unbounded) self-adjoint operator affiliated with M for which $\exp(ita) = \lambda(t)$ ($t \in \mathbb{R}$) holds. Let $\alpha > 0$, and let e_α be the spectral projection of the operator a corresponding to the interval $[0, \alpha]$. Then for each $x \in N$, one has $e_\alpha x e_\alpha \in \mathfrak{m}$ and*

$$P(e_\alpha x e_\alpha) = \int_{-\infty}^{\infty} \sigma_t^\varphi(x) \frac{1 - \cos \alpha t}{\pi t^2} dt, \quad x \in N. \tag{7}$$

Proof. It is sufficient to consider the case $x \geq 0$, so we can assume that $x = y^*y$ ($y \in N$). For $f \in L^1(\mathbb{R}) \cap L^\infty(\mathbb{R}) \cap C(\mathbb{R})$, we have

$$\begin{aligned} y\hat{f}(a) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} y\lambda(-t)f(t)dt = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \lambda(-t)\sigma_t^\varphi(y)f(t) dt \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \lambda(t)\sigma_{-t}^\varphi(y)f(-t) dt. \end{aligned}$$

³Typewriter’s note: Haagerup used the convention $\theta_s^\varphi(\lambda(t)) = e^{ist}\lambda(t)$. However, since the negative sign convention is widely accepted, we decided to change the definition.

Hence by Lemma 3.2, $\hat{f}(a)^*x\hat{f}(a) \in \mathfrak{m}$, and

$$P(\hat{f}(a)^*x\hat{f}(a)) = \int_{-\infty}^{\infty} \sigma_t^\varphi(x)|f(t)|^2 dt.$$

For $n > \frac{2}{\alpha}$, let g_n be the continuous function on \mathbb{R} for which

$$\begin{aligned} g_n(t) &= 0, & t \leq 0, t \geq \alpha, \\ g_n(t) &= 1, & t \in [\frac{1}{n}, \alpha - \frac{1}{n}], \end{aligned}$$

and for which the graph is a straight line on $[0, \frac{1}{n}]$ and $[\alpha - \frac{1}{n}, \alpha]$. Since a has no point spectrum, $g_n(a) \nearrow e_\alpha$ ($n \rightarrow \infty$). It is elementary to check that each g_n is of the form $g_n = \hat{f}_n$ for a function $f_n \in L^1(\mathbb{R}) \cap L^\infty(\mathbb{R}) \cap C(\mathbb{R})$ (use, for example, the fact that $g_n = n1_{[0, \frac{1}{n}]} * 1_{[0, \alpha - \frac{1}{n}]}$). Hence $g_n(a)^2 \in \mathfrak{m}$, and by the Plancherel Theorem, we get

$$P(g_n(a)^2) = P(\hat{f}_n(a)^*\hat{f}_n(a)) = \|f_n\|_2^2 1 = \|\hat{f}_n\|_2^2 1.$$

Since $\sup_n \|\hat{f}_n\|_2^2 = \alpha < \infty$, we have $e_\alpha \in \mathfrak{m}$ and $P(e_\alpha) = \alpha 1$. Therefore $e_\alpha M e_\alpha \subseteq \mathfrak{m}$, and the restriction of P to $e_\alpha M e_\alpha$ is a positive normal map. Hence for $x \in N$,

$$P(e_\alpha x e_\alpha) = \lim_{n \rightarrow \infty} P(g_n(a) x g_n(a)) = \lim_{n \rightarrow \infty} \int_{-\infty}^{\infty} \sigma_t^\varphi(x) |f_n(t)|^2 dt \quad (\sigma\text{-strongly}).$$

Since $\|g_n - 1_{[0, \alpha]}\|_2 \xrightarrow{n \rightarrow \infty} 0$, it follows that f_n converges in $L^2(\mathbb{R})$ to the function

$$\begin{aligned} f(t) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} 1_{[0, \alpha]}(s) e^{ist} ds \\ &= -\frac{i}{t\sqrt{2\pi}} (e^{i\alpha t} - 1). \end{aligned}$$

Hence $|f_n|^2 \xrightarrow{n \rightarrow \infty} |f|^2$ in $L^1(\mathbb{R})$, with $|f|^2(t) = \frac{1}{\pi t^2} (1 - \cos \alpha t)$ ($t \in \mathbb{R}$). Therefore (7) holds. □

LEMMA 3.4. *Let N be an injective factor of type III₁ with separable predual, φ be a faithful normal state on N and let R be the hyperfinite II₁ factor with tracial state τ . For every finite set u_1, \dots, u_n of unitaries in N and every $\varepsilon > 0$, there exist x_1, \dots, x_n in the unit ball of R , a normal unital completely positive map $T: R \rightarrow N$, such that $\psi = \varphi \circ T$ is a normal faithful state on R , and*

$$\sigma_t^\varphi \circ T = T \circ \sigma_t^\psi, \quad t \in \mathbb{R}, \tag{8}$$

$$\|T(x_k) - u_k\|_\varphi < \varepsilon, \quad k = 1, \dots, n. \tag{9}$$

Moreover, the spectrum of $h = d\psi/d\tau$ is a closed interval $[\lambda_1, \lambda_2]$, $0 < \lambda_1 < \lambda_2 < \infty$, and h has no eigenvalues.

Proof. ⁴ Let $M = N \rtimes_{\sigma^\varphi} \mathbb{R}$. By [Tak73-2], M has a normal faithful semifinite trace τ , such that

$$\tau \circ \theta_s^\varphi = e^{-s} \tau \quad (s \in \mathbb{R}).$$

The trace τ can be constructed in the following way: Let $\tilde{\varphi}$ be the dual weight of φ on M (cf. [Haa79]). Let a be the self-adjoint operator for which $\exp(ita) = \lambda(t)$ ($t \in \mathbb{R}$). Then a is affiliated with the centralizer $M_{\tilde{\varphi}}$ of $\tilde{\varphi}$ and

$$\tau = \tilde{\varphi}(e^{-a} \cdot)$$

in the sense of Pedersen-Takesaki [PT73]. By [Haa79-2], $\tilde{\varphi}$ is on the subspace \mathfrak{m} given by

$$\tilde{\varphi}(x) = \varphi \circ P(x), \quad x \in \mathfrak{m}.$$

Let $\alpha > 0$, and let $e_\alpha = 1_{[0, \alpha]}(a)$. Then by Lemma 3.3, $e_\alpha \in \mathfrak{m}$ and $P(e_\alpha) = \alpha 1$. Hence $\tilde{\varphi}|_{e_\alpha M e_\alpha}$ is a positive normal functional, and $\tilde{\varphi}(e_\alpha) = \alpha$. Finally,

$$\tau(e_\alpha) = \tilde{\varphi}(e^{-a} e_\alpha) = \int_0^\alpha e^{-t} dt = 1 - e^{-\alpha} < \infty,$$

because

$$e^{-a} e_\alpha = \int_0^\alpha e^{-\lambda} d e(\lambda),$$

where

$$a = \int_{-\infty}^\infty \lambda d e(\lambda)$$

is the spectral resolution of a , and $d\tilde{\varphi}(e(\lambda)) = d\lambda$.

Since N is of type III₁, M is a type II_∞ factor, and therefore $e_\alpha M e_\alpha$ is a II₁ factor. Moreover, the injectivity of N implies that M is also injective, so that $e_\alpha M e_\alpha$ is isomorphic to the hyperfinite factor R of type II₁ by [Con76].

CLAIM. For any $x \in N$, we have

$$\lim_{\alpha \rightarrow \infty} \left\| \frac{1}{\alpha} P(e_\alpha x e_\alpha) - x \right\|_\varphi = 0.$$

This follows from a basic property of the Fejér kernel (see e.g., [Kat68, Chapters I and VI]), but we include the proof for completeness. Let $\varepsilon > 0$. Choose $t_0 > 0$ small enough so that $\|\sigma_t^\varphi(x) - x\|_\varphi \leq \varepsilon$ for all $t \in [-t_0, t_0]$. Moreover, by $\left| \frac{1 - \cos(\alpha t)}{\pi \alpha t^2} \right| \leq \frac{2}{\pi \alpha} \cdot \frac{1}{t^2}$, we have

$$\lim_{\alpha \rightarrow \infty} \int_{|t| \geq t_0} \frac{1 - \cos(\alpha t)}{\pi \alpha t^2} dt = 0.$$

⁴Typewriter's note: Since some pages were missing from the original notes, we could not find all parts of the proofs of Lemma 3.4 and Theorem 3.1. We include the following proof for the reader's convenience.

By Lemma 3.3, we have

$$\begin{aligned} \limsup_{\alpha \rightarrow \infty} \left\| \frac{1}{\alpha} P(e_\alpha x e_\alpha) - x \right\|_\varphi &\leq \limsup_{\alpha \rightarrow \infty} \int_{-\infty}^{\infty} \|\sigma_t^\varphi(x) - x\|_\varphi \frac{1 - \cos(\alpha t)}{\pi \alpha t^2} dt \\ &\leq \varepsilon + 2\|x\|_\varphi \limsup_{\alpha \rightarrow \infty} \int_{|t| \geq t_0} \frac{1 - \cos(\alpha t)}{\pi \alpha t^2} dt \\ &= \varepsilon. \end{aligned}$$

Since $\varepsilon > 0$ is arbitrary, we obtain the conclusion.

Let $n \geq 1$, $u_1, \dots, u_n \in \mathcal{U}(N)$ and $\varepsilon, \delta > 0$ be given. By the above claim, we may choose $\alpha > 0$ large enough so that

$$\left\| \frac{1}{\alpha} P(e_\alpha u_k e_\alpha) - u_k \right\|_\varphi < \varepsilon, \quad 1 \leq k \leq n. \quad (10)$$

Define $T := \alpha^{-1} P|_{e_\alpha M e_\alpha} : R = e_\alpha M e_\alpha \rightarrow N$ and

$$\psi := \varphi \circ T = \frac{1}{\alpha} \varphi \circ P(e_\alpha \cdot e_\alpha) = \frac{1}{\alpha} \tilde{\varphi}(e_\alpha \cdot e_\alpha).$$

Then T is a normal unital completely positive map, and ψ is a normal faithful state on R . By (10) we have

$$\|T(x_k) - u_k\|_\varphi < \varepsilon, \quad 1 \leq k \leq n,$$

where $x_k := e_\alpha u_k e_\alpha$ ($1 \leq k \leq n$) are in the unit ball of R . Moreover, since $e_\alpha \in M_{\tilde{\varphi}}$ and since $\sigma_t^\varphi \circ P = P \circ \sigma_t^{\tilde{\varphi}}$ ($t \in \mathbb{R}$), we have $\sigma_t^\varphi \circ T = T \circ \sigma_t^{\tilde{\varphi}}$ ($t \in \mathbb{R}$). By construction, we have

$$h := \frac{d\psi}{d\tau} = \frac{1 - e^{-\alpha}}{\alpha} \exp(a) e_\alpha,$$

which has no atoms and the spectrum of h is a closed bounded interval in $\mathbb{R}_+^* = (0, \infty)$. \square

LEMMA 3.5. *Let $B \subset A$ be an inclusion of unital C^* -algebras and $E: A \rightarrow B$ be a unital completely positive map. Let $h \in A$ be a self-adjoint element with $\sigma(h) \subset [\lambda_1, \lambda_2]$, where $\sigma(\cdot)$ denotes the spectrum and $\lambda_1 < \lambda_2$ are reals. Then $\sigma(E(h)) \subset [\lambda_1, \lambda_2]$.*

Proof. Let $\lambda < \lambda_1$. Then $h - \lambda$ is positive and invertible. Take a nonzero $x \in A$ such that $(h - \lambda)^{\frac{1}{2}} x = 1$, so that $E((h - \lambda)^{\frac{1}{2}} x x^* (h - \lambda)^{\frac{1}{2}}) = 1$. The left hand side is dominated by $\|x\|^2 E(h - \lambda)$, whence $E(h - \lambda) \geq \|x\|^{-2} 1$, showing that $E(h) - \lambda 1$ is invertible. Thus $\lambda \notin \sigma(E(h))$. Similarly, $\sigma(E(h)) \cap (\lambda_2, \infty) = \emptyset$ holds. Therefore $\sigma(E(h)) \subset [\lambda_1, \lambda_2]$. \square

Proof of Theorem 3.1. We may assume that $0 < \varepsilon < 1$. By Lemma 3.4, there exist a normal unital completely positive map $T: R \rightarrow N$ and x_1, \dots, x_n in the unit ball of R satisfying $\sigma_t^\varphi \circ T = T \circ \sigma_t^\psi$, ($t \in \mathbb{R}$), where $\psi := \varphi \circ T: R \rightarrow N$ and

$$\|T(x_k) - u_k\|_\varphi < \frac{\varepsilon^2}{16}, \quad 1 \leq k \leq n. \quad (11)$$

Let $h := d\psi/d\tau \in R_+$. Then by Lemma 3.4, $\sigma(h) = [\lambda_1, \lambda_2]$ for some positive reals $\lambda_1 < \lambda_2$ and h does not have a point spectrum. Since $\log(\cdot)$ is continuous on $[\lambda_1, \lambda_2]$, continuous functional calculus guarantees that there exists $\delta' > 0$ such that for all $a, b \in R_+$, we have the following implication

$$\sigma(a), \sigma(b) \subset [\lambda_1, \lambda_2] \text{ and } \|a - b\| < \delta' \Rightarrow \|\log a - \log b\| < \frac{\delta}{4}. \quad (12)$$

By using the spectral decomposition of h , we can choose a partition of unity $\{p_i\}_{i=1}^\ell$ in R and $\{\mu_i\}_{i=1}^\ell$ in \mathbb{R}_+^* such that

$$\begin{aligned} \tau(p_i) &= \frac{1}{\ell}, \quad hp_i = p_i h, \\ \|(\log h)p_i - (\log \mu_i)p_i\| &< \frac{1}{4}\delta, \\ \|hp_i - \mu_i p_i\| &< \delta', \end{aligned}$$

for all ($1 \leq i \leq \ell$). Let $h_0 := \sum_{i=1}^\ell \mu_i p_i$, and we have

$$\|h - h_0\| < \delta' \quad \text{and} \quad \|\log h - \log h_0\| < \frac{1}{4}\delta. \quad (13)$$

Moreover, we may arrange $\{\mu_i\}_{i=1}^\ell$ so that $h_0 = \sum_{i=1}^\ell \mu_i p_i$ satisfies

$$\sigma(h_0) \subset [\lambda_1, \lambda_2]. \quad (14)$$

Since R is hyperfinite, there exists a type I subfactor F of R so that $p_i \in F$ ($1 \leq i \leq \ell$) and

$$\|x_k - E_F(x_k)\|_\varphi < \frac{\varepsilon^2}{16}, \quad 1 \leq k \leq n, \quad (15)$$

where $E_F: R \rightarrow F$ denotes the τ -preserving conditional expectation. Put $T_F := T|_F: F \rightarrow N$ and $y_k := E_F(x_k)$ ($1 \leq k \leq n$). Combining (11) and (15), for all $1 \leq k \leq n$, we have (use the Schwarz inequality for completely positive maps)

$$\begin{aligned} \|T_F(y_k) - u_k\|_\varphi &\leq \|T(y_k) - T(x_k)\|_\varphi + \|T(x_k) - u_k\|_\varphi \\ &\leq \|y_k - x_k\|_\psi + \|T(x_k) - u_k\|_\varphi \\ &< \frac{\varepsilon^2}{8}. \end{aligned} \quad (16)$$

Then we follow the argument of [Haa89, Lemma 6.2]. Take $v_1, \dots, v_k \in \mathcal{U}(F)$ such that

$$y_k = v_k |y_k|, \quad |y_k| := (y_k^* y_k)^{\frac{1}{2}}, \quad 1 \leq k \leq n.$$

Then again by the Schwarz inequality for completely positive maps and (16),

$$\|y_k\|_\psi \geq \|T_F(y_k)\|_\varphi > \|u_k\|_\varphi - \frac{\varepsilon^2}{8}.$$

Since $\|(y_k^* y_k)^{\frac{1}{2}}\|_\psi = \|y_k\|_\psi$ and $|y_k|^2 + (1 - |y_k|)^2 \leq 1$ (because $0 \leq |y_k| \leq 1$), we have

$$\begin{aligned} \|v_k - y_k\|_\psi^2 &= \|1 - |y_k|\|_\psi^2 \leq 1 - \| |y_k| \|_\psi^2 \\ &< 1 - \left(1 - \frac{\varepsilon^2}{8}\right)^2 \\ &< \frac{\varepsilon^2}{4}. \end{aligned}$$

Therefore since $\varepsilon^2 < \varepsilon$,

$$\begin{aligned} \|T_F(v_k) - u_k\|_\varphi &\leq \|T_F(v_k - y_k)\|_\varphi + \|T_F(y_k) - u_k\|_\varphi \\ &< \|v_k - y_k\|_\psi + \frac{\varepsilon^2}{8} \\ &< \varepsilon. \end{aligned}$$

Next, set $\chi := \tau(h_0 \cdot) \in (R_*)_+$. Note that $\sigma_t^{\chi|F} = \sigma_t^\chi|_F$ ($t \in \mathbb{R}$), since $h_0 \in F$. Then by (13), we have

$$\begin{aligned} \|h^{it} - h_0^{it}\| &= \left\| \int_0^1 \frac{d}{ds} e^{ist \log h} e^{i(1-s)t \log h_0} ds \right\| \\ &\leq \int_0^1 \|e^{ist \log h} (t \log h - t \log h_0) e^{i(1-s)t \log h_0}\| ds \\ &\leq \|\log h - \log h_0\| |t| \\ &\leq \frac{\delta |t|}{4}. \end{aligned} \tag{17}$$

On the other hand, $h_F := d\psi|_F/d\tau|_F \in F_+$ is equal to $E_F(h)$. Therefore by Lemma 3.5, $\sigma(h_F) \subset [\lambda_1, \lambda_2]$. Moreover, since $E_F(h_0) = h_0$, we have

$$\|h_F - h_0\| = \|E_F(h - h_0)\| \leq \|h - h_0\| < \delta'.$$

This shows by (12) and (14) that $\|\log h_F - \log h_0\| < \frac{\delta}{4}$. Therefore by the same argument, we have

$$\|h_F^{it} - h_0^{it}\| \leq \frac{\delta |t|}{4}, \quad t \in \mathbb{R}. \tag{18}$$

For all $t \in \mathbb{R}$ and $x \in F$,

$$\begin{aligned} \|\sigma_t^\varphi \circ T_F(x) - T_F \circ \sigma_t^{\psi|F}(x)\| &= \|T(\sigma_t^\psi(x) - \sigma_t^{\psi|F}(x))\| \\ &\leq \|\sigma_t^\psi(x) - \sigma_t^{\psi|F}(x)\| \\ &\leq \|\sigma_t^\psi(x) - \sigma_t^\chi(x)\| + \|\sigma_t^{\chi|F}(x) - \sigma_t^{\psi|F}(x)\|. \end{aligned}$$

By (17), we have

$$\begin{aligned} \|\sigma_t^\psi(x) - \sigma_t^X(x)\| &= \|h^{it}xh^{-it} - h_0^{it}xh_0^{-it}\| \\ &\leq (\|h^{it} - h_0^{it}\| + \|h^{-it} - h_0^{-it}\|)\|x\| \\ &\leq \frac{1}{2}\delta|t|\|x\|. \end{aligned}$$

Similarly, by (18),

$$\begin{aligned} \|\sigma_t^{X|F}(x) - \sigma_t^{\psi|F}(x)\| &\leq (\|h_0^{it} - h_F^{it}\| + \|h_0^{-it} - h_F^{-it}\|)\|x\| \\ &\leq \frac{1}{2}\delta|t|\|x\|. \end{aligned}$$

These altogether imply that $\|\sigma_t^\varphi \circ T_F(x) - T_F \circ \sigma_t^{\psi|F}(x)\| \leq \delta|t|\|x\|$. \square

4 \mathbb{Q} -STABLE STATES ON III_1 FACTORS

For technical reasons we shall consider a special class of normal faithful states, which we call \mathbb{Q} -stable states, because they have nice properties with respect to certain operations involving rationals.

DEFINITION 4.1. A normal faithful state φ on a von Neumann algebra N is called \mathbb{Q} -stable, if for every $m \in \mathbb{N}$, there exist m isometries $u_1, \dots, u_m \in N$ with orthogonal range projections, such that

$$\begin{aligned} \sum_{i=1}^m u_i u_i^* &= 1, \\ \varphi u_i &= \frac{1}{m} u_i \varphi, \quad i = 1, \dots, m. \end{aligned}$$

THEOREM 4.2. *Every factor of type III_1 with separable predual has a \mathbb{Q} -stable normal faithful state.*

For the proof of Theorem 4.2, we shall need two lemmas:

LEMMA 4.3. *The Araki–Woods factor R_∞ has a \mathbb{Q} -stable normal faithful state.*

Proof. Let R_λ ($0 < \lambda < 1$) be the Powers factor of type III_λ , and let φ_λ be the product state on R_λ . Then φ_λ is normal and faithful, and σ^{φ_λ} has period $-2\pi/\log \lambda$. Then the centralizer $(R_\lambda)_{\varphi_\lambda}$ is a type II_1 factor (cf. [Con73, Théorème 4.2.6]), and there exists an isometry $u \in R_\lambda$ such that

$$\sigma_t^{\varphi_\lambda}(u) = \lambda^{it}u, \quad t \in \mathbb{R}.$$

This implies that $\sigma_t^{\varphi_\lambda}(uu^*) = uu^*$ ($t \in \mathbb{R}$), i.e., $uu^* \in (R_\lambda)_{\varphi_\lambda}$. Moreover, by [Tak73, Lemma 1.6], we have

$$\varphi_\lambda u = \lambda u \varphi_\lambda,$$

and hence $\varphi_\lambda(uu^*) = (\varphi_\lambda u)(u^*) = \lambda\varphi_\lambda(u^*u) = \lambda$.

Assume now, that $\lambda = \frac{1}{m}$, $m \in \mathbb{N}, m \geq 2$. Then we can choose m equivalent orthogonal projections $p_1, \dots, p_m \in (R_\lambda)_{\varphi_\lambda}$ with sum 1, such that $p_1 = uu^*$. Next, choose partial isometries $v_1, \dots, v_m \in (R_\lambda)_{\varphi_\lambda}$ such that

$$v_i^*v_i = p_1, \quad v_iv_i^* = p_i, \quad i = 1, \dots, m.$$

Put $u_i = v_iu$, $i = 1, \dots, m$. Then u_1, \dots, u_m are m isometries in R_λ , such that $\sum_{i=1}^m u_iu_i^* = 1$, and $\varphi_\lambda u_i = \lambda u_i\varphi_\lambda$, $i = 1, \dots, m$. Put now

$$(P, \varphi) = \bigotimes_{m=2}^{\infty} (R_{\frac{1}{m}}, \varphi_{\frac{1}{m}}).$$

Then it is clear from the above computations, that φ is a \mathbb{Q} -stable normal faithful state on P (observe that it is sufficient to consider $m \geq 2$ case in Definition 4.1). Moreover, P is an ITPFI factor for which the asymptotic ratio set $r_\infty(P)$ contains $\{\frac{1}{m}; m \in \mathbb{N}\}$. Since $r_\infty(P) \cap \mathbb{R}_+$ is a closed subgroup of \mathbb{R}_+ , we have $r_\infty(P) \supseteq \mathbb{R}_+$. Therefore by Araki–Woods' Theorem [AW68, Theorem 7.6], $P \cong R_\infty$ holds. \square

LEMMA 4.4. *Let N be a factor of type III₁ with separable predual. Then there exists a normal faithful conditional expectation of N onto a subfactor P isomorphic to R_∞ .*

Proof. ⁵ We can write R_∞ as an infinite tensor product

$$R_\infty = \bigotimes_{k=1}^{\infty} (P_k, \omega_k),$$

where each P_k is a copy of the 2×2 matrices $M_2(\mathbb{C})$ and $(\omega_k)_{k=1}^{\infty}$ is a sequence of normal faithful states on $M_2(\mathbb{C})$. Let ψ be a fixed normal faithful state on N . Since N is properly infinite, we have $N \otimes M_2(\mathbb{C}) \cong N$. Moreover, by Connes–Størmer transitivity theorem [CS78], we can choose a $*$ -isomorphism $\Phi: N \otimes M_2(\mathbb{C}) \rightarrow N$ such that

$$\|(\psi \otimes \omega_1) \circ \Phi^{-1} - \psi\| < \frac{1}{2}.$$

Put $F_1 = \Phi(\mathbb{C} \otimes M_2(\mathbb{C}))$, and $\varphi_1 = (\psi \otimes \omega_1) \circ \Phi^{-1}$. Then F_1 is a type I₂ subfactor of N . Moreover, it holds that $N \cong F_1 \otimes F_1^c$, where $F_1^c = F_1' \cap N$ is the relative commutant, and $\varphi_1 = \varphi_1|_{F_1} \otimes \varphi_1|_{F_1^c}$. Moreover, we have

$$(F_1, \varphi_1|_{F_1}) \cong (P_1, \omega_1).$$

⁵Typewriter's note: this result has been extended by Haagerup–Musat [HM09, Theorem 3.5], where the authors study more general embeddings of ITPFI type III factors into type III factors as the range of normal faithful conditional expectations.

Using the same arguments to the type III₁ factor F_1^c , we can find a type I₂-subfactor $F_2 \subset F_1^c$, a normal faithful state φ'_2 on F_1^c , such that $\|\varphi'_2 - \varphi_1|_{F_1^c}\| < \frac{1}{4}$,

$$\varphi'_2 = \varphi'_2|_{F_2} \otimes \varphi'_2|_{(F_1 \otimes F_2)^c},$$

and $(F_2, \varphi'_2|_{F_2}) \cong (P_2, \omega_2)$. Thus, if we put $\varphi_2 = \varphi_1|_{F_1} \otimes \varphi'_2$, we have $\|\varphi_1 - \varphi_2\| < \frac{1}{4}$,

$$\varphi_2 = \varphi_2|_{F_1} \otimes \varphi_2|_{F_2} \otimes \varphi_2|_{(F_1 \otimes F_2)^c},$$

and

$$\begin{aligned} (F_1, \varphi_2|_{F_1}) &\cong (P_1, \omega_1), \\ (F_2, \varphi_2|_{F_2}) &\cong (P_2, \omega_2). \end{aligned}$$

Proceeding in this way, we obtain a sequence $(F_k)_{k=1}^\infty$ of mutually commuting type I₂-subfactors of N , and a sequence $(\varphi_k)_{k=1}^\infty$ of normal faithful states on N , such that

$$\|\varphi_k - \varphi_{k-1}\| < 2^{-k}, \quad k \geq 2,$$

and such that for fixed $m \in \mathbb{N}$:

$$\varphi_m = \varphi_m|_{F_1} \otimes \varphi_m|_{F_2} \otimes \dots \otimes \varphi_m|_{F_m} \otimes \varphi_m|_{(F_1 \otimes \dots \otimes F_m)^c},$$

and

$$(F_i, \varphi_m|_{F_i}) \cong (P_i, \omega_i) \quad i = 1, \dots, m.$$

Let φ be the norm limit in N_* of the sequence $(\varphi_k)_{k=1}^\infty$. Then φ is a normal state, but it can fail to be faithful. From the properties of φ_k , we have for all $m \in \mathbb{N}$,

$$\varphi = \varphi|_{F_1} \otimes \varphi|_{F_2} \otimes \dots \otimes \varphi|_{F_m} \otimes \varphi|_{(F_1 \otimes \dots \otimes F_m)^c},$$

and

$$(F_m, \varphi|_{F_m}) \cong (P_m, \omega_m).$$

Let r_k be the ratio between the largest and the smallest eigenvalues of $d\omega_k/d\text{Tr}$. Let $u \in \mathcal{U}(P_k)$. We may assume that $\omega_k = \text{Tr}(h_k \cdot)$, $h_k := \frac{1}{1+r_k} \text{diag}(r_k, 1)$ ($r_k \geq 1$). Then if $a = \begin{pmatrix} x & y \\ z & w \end{pmatrix} \in P_k$ is positive, then

$$\begin{aligned} u\omega_k u^*(a) &= \text{Tr}(h_k u^* a u) = \text{Tr}((u^* a u)^{\frac{1}{2}} h_k (u^* a u)^{\frac{1}{2}}) \\ &\geq \frac{1}{1+r_k} \text{Tr}(u^* a u) = \frac{x+w}{1+r_k} \\ &\geq r_k^{-1} \left(\frac{r_k x}{1+r_k} + \frac{w}{1+r_k} \right) \\ &= r_k^{-1} \omega_k(a). \end{aligned}$$

Similarly, $u\omega_k u^*(a) \leq r_k \omega_k(a)$ holds. This shows that

$$r_k^{-1} \omega_k \leq u\omega_k u^* \leq r_k \omega_k.$$

Hence for all $u \in \mathcal{U}(F_k)$,

$$r_k^{-1}\varphi \leq u\varphi u^* \leq r_k\varphi.$$

Thus, φ and $u\varphi u^*$ have the same support projection in N , i.e., with $e = \text{supp}(\varphi)$, we have

$$ueu^* = e, \quad u \in \mathcal{U}(F_k), \quad k \in \mathbb{N}.$$

This shows that $e \in (\bigcup_{k=1}^\infty F_k)' \cap N$. Put $G_k = eF_k$. Then $(G_k)_{k=1}^\infty$ is a sequence of commuting subfactors of eNe . Moreover, the restriction φ_e of φ to eNe is a normal faithful state on eNe , and

$$(G_1 \otimes \dots \otimes G_m, \varphi_e|_{G_1 \otimes \dots \otimes G_m}) \cong \bigotimes_{k=1}^m (P_k, \omega_k)$$

for all $m \in \mathbb{N}$. Let P be the von Neumann algebra generated by $\bigcup_{k=1}^\infty G_k$. Then

$$(P, \varphi|_P) \cong \bigotimes_{k=1}^\infty (P_k, \omega_k).$$

In particular, $P \cong R_\infty$. Moreover, since

$$\varphi_e = \varphi_e|_{G_1} \otimes \dots \otimes \varphi_e|_{G_m} \otimes \varphi_e|_{(G_1 \otimes \dots \otimes G_m)^c},$$

where $(G_1 \otimes \dots \otimes G_m)^c$ denotes the relative commutant of $\bigcup_{k=1}^m G_k$ in eNe , we have

$$\sigma_t^{\varphi_e}(G_1 \otimes \dots \otimes G_m) = G_1 \otimes \dots \otimes G_m, \quad t \in \mathbb{R}$$

for all $m \in \mathbb{N}$, and hence also $\sigma_t^{\varphi_e}(P) = P$, $t \in \mathbb{R}$. Thus by [Tak72], there exists a normal faithful conditional expectation of eNe onto P . This completes the proof, since eNe is isomorphic to N . □

Proof of Theorem 4.2. Let N be a type III₁ factor with separable predual. By Lemmata 4.3 and 4.4, we can choose a normal faithful conditional expectation E of N onto a subfactor P of N isomorphic to R_∞ . Moreover, we can choose a \mathbb{Q} -stable normal faithful state ω on P . Put $\varphi = \omega \circ E$. Then it follows from the bimodule property of conditional expectations [Tom58, Theorem 1] that φ is a \mathbb{Q} -stable normal faithful state on N . □

THEOREM 4.5. *Let φ be a \mathbb{Q} -stable normal faithful state on a von Neumann algebra N , let $m \in \mathbb{N}$, and let q_1, \dots, q_m be m positive rational numbers with sum 1. Then there exists a type I _{m} subfactor F of N , such that*

- (a) $\varphi = \varphi|_F \otimes \varphi|_{F^c}$.
- (b) $\varphi|_{F^c}$ is \mathbb{Q} -stable.
- (c) $d\varphi|_F/d\text{Tr}_F$ has eigenvalues (q_1, \dots, q_m) .

Here, Tr_F denotes the trace on F for which $\text{Tr}_F(1) = m$.

We prove first:

LEMMA 4.6. *Let φ be a \mathbb{Q} -stable normal faithful state on a von Neumann algebra N , and let q_1, \dots, q_m be positive rational numbers with sum 1. Then there exist isometries $u_1, \dots, u_m \in N$ with orthogonal ranges, such that*

$$\sum_{i=1}^m u_i u_i^* = 1, \\ \varphi u_i = q_i u_i \varphi, \quad i = 1, \dots, m.$$

Proof. We can choose integers $p, p_1, \dots, p_m \in \mathbb{N}$ such that

$$q_i = \frac{p_i}{p}, \quad i = 1, \dots, m.$$

Note that $\sum_{i=1}^m p_i = p$. By Definition 4.1, for each $i \in \{1, \dots, m\}$ we can choose p_i isometries v_{i1}, \dots, v_{ip_i} in N with orthogonal ranges, such that

$$\sum_{j=1}^{p_i} v_{ij} v_{ij}^* = 1 \quad \text{and} \quad \varphi v_{ij} = \frac{1}{p_i} v_{ij} \varphi, \quad j = 1, \dots, p_i.$$

Moreover, since the set $\{(i, j); 1 \leq i \leq m, 1 \leq j \leq p_i\}$ contains $\sum_{i=1}^m p_i = p$ elements, we can also find isometries $w_{ij} \in N$ ($1 \leq i \leq m, 1 \leq j \leq p_i$) with orthogonal ranges, such that

$$\sum_{i=1}^m \sum_{j=1}^{p_i} w_{ij} w_{ij}^* = 1 \quad \text{and} \quad \varphi w_{ij} = \frac{1}{p} w_{ij} \varphi, \quad 1 \leq i \leq m, 1 \leq j \leq p_i.$$

Put now

$$u_i := \sum_{j=1}^{p_i} w_{ij} v_{ij}^*, \quad i = 1, \dots, m.$$

Then

$$u_i^* u_i = \sum_{j=1}^{p_i} v_{ij} v_{ij}^* = 1, \\ \sum_{i=1}^m u_i u_i^* = \sum_{i=1}^m \sum_{j=1}^{p_i} w_{ij} w_{ij}^* = 1,$$

and since $\varphi w_{ij} = \frac{1}{p} w_{ij} \varphi$ and $v_{ij}^* \varphi = \frac{1}{p_i} \varphi v_{ij}^*$ for all (i, j) , we get

$$\varphi u_i = \sum_{j=1}^{p_i} \varphi w_{ij} v_{ij}^* = \sum_{j=1}^{p_i} \frac{p_i}{p} w_{ij} v_{ij}^* \varphi = q_i u_i \varphi.$$

This proves Lemma 4.6. □

Proof of Theorem 4.5. Choose m isometries $u_1, \dots, u_m \in N$ satisfying the conditions in Lemma 4.6. We can define a $*$ -isomorphism Φ of $N \otimes M_m(\mathbb{C})$ onto N by

$$\Phi \left(\sum_{i,j=1}^m x_{ij} \otimes e_{ij} \right) := \sum_{i,j=1}^m u_i x_{ij} u_j^*,$$

where $(e_{ij})_{i,j=1}^m$ are the matrix units in $M_m(\mathbb{C})$. Then using $\varphi u_i = \lambda_i u_i \varphi$, we get

$$\begin{aligned} (\varphi \circ \Phi) \left(\sum_{i,j=1}^m x_{ij} \otimes e_{ij} \right) &= \sum_{i,j=1}^m \varphi(u_i x_{ij} u_j^*) \\ &= \sum_{i,j=1}^m q_i \varphi(x_{ij} u_j^* u_i) \\ &= \sum_{i=1}^m q_i \varphi(x_{ii}). \end{aligned}$$

Hence

$$\varphi \circ \Phi = \varphi \otimes \omega,$$

where ω is the state on $M_m(\mathbb{C})$ for which

$$\frac{d\omega}{d\text{Tr}} = \begin{pmatrix} q_1 & 0 & \cdots & 0 \\ 0 & q_2 & & \vdots \\ \vdots & & \ddots & 0 \\ 0 & \cdots & 0 & q_m \end{pmatrix}.$$

Put now $F := \Phi(\mathbb{C} \otimes M_m(\mathbb{C}))$. Then the relative commutant of F in N is $\Phi(N \otimes \mathbb{C})$. Since $\varphi \circ \Phi = \varphi \otimes \omega$, φ itself is a tensor product state with respect to the decomposition

$$N = F \cdot F^c \cong F \otimes F^c.$$

Moreover, $d\varphi|_F/d\text{Tr}_F$ has eigenvalues (q_1, \dots, q_m) . Let Φ_0 be the isomorphism of N onto F^c given by

$$\Phi_0(x) = \Phi(x \otimes 1), \quad x \in N.$$

Then $\varphi|_{F^c} \circ \Phi_0 = \varphi$. Therefore $\varphi|_{F^c}$ is a \mathbb{Q} -stable normal faithful state on F^c . \square

5 PROOF OF MAIN THEOREM

In this section we prove the main result of the paper:

THEOREM 5.1. *Every injective factor N of type III_1 on a separable Hilbert space is isomorphic to the Araki–Woods factor R_∞ .*

We need preparations. In this section, for each von Neumann algebra N , we fix a standard form $(N, H, J, \mathcal{P}^\natural)$. For each $\varphi \in (N_*)_+$, we denote by ξ_φ the unique representing vector in \mathcal{P}^\natural [Haa75].

LEMMA 5.2. *Let N be a properly infinite factor with separable predual and with a normal faithful state φ , let F be a finite dimensional σ^φ -invariant subfactor of N , and let $T: F \rightarrow N$ be a unital completely positive map, such that $\varphi \circ T = \varphi$ and*

$$\|\sigma_t^\varphi \circ T - T \circ \sigma_t^{\varphi|_F}\| \leq \delta|t|, \quad t \in \mathbb{R}, \quad (19)$$

where $\delta > 0$ is a constant. Then there exists a norm-continuous map $a: \mathbb{R} \rightarrow N$ such that

$$(a) \quad \int_{-\infty}^{\infty} a(t)^* a(t) dt = 1 \quad (\sigma\text{-strongly}),$$

$$(b) \quad \int_{-\infty}^{\infty} e^{-t} \varepsilon_{F, \varphi}(a(t)a(t)^*) dt = 1 \quad (\sigma\text{-strongly}),$$

$$(c) \quad \int_{-\infty}^{\infty} \|a(t)\xi_\varphi - e^{-t/2}\xi_\varphi a(t)\|^2 dt < \frac{\delta}{8},$$

$$(d) \quad \left\| T(x) - \int_{-\infty}^{\infty} a(t)^* x a(t) dt \right\| \leq \delta^{\frac{1}{2}} \|x\|, \quad x \in F,$$

where $\varepsilon_{F, \varphi}$ is the normal faithful conditional expectation of N onto F that leaves the state φ invariant.

Proof. Let f be the function

$$f(t) := (\pi\delta)^{-\frac{1}{4}} \exp\left(-\frac{1}{2\delta}t^2\right), \quad t \in \mathbb{R},$$

and let g be the Fourier-transformed of f :

$$g(s) := \left(\frac{\delta}{\pi}\right)^{\frac{1}{4}} \exp\left(-\frac{\delta}{2}s^2\right), \quad s \in \mathbb{R}.$$

Note that

$$\int_{-\infty}^{\infty} f(t)^2 dt = \int_{-\infty}^{\infty} g(s)^2 ds = 1.$$

By [Haa85, Proposition 2.1], there exists an operator $a \in N$ such that

$$T(x) = a^* x a, \quad x \in F.$$

In particular, $a^*a = 1$, i.e., a is an isometry. Put

$$a(t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-is(t-\delta/4)} g(s) \sigma_s^\varphi(a) ds, \quad t \in \mathbb{R}.$$

Since $t \mapsto e^{-is(t-\delta/4)} g(s)$ is a continuous map from \mathbb{R} to $L^1(\mathbb{R})$, the map $t \mapsto a(t)$ is a norm-continuous map from \mathbb{R} to N . Using the Plancherel formula in $L^2(\mathbb{R}, H)$, we get

$$\int_{-\infty}^{\infty} \|a(t)\xi\|^2 dt = \int_{-\infty}^{\infty} g(s)^2 \|\sigma_s^\varphi(a)\xi\|^2 ds = \|\xi\|^2$$

for all $\xi \in H$. Hence

$$\int_{-\infty}^{\infty} a(t)^* a(t) dt = 1 \quad (\sigma\text{-weakly}).$$

Since the convergence of the integral is monotone, we get (a). Using again the Plancherel formula, we get for $\xi, \eta \in H$ and $x \in F$,

$$\begin{aligned} \int_{-\infty}^{\infty} \langle xa(t)\xi, a(t)\eta \rangle dt &= \int_{-\infty}^{\infty} g(s)^2 \langle x\sigma_s^\varphi(a)\xi, \sigma_s^\varphi(a)\eta \rangle ds \\ &= \int_{-\infty}^{\infty} g(s)^2 \langle \sigma_s^\varphi \circ T \circ \sigma_{-s}^\varphi(x)\xi, \eta \rangle ds. \end{aligned}$$

Hence for $x \in F$,

$$\int_{-\infty}^{\infty} a(t)^* xa(t) dt = \int_{-\infty}^{\infty} g(s)^2 \sigma_s^\varphi \circ T \circ \sigma_{-s}^\varphi(x) ds. \quad (20)$$

Note that the left hand side of (20) converges σ -strongly, because $F = \text{span}(F_+)$ and for $x \in F_+$, the integral converges σ -weakly and the convergence is monotone. Therefore by (19), for each $x \in F$ we get

$$\begin{aligned} \left\| T(x) - \int_{-\infty}^{\infty} a(t)^* xa(t) dt \right\| &\leq \delta \|x\| \int_{-\infty}^{\infty} |s| g(s)^2 ds \\ &= \left(\frac{\delta}{\pi} \right)^{\frac{1}{2}} \|x\| \\ &\leq \delta^{\frac{1}{2}} \|x\|. \end{aligned}$$

This proves (d). Since $g(s)$ has the analytic extension to the function $g: \mathbb{C} \rightarrow \mathbb{C}$, and since the integrals

$$\int_{-\infty}^{\infty} |g(s + iu)| ds = \left(\frac{4\pi}{\delta} \right)^{\frac{1}{4}} e^{\frac{\delta}{2}u^2}, \quad u \in \mathbb{R}$$

are uniformly bounded for u on bounded subsets of \mathbb{R} , it follows that $a(t)$ is analytic with respect to σ^φ (in the sense of [PT73]) and that

$$\sigma_\alpha^\varphi(a(t)) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^\infty e^{-i(s-\alpha)(t-\frac{\delta}{4})} g(s-\alpha) \sigma_s^\varphi(a) ds,$$

for all $\alpha \in \mathbb{C}$. To prove (c), we use the equality

$$\xi_\varphi a(t) = J_\varphi a(t)^* \xi_\varphi = \Delta_{\frac{1}{\varphi}} a(t) \xi_\varphi = \sigma_{-i/2}^\varphi(a(t)) \xi_\varphi.$$

Hence

$$e^{-t/2} \xi_\varphi a(t) = \frac{e^{-\frac{\delta}{8}}}{\sqrt{2\pi}} \int_{-\infty}^\infty e^{-is(t-\frac{\delta}{4})} g(s+\frac{i}{2}) \sigma_s^\varphi(a) \xi_\varphi ds.$$

Using the Plancherel formula, we get

$$\int_{-\infty}^\infty \|a(t)\xi_\varphi - e^{-t/2}\xi_\varphi a(t)\|^2 dt = \int_{-\infty}^\infty |g(s) - e^{-\frac{\delta}{8}}g(s+\frac{i}{2})|^2 \|a\xi_\varphi\|^2 ds.$$

On the other hand, $g(s+\frac{i}{2})$ is the Fourier–Plancherel transformed of $e^{t/2}f(t)$. Therefore the above integral is equal to

$$\int_{-\infty}^\infty f(t)^2 \left(1 - e^{-\frac{\delta}{8}+\frac{t}{2}}\right)^2 dt.$$

It is easy to compute that for $\gamma \in \mathbb{R}$,

$$\int_{-\infty}^\infty f(t)^2 e^{\gamma t} dt = \exp(\frac{1}{4}\gamma^2\delta).$$

Therefore

$$\begin{aligned} \int_{-\infty}^\infty f(t)^2 \left(1 - e^{-\frac{\delta}{8}+\frac{t}{2}}\right)^2 dt &= 2(1 - e^{-\frac{\delta}{16}}) \\ &< \frac{\delta}{8}. \end{aligned}$$

This proves (c). Put now

$$A(t) := e^{-t}\varepsilon_{F,\varphi}(a(t)a(t)^*), \quad t \in \mathbb{R}.$$

Since $\varepsilon_{F,\varphi}$ is a normal faithful conditional expectation of N onto F that leaves φ invariant, we have for $x \in F$, that

$$\varphi(A(t)x) = e^{-t}\varphi(a(t)a(t)^*x).$$

By the KMS-condition, it follows that if $a, b \in N$ and a is σ^φ -analytic, then

$$\varphi(ab) = \varphi(b\sigma_{-i}^\varphi(a))$$

(cf. [Haa79, Theorem 3.2]). Hence for $x \in F$,

$$\varphi(A(t)x) = e^{-t}\varphi(a(t)^*x\sigma_{-i}^\varphi(a(t))).$$

Using

$$e^{-t}\sigma_{-i}^\varphi(a(t)) = \frac{e^{-\frac{\delta}{4}}}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-is(t-\frac{\delta}{4})} g(s+i)\sigma_s^\varphi(a) ds,$$

we get by the Plancherel formula, that

$$\begin{aligned} \int_{-\infty}^{\infty} \varphi(A(t)x) dt &= \int_{-\infty}^{\infty} \langle x(e^{-t}\sigma_{-i}^\varphi(a(t)))\xi_\varphi, a(t)\xi_\varphi \rangle dt \\ &= e^{-\frac{\delta}{4}} \int_{-\infty}^{\infty} g(s+i)\overline{g(s)} \langle x\sigma_s^\varphi(a)\xi_\varphi, \sigma_s^\varphi(a)\xi_\varphi \rangle ds. \end{aligned}$$

Since $\varphi \circ T = \varphi$, it holds that

$$\langle x\sigma_s^\varphi(a)\xi_\varphi, \sigma_s^\varphi(a)\xi_\varphi \rangle = \varphi \circ \sigma_s^\varphi \circ T \circ \sigma_{-s}^\varphi(x) = \varphi(x),$$

Hence

$$\int_{-\infty}^{\infty} \varphi(A(t)x) dt = e^{-\frac{\delta}{4}}\varphi(x) \int_{-\infty}^{\infty} g(s+i)\overline{g(s)} ds.$$

Since $g(s+i)$ is the Fourier–Plancherel transformed of $f(t)e^t$, we get

$$\int_{-\infty}^{\infty} g(s+i)\overline{g(s)} ds = \int_{-\infty}^{\infty} |f(t)|^2 e^t dt = e^{\frac{\delta}{4}}.$$

Since F is finite-dimensional and φ is faithful on F , every $\psi \in F_*$ is of the form $\varphi(\cdot x)$, $x \in F$. This shows that

$$\int_{-\infty}^{\infty} \psi(A(t)) dt = \psi(1), \quad \psi \in F_*,$$

that is, we have

$$\int_{-\infty}^{\infty} A(t) dt = 1 \quad (\sigma\text{-weakly}).$$

This proves (b). □

LEMMA 5.3. *Let N, φ, F and $\varepsilon_{F,\varphi}$ be as in Lemma 5.2. Let $\lambda > 0$ and assume that c_1, \dots, c_s are operators in $F^c = F' \cap N$ such that*

$$\begin{aligned} \varphi c_i &= \lambda c_i \varphi, \quad i = 1, \dots, s, \\ \sum_{i=1}^s c_i^* c_i &= 1. \end{aligned}$$

Then for all $x \in N$,

$$\varepsilon_{F,\varphi} \left(\sum_{i=1}^s c_i x c_i^* \right) = \lambda \varepsilon_{F,\varphi}(x).$$

Proof. It is sufficient to check the formula for $x \in N$ of the form $x = ab$, $a \in F$, $b \in F^c$. For $z \in F^c$, $\varepsilon_{F,\varphi}(z)$ commutes with every element in F . Hence $\varepsilon_{F,\varphi}(z)$ is a scalar multiple of the identity. Using that $\varepsilon_{F,\varphi}$ leaves φ invariant, we get

$$\varepsilon_{F,\varphi}(z) = \varphi(z)1, \quad z \in F^c.$$

Therefore

$$\begin{aligned} \varepsilon_{F,\varphi} \left(\sum_{i=1}^s c_i x c_i^* \right) &= \varepsilon_{F,\varphi} \left(a \left(\sum_{i=1}^s c_i b c_i^* \right) \right) \\ &= \varphi \left(\sum_{i=1}^s c_i b c_i^* \right) a \\ &= \lambda \varphi \left(\sum_{i=1}^s b c_i^* c_i \right) a \\ &= \lambda \varphi(b) a \\ &= \lambda \varepsilon_{F,\varphi}(x). \end{aligned}$$

□

LEMMA 5.4. *Let φ be a \mathbb{Q} -stable normal faithful state on an injective factor N of type III_1 with separable predual. Let $u_1, \dots, u_n \in \mathcal{U}(N)$, let $\delta > 0$. Then there exist a finite dimensional σ^φ -invariant subfactor F of N and unitaries $v_1, \dots, v_n \in \mathcal{U}(F)$, such that for every σ -strong neighborhood \mathcal{V} of 0 in N , there exists a finite set b_1, \dots, b_r of operators in N with the following properties:*

- (a) $\sum_{i=1}^r b_i^* b_i \in 1 + \mathcal{V}$ and $\sum_{i=1}^r b_i^* b_i \leq 1$.
- (b) $\varepsilon_{F,\varphi} \left(\sum_{i=1}^r b_i b_i^* \right) \in 1 + \mathcal{V}$ and $\varepsilon_{F,\varphi} \left(\sum_{i=1}^r b_i b_i^* \right) \leq 1$.
- (c) $\sum_{i=1}^r \|b_i \xi_\varphi - \xi_\varphi b_i\|^2 < \delta$.
- (d) $\sum_{i=1}^r \|b_i u_k - v_k b_i\|_\varphi^2 < \delta$, $k = 1, \dots, n$.

Proof. Put $\delta_1 = \min(\delta^2/16, \delta)$. By Theorem 3.1, there exist $m \in \mathbb{N}$, a unital completely positive map $T_0: M_m(\mathbb{C}) \rightarrow N$ and unitaries $w_1, \dots, w_n \in M_m(\mathbb{C})$ such that $\psi := \varphi \circ T_0 \in M_m(\mathbb{C})_*$ satisfies

$$\begin{aligned} \|\sigma_t^\varphi \circ T_0 - T_0 \circ \sigma_t^\psi\| &\leq \frac{\delta_1}{2}|t|, \quad t \in \mathbb{R}, \\ \|T_0(w_k) - u_k\|_\varphi &< \varepsilon, \quad k = 1, \dots, n. \end{aligned}$$

Let $\{q'_1, \dots, q'_m\}$ be the spectrum of $d\psi/d\text{Tr} \in M_m(\mathbb{C})_+$ where the multiplicity is taken into account. Let $\{q_1, \dots, q_m\}$ be positive rationals with sum 1, and let χ on $M_m(\mathbb{C})_+$ such that $d\chi/d\text{Tr}$ has the same spectral projections as $d\psi/d\text{Tr}$ but the eigenvalues replaced by $\{q_1, \dots, q_m\}$. Since $\|e^{ia} - e^{ib}\| \leq \|a - b\|$ for self-adjoint operators a, b , (cf. (17) in Theorem 3.1), we may arrange q_i 's so that the following inequality holds:

$$\|\sigma_t^\psi - \sigma_t^\chi\|_{M_m(\mathbb{C})} \leq \frac{\delta_1}{2}|t|, \quad t \in \mathbb{R}.$$

Since φ is \mathbb{Q} -stable and q_i 's are rationals, by Theorem 4.5, there exists a finite-dimensional subfactor $F \subset N$ and a state-preserving $*$ -isomorphism $\Phi: (M_m(\mathbb{C}), \chi) \rightarrow (F, \varphi|_F)$ such that $\varphi|_{F^c}$ is \mathbb{Q} -stable. Define $T := T_0 \circ \Phi^{-1}: F \rightarrow N$ and $v_k := \Phi_0(w_k) \in \mathcal{U}(F) (1 \leq k \leq n)$. Then if $x = \Phi(y)$ ($y \in M_m(\mathbb{C})$) and $t \in \mathbb{R}$, we have

$$\begin{aligned} \|\sigma_t^\varphi \circ T(x) - T \circ \sigma_t^{\varphi|_F}(x)\| &= \|\sigma_t^\varphi \circ T_0(y) - T_0 \circ \Phi^{-1} \circ \sigma_t^{\varphi|_F} \circ \Phi(y)\| \\ &= \|\sigma_t^\varphi \circ T_0(y) - T_0 \circ \sigma_t^\chi(y)\| \\ &\leq \|\sigma_t^\varphi \circ T_0(y) - T_0 \circ \sigma_t^\psi(y)\| + \|T(\sigma_t^\psi(y) - \sigma_t^\chi(y))\| \\ &\leq \delta_1|t|\|y\|. \end{aligned}$$

Therefore we obtain

$$\begin{aligned} T(1) &= 1, \quad \varphi \circ T = \varphi|_F, \\ \|\sigma_t^\varphi \circ T - T \circ \sigma_t^{\varphi|_F}\| &\leq \delta_1|t|, \quad t \in \mathbb{R}, \\ \|T(v_k) - u_k\|_\varphi &< \delta_1^{\frac{1}{2}}, \quad k = 1, \dots, n. \end{aligned}$$

Choose now a norm-continuous function $t \mapsto a(t)$ of \mathbb{R} into N , such that the conditions (a), (b), (c) and (d) in Lemma 5.2 are satisfied with respect to δ_1 instead of δ . Then using (d), we have

$$\|u_k - \int_{-\infty}^{\infty} a(t)^* v_k a(t) dt\|_\varphi < 2\delta_1^{\frac{1}{2}} \leq \frac{\delta}{2}$$

for $k = 1, \dots, n$. Using that $\int_{-\infty}^{\infty} a(t)^* a(t) dt = 1$, it follows that

$$\begin{aligned} \int_{-\infty}^{\infty} \|a(t)u_k - v_k a(t)\|_\varphi^2 dt &= 2 - 2\text{Re} \int_{-\infty}^{\infty} \langle a(t)u_k \xi_\varphi, v_k a(t) \xi_\varphi \rangle dt \\ &= 2 - 2\text{Re} \left(\langle u_k \xi_\varphi, \int_{-\infty}^{\infty} a(t)^* v_k a(t) \xi_\varphi dt \rangle \right) \\ &\leq 2 \|u_k - \int_{-\infty}^{\infty} a(t)^* v_k a(t) dt\|_\varphi \\ &< \delta. \end{aligned}$$

Let now \mathcal{V} be a σ -strong neighborhood of 0 in N . It is no loss of generality to assume that \mathcal{V} is open. For sufficiently large $\gamma \in \mathbb{R}_+$, we have:

- (a') $\int_{-\gamma}^{\gamma} a(t)^* a(t) dt \in 1 + \mathcal{V}$ and $\int_{-\gamma}^{\gamma} a(t)^* a(t) dt \leq 1$.
- (b') $\int_{-\gamma}^{\gamma} e^{-t} \varepsilon_{F,\varphi}(a(t)a(t)^*) dt \in 1 + \mathcal{V}$ and $\int_{-\gamma}^{\gamma} e^{-t} \varepsilon_{F,\varphi}(a(t)a(t)^*) dt \leq 1$.
- (c') $\int_{-\gamma}^{\gamma} \|a(t)\xi_{\varphi} - e^{-t/2}\xi_{\varphi}a(t)\|^2 dt < \frac{\delta_1}{8} \leq \delta$.
- (d') $\int_{-\gamma}^{\gamma} \|a(t)u_k - v_k a(t)\|_{\varphi}^2 dt < \delta$.

Since $t \mapsto a(t)$ is norm-continuous, we can approximate (in norm) the above N -valued Riemann integrals over $[-\gamma, \gamma]$ to get the following statements: there exists an $h_0 > 0$ such that when $0 < h < h_0$, the operators

$$a_j = h^{-\frac{1}{2}} a(jh), \quad j \in \mathbb{Z}$$

satisfy the following relations:

- (a'') $\sum_{j=-p}^p a_j^* a_j \in 1 + \mathcal{V}$.
- (b'') $\sum_{j=-p}^p e^{-jh} \varepsilon_{F,\varphi}(a_j a_j^*) \in 1 + \mathcal{V}$.
- (c'') $\sum_{j=-p}^p \|a_j \xi_{\varphi} - e^{-\frac{1}{2}jh} \xi_{\varphi} a_j\|^2 < \delta$.
- (d'') $\sum_{j=-p}^p \|a_j u_k - v_k a_j\|_{\varphi}^2 dt < \delta$,

where p is the largest integer smaller than γ/h_0 . Moreover, since the Riemann sum is norm-convergent, by multiplying a scalar $c > 0$ to a_j 's which is sufficiently close to 1 if necessary, we may moreover assume that

$$\sum_{j=-p}^p a_j^* a_j \leq 1 \tag{21}$$

$$\sum_{j=-p}^p e^{-jh} \varepsilon_{F,\varphi}(a_j a_j^*) \leq 1. \tag{22}$$

Choose now $h \in (0, h_0)$, such that $\exp(h) \in \mathbb{Q}$. This implies that the numbers $q_j = e^{-jh}$, $j \in \mathbb{Z}$ are rational. Since the restriction of φ to F^c is \mathbb{Q} -stable, there exists for each $j \in \mathbb{Z}$ a finite set of operators $c_{j1}, \dots, c_{js(j)}$ in F^c such that

$$\varphi c_{ji} = e^{-jh} c_{ji} \varphi, \quad i = 1, \dots, s(j)$$

and

$$\sum_{i=1}^{s(j)} c_{ji}^* c_{ji} = 1.$$

Here we use Lemma 4.6 together with the fact that $\varphi = \varphi|_F \otimes \varphi|_{F^c}$. Put

$$b_{ji} = c_{ji} a_j, \quad |j| \leq p, \quad 1 \leq i \leq s(j).$$

Then by (21),

$$(a''') \quad \sum_{j=-p}^p \sum_{i=1}^{s(j)} b_{ji}^* b_{ji} = \sum_{j=-p}^p a_j^* a_j \in 1 + \mathcal{V} \quad \text{and} \quad \sum_{j=-p}^p \sum_{i=1}^{s(j)} b_{ji}^* b_{ji} \leq 1,$$

and by (22) and Lemma 5.3,

$$(b''') \quad \varepsilon_{F,\varphi} \left(\sum_{j=-p}^p \sum_{i=1}^{s(j)} b_{ji} b_{ji}^* \right) = \sum_{j=-p}^p e^{-jh} \varepsilon_{F,\varphi}(a_j a_j^*) \in 1 + \mathcal{V},$$

and $\varepsilon_{F,\varphi} \left(\sum_{j=-p}^p \sum_{i=1}^{s(j)} b_{ji} b_{ji}^* \right) \leq 1.$

The equality $\varphi c_{ji} = e^{-jh} c_{ji} \varphi$ implies that

$$\xi_\varphi c_{ji} = e^{-\frac{1}{2}jh} c_{ji} \xi_\varphi.$$

Therefore

$$(c''') \quad \sum_{j=-p}^p \sum_{i=1}^{s(j)} \|b_{ji} \xi_\varphi - \xi_\varphi b_{ji}\|^2 = \sum_{j=-p}^p \sum_{i=1}^{s(j)} \|c_{ji}(a_j \xi_\varphi - e^{-\frac{1}{2}jh} \xi_\varphi a_j)\|^2$$

$$= \sum_{j=-p}^p \|a_j \xi_\varphi - e^{-\frac{1}{2}jh} \xi_\varphi a_j\|^2$$

$$< \varepsilon.$$

Finally, using that $v_k \in F$ and $c_{ji} \in F^c$, we get

$$(d''') \quad \sum_{j=-p}^p \sum_{i=1}^{s(j)} \|b_{ji} u_k - v_k b_{ji}\|_\varphi^2 = \sum_{j=-p}^p \sum_{i=1}^{s(j)} \|c_{ji}(a_j u_k - v_k a_j)\|_\varphi^2$$

$$= \sum_{j=-p}^p \|a_j u_k - v_k a_j\|_\varphi^2$$

$$< \delta.$$

This completes the proof of Lemma 5.4. □

In the proof of the following lemma, it is essential that injective type III₁ factors (on a separable Hilbert space) have trivial bicentralizers.

LEMMA 5.5. *Let φ be a \mathbb{Q} -stable normal faithful state on an injective factor N of type III₁ with separable predual. Let $u_1, \dots, u_n \in \mathcal{U}(N)$, and let $\delta > 0$. Then there exists a finite dimensional σ^φ -invariant subfactor F of N and $v_1, \dots, v_n \in \mathcal{U}(F)$, such that for every σ -strong neighborhood \mathcal{V} of 0 in N , there exists a finite set a_1, \dots, a_p of operators in N with the following properties:*

- (a) $\sum_{i=1}^p a_i^* a_i \in 1 + \mathcal{V}$ and $\sum_{i=1}^p a_i^* a_i \leq 1$.
- (b) $\sum_{i=1}^p a_i a_i^* \in 1 + \mathcal{V}$ and $\sum_{i=1}^p a_i a_i^* \leq 1$.
- (c) $\sum_{i=1}^p \|a_i \xi_\varphi - \xi_\varphi a_i\|^2 < \delta$.
- (d) $\sum_{i=1}^p \|a_i u_k - v_k a_i\|_\varphi^2 < \delta, \quad k = 1, \dots, n$.

Proof. Choose an F and $v_1, \dots, v_n \in \mathcal{U}(F)$ satisfying the properties of Lemma 5.4 with respect to $(u_1, \dots, u_n, \delta)$, and let \mathcal{V} be a σ -strongly open neighborhood of 0 in N . By Lemma 5.4, there exists $b_1, \dots, b_r \in N$ such that

- (a') $\sum_{i=1}^r b_i^* b_i \in 1 + \mathcal{V}$ and $\sum_{i=1}^r b_i^* b_i \leq 1$.
- (b') $\varepsilon_{F, \varphi} \left(\sum_{i=1}^r b_i b_i^* \right) \in 1 + \mathcal{V}$ and $\varepsilon_{F, \varphi} \left(\sum_{i=1}^r b_i b_i^* \right) \leq 1$.
- (c') $\sum_{i=1}^r \|b_i \xi_\varphi - \xi_\varphi b_i\|^2 < \delta$.
- (d') $\sum_{i=1}^r \|b_i u_k - v_k b_i\|_\varphi^2 < \delta, \quad k = 1, \dots, n$.

Let $\delta' > 0$ and h denote the operator $\sum_{i=1}^r b_i b_i^*$. Since $B_\varphi = \mathbb{C}1$, by Proposition 2.6, we have

$$\varepsilon_{F, \varphi}(h) \in \overline{\text{conv}}\{whw^*; w \in \mathcal{U}(F^c), \|w\xi_\varphi - \xi_\varphi w\| < \delta'\}. \quad (23)$$

Here, $\overline{\text{conv}}$ in (23) denotes the σ -strong closure. Hence there exist $w_1, \dots, w_s \in \mathcal{U}(F^c)$, and scalars $\lambda_1, \dots, \lambda_s \in \mathbb{R}_+$, with sum 1, such that

$$\|w_j \xi_\varphi - \xi_\varphi w_j\| < \delta', \quad j = 1, \dots, s$$

and

$$\sum_{j=1}^s \lambda_j w_j h w_j^* \in 1 + \mathcal{V}.$$

Put

$$a_{ij} := \lambda_j^{\frac{1}{2}} w_j b_i, \quad i = 1, \dots, r, \quad j = 1, \dots, s.$$

Then

$$(a'') \quad \sum_{i=1}^r \sum_{j=1}^s a_{ij}^* a_{ij} = \sum_{i=1}^r b_i^* b_i \in 1 + \mathcal{V} \quad \text{and} \quad \sum_{i=1}^r \sum_{j=1}^s a_{ij}^* a_{ij} \leq 1.$$

$$(b'') \quad \sum_{i=1}^r \sum_{j=1}^s a_{ij} a_{ij}^* = \sum_{j=1}^s \lambda_j w_j h w_j^* \in 1 + \mathcal{V} \quad \text{and} \quad \sum_{i=1}^r \sum_{j=1}^s a_{ij} a_{ij}^* \leq 1.$$

Moreover, using

$$a_{ij} \xi_\varphi - \xi_\varphi a_{ij} = \lambda_j^{\frac{1}{2}} w_j (b_i \xi_\varphi - \xi_\varphi b_i) + \lambda_j^{\frac{1}{2}} (w_j \xi_\varphi - \xi_\varphi w_j) b_i,$$

we obtain

$$\begin{aligned} (c'') \quad & \left(\sum_{i=1}^r \sum_{j=1}^s \|a_{ij} \xi_\varphi - \xi_\varphi a_{ij}\|^2 \right)^{\frac{1}{2}} \leq \\ & \leq \left(\sum_{i=1}^r \sum_{j=1}^s \lambda_j \|b_i \xi_\varphi - \xi_\varphi b_i\|^2 \right)^{\frac{1}{2}} + \delta' \left(\sum_{i=1}^r \sum_{j=1}^s \lambda_j \|b_i\|^2 \right)^{\frac{1}{2}} \\ & = \left(\sum_{i=1}^r \|b_i \xi_\varphi - \xi_\varphi b_i\|^2 \right)^{\frac{1}{2}} + \delta' \left(\sum_{i=1}^r \|b_i\|^2 \right)^{\frac{1}{2}}. \end{aligned}$$

Finally, since $v_k \in F$ and $w_j \in F^c$, we have

$$\begin{aligned} (d'') \quad & \sum_{i=1}^r \sum_{j=1}^s \|a_{ij} u_k - v_k a_{ij}\|_\varphi^2 = \sum_{i=1}^r \sum_{j=1}^s \lambda_j \|w_j (b_i u_k - v_k b_i)\|_\varphi^2 \\ & = \sum_{i=1}^r \|b_i u_k - v_k b_i\|_\varphi^2 \\ & < \delta. \end{aligned}$$

Since $\delta' > 0$ was arbitrary (independent of δ, \mathcal{V} , and b_1, \dots, b_r), we can assume that

$$\left(\sum_{i=1}^r \|b_i \xi_\varphi - \xi_\varphi b_i\|^2 \right)^{\frac{1}{2}} + \delta' \left(\sum_{i=1}^r \|b_i\|^2 \right)^{\frac{1}{2}} < \delta^{\frac{1}{2}}.$$

This proves Lemma 5.5. □

LEMMA 5.6. *Let N be an injective factor of type III_1 with separable predual, and let φ be a \mathbb{Q} -stable normal faithful state on N . Let $u_1, \dots, u_n \in \mathcal{U}(N)$ and let $\varepsilon > 0$. Then there exist a σ^φ -invariant finite dimensional subfactor F of N , $v_1, \dots, v_n \in \mathcal{U}(F)$ and a unitary $w \in \mathcal{U}(N)$ such that*

$$\|w\xi_\varphi - \xi_\varphi w\| < \varepsilon,$$

and

$$\|w^*v_k w - u_k\|_\varphi < \varepsilon, \quad k = 1, \dots, n.$$

Proof. Let $\delta(n, \varepsilon) > 0$ be the function in Theorem 2.8, and put $\delta_1 = \frac{1}{16}\delta(n + 1, \varepsilon/2)$. Choose F and $v_1, \dots, v_n \in \mathcal{U}(F)$, such that the conditions of Lemma 5.5 are satisfied with respect to $(u_1, \dots, u_n, \delta_1)$. Put

$$\xi_k = u_k \xi_\varphi, \quad \eta_k = v_k \xi_\varphi, \quad k = 1, \dots, n.$$

For every σ -strong neighborhood \mathcal{V} of 0 in N , there exist $a_1, \dots, a_p \in N$, such that (a), (b), (c) and (d) in Lemma 5.5 are satisfied. Since

$$a_i \xi_k - \eta_k a_i = (a_i u_k - v_k a_i) \xi_\varphi + v_k (a_i \xi_\varphi - \xi_\varphi a_i),$$

we have

$$\begin{aligned} \left(\sum_{i=1}^p \|a_i \xi_k - \eta_k a_i\|^2 \right)^{\frac{1}{2}} &\leq \left(\sum_{i=1}^p \|a_i u_k - v_k a_i\|_\varphi^2 \right)^{\frac{1}{2}} + \left(\sum_{i=1}^p \|a_i \xi_\varphi - \xi_\varphi a_i\|^2 \right)^{\frac{1}{2}} \\ &< 2\delta_1^{\frac{1}{2}}. \end{aligned}$$

Moreover,

$$\left(\sum_{i=1}^p \|a_i \xi_\varphi - \xi_\varphi a_i\|^2 \right)^{\frac{1}{2}} < \delta_1^{\frac{1}{2}} < 2\delta_1^{\frac{1}{2}}.$$

Since $\sum_{i=1}^p a_i^* a_i \in 1 + \mathcal{V}$, $\sum_{i=1}^p a_i^* a_i \leq 1$, $\sum_{i=1}^p a_i a_i^* \in 1 + \mathcal{V}$ and $\sum_{i=1}^p a_i a_i^* \leq 1$, the two $(n+1)$ -tuples $(\xi_1, \dots, \xi_n, \xi_\varphi)$ and $(\eta_1, \dots, \eta_n, \xi_\varphi)$ satisfies the conditions of Remark 2.9 with $4\delta_1^{\frac{1}{2}}$ instead of δ , so that the two $(n+1)$ -tuples are $16\delta_1$ -related, or equivalently they are $\delta(n+1, \frac{\varepsilon}{2})$ -related in the sense of Remark 2.9. Hence by Theorem 2.8, there exists a unitary operator $w \in \mathcal{U}(N)$, such that

$$\|w\xi_k - \eta_k w\| < \frac{\varepsilon}{2}, \quad k = 1, \dots, n.$$

and

$$\|w\xi_\varphi - \xi_\varphi w\| < \frac{\varepsilon}{2}.$$

Therefore

$$\begin{aligned} \|w^*v_k w - u_k\|_\varphi &= \|w^*(v_k w - w u_k)\xi_\varphi\| \\ &= \|(w u_k - v_k w)\xi_\varphi\| \\ &= \|(w\xi_k - \eta_k w) + v_k(\xi_\varphi w - w\xi_\varphi)\| \\ &< \varepsilon, \end{aligned}$$

which completes the proof of Lemma 5.6. □

Now we are ready to prove the main theorem of the paper.

Proof of Theorem 5.1. By [AW68, Theorem 7.6], it is sufficient to show that N is an ITPFI-factor. Let φ be a \mathbb{Q} -stable normal faithful state on N , let $u_1, \dots, u_n \in \mathcal{U}(N)$, and let $\varepsilon > 0$. Choose now F , $v_1, \dots, v_n \in \mathcal{U}(F)$ and $w \in \mathcal{U}(N)$ as in Lemma 5.6. Put

$$w_k = w^* v_k w, \quad k = 1, \dots, n.$$

Then $F_1 := w^* F w$ is a finite-dimensional subfactor of N , $w_1, \dots, w_n \in \mathcal{U}(F_1)$ and

$$\|w_k - u_k\|_\varphi < \varepsilon, \quad k = 1, \dots, n.$$

Hence if we put $\varphi_1 = w^* \varphi w$, then by $\varphi = \varphi|_F \otimes \varphi|_{F^c}$, we have

$$\varphi_1 = \varphi_1|_{F_1} \otimes \varphi_1|_{F_1^c}.$$

Since the representing vector of φ_1 in \mathcal{P}_N^{\natural} is $w^* \xi_\varphi w$, we have

$$\begin{aligned} \|\varphi - \varphi_1\| &\leq \|\xi_\varphi - w^* \xi_\varphi w\| \|\xi_\varphi + w^* \xi_\varphi w\| \\ &\leq 2 \|w \xi_\varphi - \xi_\varphi w\| \\ &< 2\varepsilon. \end{aligned}$$

This shows that φ satisfies the product condition in Proposition 2.2, and thus N is an ITPFI factor. \square

REFERENCES

- [Ara74] H. Araki, Some properties of modular conjugation operator of von Neumann algebras and non-commutative Radon-Nikodym theorem with a chain rule, *Pac. J. Math.* 50 (1974), 309–354.
- [AW68] H. Araki and E. J. Woods, A classification of factors, *Publ. Res. Inst. Math. Sci. Ser. A.* 4 (1968), 51–130.
- [Con72] A. Connes, Une classification des facteurs de type III (French), *C. R. Acad. Sci. Paris Ser. A-B* 275 (1972), 523–525.
- [Con73] A. Connes, Une classification des facteurs de type III (French), *Ann. Sci. Ec. Norm. Sup.* 6 (1973), 133–252.
- [Con74] A. Connes, Almost periodic states and factors of type III₁, *J. Funct. Anal.* 16 (1974), 415–445.
- [Con76] A. Connes, Classification of injective factors, *Ann. Math.* 104 (1976), 73–115.
- [Con85] A. Connes, Factors of type III₁, property L'λ, and closure of inner automorphisms. *J. Operator Theory* 14 (1985), 189–211.
- [CS78] A. Connes, E. Størmer, Homogeneity of the state space of factors of type III₁, *J. Funct. Anal.* 28 (1978), 187–196.

- [CT77] A. Connes and M. Takesaki, The flow of weights on factors of type III, *Tohoku Math. J.* 29 (1977), 473–575.
- [CW85] A. Connes and E. J. Woods, Approximately transitive flows and ITPFI factors, *Ergod. Th. Dynam. Sys.* 5 (1985), 203–236.
- [Haa75] U. Haagerup, The standard form of von Neumann algebras, *Math. Scand.* 37 (1975), 271–283.
- [Haa79] U. Haagerup, Operator-valued weights in von Neumann algebras. I, *J. Funct. Anal.* 32 (1979), 175–206.
- [Haa79-2] U. Haagerup, Operator-valued weights in von Neumann algebras. II, *J. Funct. Anal.* 33 (1979), 339–361.
- [Haa85] U. Haagerup, A new proof of the equivalence of injectivity and hyperfiniteness for factors on a separable Hilbert space, *J. Funct. Anal.* 62 (1985), 160–201.
- [Haa87] U. Haagerup, Connes’ bicentralizer problem and uniqueness of the injective factor of type III₁. *Acta Math.* 158 (1987), 95–148.
- [Haa89] U. Haagerup, The injective factors of type III_λ, 0 < λ < 1, *Pacific. J. Math.* 137 (1989), 265–310.
- [HM09] U. Haagerup and M. Musat, Classification of hyperfinite factors up to completely bounded isomorphism of their preduals, *J. reine angew. Math.* 630 (2009), 141–176.
- [Kat68] Y. Katznelson, An introduction to harmonic analysis, John Wiley & sons (1968).
- [PT73] G. K. Pedersen, M. Takesaki, The Radon-Nikodym theorem for von Neumann algebras, *Acta Math.* 130 (1973), 53–87.
- [Tak72] M. Takesaki, Conditional expectations in von Neumann algebras, *J. Funct. Anal.* 9 (1972), 306–321.
- [Tak73] M. Takesaki, The structure of von Neumann algebras with a homogeneous periodic state, *Acta. Math.* 131 (1973), 79–121.
- [Tak73-2] M. Takesaki, Duality for crossed products and the structure of von Neumann algebras of type III, *Acta. Math.* 131, (1973), 249–310.
- [Tom58] J. Tomiyama, On the projection of norm one in W*-algebras, III, *Tohoku Math. J.*, 10 (1958), 204–209.

Uffe Haagerup
Department of Mathematics
and Computer Science
The University of Southern
Denmark
Campusvej 55, DK-5230
Odense M, Denmark