

**NASA TECHNICAL NOTE**



**NASA TN D-2460**  
C.1

**NASA TN D-2460**

LOAN COPY: RETU  
AFWL (WLIL-2  
KIRTLAND AFB, N

0079546



TECH LIBRARY KAFB, NM

# LOW-COVERAGE HEATS OF ADSORPTION

III — ALKALI METAL IONS ON TUNGSTEN;  
ATOM-METAL INTERACTION THEORY

*by Harold E. Neustadter and Keung P. Luke*

*Lewis Research Center  
Cleveland, Ohio*



LOW-COVERAGE HEATS OF ADSORPTION

III - ALKALI METAL IONS ON TUNGSTEN;  
ATOM-METAL INTERACTION THEORY

By Harold E. Neustadter and Keung P. Luke

Lewis Research Center  
Cleveland, Ohio

NATIONAL AERONAUTICS AND SPACE ADMINISTRATION

---

For sale by the Office of Technical Services, Department of Commerce,  
Washington, D.C. 20230 -- Price \$0.50

# LOW-COVERAGE HEATS OF ADSORPTION

## III - ALKALI METAL IONS ON TUNGSTEN;

### ATOM-METAL INTERACTION THEORY

by Harold E. Neustadter and Keung P. Luke

Lewis Research Center

#### SUMMARY

The heat of adsorption of an ion on a metal (in the limit of zero coverage) is expressed as the superposition of a van der Waals type interaction and an image charge type interaction. The reported experimental values of the heat of adsorption of sodium, rubidium, and cesium ions on a tungsten substrate are then compared with results of a theoretical evaluation based on the above model.

#### INTRODUCTION

Considerable interest in alkali metal plasma for power generation and space propulsion has prompted extensive studies of adsorption of alkali metals on various substrates. A requirement for an adequate adsorption theory is the

specification of the heat of adsorption in the limit of zero coverage. This report proposes an equation for calculating the heat of adsorption  $\phi$  of an ion on a metal at coverages sufficiently low to allow neglect of interactions between neighboring adsorbed ions. Results of calculations based on this equation are compared with experimental values for sodium, rubidium, and cesium ions adsorbed on a tungsten substrate (table I). Note that attractive energies are denoted as positive.

TABLE I. - EXPERIMENTAL ZERO-COVERAGE

#### HEATS OF ADSORPTION

Source	Sodium	Rubidium	Cesium
	Heat of adsorption, $\phi$ , ev		
Bosworth (ref. 1)	2.50		
Hughes and Levenstein (ref. 2)		1.90	
de Boer and Veenemans (ref. 3)			<sup>a</sup> 2.36
Taylor and Langmuir (ref. 4)			<sup>a</sup> 2.04
Becker (ref. 5)			2.19
Scheer and Fine (ref. 6)			2.04

<sup>a</sup>Not measured directly, but calculated from the assumption that all cesium adsorbed on tungsten exists only as ions.

#### SYMBOLS

$a_0$  Bohr radius

$e$  electron charge

- $k_m$  defined by eq. (5)  
 $n_o$  free electron density of metal (tungsten)  
 $R$  distance from metal surface to center of ion  
 $\alpha$  electronic polarizability  
 $\beta$  constant,  $\approx 0.33$   
 $\kappa$  defined by eq. (3)  
 $\phi$  heat of adsorption of ion in limit of zero coverage  
 $\phi_i$  defined by eq. (2)  
 $\phi_w$  defined by eq. (4)

#### ANALYSIS

A model commonly used to explain the adsorptive bond approximates the adsorbed ion as a point charge and utilizes classical image-charge theory. This is an oversimplification in that (1) it neglects the contribution of the electronic distribution of the ion and (2) classical image-charge theory breaks down in the immediate region of the metal surface.

The adsorption energies can be treated as arising from the superposition of (1) the interaction  $\phi_i$  of the net ionic charge with the metal and (2) a van der Waals type interaction  $\phi_w$  of the instantaneous dipole moment of the ion with the metal. Thus,

$$\phi = \phi_i + \phi_w \quad (1)$$

For  $\phi_i$  a quantum mechanically corrected image-charge expression must be used, since the ions are 1/2 to 2 angstroms from the metal surface. Sachs and Dexter (ref. 7) have proposed such an expression:

$$\phi_i = \frac{e^2}{4R} - \beta \kappa^{1/2} \frac{e^2}{4R^2} \quad (2)$$

where  $R$  is the distance from the surface to the center of the ion,  $\beta$  is a constant, and

$$\kappa = -\frac{1}{3} \left( \frac{3}{8\pi} \right)^{1/3} n_o^{-2/3} + \frac{(2\pi)^2}{6} \left( \frac{3}{8\pi} \right)^{2/3} a_o n_o^{-1/3} \quad (3)$$

in which  $n_o$  is the free electron density of the metal, and  $a_o$  is the Bohr radius.

The energy arising from the polarization of an adsorbed particle has been studied by Prosen and Sachs (ref. 8) and is given by<sup>1</sup>

$$\phi_w = \frac{\alpha e^2 \pi k_m}{(2\pi)^3} \frac{\ln(2k_m R)}{R^2} \quad (4)$$

where  $\alpha$  is the electronic dipole polarizability of the adsorbed ion (table II), and

$$k_m = (3\pi^2 n_0)^{1/3} \quad (5)$$

TABLE II. - ELECTRONIC POLARIZABILITY

Source	Sodium ion	Rubidium ion	Cesium ion
	Electronic polarizability, $\alpha$ , $\text{\AA}^3$		
Sternheimer (ref. 10)	<sup>a</sup> 0.152	<sup>a</sup> 2.920	<sup>a</sup> 5.60
Fajan and Joos (ref. 11)	.196	1.56	2.56
Born and Heisenberg (ref. 12)	.210	1.81	2.79
Pauling (ref. 13)	.189	1.41	2.44
Tessman, Kahn, and Shockley (ref. 14)	.255	1.797	3.137

<sup>a</sup>Calculated.

The complete expression for the heat of adsorption of ions on a metal in the limit of zero coverage is obtained by adding equations (2) and (4):

$$\phi = \frac{e^2}{4R} - \beta \kappa^{1/2} \frac{e^2}{4R^2} + \frac{\alpha e^2 \pi k_m^2}{(2\pi)^3} \frac{\ln(2k_m R)}{R^2} \quad (6)$$

Implicit in equation (6) and the choice of ionic values of polarizability is the assumption that a particle that desorbs as an ion has the properties of an ion prior to desorption. Hughes (ref. 15) in his adsorption-lifetime studies has found evidence to support this.

To apply equation (6), it is necessary to evaluate the constant  $\beta$  and  $R$ , the distance from the nucleus of the ion to the metal surface. The location of this "surface" is not well defined, however. Physically it is that plane on which all excess electrons would collect. The constant  $\beta$  depends on the choice of wave function used in deriving equation (2). Sachs and Dexter estimated  $\beta$  to be 0.46; however, their approximations appear to make this an upper limit. That this value of  $\beta$  is too high has been substantiated by others in two instances (ref. 16). Since  $\beta$  is not uniquely determined, this report evaluates it by matching the calculated value of  $\phi$  for sodium adsorption to the reported experimental value. In the course of evaluating  $\beta$ ,  $R$  will also be determined. The value of  $\phi$  for sodium was selected because the electronic polarizability of the sodium ion is very small (table II). Thus,

<sup>1</sup>The choice of equation (4) to express the van der Waals type interaction is based on the results of an earlier study by the authors of this report (ref. 9).

the contribution of  $\phi_w$  to the heat of adsorption of sodium ions on tungsten is small ( $\phi_w \cong 0.04$  ev) and one can approximate

$$\phi(\text{Na}^+ - \text{W}) = \phi_i = 2.46 \text{ ev} \quad (7)$$

At equilibrium the sodium ion will be at that  $R$  value for which  $\phi_i$  is a maximum. Taking

$$d\phi_i/dR = 0 \quad (8)$$

gives the result

$$R = 2\beta\kappa^{1/2} \quad (9)$$

For tungsten (lattice constant of 3.16 Å, one conduction electron per atom)  $\kappa^{1/2} = 1.1$ , and from equation (9)  $R = 2.2\beta$ . The constant  $\beta$  can now be evaluated by using this result in equations (7) and (2), the final results<sup>2</sup> being  $\beta \cong 0.33$  and  $R = 0.73$  Å.

The use of a hard sphere model, the assumption of adsorption taking place on the (110) plane of tungsten, and the value  $R = 0.73$  Å, locate the metal surface at 0.93 Å from the plane through the centers of the outermost layer of




Figure 1. - Sodium ion adsorbed on a (110) plane of tungsten. (Dimensions in angstroms.)

atoms (see fig. 1). This is 71 percent of the distance from the center of the surface layer of atoms to the outermost tangent of the surface layer of atoms, which is a physically reasonable result.

Sufficient evidence justifies the assumption that for tungsten emitters the emitting surfaces were essentially (110) planes (refs. 4, 5, and 17). The distance  $R$  was also calculated for rubidium and cesium ions on tungsten by using the same hard sphere model, and assuming the value of  $\beta$ , and the location of the surface to be independent of the nature of the adsorbed ion.

Because  $\alpha$  is not uniquely determined (table II), the calculation of  $\phi$  was performed for both the most recently reported calculated values, and the average of all reported experimental values of  $\alpha$ . The results are listed in

<sup>2</sup>This value of  $\beta$  is slightly lower than the upper limit  $\beta = 0.46$  suggested by Sachs and Dexter (ref. 7) and considerably higher than the value  $\beta = 0.09$  used by Cutler and Gibbons (ref. 16), which is reasonable.

table III. In general, the calculated values of  $\phi$  are in reasonable agreement with the experimental values. That the calculations based on the calculated values of  $\alpha$  show a slightly lower value for the cesium ion - tungsten

TABLE III. - APPLICATION OF EQUATION (6) TO ADSORPTION

OF ALKALI METAL IONS ON TUNGSTEN

Ion	Electronic polarizability, $\alpha, \text{A}^3$	Distance from metal surface to center of ion, $R, \text{A}$	Interaction energy, $\Phi_i, \text{ev}$	Heat of adsorption, $\phi, \text{ev}$	
				Calculated	Experimental
Sodium	<sup>a</sup> 0.200	<sup>a</sup> 0.73	<sup>a</sup> 2.460	2.51	2.50
	<sup>b</sup> 0.152			2.50	
Rubidium	<sup>a</sup> 1.899	<sup>a</sup> 1.44	<sup>a</sup> 1.875	2.13	1.90
	<sup>b</sup> 2.920			2.27	
Cesium	<sup>a</sup> 3.305	<sup>a</sup> 1.69	<sup>a</sup> 1.660	2.03	2.04 to 2.19
	<sup>b</sup> 5.600			2.29	

<sup>a</sup>Average of value listed in table II.

<sup>b</sup>Calculated.

adsorption than for the rubidium ion - tungsten adsorption is particularly interesting. This is in agreement with the experimental results. While the major bonding is due to the image-force of the net charge of the ion, the polarization effect cannot be neglected. It is primarily the difference in polarizabilities that accounts for the heat of adsorption of cesium on tungsten being lower than that of rubidium on tungsten when the calculated values of  $\alpha$  are used.

CONCLUDING REMARKS

While equation (6) is not exact (the polarization term considers only dipole-dipole interactions), the results obtained (table III) are generally good. On the basis of the calculations performed, the theory presented is adequate to explain the phenomena of adsorption of alkali metal ions on tungsten in the limit of zero coverage and emphasizes that the effect arising from the polarizability of the adsorbed ion should not be neglected.

Unfortunately, the experimental values of  $\phi$  had to be collected from a variety of experimental techniques and laboratory conditions of numerous workers. It must also be recognized that except for the cesium adsorption there is only one reported value of  $\phi$  for each system. The results of this study must necessarily be considered within the framework of these limitations.

Lewis Research Center  
National Aeronautics and Space Administration  
Cleveland, Ohio, May 27, 1964

## REFERENCES

1. Bosworth, Richard Charles Leslie: The Mobility of Sodium on Tungsten. Proc. Roy. Soc. (London), ser. A, vol. 150, 1935, pp. 58-76.
2. Hughes, F. L., and Levinstein, H.: Mean Adsorption Lifetime of Rb on Etched Tungsten Single Crystals: Ions. Phys. Rev., 113, no. 4, Feb. 15, 1959, pp. 1029-1035.
3. de Boer, J. H., and Veenemans, C. F.: Adsorption of Alkali Metals on Metal Surfaces. Chemical Physica, vol. 1, 1934, pp. 753-762.
4. Taylor, John Bradshaw, and Langmuir, Irving: The Evaporation of Atoms, Ions and Electrons from Caesium Films on Tungsten. Phys. Rev., vol. 44, no. 6, Sept. 15, 1933, pp. 423-458.
5. Becker, J. A.: Adsorption on Metal Surfaces and Its Bearing on Catalysis. Advances in Catalysis, vol. 7, Academic Press, 1955.
6. Sheer, Milton D., and Fine, Joseph: Kinetics of  $\text{Cs}^+$  Desorption from Tungsten. Jour. Chem. Phys., vol. 37, no. 1, July 1962, pp. 107-113.
7. Sachs, R. G., and Dexter, D. L.: Quantum Limits of the Electrostatic Image Charge Theory. Jour. Appl. Phys., vol. 21, 1950, pp. 1304-1308.
8. Prosen, E. J. R., and Sachs, R. G.: The Interaction Between a Molecule and a Metal Surface. Phys. Rev., vol. 61, Jan. 1-15, 1942, pp. 65-73.
9. Neustadter, Harold E., and Luke, Kueng P.: Low-Coverage Heats of Adsorption. I - Alkali Metal Atoms on Tungsten; Atom-Metal Interaction Theory. NASA TN D-2430, 1964.
10. Sternheimer, R. M.: Electronic Polarizabilities of Ions. Phys. Rev., vol. 115, no. 5, Sept. 1, 1959, pp. 1198-1206.
11. Fajans, K., and Joos, G.: Molrefraktion von Ionen und Molekülen im Lichte der Atomstruktur. (Mass Refraction of Ions and Molecules in the Light of the Atomic Structure.) Zs. Physik, bd. 23, 1924, pp. 1-46.
12. Born, M., and Heisenberg, W.: Über den Einfluss der Deformierbarkeit der Ionen auf Optische und Chemische Konstanten. (Effect of Ion Deformability on Optical and Chemical Constants.) Zs. Physik, bd. 23, 1924, pp. 388-410.
13. Pauling, Linus: The Theoretical Prediction of the Physical Properties of Many-Electron Atoms and Ions. Mole Refraction, Diamagnetic Susceptibility and Extension in Space. Proc. Roy. Soc. (London), ser. A, vol. 114, 1927, pp. 181-211. Chemical Abstracts, vol. 25, June-Oct. 1927.
14. Tessman, Jack R., Kahn, A. H., and Shockley, William: Electronic Polarizabilities of Ions in Crystals. Phys. Rev., vol. 29, no. 24, Nov. 15, 1953, pp. 890-895.



15. Hughes, F. L.: Mean Adsorption Lifetime of Rb on Etched Tungsten Single Crystals: Neutrals. Phys. Rev., vol. 113, no. 4, Feb. 15, 1959, pp. 1036-1038.
16. Cutler, P. H., and Gibbons, J. J.: Model for the Surface Potential Barrier and the Periodic Deviations in the Schottky Effect. Phys. Rev., vol. 111, no. 2, July 15, 1958, pp. 394-402.
17. Hughes, F. L., Levinstein, H., and Kaplan, R.: Surface Properties of Etched Tungsten Single Crystals. Phys. Rev., vol. 113, no. 4, Feb. 15, 1959, pp. 1023-1028.

2/17/85  
J

*"The aeronautical and space activities of the United States shall be conducted so as to contribute . . . to the expansion of human knowledge of phenomena in the atmosphere and space. The Administration shall provide for the widest practicable and appropriate dissemination of information concerning its activities and the results thereof."*

—NATIONAL AERONAUTICS AND SPACE ACT OF 1958

## NASA SCIENTIFIC AND TECHNICAL PUBLICATIONS

**TECHNICAL REPORTS:** Scientific and technical information considered important, complete, and a lasting contribution to existing knowledge.

**TECHNICAL NOTES:** Information less broad in scope but nevertheless of importance as a contribution to existing knowledge.

**TECHNICAL MEMORANDUMS:** Information receiving limited distribution because of preliminary data, security classification, or other reasons.

**CONTRACTOR REPORTS:** Technical information generated in connection with a NASA contract or grant and released under NASA auspices.

**TECHNICAL TRANSLATIONS:** Information published in a foreign language considered to merit NASA distribution in English.

**TECHNICAL REPRINTS:** Information derived from NASA activities and initially published in the form of journal articles.

**SPECIAL PUBLICATIONS:** Information derived from or of value to NASA activities but not necessarily reporting the results of individual NASA-programmed scientific efforts. Publications include conference proceedings, monographs, data compilations, handbooks, sourcebooks, and special bibliographies.

*Details on the availability of these publications may be obtained from:*

SCIENTIFIC AND TECHNICAL INFORMATION DIVISION  
NATIONAL AERONAUTICS AND SPACE ADMINISTRATION  
Washington, D.C. 20546