

RESEARCH ARTICLE

Open Access



BOTRYTIS-INDUCED KINASE1, a plasma membrane-localized receptor-like protein kinase, is a negative regulator of phosphate homeostasis in *Arabidopsis thaliana*

Huijuan Zhang^{1,2}, Lei Huang², Yongbo Hong² and Fengming Song^{2*}

Abstract

Background: Plants have evolved complex coordinated regulatory networks to cope with deficiency of phosphate (Pi) in their growth environment; however, the detailed molecular mechanisms that regulate Pi sensing and signaling pathways are not fully understood yet. We report here that the involvement of Arabidopsis BIK1, a plasma membrane-localized receptor-like protein kinase that plays critical role in immunity, in Pi starvation response.

Results: qRT-PCR analysis revealed that expression of *BIK1* was induced by Pi starvation and GUS staining indicated that the *BIK1* promoter activity was detected in root, stem and leaf tissues of plants grown in Pi starvation condition, demonstrating that *BIK1* is responsive to Pi starvation stress. The *bik1* plants accumulated higher Pi content in root and leaf tissues and exhibited altered root architecture such as shorter primary roots, longer and more root hairs and lateral roots, as compared with those in the wild type plants, when grown under Pi sufficient and deficient conditions. Increased anthocyanin content and acid phosphatase activity, reduced accumulation of reactive oxygen species and downregulated expression of Pi starvation-induced genes including *PHR1*, *WRKY75*, *AT4*, *PHT1;2* and *PHT1;4* were observed in *bik1* plants grown under Pi deficient condition. Furthermore, the expression of *PHO2* was downregulated while the expression of *miRNA399a* and *miRNA399d*, which target to *PHO2*, was upregulated in *bik1* plants, compared to the wild type plants, when grown under Pi deficient condition.

Conclusion: Our results demonstrate that *BIK1* is a Pi starvation-responsive gene that functions as a negative regulator of Pi homeostasis in Arabidopsis.

Keywords: BOTRYTIS-INDUCED KINASE1 (BIK1), Phosphate starvation stress, Phosphate starvation response, Root architecture

Background

Phosphate (Pi) is one of the indispensable macronutrients to plants for growth, development and reproduction. Pi deficiency (-Pi) is one of the main limiting factors for increasing crop yield and improving quality because of low bioavailability of Pi in soil [1–4]. To cope with Pi deficiency, plants develop a series of tightly controlled adaptive responses including external developmental alterations of

increasing Pi absorption and internal metabolic, physiological and biochemical alterations of reducing Pi usage [4–6]. To increase Pi uptake under Pi depletion stress, plants adapt to modulate the root architecture bearing more and longer lateral roots as well as denser root hairs, which enable the roots to explore the Pi resources in soil [7–12]. To reduce Pi consumption under Pi deficiency condition, plants often modulate metabolisms to maintain intracellular Pi homeostasis by reducing metabolic consumption of Pi [1, 3, 13–15] and degrading compounds to release Pi [16]. Furthermore, a series of physiological

* Correspondence: fmsong@zju.edu.cn

²National Key Laboratory for Rice Biology, Institute of Biotechnology, Zhejiang University, Hangzhou 310058, People's Republic of China
Full list of author information is available at the end of the article

and biochemical adaptations including the induction and secretion of phosphatases and organic acids and accumulation of protective metabolites such as anthocyanin help augment the availability of both endogenous and exogenous Pi [1, 4, 17, 18].

Genetic, physiological and biochemical studies in *Arabidopsis* have demonstrated that the acquisition, allocation, and metabolism of Pi are highly regulated processes and require the concerted action of multiple membrane Pi transport systems [19–23]. Among five distinct classes of proteins possessing Pi transport activity, the plastidic Pi translocator group function as antiport systems, whereas the other four Pi transporter families, named the PHOSPHATE TRANSPORTER1 (Pht1), Pht2, Pht3 and Pht4, contribute to the acquisition, allocation and remobilization of Pi [24–27]. Most of the plasma membrane-localized high affinity transporters in the Pht1 family mediate Pi acquisition from external environment [1, 24, 28–30]. Once Pi is transported into root epidemic cells, translocation and allocation of Pi within the plants and cells are key steps in maintaining Pi homeostasis at cell and whole plant levels. The low affinity transporters in the Pht2 and Pht4 families are thought to participate mainly in Pi transfer across internal cellular membranes and thus allocate Pi in different compartments of the cells [25, 27, 31–33]. Pht1;5, a Pht1 family member, PHO1, Pht1;8 and Pht1;9 were shown to play critical roles in systemic regulation of Pi homeostasis, e.g. mobilization of Pi from source to sink organs in accordance with the Pi status of the plant [34–36].

Functional characterization of genes in a number of mutants with altered response to Pi depletion have led to the identification of several different regulatory mechanisms controlling the adaptive responses to Pi starvation. These regulatory mechanisms include transcriptional regulation by transcription factors such as PHR1, WRKY6, WRKY42, WRKY45, WRKY75, ZAT6, bHLH32, ERF070 and MYB62 [37–47], posttranscriptional regulation by microRNAs including miRNA399 [48–53], posttranslational regulation by protein modifications such as sumoylation of SIZ1 [54], phosphorylation of Pht1.1 and activation of MKK9-MPK3/MPK6 module [55, 56], deubiquitination of UBP14 [57] and chromatin histone modification and epigenetic [46, 58]. Furthermore, it was also demonstrated that Pi starvation response cross-talks signaling mediated by different hormones such as auxin [10, 15, 59–61], cytokinin [10, 15, 62, 63], ethylene [9, 10, 64–66] and gibberellin [67]. Although great advance on the adaptive response to Pi starvation has been made during the last decade, the molecular mechanism that regulates these adaptive processes has yet to be elucidated in detail.

The *Arabidopsis thaliana* *BOTRYTIS-INDUCED KINASE1* (*BIK1*) encodes a plasma membrane-localized receptor-like protein kinase and plays critical

roles in *Botrytis cinerea* resistance [68]. Recent studies have shown that BIK1 interacts with receptors for pathogen- or damage-associated molecular patterns such as FLS2 and PEPRs to regulate immune response against different types of pathogens [69–73] and defense response to insect pests [74]. The *bik1* mutant plants showed an altered root architecture [68], similar to morphological phenotypes often seen in mutants with Pi starvation response [75], indicating a possible involvement of *BIK1* in Pi starvation response in *Arabidopsis*. Therefore, we investigated whether BIK1 functions in Pi starvation response and our results demonstrate that BIK1 plays a role in regulation of Pi homeostasis in *Arabidopsis*.

Results

Responsiveness of *BIK1* to Pi starvation

When grown on MS medium under normal conditions, the *bik1* plants produced shorter primary roots and longer and significantly more root hairs and lateral roots than WT plants [68], which is reminiscent of the mutants with defects in Pi nutrition [75]. These observations led us to examine whether BIK1 has a function in Pi starvation response. We first examined whether *BIK1* is responsive to Pi starvation stress by analyzing the expression patterns of *BIK1* in seedlings grown under +Pi and -Pi conditions. As shown in Fig. 1a, expression of *BIK1* was detected in roots, shoots and leaves of seedlings grown under +Pi condition and no significant changes in *BIK1* expression was observed during the experiment period. However, expression level of *BIK1* was markedly induced with similar patterns in roots, shoots and leaves of seedlings after transferring to medium without Pi supplement (Fig. 1a). The transcript levels of *BIK1* in seedlings grown under -Pi condition increased at 12 h and peaked at 24 and 48 h after transferring, leading to 7.5 ~ 11.2 folds of increases over those in seedlings grown under +Pi condition (Fig. 1a). To gain further information on spatial expression of the *BIK1* gene in response to Pi starvation, we generated *BIK1_{pro}::GUS* transgenic lines and compared the GUS staining patterns in T2 seedlings during Pi starvation stress. Slight GUS staining was observed in leaves and roots of the *BIK1_{pro}::GUS* seedlings at 0 h after transferred to medium without Pi supplement (Fig. 1b), indicating a basal expression of *BIK1* in leaf and root tissues, similar to the results from RT-PCR. At 1 day after transferred to medium without Pi supplement, significant GUS staining was easily seen in roots, shoot and leaves of the *BIK1_{pro}::GUS* seedlings (Fig. 1b). qRT-PCR analysis showed 7.3 and 9.7 folds of increases in the expression levels of *GUS* gene in root and leaf tissues of the *BIK1_{pro}::GUS*





Fig. 1 Responsiveness of *BIK1* to Pi starvation. **a** Expression changes of *BIK1* in different tissues of WT plants under +Pi and -Pi conditions. Seven-day-old seedlings grown hydroponically under normal Pi condition were transferred to medium supplemented with Pi (+Pi, 250 μ M) or without Pi (-Pi). Roots, shoots and leaves were collected for analysis of *BIK1* expression by qRT-PCR at indicated time points after transferring. **b** Detection of *BIK1* promoter activity in *BIK1_{pro}::GUS* seedlings by GUS staining. **c** Expression changes of *GUS* gene in root and leaf tissues of *BIK1_{pro}::GUS* seedlings under -Pi condition. Seven-day-old *BIK1_{pro}::GUS* seedlings grown in normal medium were transferred to medium without Pi supplement and samples were collected at 0 and 1 day after transferring for GUS staining and analysis of gene expression. Data were normalized with the transcript level of *UBQ10* and relative expression levels were shown as folds of the *UBQ10* level. Data presented are the means \pm SD from three independent experiments and * above the error bars indicate significant differences at $p < 0.05$ level between the +Pi and -Pi conditions

seedlings at 1 day after transferring to -Pi condition (Fig. 1c). Notably, GUS staining was clearly observed in vascular tissues of roots, shoots and leaves of the *BIK1_{pro}::GUS* seedlings subjected to Pi starvation (Fig. 1b). Together, data from RT-PCR and GUS staining demonstrate that *BIK1* is responsive to Pi starvation.

Increased Pi concentration in *bik1* plants

We next examined whether loss of *BIK1* function affects Pi homeostasis in *bik1* plants. Total Pi contents in leaves and roots of WT and *bik1* plants grown hydroponically under +Pi (250 μ M) and -Pi conditions for 27 days were measured. As shown in Fig. 2, a significant increase in total Pi contents was observed in leaves and roots of



bik1 plants as compared to WT plants under both +Pi and -Pi conditions. When grown under +Pi condition, total Pi contents in roots and leaves of *bik1* plants were 0.43 and 1.12 times higher over those in WT plants, respectively (Fig. 2a and b). Similarly, when grown under -Pi condition, total Pi contents in roots and leaves of *bik1* plants were 0.55 and 1.05 times higher than those of WT plants, respectively (Fig. 2a and b). These data indicate that BIK1 has a function either in Pi uptake or in the transfer of Pi from the roots to the leaves.

Altered root architecture in *bik1* plants

Root architecture plays important roles in maintaining Pi homeostasis in plants [75]. We compared the root architecture of WT and *bik1* seedlings grown under Pi normal (+Pi) and starvation (-Pi) conditions. When compared with WT seedlings, the *bik1* seedlings grown under +Pi condition showed shorter primary root and more lateral roots (Fig. 3a, c and d), and this trend was much evident in the *bik1* seedlings grown under -Pi condition (Fig. 3a, c and d). When grown under +Pi and -Pi conditions, the elongation rate of the primary and lateral roots of the *bik1* seedlings was markedly reduced as compared with WT seedlings (Fig. 3e and f). Furthermore, the *bik1* seedlings grown under +Pi and -Pi conditions showed more and longer root hairs than WT seedlings (Fig. 3b, g and h). These data indicate that BIK1 plays an important role in regulating development of root architecture.

Increased anthocyanin accumulation and acid phosphatase activity in *bik1* plants

Accumulation of anthocyanin and increased secretion of acid phosphatases are characteristic symptoms in plants under Pi starvation conditions [3]. We thus examined

and compared the accumulation of anthocyanin and activity of acid phosphatase between WT and *bik1* plants grown under +Pi and -Pi conditions. When grown on medium under -Pi condition for 5 days, the leaves of the *bik1* seedlings turned to be purple while WT seedlings still kept green (Fig. 4a). The anthocyanin contents in *bik1* seedlings grown under +Pi and -Pi conditions were much higher than those in WT seedlings, showing 32 % and 105 % increase, respectively (Fig. 4b). No visible staining and significant changes in acid phosphatase activity were detected in WT and *bik1* seedlings grown +Pi condition (Fig. 4c and d). When grown under -Pi condition, roots of the *bik1* seedlings are stained deep blue while the staining of roots of WT seedlings is much lighter (Fig. 4c). Similarly, activity of acid phosphatase in roots of *bik1* seedlings is much higher than that in WT seedlings, grown under -Pi condition, showing an increase of ~65 % (Fig. 4d). These results indicate that the *bik1* seedlings secrete higher levels of acid phosphatase than WT seedlings under Pi starvation stress. Collectively, these results suggest that loss of BIK1 function results in typical Pi starvation responses as revealed by the increased anthocyanin contents and acid phosphatase activity.

Reduced ROS accumulation in *bik1* plants under Pi starvation condition

ROS, generated and accumulated in plant response to different types of stresses, has been implicated in many biological processes including biotic and abiotic responses [76, 77]. Therefore, we examined whether mutation in BIK1 gene affects the balance of ROS and hence leads to accumulation of ROS in plants during Pi starvation stress. In NBT staining of superoxide anion, no significant difference was detected between WT and *bik1* plants grown under +Pi condition; however, significant accumulation of superoxide anion in leaves, stem and roots of WT and *bik1* plants grown under -Pi condition was observed (Fig. 5a). Notably, accumulation of superoxide anion in WT plants was much higher than that in *bik1* plants grown under -Pi condition (Fig. 5a). Quantification of H₂O₂ contents revealed a significant lower level of H₂O₂ accumulated in *bik1* plants than that in WT plants when grown under +Pi condition (Fig. 5b). Accumulation of H₂O₂ in WT and *bik1* plants grown under -Pi condition was markedly increased, leading to 95 and 78 % of increases as compared with those in plants grown under +Pi condition, respectively (Fig. 5b). Under -Pi condition, the content of H₂O₂ in *bik1* plants was approximately 54 % of that in WT plants (Fig. 5b). These results indicate that loss of BIK1 function led to reduced accumulation of ROS in *bik1* plants under -Pi starvation stress.




Fig. 3 (See legend on next page.)

(See figure on previous page.)

Fig. 3 Altered root architecture in *bik1* seedlings grown under +Pi and -Pi conditions. Seven-day-old seedlings grown under normal condition were transferred into fresh medium supplemented with (+Pi, 250 μ M) or without Pi (-Pi) and grown for another 7 days. **a** Root architecture in WT and *bik1* seedlings. **b** Root hairs in WT and *bik1* seedlings. Fragments of 5 mm from root tips were presented. **c** to **h** Comparative parameters for length of primary roots (**c**), numbers of lateral roots (**d**), elongation of primary roots (**e**) and lateral roots (**f**), and numbers (**g**) and length (**h**) of root hairs of WT and *bik1* seedlings grown under +Pi and -Pi condition. Data presented are the means \pm SD from three independent experiments and different letters above the columns indicate significant differences at $p < 0.05$ level

Altered expression of Pi starvation-induced genes and miRNA399 in *bik1* plants

Expression of some well-characterized Pi starvation-induced genes was analyzed and compared between WT and *bik1* seedlings that had been transferred into medium with Pi (+Pi) or without Pi (-Pi) for 5 days. When grown under +Pi condition, the expression levels of *AT4* [78], *WRKY75* [39], *PHT1;2* and *PHT1;4* [24, 28] in *bik1* seedlings were comparable to those in WT seedlings, while the expression of *PHR1* [37] in *bik1* seedlings was reduced (Fig. 6a). The expression levels of the Pi starvation-induced genes tested were markedly increased in WT and *bik1* plants grown under -Pi condition (Fig. 6a). However, significant reductions in the expression levels of these Pi starvation-induced genes was observed in *bik1* plants, as compared with those in WT plants, under -Pi condition (Fig. 6a). These results indicate that mutation in

BIK1 leads to reduced expression of the Pi starvation-induced genes in plants grown under -Pi condition.

It was found that expression of *PHO2*, encoding a ubiquitin-conjugating E2 enzyme, was suppressed by *miRNA399*, which controls Pi homeostasis in plants and whose expression is up-regulated by Pi starvation [48–52]. To examine whether *PHO2/miRNA399* links to increased accumulation of Pi in *bik1* plants, we analyzed the changes in levels of *PHO2* expression and two *miRNA399* primary transcripts, *miRNA399a* and *miRNA399d*, in *bik1* seedlings grown under +Pi and -Pi conditions. The expression level of *PHO2* and the transcript levels of *miRNA399a* and *miRNA399d* were comparable in WT and *bik1* seedlings grown under +Pi condition. However, the expression levels of *PHO* in WT and *bik1* seedlings under -Pi condition were significantly reduced whereas the transcript levels of *miRNA399a* and *miRNA399d* in WT and *bik1* seedlings grown




Fig. 4 Increased levels of anthocyanin accumulation and acid phosphatases activity in *bik1* plants. **a** and **b** Measurements of anthocyanin contents. Seven-day-old WT and *bik1* seedlings grown on 1/2 MS were transferred into MS medium supplemented with Pi (+Pi, 1 mM) or without Pi (-Pi). Photos were taken and anthocyanin contents were measured at 5 days after treatment. **c** and **d** Detection of acid phosphatase activity. Ten-day-old seedlings grown in liquid medium were transferred to fresh medium supplemented with Pi (+Pi, 1 mM) or without Pi (-Pi) and covered with a layer of 0.008 % BCIP-containing agarose. Photos were taken and the activity of acid phosphatase in roots was estimated at 5 days after treatment. Data presented are the means \pm SD from three independent experiments and different letters above the columns indicate significant differences at $p < 0.05$ level between WT and *bik1* plants grown under same condition




Fig. 5 Reduced accumulation of ROS in *bik1* plants. Four-week-old seedlings grown under + Pi (250 μM) and -Pi conditions were collected for detection of superoxide anion and measurement of H₂O₂. **a** NBT staining of superoxide anion in WT and *bik1* plants grown under + Pi and -Pi conditions. **b** Quantification of H₂O₂ contents in WT and *bik1* plants grown under + Pi and -Pi conditions. Data presented are the means ± SD from three independent experiments and different letters above the columns indicate significant differences at *p* < 0.05 level between WT and *bik1* plants grown under same condition




Fig. 6 Down-regulated expression of Pi starvation-induced genes in *bik1* plants. **a** Expression patterns of Pi starvation-induced genes in WT and *bik1* seedlings grown under + Pi and -Pi conditions. Seven-day-old seedlings grown under normal Pi condition were treated with Pi (+Pi, 250 μM) or without Pi (-Pi) for 5 days and total RNA were extracted from root tissues at 5 days after treatment. **b** Expression patterns of *PHO2* and *miR399* in WT and *bik1* seedlings grown under + Pi and -Pi conditions. Seven-day-old seedlings grown under normal Pi condition were transferred to medium supplemented with Pi (+Pi, 250 μM) or without Pi (-Pi) and samples were collected at 48 h after treatment. Expression of Pi starvation-responsive genes and miR399s was analyzed by qRT-PCR using specific primers and data were normalized with the transcript level of *UBQ10* as an internal control. Data presented are the means ± SD from three independent experiments and different letters above the columns indicate significant differences at *p* < 0.05 level

under $-P_i$ condition were markedly increased, as compared with the corresponding seedlings under $+P_i$ condition (Fig. 6b). A further reduction in the expression of *PHO2* and a further increase in the transcripts of *miRNA399a* and *miRNA399d* in *bik1* seedlings was observed as compared with those in WT seedlings grown under $-P_i$ condition (Fig. 6b). Together, these data indicate that the *PHO2/miRNA399* plays a role in regulating P_i accumulation in *bik1* plants under $-P_i$ condition.

Discussion

The maintenance of cellular P_i homeostasis in plants involves complicated regulatory mechanisms. Several studies have demonstrated that posttranslational modifications such as phosphorylation, sumoylation and ubiquitination of regulatory proteins play critical roles in P_i starvation responses [54, 55, 57]. Recently, genome-wide co-expression analysis identifies dozens of protein kinases as important regulators of P_i deficiency-induced root hair remodeling [79, 80]. It was also found that *in vivo* phosphorylation activation of PPC1 (phosphoenolpyruvate carboxylase 1) is involved in the metabolic adaptations of P_i starvation [81] and activation of MKK9-MPK3/MPK6 enhances phosphate acquisition in Arabidopsis [56]. These findings indicate an important role for protein kinases in regulating P_i starvation response. In this study, we found that BIK1, a plasma membrane-localized receptor-like protein kinase [68], plays an important role in modulating P_i starvation responses and functions as a negative regulator of P_i homeostasis in Arabidopsis plants. Thus, evidence presented in the current study renders BIK1 for a novel function in P_i starvation response, in addition to its previously reported functions in immunity [68–73].

Several lines of evidence presented in this study support that BIK1 functions in P_i starvation response. Earlier study has shown that expression of *BIK1* could be induced by infection with *B. cinerea* as well as treatments with some well-known defense signaling molecules [68]. Our qRT-PCR analysis of *BIK1* expression and determination of *BIK1* promoter activity in stable *BIK1_{pro}::GUS* transgenic seedlings demonstrate that *BIK1* can be induced by P_i starvation and P_i starvation-induced expression of *BIK1* in root and shoot tissues initiated earlier than in leaf tissues (Fig. 1a). The pattern of *BIK1* expression induced by P_i starvation is similar to most of the P_i starvation-induced genes identified so far, like *WKRY75*, *ERF070* and *ZAT6* [39, 40, 44]. Importantly, promoter activity analysis revealed that GUS staining driven by the *BIK1* promoter was initiated in vascular tissues of root (Fig. 1b) that is the primary organ that responds to P_i starvation, further confirming the responsiveness of *BIK1* to P_i starvation. Thus, it is likely that *BIK1* can respond rapidly to the altered P_i status of plants under P_i starvation condition. It was

particularly noteworthy that BIK1 is a plasma membrane-localized receptor-like protein kinase and has been shown to be phosphorylated by other kinases (e.g. BAK1 and FLS2) for its full activity in plant immune responses [69, 70]. In this context, it is thus possible that BIK1 is phosphorylated by unknown kinase during P_i starvation stress and phosphorylated BIK1 acts in P_i starvation response, in addition to its transcriptional regulation upon P_i starvation.

The primary root, lateral root, and root hairs are three main components of the root architecture that is critical to absorption of P_i from soil. Low P availability can drastically alter the root architecture by switching the indeterminate growth to determinate growth to promote lateral root growth [75]. Generally, not only the primary roots but also almost all mature lateral roots enter into the determinate developmental program under low P_i condition [82]. The *bik1* seedlings showed a significant change in root architecture and the primary root length of *bik1* plants was significantly decreased (Fig. 6). This is similar to a common phenomenon observed that the primary root length is decreased significantly due to the P_i starvation-induced determinate growth in primary root [82]. In addition to the change of the primary root, the *bik1* plants showed increased number and length of lateral roots and root hairs (Fig. 6), which increase root surface contacting an increased soil volume to explore P_i availability in soil [75]. Similar root architectures were observed in the *bik1* plants under $+P_i$ and $-P_i$ conditions (Fig. 6). Thus, it is likely that the effect of BIK1 on the development of root architecture is independent of the P_i status in plants and P_i availability in soil. Nevertheless, the characteristic root architecture observed in *bik1* plants suggests that BIK1 is a negative regulator of lateral root and root hair development. This is in agreement with previous observations that the *bik1* plants showed some defects in growth and development, e.g. weak stem strength, early flowering and less seed setting [68]. Together, these data further demonstrate that BIK1 is required for normal plant growth and development [68]. It will be interesting to investigate whether the growth and developmental defects in *bik1* plants is due to an activated P_i starvation response and if this is the case, the results obtained will further support a crosstalk between P_i starvation response and plant growth/development. It was previously found that overexpression of *ZAT6* retards growth and results in typical P_i starvation responses [40].

Accompanied with high contents of total P_i in roots and leaves of *bik1* plants under $+P_i$ and $-P_i$ conditions (Fig. 2) are the significant accumulation of anthocyanin (Fig. 4a and b), increased activity of acid phosphatase (Fig. 4a and b) and alterations in expression of P_i

starvation-induced genes (Fig. 6). Collectively, these data imply that, whatever grown under + Pi condition or under -Pi condition, a Pi starvation response including those of physiological, molecular and metabolic changes is activated in *bik1* plants. Increased activity of acid phosphatases in *bik1* plants grown under -Pi condition (Fig. 3b) not only represents a characteristic Pi starvation response [83] but also can release more Pi available for absorption by the roots. This is supported by the findings that mutations in genes encoding for purple acid phosphatases affect markedly the uptake of Pi from exogenous sources [84–86]. Because expression of *BIK1* was rapidly induced by Pi starvation (Fig. 1), it seems possible that *BIK1* is involved in Pi uptake by regulating the activity of acid phosphatase. Similar observations were also obtained for some Pi starvation response regulators such as *PHR1* and *ZAT6*, which have been shown to affect activity of acid phosphatase and Pi uptake in transgenic plants with altered expression of these genes [40, 85, 87, 88]. Interestingly, expression of *Pht1;2* and *Pht1;4*, encoding for high-affinity Pi transporters [24, 28], *WRKY75*, encoding a WRKY transcriptional factor involved in Pi acquisition [39], *At4*, a member of the Mt4/TPSI1 gene family involved in Pi distribution [78], and *PHR1*, encoding a MYB transcriptional factor involved in Pi starvation response signaling [37], was suppressed significantly in roots of *bik1* plants under -Pi condition (Fig. 6a). These data demonstrate that *BIK1* has global effect on a set of Pi starvation-induced genes that are involved in Pi acquisition and mobilization. Notably, the expression levels of the tested Pi starvation-responsive gene except *PHR1* in *bik1* plants were slightly reduced as compared to those in WT plants under + Pi condition (Fig. 6a), indicating a possibility that these Pi starvation-responsive genes may be also affected by mutation of *BIK1* itself. Particularly, the expression of *PHR1* in *bik1* plants grown under + Pi condition was significantly reduced as compared to that in WT plants (Fig. 6a), leading to an open question whether *BIK1* acts upstream of *PHR1* in regulating the Pi starvation signaling. This can be clarified by phenotyping and analysis of *PHR1* expression in *BIK1*-overexpressing plants and/or detailed examination of biochemical and genetic requirement of *BIK1* for Pi starvation response. It is currently difficult to link the function of *BIK1* to Pi uptake or its root-to-leaf translocation as Pi uptake and transportation in WT plants is also affected under -Pi condition (Fig. 2). This can further be explained by the reduced expression of these tested Pi starvation-induced genes in root of *bik1* plants as compared to those in WT plants under -Pi condition (Fig. 6). Furthermore, altered expression patterns of *PHO2* and *miRNA399a/d*, which are thought to be involved in systemic signaling of Pi starvation response and Pi distribution in plants [49, 52], were also observed in *bik1* plants

grown under + Pi and -Pi conditions (Fig. 6b). Thus, it is likely that *BIK1* is involved in maintaining Pi homeostasis in whole plants. Taken together, these data support an idea that *BIK1* is a negative regulator of Pi starvation responses, probably through affecting development of root architecture and a series of physiological and biochemical events related to Pi acquisition, mobilization and translocation. It is reasonable to speculate that, under Pi starvation stress, the *bik1* plants may experience a reduced Pi content as the WT plants but they can uptake efficiently Pi from growth environment with their significantly increased root surface area, leading to an increased Pi content in root and leaf tissues of *bik1* plants (Fig. 2). However, the detailed mechanism that *BIK1* functions in Pi starvation response remains to be explored further. Particularly, characterization of the targets that are phosphorylated by *BIK1* will be greatly helpful in elucidating the early signaling events that determine Pi starvation response.

ROS not only are toxic compounds produced in plant response to various stresses but also plays an integral role as signaling molecules in regulation of numerous biological processes such as growth, development, and responses to biotic and/or abiotic stress [89–91]. The involvement of ROS in Pi starvation response has been established recently. In this study, we found that both WT and *bik1* plants grown under -Pi condition accumulated more ROS, represented by H_2O_2 and superoxide anion, than those in plants grown under + Pi condition (Fig. 5). This is similar to the previous observations that H_2O_2 concentrations in roots increased upon Pi deprivation [92]. Notably, the accumulation of superoxide anion in WT and *bik1* plants under + Pi condition was comparable, the accumulation of superoxide anion in *bik1* plants was significantly reduced as compared to that in WT plants under -Pi condition (Fig. 5). By contrast, the *bik1* plants accumulated lower level of H_2O_2 than WT plants grown under + Pi condition (Fig. 5), which is similar to the histochemical staining in soil-grown *bik1* plants [68]. Thus, it is likely that *BIK1* affects the accumulation of superoxide anion and H_2O_2 in different ways: loss of *BIK1* function suppressed, at least partially, the Pi starvation-induced accumulation of superoxide anion while *BIK1* regulates directly the accumulation of H_2O_2 in plants under normal condition. This may also imply different functions of superoxide anion and H_2O_2 in Pi starvation response. On the other hand, increased accumulation of ROS in plants under -Pi condition might be one of the stress responses as those in response to other abiotic stress [90]. It was recently shown that localization pattern of ROS accumulated in root during Pi starvation stress is critical to shape the root architecture [93–96]. The reduced accumulation of ROS in *bik1* plants in relative to those in WT plants (Fig. 5), which is similar to the observation

that loss of *BIK1* function impaired the PAMP-induced ROS burst in immunity [97, 98], may attribute to the altered root architecture of *bik1* plants (Fig. 3). However, detailed analysis of ROS localization and the possible mechanism regulating ROS generation in root of *bik1* plants will provide new insights into the connection between *BIK1* and ROS in development of root architecture.

Conclusions

In summary, this study demonstrates that *BIK1* is a Pi starvation-responsive gene and functions as a negative regulator of Pi homeostasis in *Arabidopsis*. This not only renders a novel function for *BIK1* but also strengthens our understanding of post-transcriptional regulation during Pi starvation responses in plants. Considering that *BIK1* is a plasma membrane-localized receptor-like protein kinase, it thus may play a role in sensing and/or processing of the Pi starvation signal during early stage of Pi starvation stress.

Methods

Plant materials and growth condition

Arabidopsis thaliana ecotype Columbia (Col-0) and *bik1* (provided by Dr. Tesfaye Mengiste, Purdue University, USA) mutant [68] were used in this study. Seeds were surface sterilized with 70 % ethanol, washed with sterilized distill water and vernalized at 4 °C for 2 days before germination. Hydroponic, solid medium and liquid medium cultivation were used for different purpose of experiments. For solid medium cultivation, the basic medium used contained 2.06 mM NH_4NO_3 , 1.88 mM KNO_3 , 0.31 mM MgSO_4 , 0.1 mM MnSO_4 , 0.03 mM ZnSO_4 , 0.1 mM CuSO_4 , 0.3 mM CaCl_2 , 5.0 mM KI , 0.1 mM CoCl_2 , 0.1 mM FeSO_4 ; 0.1 mM EDTA , 0.1 mM H_3BO_3 , 1 mM $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$, 3 g/L sucrose, 10 g/L agar, pH 5.8. When treated for Pi sufficiency (+Pi) and Pi deficiency (-Pi), the medium was supplemented with 1 mM KH_2PO_4 , 10 μM KH_2PO_4 and 0.99 mM KCl . For hydroponic cultivation, seedlings at 5~7-leaf stage were transferred to 1/2 Hoagland solution for short adaption and then transferred to hydroponic solution containing 250 μM Pi or no Pi [99]. For liquid medium cultivation, seeds were dispensed in 1/2 MS medium without agar and rinsed with sterilized distilled water at 7 days. Seedlings were then transferred in MS liquid medium with Pi (1 mM) or without Pi and allowed to grow with shaking at 85 rpm. Plants were grown in a growth room under a 16 h light (100 $\mu\text{mol} \cdot \text{s}^{-1} \cdot \text{m}^{-2}$ photons $\text{m}^{-2} \text{sec}^{-1}$ of intensity) and 8 h dark cycle at 22 ± 2 °C with 60 % relative humidity.

Measurement of root system architecture

Seven-day-old seedlings grown on 1/2 MS medium were transferred to Petri dishes and treated for + Pi or -Pi for another 7 days. Length of main root, number and length of the lateral root were measured. Total numbers of root hairs in a 5 mm region from root tip were recorded. Data were recorded from 15 individual plants from each treatment.

Anthocyanin analysis

Measurement of anthocyanin was performed as described previously [100]. Briefly, seedlings grown under + Pi or -Pi condition in solid medium were put into 20 mL extraction solution (propanol : HCl : H_2O = 18:1:81, V/V/V) and boiled for 90 s. The mixtures were kept overnight in dark at room temperature, centrifuged for 40 min at 5000 \times g, and the absorbance (A) was measured at 535 nm and 650 nm. The A_{535} values were corrected with the A_{650} values using formula $A_{535} = A_{535} - 2.2 \times A_{650}$. Anthocyanin contents were calculated according to a standard curve prepared with the same protocol.

Quantification of total Pi content

Total Pi content was quantified according to the U.S. Environmental Protection Agency Method 365.2 with minor modifications [39]. Briefly, samples (~50 mg/sample) were flamed to ash after recording of their dry weights and then 100 μL of concentrated HCl was added. Ten microliters of the mixture were drawn and diluted into 790 μL of water, followed by addition of 200 μL assay solution (4.8 mM NH_4MoO_4 , 2.5 N H_2SO_4 , and 35 μM ascorbic acid). The reactions were incubated at 45 °C for 20 min and the absorbance at 650 nm was measured spectrophotometrically. Contents of total Pi were calculated according to the Pi standard curve prepared with the same procedure.

Quantification and staining of acid phosphatase activity

Activity of acid phosphatase was quantified using the pNPP hydrolysis assay according to a previously described method [101]. Briefly, 30 mg of samples were grounded, transferred to Eppendorf tubes and then spin for 10 min at 2000 \times g. Reactions containing 100 μL supernatant, 100 μL *p*-nitrophenol sodium phosphate and 2.8 mL buffer were kept at 30 °C for 10 min, with shaking occasionally, and terminated by addition of 1 mL 0.5 M NaOH. The absorbance of the reactions was determined spectrophotometrically at 400 nm and the activity of acid phosphatase was calculated from the production of *p*-nitrophenol. Total protein was estimated using Bradford's reagent and the total acid phosphatase activity was expressed as mU/mg protein. Staining of acid phosphatase activity was performed as

described by Tomscha et al. [102] with minor modifications. Ten-day-old seedlings grown in liquid medium were rinsed in $-Pi$ medium, transferred to fresh medium supplemented with Pi (1 mM) or without Pi and then covered with a layer of 0.05 % agarose solution containing 0.008 % 5-bromo-4-chloro-3-indolyl phosphate (BCIP) [103].

Detection and quantification of H_2O_2

For quantification of H_2O_2 , 30 mg samples from 4-week-old plants were completely ground, followed by addition of 200 μ L 20 mM K_2HPO_4 (pH6.5) phosphate buffer. Quantification of H_2O_2 was performed using a commercial kit (Jiancheng Bioengineering Institute, Nanjing, China) according to the manufacturer's recommendation. Detection of superoxide anion was performed by the nitroblue tetrazolium (NBT) staining [104]. Four-week-old seedlings were vacuum infiltrated with 2 ml of 10 mM potassium phosphate buffer (pH 7.5) containing 10 mM NaN_3 and 0.1 % NBT for 30 min, cleared by boiling in 96 % ethanol, remained in 50 % ethanol before taking photos.

qRT-PCR analysis of gene expression

Total RNA was extracted using TRIZOL reagent (Invitrogen, Shanghai, China) according to the manufacturer's instructions. First strand cDNA was synthesized from 500 ng of total RNA using SuperScript III Kit (Invitrogen, Shanghai, China). The qPCR reactions contained 12.5 μ L SYBR Premix Ex TaqTM (TaKaRa, Dalian, China), 0.1 μ g cDNA and 7.5 pmol of each of gene-specific primers in 25 μ L and were conducted on a CFX96 real-time PCR system (BioRad, Hercules, CA, USA). Gene-specific primers used were as followings: BIK1-q-F, 5'-ACT TAT GGG TAC GCC GCG CCT GAG T-3'; BIK1-q-R, 5'-GGC ACG GAC CAC TTG GTC CA-3'; GUS-q-1F, 5'-AGG TGC ACG GGA ATG TTT CG-3'; GUS-q-1R, 5'-TGT GAG AGT CGC AGA ACA TT-3'; PHR1-q-1F, 5'-GTG ATT GGC ATG AAT GGG CTG AC-3'; PHR1-q-1R, 5'-CGC AAT TCC ACA GAC GGA GAA GG-3'; AT4-q-1F, 5'-GAT CGA AGT TGC CCA AAC GA -3'; AT4-q-1R, 5'-GAG CGA TGA AGA TTG CAT GAA G-3'; WRKY75-q-1F, 5'-GAG AAA TCC ACC GAA AAC TTC GAG CAT AT-3'; WRKY75-q-1R, 5'-GCA TGG TTT TTC TTT TCA ACA CAC GTA AAA TGT A-3'; PHT1.2-q-1F, 5'-AGG GCA AGT CCC TCG AAG AAC T-3'; PHT1.2-q-1R, 5'-ATC AAA CAA ACC ACA AAC AAC TCC ACA T-3'; PHT1.4-q-1F, 5'-TTG CTC CTA ATT TTC CTG ATG CT -3'; PHT1.4-q-1R, 5'-TGT GCC GGC CGA AAT CT-3'. Pri-miRNA399a-1F, 5'-TGG CAG GAA ACC ATT ACT TAG ATC T-3'; Pri-miRNA399a-1R, 5'-TCA CTA ATT AAA AGC AAT GCA TAA AGA GA-3'; Pri-miRNA399d-1F, 5'-TTA CTG GGC GAA

TAC TCC TAT GG-3'; Pri-miRNA399d-1R, 5'-ATT TTA CTT GCA TAT CTA GCC AAT GC-3'; PHO2-q-1F, 5'-AGG TTT GAA GCT CCA CCC TCA-3'; PHO2-q-1R, 5'-CCC AAG ATG TGA TTG GAG TTC C-3'; UBQ10-q-1F, 5'-GGC CTT GTA TAA TCC CTG ATG AAT AAG-3'; UBQ10-q-1R, 5'-AAA GAG ATA ACA GGA ACG GAA ACA TAG T-3'. Relative gene expression levels were calculated using $2^{-\Delta\Delta CT}$ method with three independent biological replicates.

Generation of *BIK1_{pro}::GUS* transgenic line and GUS staining

A 2 kb sequence upstream of the *BIK1* start codon was PCR amplified using primers AtBIK1-GUS-1F (5'-ATA **CTG CAG CTT GTT** GAT TGA TTA ATA GAT TAC C-3, a *Pst*I site in bold) and AtBIK1-GUS-1R (5'-GCC **GGA TCC** AGA ACT GAA GCA AGA ACC CAT C-3', a *Bam*HI site in bold) and cloned into vector pCAM-BIA1301. Transformation of wild-type Col-0 plants was performed using the floral dip infiltration method mediated by *Agrobacterium tumefaciens* strain GV3101. Plants of T2 generations from kanamycin-resistant transformants were used for GUS histochemical staining [105].

Abbreviations

+Pi, Pi sufficiency; *B. cinerea*, *Botrytis cinerea*; BCIP, 5-bromo-4-chloro-3-indolyl phosphate; *BIK1*, *Botrytis-induced kinase1*; MKK, mitogen-activated protein kinase; MPK, mitogen-activated protein kinase; NBT, nitroblue tetrazolium; PAMP, pathogen associated molecular pattern; *Pht1*, *PHOSPHATE TRANSPORTER1*; Pi, phosphate; $-Pi$, Pi deficiency; pNPP, *p*-nitrophenol sodium phosphate; *PPC1*, *phosphoenolpyruvate carboxylase 1*; qRT-PCR, quantitative reverse transcription PCR; ROS, reactive oxygen species; WT, wild type

Acknowledgement

We are grateful to Dr. Tesfaye Mengiste, Purdue University, USA, for providing the Arabidopsis WT and *bik1* seeds used in this study.

Funding

This work was supported by the National Natural Foundation of Sciences (no. 31272028) and the Ph.D. Program Foundation of Ministry of Education of China (no. 20120101110070).

Availability of supporting data

All data supporting the results of this article are included within the article.

Authors' contributions

HZ and FS designed the study and all experiments; HZ, LH and YH conducted all the molecular biology and physiology experiments; HZ and FS analyzed the data; FS drafted the manuscript with HZ. All authors have read and approved the final version of the manuscript.

Competing interests

The authors declare that they have no competing interests.

Consent for publication

Not applicable.

Ethics approval and consent to participate

Not applicable. No human or animals were involved in this study.

Author details

¹College of Life Science, Taizhou University, Taizhou, Zhejiang 318001, People's Republic of China. ²National Key Laboratory for Rice Biology, Institute of Biotechnology, Zhejiang University, Hangzhou 310058, People's Republic of China.

Received: 27 January 2016 Accepted: 28 June 2016

Published online: 07 July 2016

References

- Raghothama KG. Phosphate transport and signaling. *Curr Opin Plant Biol.* 2000;3:182–7.
- Lynch JP, Brown KM. Topsoil foraging—an architectural adaptation of plants to low phosphorus availability. *Plant Soil.* 2001;237:225–37.
- Abel S, Ticconi CA, Delatorre CA. Phosphate sensing in higher plant. *Physiol Plant.* 2002;115:1–8.
- Ticconi CA, Abel S. Short on phosphate: plant surveillance and countermeasures. *Trends Plant Sci.* 2004;9:548–55.
- Bucher M, Rausch C, Daram P. Molecular and biochemical mechanisms of phosphorus uptake into plants. *J Plant Nutr Soil Sci.* 2001;164:209–17.
- Raghothama KG, Karthikeyan AS. Phosphate acquisition. *Plant Soil.* 2005;274:37–49.
- Williamson LC, Ribrioux SPCP, Fitter AH, Leyser HMO. Phosphate availability regulates root system architecture in *Arabidopsis*. *Plant Physiol.* 2001;126:875–82.
- Ma Z, Bielenberg DG, Brown KM, Lynch JP. Regulation of root hair density by phosphorus availability in *Arabidopsis thaliana*. *Plant Cell Environ.* 2001;24:459–67.
- Schmidt W, Schikora A. Different pathways are involved in phosphate and iron stress-induced alterations of root epidermal cell development. *Plant Physiol.* 2001;125:2078–84.
- Lopez-Bucio J, Hernández-Abreu E, Sánchez-Calderón L, Nieto-Jacobo MF, Simpson J, Herrera-Estrella L. Phosphate availability alters architecture and causes changes in hormone sensitivity in the *Arabidopsis* root system. *Plant Physiol.* 2002;129:244–56.
- López-Bucio J, Cruz-Ramírez A, Herrera-Estrella A. The role of nutrient availability in regulating root architecture. *Curr Opin Plant Biol.* 2003;6:280–7.
- Niu YF, Chai RS, Jin GL, Wang H, Tang CX, Zhang YS. Responses of root architecture development to low phosphorus availability: a review. *Ann Bot.* 2013;112:391–408.
- Mimura T. Regulation of phosphate transport and homeostasis in plant cells. *Int Rev Cytol.* 1999;191:149–200.
- Vance CP, Uhde-Stone C, Allan DL. Phosphorus acquisition and use: critical adaptations by plants for securing a nonrenewable resource. *New Phytol.* 2003;157:423–47.
- Kobayashi K, Masuda T, Takamiya K, Ohta H. Membrane lipid alteration during phosphate starvation is regulated by phosphate signaling and auxin/cytokinin cross-talk. *Plant J.* 2006;47:238–48.
- Oropeza-Aburto A, Cruz-Ramírez A, Acevedo-Hernández GJ, Pérez-Torres CA, Caballero-Pérez J, Herrera-Estrella L. Functional analysis of the *Arabidopsis* *PLDZ2* promoter reveals an evolutionarily conserved low-Pi-responsive transcriptional enhancer element. *J Exp Bot.* 2012;63:2189–202.
- Rausch C, Bucher M. Molecular mechanisms of phosphate transport in plants. *Planta.* 2002;216:23–37.
- López-Arredondo DL, Leyva-González MA, González-Morales SI, López-Bucio J, Herrera-Estrella L. Phosphate nutrition: improving low-phosphate tolerance in crops. *Annu Rev Plant Biol.* 2014;65:95–123.
- Franco-Zorrilla JM, González E, Bustos R, Linhares F, Leyva A, Paz-Ares J. The transcriptional control of plant responses to phosphate limitation. *J Exp Bot.* 2004;55:285–93.
- Lin WY, Lin SI, Chiou TJ. Molecular regulators of phosphate homeostasis in plants. *J Exp Bot.* 2009;60:1427–38.
- Rouached H, Arpat AB, Poirier Y. Regulation of phosphate starvation responses in plants: signaling players and cross-talks. *Mol Plant.* 2010;3:288–99.
- Jain A, Nagarajan VK, Raghothama KG. Transcriptional regulation of phosphate acquisition by higher plants. *Cell Mol Life Sci.* 2012;69:3207–24.
- Zhang Z, Liao H, Lucas WJ. Molecular mechanisms underlying phosphate sensing, signaling, and adaptation in plants. *J Integr Plant Biol.* 2014;56:192–220.
- Muchhal US, Pardo JM, Raghothama KG. Phosphate transporters from the higher plant *Arabidopsis thaliana*. *Proc Natl Acad Sci U S A.* 1996;93:10519–23.
- Daram P, Brunner S, Rausch C, Steiner C, Amrhein N, Bucher M. Pht2;1 encodes a low-affinity phosphate transporter from *Arabidopsis*. *Plant Cell.* 1999;11:2153–66.
- Knappe S, Flügge UI, Fischer K. Analysis of the plastidic phosphate translocator gene family in *Arabidopsis* and identification of new phosphate translocator-homologous transporters, classified by their putative substrate-binding site. *Plant Physiol.* 2003;131:1178–90.
- Guo B, Irigoyen S, Fowler TB, Versaw WK. Differential expression and phylogenetic analysis suggest specialization of plastid-localized members of the PHT4 phosphate transporter family for photosynthetic and heterotrophic tissues. *Plant Signal Behav.* 2008;3:784–90.
- Shin H, Shin HS, Dewbre GR, Harrison MJ. Phosphate transport in *Arabidopsis*: Pht1;1 and Pht1;4 play a major role in phosphate acquisition from both low- and high-phosphate environments. *Plant J.* 2004;39:629–42.
- Nussaume L, Kanno S, Javot H, Marin E, Pochon N, Ayadi A, Nakanishi TM, Thibaud MC. Phosphate import in plants: Focus on the PHT1 transporters. *Front Plant Sci.* 2011;2:83.
- Remy E, Cabrito TR, Batista RA, Teixeira MC, Sá-Correia I, Duque P. The Pht1;9 and Pht1;8 transporters mediate inorganic phosphate acquisition by the *Arabidopsis thaliana* root during phosphorus starvation. *New Phytol.* 2012;195:356–71.
- Cubero B, Nakagawa Y, Jiang XY, Miura KJ, Li F, Raghothama KG, Bressan RA, Hasegawa PM, Pardo JM. The phosphate transporter PHT4;6 is a determinant of salt tolerance that is localized to the Golgi apparatus of *Arabidopsis*. *Mol Plant.* 2009;2:535–52.
- Hassler S, Lemke L, Jung B, Möhlmann T, Krüger F, Schumacher K, Espen L, Martinioia E, Neuhaus HE. Lack of the Golgi phosphate transporter PHT4;6 causes strong developmental defects, constitutively activated disease resistance mechanisms and altered intracellular phosphate compartmentation in *Arabidopsis*. *Plant J.* 2012;72:732–44.
- Liu J, Yang L, Luan M, Wang Y, Zhang C, Zhang B, et al. A vacuolar phosphate transporter essential for phosphate homeostasis in *Arabidopsis*. *Proc Natl Acad Sci U S A.* 2015;112:E6571–8.
- Hamburger D, Rezzonico E, MacDonald-Comber Petétot J, Somerville C, Poirier Y. Identification and characterization of the *Arabidopsis* *PHO1* gene involved in phosphate loading to the xylem. *Plant Cell.* 2002;14:889–902.
- Nagarajan VK, Jain A, Poling MD, Lewis AJ, Raghothama KG, Smith AP. *Arabidopsis* Pht1;5 mobilizes phosphate between source and sink organs and influences the interaction between phosphate homeostasis and ethylene signaling. *Plant Physiol.* 2011;156:1149–63.
- Lapis-Gaza HR, Jost R, Finnegan PM. *Arabidopsis* PHOSPHATE TRANSPORTER1 genes PHT1;8 and PHT1;9 are involved in root-to-shoot translocation of orthophosphate. *BMC Plant Biol.* 2014;14:334.
- Rubio V, Linhares F, Solano R, Martín AC, Iglesias J, Leyva, Paz-Ares J. A conserved MYB transcription factor involved in phosphate starvation signaling both in vascular plants and in unicellular algae. *Genes Dev.* 2001;15:2122–33.
- Chen ZH, Nimmo GA, Jenkins GI, Nimmo HG. BHLH32 modulates several biochemical and morphological processes that respond to Pi starvation in *Arabidopsis*. *Biochem J.* 2007;405:191–8.
- Devaiah BN, Karthikeyan AS, Raghothama KG. WRKY75 transcription factor is a modulator of phosphate acquisition and root development in *Arabidopsis*. *Plant Physiol.* 2007;143:1789–801.
- Devaiah BN, Nagarajan VK, Raghothama KG. Phosphate homeostasis and root development in *Arabidopsis* are synchronized by the zinc finger transcription factor ZAT6. *Plant Physiol.* 2007;145:147–59.
- Devaiah BN, Madhuvantri R, Karthikeyan AS, Raghothama KG. Phosphate starvation responses and gibberellic acid biosynthesis are regulated by the MYB62 transcription factor in *Arabidopsis*. *Mol Plant.* 2009;2:43–58.
- Chen YF, Li LQ, Xu Q, Kong YH, Wang H, Wu WH. The WRKY6 transcription factor modulates PHOSPHATE1 expression in response to low Pi stress in *Arabidopsis*. *Plant Cell.* 2009;21:3554–66.
- Wang H, Xu Q, Kong YH, Chen Y, Duan JY, Wu WH, Chen YF. *Arabidopsis* WRKY45 transcription factor activates PHOSPHATE TRANSPORTER1;1 expression in response to phosphate starvation. *Plant Physiol.* 2014;164:2020–9.
- Ramaiah M, Jain A, Raghothama KG. Ethylene Response Factor070 regulates root development and phosphate starvation-mediated responses. *Plant Physiol.* 2014;164:1484–98.
- Su T, Xu Q, Zhang FC, Chen Y, Li LQ, Wu WH, Chen YF. WRKY42 modulates phosphate homeostasis through regulating phosphate translocation and acquisition in *Arabidopsis*. *Plant Physiol.* 2015;167:1579–91.
- Chen CY, Wu K, Schmidt W. The histone deacetylase HDA19 controls root cell elongation and modulates a subset of phosphate starvation responses in *Arabidopsis*. *Sci Rep.* 2015;5:15708.

47. Pant BD, Burgos A, Pant P, Cuadros-Inostroza A, Willmitzer L, Scheible WR. The transcription factor PHR1 regulates lipid remodeling and triacylglycerol accumulation in *Arabidopsis thaliana* during phosphorus starvation. *J Exp Bot.* 2015;66:1907–18.
48. Fujii H, Chiou TZ, Lin S-I, Aung K, Zhu J-K. A miRNA involved in phosphate-starvation response in *Arabidopsis*. *Curr Biol.* 2005;15:2038–43.
49. Bari R, Pant BD, Stitt M, Scheible WR. PHO2, MicroRNA399, and PHR1 define a phosphate-signaling pathway in plants. *Plant Physiol.* 2006;141:988–99.
50. Aung K, Lin SI, Wu CC, Huang YT, Su CL, Chiou TZ. *pho2*, a phosphate overaccumulator, is caused by a nonsense mutation in a microRNA399 target gene. *Plant Physiol.* 2006;141:1000–11.
51. Chiou TJ, Aung K, Lin SI, Wu CC, Chiang SF, Sua CL. Regulation of phosphate homeostasis by microRNA in *Arabidopsis*. *Plant Cell.* 2006;18:412–21.
52. Lin SI, Chiang SF, Lin WY, Chen JW, Tseng CY, Wu PC, Chiou TJ. Regulatory network of microRNA399 and PHO2 by systemic signaling. *Plant Physiol.* 2008;147:732–46.
53. Hsieh LC, Lin SI, Shih AC, Chen JW, Lin WY, Tseng CY, Li WH, Chiou TJ. Uncovering small RNA-mediated responses to phosphate deficiency in *Arabidopsis* by deep sequencing. *Plant Physiol.* 2010;151:2120–32.
54. Miura K, Rus A, Sharkhuu A, Yokoi S, Karthikeyan AS, Raghothama KG, Baek D, Koo YD, Jin JB, Bressan RA, Yun DJ, Hasegawa PM. The *Arabidopsis* SUMO E3 ligase SIZ1 controls phosphate deficiency responses. *Proc Natl Acad Sci U S A.* 2005;102:7760–5.
55. Bayle V, Arrighi JF, Creff A, Nespoulous C, Vialaret J, Rossignol M, Gonzalez E, Paz-Ares J, Nussaume L. *Arabidopsis thaliana* high-affinity phosphate transporters exhibit multiple levels of posttranslational regulation. *Plant Cell.* 2011;23:1523–35.
56. Lei L, Li Y, Wang Q, Xu J, Chen Y, Yang H, et al. Activation of MKK9-MPK3/MPK6 enhances phosphate acquisition in *Arabidopsis thaliana*. *New Phytol.* 2014;203:1146–60.
57. Li WF, Perry PJ, Prafulla NN, Schmidt W. Ubiquitin-specific protease 14 (UBP14) is involved in root responses to phosphate deficiency in *Arabidopsis*. *Mol Plant.* 2010;3:212–23.
58. Yong-Villalobos L, González-Morales SI, Wrobel K, Gutiérrez-Alanis D, Cervantes-Peréz SA, Hayano-Kanashiro C, Oropeza-Aburto A, Cruz-Ramírez A, Martínez O, Herrera-Estrella L. Methylome analysis reveals an important role for epigenetic changes in the regulation of the *Arabidopsis* response to phosphate starvation. *Proc Natl Acad Sci U S A.* 2015;112:E7293–302.
59. López-Bucio J, Hernández-Abreu E, Sánchez-Calderón L, Pérez-Torres A, Rampey RA, Bartel B, Herrera-Estrella L. An auxin transport independent pathway is involved in phosphate stress-induced root architectural alterations in *Arabidopsis*. Identification of BIG as a mediator of auxin in pericycle cell activation. *Plant Physiol.* 2005;137:681–91.
60. Nacry P, Canivenc G, Muller B, Azmi A, Van Onckelen H, Rossignol M, Doumas P. A role for auxin redistribution in the responses of the root system architecture to phosphate starvation in *Arabidopsis*. *Plant Physiol.* 2005;138:2061–74.
61. Grieneisen VA, Xu J, Mar'ee AFM, Hogeweg P, Scheres B. Auxin transport is sufficient to generate a maximum and gradient guiding root growth. *Nature.* 2007;449:1008–13.
62. Martin AC, del Pozo JC, Iglesias J, Rubio V, Solano R, de la Pena A, Paz-Ares J. Influence of cytokinins on the expression of phosphate starvation responsive genes in *Arabidopsis*. *Plant J.* 2000;24:559–67.
63. Wang X, Yi K, Tao Y, Wang F, Wu Z, Jiang D, Chen X, Zhou L, Wu P. Cytokinin represses phosphate-starvation response through increasing of intracellular phosphate level. *Plant Cell Environ.* 2006;29:1924–35.
64. Ma Z, Baskin TI, Brown KM, Lynch JP. Regulation of root elongation under phosphorus stress involves changes in ethylene responsiveness. *Plant Physiol.* 2003;131:1381–90.
65. Zhang Y, Lynch JP, Brown K. Ethylene and phosphorus availability have interacting yet distinct effects on root hair development. *J Exp Bot.* 2003;54:2351–61.
66. Nagarajan VK, Smith AP. Ethylene's role in phosphate starvation signaling: more than just a root growth regulator. *Plant Cell Physiol.* 2012;53:277–86.
67. Jiang CF, Gao XH, Liao LL, Harberd NP, Fu XD. Phosphate starvation root architecture and anthocyanin accumulation responses are modulated by the gibberellin-DELLA signaling pathway in *Arabidopsis*. *Plant Physiol.* 2007;145:1460–70.
68. Veronese P, Nakagami H, Bluhm B, Abuqamar S, Chen X, Salmeron J, Dietrich RA, Hirt H, Mengiste T. The membrane-anchored BOTRYTIS-INDUCED KINASE1 plays distinct roles in *Arabidopsis* resistance to necrotrophic and biotrophic pathogens. *Plant Cell.* 2006;18:257–73.
69. Lu D, Wu S, Gao X, Zhang Y, Shan L, He P. A receptor-like cytoplasmic kinase, BIK1, associates with a flagellin receptor complex to initiate plant innate immunity. *Proc Natl Acad Sci U S A.* 2010;107:496–501.
70. Zhang J, Li W, Xiang T, Liu Z, Laluk K, Ding X, Dietrich RA, Hirt H, Mengiste T. Receptor-like cytoplasmic kinases integrate signaling from multiple plant immune receptors and are targeted by a *Pseudomonas syringae* effector. *Cell Host Microbe.* 2010;7:290–301.
71. Laluk K, Luo H, Chai M, Dhawan R, Lai Z, Mengiste T. Biochemical and genetic requirements for function of the immune response regulator BOTRYTIS-INDUCED KINASE1 in plant growth, ethylene signaling, and PAMP-triggered immunity in *Arabidopsis*. *Plant Cell.* 2011;23:2831–49.
72. Liu Z, Wu Y, Yang F, Zhang Y, Chen S, Xie Q, Tian X, Zhou JM. BIK1 interacts with PEPRs to mediate ethylene-induced immunity. *Proc Natl Acad Sci U S A.* 2013;110:6205–10.
73. Lin W, Lu D, Gao X, Jiang S, Ma X, Wang Z, Mengiste T, He P, Shan L. Inverse modulation of plant immune and brassinosteroid signaling pathways by the receptor-like cytoplasmic kinase BIK1. *Proc Natl Acad Sci U S A.* 2013;110:12114–9.
74. Lei J, A Finlayson S, Salzman RA, Shan L, Zhu-Salzman K. BOTRYTIS-INDUCED KINASE1 modulates *Arabidopsis* resistance to green peach aphids via PHYTOALEXIN DEFICIENT4. *Plant Physiol.* 2014;165:1657–70.
75. Péret B, Clément M, Nussaume L, Desnos T. Root developmental adaptation to phosphate starvation: better safe than sorry. *Trends Plant Sci.* 2011;16:442–50.
76. Mittler R, Vanderauwera S, Gollery M, Van Breusegem F. Reactive oxygen gene network of plants. *Trends Plant Sci.* 2004;9:490–8.
77. Foyer CH, Noctor G. Redox homeostasis and antioxidant signaling: a metabolic interface between stress perception and physiological responses. *Plant Cell.* 2005;17:1866–75.
78. Shin H, Shin HS, Chen R, Harrison MJ. Loss of *At4* function impacts phosphate distribution between the roots and the shoots during phosphate starvation. *Plant J.* 2006;45:712–26.
79. Misson J, Raghothama KG, Jain A, Jouhet J, Block MA, Bligny R, Ortet P, Creff A, Somerville S, Rolland N, et al. A genome-wide transcriptional analysis using *Arabidopsis thaliana* Affymetrix gene chips determined plant responses to phosphate deprivation. *Proc Natl Acad Sci U S A.* 2005;102:11934–9.
80. Lan P, Li W, Schmidt W. Genome-wide co-expression analysis predicts protein kinases as important regulators of phosphate deficiency-induced root hair remodeling in *Arabidopsis*. *BMC Genomics.* 2013;14:210.
81. Gregory A, Hurlley BA, Tran HT, Valentine AJ, She YM, Knowles VL, Plaxton WC. In vivo regulatory phosphorylation of the phosphoenolpyruvate carboxylase AtPPC1 in phosphate-starved *Arabidopsis thaliana*. *Biochem J.* 2009;420:57–65.
82. Sanchez-Calderon L, Lopez-Bucio J, Chacon-Lopez A, Cruz-Ramirez A, Nieto-Jacobo F, Dubrovsky JG, Herrera-Estrella L. Phosphate starvation induces a determinate development program in root of *Arabidopsis thaliana*. *Plant Cell Physiol.* 2005;46:174–84.
83. Plaxton WC, Tran HT. Metabolic adaptations of phosphate starved plants. *Plant Physiol.* 2011;156:1006–15.
84. Hurlley BA, Tran HT, Marty NJ, Park J, Snedden WA, Mullen RT, Mullen RT, Plaxton WC. The dual-targeted purple acid phosphatase isozyme AtPAP26 is essential for efficient acclimation of *Arabidopsis* to nutritional phosphate deprivation. *Plant Physiol.* 2010;153:1112–22.
85. Robinson WD, Park J, Tran HT, Del Vecchio HA, Ying S, Zins JL, Patel K, McKnight TD, Plaxton WC. The secreted purple acid phosphatase isozymes AtPAP12 and AtPAP26 play a pivotal role in extracellular phosphate-scavenging by *Arabidopsis thaliana*. *J Exp Bot.* 2012;63:6531–42.
86. Del Vecchio HA, Ying S, Park J, Knowles VL, Kanno S, Tanoi K, et al. The cell wall-targeted purple acid phosphatase AtPAP25 is critical for acclimation of *Arabidopsis thaliana* to nutritional phosphorus deprivation. *Plant J.* 2014;80:569–81.
87. Tran HT, Plaxton WC. Proteomic analysis of alterations in the secretome of *Arabidopsis thaliana* suspension cells subjected to nutritional phosphate deficiency. *Proteomics.* 2008;8:4317–26.
88. Tran HT, Qian W, Hurlley BA, She YM, Wang D, Plaxton WC. Biochemical and molecular characterization of AtPAP12 and AtPAP26: the predominant

- purple acid phosphatase isozymes secreted by phosphate-starved *Arabidopsis thaliana*. *Plant Cell Environ.* 2010;33:1789–803.
89. Torres MA. ROS in biotic interactions. *Physiol Plant.* 2010;138:414–29.
 90. Suzuki N, Koussevitzky S, Mittler R, Miller G. ROS and redox signalling in the response of plants to abiotic stress. *Plant Cell Environ.* 2012;35:259–70.
 91. Baxter A, Mittler R, Suzuki N. ROS as key players in plant stress signalling. *J Exp Bot.* 2013;65:1229–40.
 92. Shin R, Berg RH, Schachtman DP. Reactive oxygen species and root hairs in *Arabidopsis* root response to nitrogen, phosphorus and potassium deficiency. *Plant Cell Physiol.* 2005;46:1350–7.
 93. Foreman J, Demidchik V, Bothwell JH, Mylona P, Miedema H, Torres MA, Linstead P, Costa S, Brownlee C, Jones JD, Davies JM, Dolan L. Reactive oxygen species produced by NADPH oxidase regulate plant cell growth. *Nature.* 2003;422:442–6.
 94. Rentel MC, Lecourieux D, Ouaked F, Usher SL, Petersen L, Okamoto H, Knight H, Peck SC, Grierson CS, Hirt H, Knight MR. OX11 kinase is necessary for oxidative burst-mediated signalling in *Arabidopsis*. *Nature.* 2004;427:858–61.
 95. Tyburski J, Dunajska K, Tretyn A. Reactive oxygen species localization in roots of *Arabidopsis thaliana* seedlings under phosphate deficiency. *Plant Growth Reg.* 2009;59:27–36.
 96. Chacón-López A, Ibarra-Laclette E, Sánchez-Calderón L, Gutiérrez-Alanis D, Herrera-Estrella L. Global expression pattern comparison between *low phosphorus insensitive 4* and WT *Arabidopsis* reveals an important role of reactive oxygen species and jasmonic acid in the root tip response to phosphate starvation. *Plant Signal Behav.* 2011;6:382–92.
 97. Kadota Y, Sklenar J, Derbyshire P, Stransfeld L, Asai S, Ntoukakis V, Jones JD, Shirasu K, Menke F, Jones A, Zipfel C. Direct regulation of the NADPH oxidase RBOHD by the PRR-associated kinase BIK1 during plant immunity. *Mol Cell.* 2014;54:43–55.
 98. Li L, Li M, Yu L, Zhou Z, Liang X, Liu Z, Jones JD, Shirasu K, Menke F, Jones A, Zipfel C. The FLS2-associated kinase BIK1 directly phosphorylates the NADPH oxidase RbohD to control plant immunity. *Cell Host Microbe.* 2014;15:329–38.
 99. Karthikeyan AS, Varadarajan DK, Mukatira UT, D'Urzo MP, Damz B, Raghothama KG. Regulated expression of *Arabidopsis* phosphate transporters. *Plant Physiol.* 2002;130:221–33.
 100. Lange H, Shropshire W, Mohr H. An analysis of phytochrome-mediated anthocyanin synthesis. *Plant Physiol.* 1971;47:649–55.
 101. Richardson AE, Hadobas PA, Hayes JE. Extracellular secretion of *Aspergillus* phytase from *Arabidopsis* roots enables plants to obtain phosphorus from phytate. *Plant J.* 2001;25:641–9.
 102. Tomscha JL, Trull MC, Deikman J, Lynch JP, Guitinan MJ. Phosphatase under-producer mutants have altered phosphorus relations. *Plant Physiol.* 2004;135:334–45.
 103. Trull MC, Deikman J. An *Arabidopsis* mutant missing one acid phosphatase isoform. *Planta.* 1998;206:544–50.
 104. Doke N. Involvement of superoxide anion generation in the hypersensitive response of potato tuber tissue to infection with an incompatible race of *Phytophthora infestans* and to the hyphal wall components. *Physiol Plant Pathol.* 1983;23:345–57.
 105. Jefferson RA, Kavanagh TA, Bevan MW. GUS fusions: beta-glucuronidase as a sensitive and versatile gene fusion marker in higher plants. *EMBO J.* 1987;6:3901–7.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

