



PUBLISHED FOR SISSA BY SPRINGER

RECEIVED: June 26, 2012

ACCEPTED: August 8, 2012

PUBLISHED: August 23, 2012

S-duality in twistor space

Sergei Alexandrov^{a,b} and Boris Pioline^{c,d}

^a *Université Montpellier 2, Laboratoire Charles Coulomb UMR 5221,
F-34095, Montpellier, France*

^b *CNRS, Laboratoire Charles Coulomb UMR 5221,
F-34095, Montpellier, France*

^c *CERN PH-TH, Case C01600, CERN,
CH-1211 Geneva 23, Switzerland*

^d *Laboratoire de Physique Théorique et Hautes Energies, CNRS UMR 7589,
Université Pierre et Marie Curie, 4 place Jussieu, 75252 Paris cedex 05, France*

E-mail: salexand@univ-montp2.fr, boris.pioline@cern.ch

ABSTRACT: In type IIB string compactifications on a Calabi-Yau threefold, the hypermultiplet moduli space \mathcal{M}_H must carry an isometric action of the modular group $SL(2, \mathbb{Z})$, inherited from the S-duality symmetry of type IIB string theory in ten dimensions. We investigate how this modular symmetry is realized at the level of the twistor space of \mathcal{M}_H , and construct a general class of $SL(2, \mathbb{Z})$ -invariant quaternion-Kähler metrics with two commuting isometries, parametrized by a suitably covariant family of holomorphic transition functions. This family should include \mathcal{M}_H corrected by D3-D1-D(-1)-instantons (with five-brane corrections ignored) and, after taking a suitable rigid limit, the Coulomb branch of five-dimensional $\mathcal{N} = 2$ gauge theories compactified on a torus, including monopole string instantons. These results allow us to considerably simplify the derivation of the mirror map between type IIA and IIB fields in the sector where only D1-D(-1)-instantons are retained.

KEYWORDS: D-branes, Superstring Vacua, Supersymmetric Effective Theories

ARXIV EPRINT: [1206.1341](https://arxiv.org/abs/1206.1341)

Contents

1	Introduction	1
2	Hypermultiplet moduli space in CY string vacua	3
2.1	Classical metric	4
2.2	Quantum corrections	6
2.3	Twistorial description of \mathcal{M}_H	7
3	S-duality and D1-D(-1)-instantons	9
3.1	S-duality in the classical twistor space	10
3.2	Type IIB twistorial construction of D1-D(-1)-instanton corrections	11
3.3	The key observations	12
3.4	The quantum mirror map revisited	14
3.5	Equivalence between IIA and IIB constructions	16
4	S-duality in twistor space with two isometries	17
5	Discussion	22
A	Twistorial description of quaternion-Kähler manifolds	23
B	S-duality invariance at linear order	25

1 Introduction

String vacua with $\mathcal{N} = 2$ supersymmetry in four dimensions offer a fascinating vantage point on the non-perturbative spectrum and symmetries of string theory. In the low energy approximation, the dynamics around these vacua is described by the metric on the moduli space parametrized by the massless scalar fields arising after compactification from ten to four dimensions. This space factorizes into the direct product of the vector multiplet (VM) moduli space \mathcal{M}_V and the hypermultiplet (HM) moduli space \mathcal{M}_H [1], which encode distinct properties of the internal manifold. Whereas the geometry of \mathcal{M}_V has been under control for many years, a complete description of the geometry of \mathcal{M}_H at the non-perturbative level is still missing. This is in part due to the fact that the metric on \mathcal{M}_H must be quaternion-Kähler (QK) [2], a property which is far more complicated to enforce than special Kähler geometry in the vector multiplet sector.

Nevertheless, remarkable progress was achieved in recent years in the context of type II strings compactified on a Calabi-Yau (CY) threefold, based on the development of twistorial methods for quaternion-Kähler geometry (see [3] for an extensive review). This approach was developed in the mathematical literature in [4–6] along with projective superspace

methods in the physics literature [7–10], and further tailored for string theory applications in [11, 12]. By lifting the QK space to its twistor space, a $\mathbb{C}P^1$ bundle endowed with canonical complex and contact structures, this method provides an efficient parametrization of the quaternion-Kähler metric in terms of a set of holomorphic transition functions between local Darboux coordinate systems, which play a role similar to the holomorphic prepotential of special Kähler geometry. Using this formalism, it was understood how to describe various quantum corrections to the geometry of \mathcal{M}_H consistently with supersymmetry, starting with D1-D(-1)-instantons [13], further including all D-brane instantons at the fully non-linear level¹ [14, 15] and finally, NS5-brane instantons in the linear approximation [16] (see also [17–24] for closely related work). These advances took place in parallel with related developments in the context of rigid $\mathcal{N} = 2$ field theories [25–28].

Unlike the calculation of the one-loop correction to \mathcal{M}_H , which was found by a scattering amplitude calculation [29–31], the above results on instanton corrections were obtained by postulating certain symmetries and dualities. Among those, the isometric action of the modular group $SL(2, \mathbb{Z})$, inherited from S-duality of type IIB string theory in ten dimensions, played a central role [13, 16]. Nevertheless, it has not been verified so far that the type IIA construction of the D-instanton corrections in [14, 15] was in fact consistent with S-duality of the mirror formulation. In particular, in the limit where only D(-1), D1 and D3 instantons are retained, the metric on \mathcal{M}_H must be modular invariant, which provides a non-trivial constraint on the generalized Donaldson-Thomas invariants governing such D-instantons.

One of the difficulties in demonstrating S-duality is an inherent ambiguity in the twistor construction, namely the fact that the Darboux coordinates are only defined up to a local complex contact transformation. Thus, while any isometry of \mathcal{M}_H can always be lifted to a holomorphic action in twistor space, the Darboux coordinates need only be covariant up to a local complex contact transformation. Thus, the condition for S-duality invariance is that the set of holomorphic transition functions should commute with S-duality, up to local complex transformations in each patch.

In this work we give a solution to this problem and provide a twistorial construction of a general class of QK metrics preserving two continuous commuting isometries, which is explicitly $SL(2, \mathbb{Z})$ -invariant and parametrized by a suitably covariant family of transition functions $G_{m,n}$. Physically, the two isometries correspond to the unbroken Peccei-Quinn symmetries in the absence of D5 and NS5-brane instantons. For vanishing $G_{m,n}$, these metrics reduce to the local c-map metrics [32, 33], which arise in the weak coupling, large volume limit of type IIB string theory. Thus, this class of the metrics should contain the quantum corrected HM moduli space in the limit where fivebrane instantons are exponentially suppressed compared to D(-1), D1, D3 instantons. After taking a rigid limit, this class should also contain the hyperkähler metric on the Coulomb branch of five-dimensional $\mathcal{N} = 2$ gauge theories compactified on a torus T^2 , where the modular group of the torus

¹The holomorphic transition functions specified in [14, 15] uniquely determine the metric on \mathcal{M}_H , provided the generalized Donaldson-Thomas invariants of the CY threefold are known. It is in general a formidable task to compute these invariants.

plays the role of S-duality. Thus, this construction should provide constraints on the contributions of the monopole string instantons considered in [27].

To motivate this construction, we first reconsider the much simpler case of $SL(2, \mathbb{Z})$ -invariant toric QK spaces, i.e. $4n$ -dimensional spaces \mathcal{M}_H with $n+1$ commuting isometries. Physically, this case arises when only D1-D(-1)-contributions (or equivalently (p, q) -string instantons) are retained and, just like the undeformed c -map metric, is amenable to the Legendre transform construction. This case was studied in detail in [34], where a set of *ad hoc* local contact transformations was designed that bring the general type IIA construction of [14] into a manifestly $SL(2, \mathbb{Z})$ -invariant form. This invariance was then exploited in order to obtain instanton corrections to the ‘mirror map’, relating the type IIB fields (covariant under S-duality) with the type IIA fields (covariant under monodromies). However, the geometric meaning of both the contact transformation and the quantum mirror map, had remained unclear. Here we illuminate the *ad hoc* construction of [34], exposing the inner workings of S-duality in the twistor space, and considerably streamlining the derivation of the mirror map.

Having recast the construction of [34] in a manifestly S-duality invariant fashion, the extension to S-duality invariant QK metrics with only two commuting isometries is then relatively straightforward, despite the fact that such metrics are no longer amenable to the Legendre transform construction. The key ingredient is the non-linear condition (4.4) on the holomorphic transition functions $G_{m,n}$, which ensures the $SL(2, \mathbb{Z})$ -invariance of the QK metric. Identifying the functions $G_{m,n}$ which are relevant in the context of D3-brane instanton corrections to the the HM moduli space lies beyond the scope of this paper. In subsequent work [35] we provide a detailed analysis of the modular properties of the type IIA twistorial construction of [14], uncovering how S-duality is realized in this framework.

The organization of the paper is as follows. In section 2 we provide a brief review of the HM moduli space both at classical and quantum level. In particular, we explain how its QK geometry, corrected by D2-instantons of type IIA theory, is encoded in the twistor data consisting of a covering of the twistor fiber and an associated set of holomorphic transition functions. Then in section 3 we study the realization of S-duality in the dual type IIB formulation keeping contributions only from D1-D(-1)-instantons. By refining the construction of [34], we demonstrate how S-duality and the quantum mirror map follow from the simple transformation property (3.15) of the transition functions encoding instanton corrections. In section 4, we turn to the more general case of QK spaces with two commuting isometries, and provide sufficient conditions on the covering of the twistor sphere and holomorphic transition functions which ensure the $SL(2, \mathbb{Z})$ -invariance of the metric. We conclude with some open questions in section 5. Two appendices contain a review of the twistorial description of QK spaces and a proof of the S-duality invariance of the construction of section 4 in the linear approximation.

2 Hypermultiplet moduli space in CY string vacua

In this section, we briefly recall some basic facts about the hypermultiplet moduli space \mathcal{M}_H in type II string theory compactified on a CY threefold. Since mirror symmetry

identifies type IIA compactified on a Calabi-Yau threefold \mathfrak{Y} and type IIB compactified on the mirror threefold $\hat{\mathfrak{Y}}$, the moduli space \mathcal{M}_H has two equivalent, dual descriptions. On either side, it governs the dynamics of²

1. the four-dimensional dilaton $r = e^\phi = 1/g_4^2$ or its ten-dimensional cousin g_s ;
2. the complex moduli $z^a = b^a + it^a$ parametrizing the moduli space $\mathcal{K}_C(\mathfrak{Y})$ of complex structures on \mathfrak{Y} or the complexified Kähler structure of $\mathcal{K}_K(\hat{\mathfrak{Y}})$ — these two spaces also describe the VM moduli space of the dual type II theory compactified on the same CY threefold, and are identified by classical mirror symmetry;
3. the RR scalars $\zeta^\Lambda, \tilde{\zeta}_\Lambda$ or $c^0, c^a, \tilde{c}_a, \tilde{c}_0$ obtained as integrals of RR gauge potentials over a symplectic basis of cycles in $H_3(\mathfrak{Y}, \mathbb{Z})$ or $H_{\text{even}}(\hat{\mathfrak{Y}}, \mathbb{Z})$;
4. the NS axion σ or ψ , dual to the NS 2-form B in four dimensions.

On the type IIB side, the ten-dimensional string coupling $\tau_2 \equiv 1/g_s$ and the RR axion $\tau_1 \equiv c^0$ combine into the ten-dimensional axio-dilaton field $\tau = \tau_1 + i\tau_2$. As a result, the field basis $(\tau, b^a, t^a, c^a, \tilde{c}_a, \tilde{c}_0, \psi)$ has simple transformation rules under S-duality. In contrast, the field basis $(\phi, z^a, \zeta^\Lambda, \tilde{\zeta}_\Lambda, \sigma)$ transform naturally under monodromies, which are manifest on the type IIA side as they correspond to a change of symplectic basis in $H_3(\mathfrak{Y}, \mathbb{Z})$. Thus, we refer to the latter field basis as the type IIA coordinates, and to the former as the type IIB coordinates.

2.1 Classical metric

Supersymmetry constrains the moduli space \mathcal{M}_H to carry a quaternion-Kähler metric with negative scalar curvature [2]. At classical level the metric on \mathcal{M}_H can be obtained from the moduli space $\mathcal{K}_C(\mathfrak{Y})$ of complex structures on \mathfrak{Y} , or equivalently the moduli space $\mathcal{K}_K(\hat{\mathfrak{Y}})$ of Kähler structures on $\hat{\mathfrak{Y}}$, via the local c -map construction [32, 33]. The space $\mathcal{K}_C(\mathfrak{Y}) = \mathcal{K}_K(\hat{\mathfrak{Y}})$ is a special Kähler manifold characterized by a holomorphic prepotential $F(X^\Lambda)$, homogeneous of degree 2 in the special complex coordinates X^Λ such that $X^a/X^0 = z^a = b^a + it^a$. Thus, the classical metric is completely determined by the prepotential and is given by [33]

$$\begin{aligned}
 ds_{\text{cl}}^2 = & \frac{dr^2}{r^2} + 4ds_{\mathcal{S}\mathcal{K}}^2 - \frac{1}{2r} \left(d\tilde{\zeta}_\Lambda - \bar{\mathcal{N}}_{\Lambda\Sigma} d\zeta^\Sigma \right) \text{Im} \mathcal{N}^{\Lambda\Lambda'} \left(d\tilde{\zeta}_{\Lambda'} - \mathcal{N}_{\Lambda'\Sigma'} d\zeta^{\Sigma'} \right) \\
 & + \frac{1}{16r^2} \left(d\sigma + \tilde{\zeta}_\Lambda d\zeta^\Lambda - \zeta^\Lambda d\tilde{\zeta}_\Lambda \right)^2,
 \end{aligned}
 \tag{2.1}$$

where $ds_{\mathcal{S}\mathcal{K}}^2$ is the metric on $\mathcal{K}_C(\mathfrak{Y}) = \mathcal{K}_K(\hat{\mathfrak{Y}})$ with Kähler potential $\mathcal{K} = -\log[i(\bar{X}^\Lambda F_\Lambda - X^\Lambda \bar{F}_\Lambda)]$,

$$\mathcal{N}_{\Lambda\Lambda'} = \bar{\tau}_{\Lambda\Lambda'} + 2i \frac{[\text{Im} \tau \cdot X]_\Lambda [\text{Im} \tau \cdot X]_{\Lambda'}}{X^\Sigma \text{Im} \tau_{\Sigma\Sigma'} X^{\Sigma'}}, \quad \tau_{\Lambda\Sigma} \equiv \partial_{X^\Lambda} \partial_{X^\Sigma} F(X).
 \tag{2.2}$$

²Our conventions are such that the indices Λ, Σ, \dots (resp. a, b, \dots) run from 0 (resp. 1) till $h^{2,1}(\mathfrak{Y}) = h^{1,1}(\hat{\mathfrak{Y}})$.

Here we expressed the metric in terms of the type IIA fields where it takes an especially simple form. In particular, it is explicitly invariant under symplectic transformations and the continuous shifts generating the Heisenberg algebra

$$T_{\eta^\Lambda, \tilde{\eta}_\Lambda, \kappa} : (\zeta^\Lambda, \tilde{\zeta}_\Lambda, \sigma) \mapsto (\zeta^\Lambda + \eta^\Lambda, \tilde{\zeta}_\Lambda + \tilde{\eta}_\Lambda, \sigma + 2\kappa - \tilde{\eta}_\Lambda \zeta^\Lambda + \eta^\Lambda \tilde{\zeta}_\Lambda). \quad (2.3)$$

Moreover, the symplectic vector (X^Λ, F_Λ) is identified as the integral of the holomorphic three-form on \mathfrak{Y} along a symplectic basis in $H_3(\mathfrak{Y}, \mathbb{Z})$, while $\mathcal{N}_{\Lambda\Sigma}$ and $\tau_{\Lambda\Sigma}$ are the Weil and Griffiths period matrices on the intermediate Jacobian $H^3(\mathfrak{Y}, \mathbb{R})/H^3(\mathfrak{Y}, \mathbb{Z})$, respectively.

To rewrite the metric (2.1) in type IIB variables, we use the fact that the holomorphic prepotential can be written, in the limit where the volume of the mirror threefold $\hat{\mathfrak{Y}}$ is infinite, as³

$$F^{\text{cl}}(X^\Lambda) = -\kappa_{abc} \frac{X^a X^b X^c}{6X^0}, \quad (2.4)$$

where κ_{abc} is the triple intersection product in $H_4(\hat{\mathfrak{Y}}, \mathbb{Z})$. Moreover, the type IIA fields are related to the type IIB ones via the following classical ‘mirror map’ [36]

$$\begin{aligned} r &= \frac{\tau_2^2}{2} \mathcal{V}, & z^a &= b^a + it^a, & \zeta^0 &= \tau_1, & \zeta^a &= -(c^a - \tau_1 b^a), \\ \tilde{\zeta}_a &= \tilde{c}_a + \frac{1}{2} \kappa_{abc} b^b (c^c - \tau_1 b^c), & \tilde{\zeta}_0 &= \tilde{c}_0 - \frac{1}{6} \kappa_{abc} b^a b^b (c^c - \tau_1 b^c), \\ \sigma &= -2(\psi + \frac{1}{2} \tau_1 \tilde{c}_0) + \tilde{c}_a (c^a - \tau_1 b^a) - \frac{1}{6} \kappa_{abc} b^a c^b (c^c - \tau_1 b^c), \end{aligned} \quad (2.5)$$

where \mathcal{V} is the volume of the threefold $\hat{\mathfrak{Y}}$ in string units,

$$\mathcal{V} = \frac{1}{6} \int_{\hat{\mathfrak{Y}}} J \wedge J \wedge J = \frac{1}{6} \kappa_{abc} t^a t^b t^c. \quad (2.6)$$

The virtue of the coordinate transformation (2.5) is to make manifest the invariance of the metric (2.1) under the action of $\text{SL}(2, \mathbb{Z})$ [30, 36]⁴

$$\begin{aligned} \tau &\mapsto \frac{a\tau + b}{c\tau + d}, & t^a &\mapsto t^a |c\tau + d|, & \tilde{c}_a &\mapsto \tilde{c}_a, \\ \begin{pmatrix} c^a \\ b^a \end{pmatrix} &\mapsto \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} c^a \\ b^a \end{pmatrix}, & \begin{pmatrix} \tilde{c}_0 \\ \psi \end{pmatrix} &\mapsto \begin{pmatrix} d & -c \\ -b & a \end{pmatrix} \begin{pmatrix} \tilde{c}_0 \\ \psi \end{pmatrix}, & ad - bc &= 1, \end{aligned} \quad (2.7)$$

which corresponds to the S-duality symmetry of type IIB supergravity in ten dimensions. The metric (2.1) is in fact invariant under the continuous action of $\text{SL}(2, \mathbb{R})$, but as we now recall, this invariance is broken by quantum corrections.

³In general, the cubic prepotential (2.4) must be supplemented by a quadratic term $\frac{1}{2} A_{\Lambda\Sigma} X^\Lambda X^\Sigma$ which is necessary for consistency with charge quantization. This term can always be removed by working in a non-integer symplectic basis, which we assume in this work.

⁴In [16] it was found that the transformation of \tilde{c}_a must contain an additional constant shift proportional to the second Chern class $c_{2,a}$ of $\hat{\mathfrak{Y}}$, which is necessary for consistency with charge quantization. This subtlety is closely related to the quadratic correction to the prepotential mentioned in the previous footnote, and can be safely ignored for the purposes of this paper.

2.2 Quantum corrections

Away from the classical, large volume limit, the HM moduli space receives two types of quantum corrections. The first type, namely α' -corrections, occur only in the type IIB formulation. These effects preserve the c -map form (2.1) of the metric, and simply correct the holomorphic prepotential (2.4) into $F = F^{\text{cl}} + F^{\text{w.s.}}$ where [37, 38]

$$F^{\text{w.s.}}(X^\Lambda) = \chi_{\hat{\mathfrak{Y}}} \frac{\zeta(3)(X^0)^2}{2(2\pi i)^3} - \frac{(X^0)^2}{(2\pi i)^3} \sum_{q_a \gamma^a \in H_2^+(\hat{\mathfrak{Y}})} n_{q_a}^{(0)} \text{Li}_3 \left(e^{2\pi i q_a X^a / X^0} \right). \quad (2.8)$$

The first term, proportional to the Euler number $\chi_{\hat{\mathfrak{Y}}}$ of $\hat{\mathfrak{Y}}$, corresponds to a perturbative correction in the α' -expansion around the large volume limit. The second term corresponds to a sum of worldsheet instantons, labeled by their effective homology class $q_a \gamma^a \in H_2^+(\hat{\mathfrak{Y}}, \mathbb{Z})$ (i.e. $q_a \in \mathbb{Z}^+$ for all a , not all q_a 's vanishing simultaneously), and weighted by the genus zero Gopakumar-Vafa invariants $n_{q_a}^{(0)} \in \mathbb{Z}$. These effects contribute through the tri-logarithm function $\text{Li}_3(x) = \sum_{m=1}^{\infty} m^{-3} x^m$, which takes into account multi-covering effects. Note that the last two terms in (2.8) may be combined by including the zero class $q_a = 0$ in the sum and setting $n_0^{(0)} = -\chi_{\hat{\mathfrak{Y}}}/2$. The metric on \mathcal{M}_H including the α' -corrections is expressed as in (2.1) where F denotes now the full prepotential.

In contrast, the second type of quantum corrections, namely corrections in the string coupling g_s , take the metric outside the class of c -map metrics. At the perturbative level, it can be argued that the only non-trivial correction occurs at one-loop, and is determined solely by the Euler number $\chi_{\mathfrak{Y}} = -\chi_{\hat{\mathfrak{Y}}}$ [12, 29, 30, 39]. Its effect on the metric was explicitly evaluated in [39, 40], and will be described in twistorial terms in section 2.3.

At the non-perturbative level, the corrections to \mathcal{M}_H split again in two types. The first type corresponds, on the type IIA side, to D2-branes wrapping special Lagrangian cycles in \mathfrak{Y} , or, on the type IIB side, to D5-D3-D1-D(-1)-branes wrapping algebraic cycles of the corresponding dimension (or more generally, coherent sheaves on $\hat{\mathfrak{Y}}$). In either case, the correction takes the schematic form [41]

$$\delta ds^2|_{\text{D2}} \sim \Omega(\gamma, z^a) e^{-2\pi|Z_\gamma|/g_s - 2\pi i(q_\Lambda \zeta^\Lambda - p^\Lambda \tilde{\zeta}_\Lambda)}, \quad (2.9)$$

where the charge vector $\gamma = (p^\Lambda, q_\Lambda) = (p^0, p^a, q_a, q_0)$ takes value in $H_3(\mathfrak{Y}, \mathbb{Z})$ or $H_{\text{even}}(\hat{\mathfrak{Y}}, \mathbb{Z})$, the central charge function ($z^0 \equiv 1$)

$$Z_\gamma(z^a) = q_\Lambda z^\Lambda - p^\Lambda F_\Lambda(z^a) \quad (2.10)$$

measures the area of the supersymmetric cycle in the homology class γ , and $\Omega(\gamma, z^a)$ is the corresponding Donaldson-Thomas invariant, which counts (with signs) the number of supersymmetric cycles with charge γ . Due to their dependence on the RR axions $\zeta^\Lambda, \tilde{\zeta}_\Lambda$, or equivalently $\tau_1, c^a, \tilde{c}_a, \tilde{c}_0$, the D-instanton effects (2.9) break the continuous Heisenberg symmetries (2.3) to a discrete subgroup. In the large volume limit on the type IIB side, this breaking occurs in a hierarchical fashion, namely translations along τ_1 are broken at leading order, translations along c^a at order $\mathcal{O}(e^{-\nu^{1/3}})$, translations along \tilde{c}_a at order $\mathcal{O}(e^{-\nu^{2/3}})$ and finally translations along \tilde{c}_0 at order $\mathcal{O}(e^{-\nu})$.

Finally, the last type of corrections is generated by Euclidean NS5-branes wrapping the whole CY threefold. They contribute to the metric schematically as [41]

$$\delta ds^2|_{\text{NS5}} \sim e^{-2\pi|k|\mathcal{V}/g_s^2 - i\pi k\sigma}, \tag{2.11}$$

and break the last continuous isometry along σ (or ψ) which remained after inclusion of all D-instanton effects. Just like D5-branes, they are suppressed as $\mathcal{O}(e^{-\mathcal{V}})$ in the large volume limit.

What is the fate of S-duality at the quantum level? Firstly, the continuous $\text{SL}(2, \mathbb{R})$ symmetry (2.7) is certainly broken by the α' -corrections in (2.8). However, this symmetry descends from the usual S-duality action in ten-dimensional string theory, and there is overwhelming evidence that the modular subgroup $\text{SL}(2, \mathbb{Z})$ remains a symmetry at the quantum level (see [36, 42] and citations thereof). It is a priori unclear whether this symmetry should remain intact after compactification on a Calabi-Yau three-fold $\hat{\mathfrak{Y}}$, but this is supported by the fact that upon compactifying on S^1 , T-dualizing to type IIA string theory and lifting to M-theory, this action becomes the usual large diffeomorphisms of the torus in M-theory on $\hat{\mathfrak{Y}} \times T^2$. Thus, it is natural to assume that quantum corrections should preserve an isometric action of the discrete group $\text{SL}(2, \mathbb{Z})$.

Furthermore, from (2.7) it is possible to distinguish two sectors which should be preserved by S-duality. The first one is obtained by setting all fivebrane charges to zero, i.e. ignoring effects of order $\mathcal{O}(e^{-\mathcal{V}})$ in the large volume limit, leaving only D3-D1-F1-D(-1)-instantons. As noted above, in this approximation the moduli space has two continuous isometries along ψ and \tilde{c}_0 . The second sector arises by further setting the D3-brane charge $p^a = 0$, i.e. ignoring effects of order $\mathcal{O}(e^{-\mathcal{V}^{2/3}})$ or smaller in the large volume limit, leaving only D1-F1-D(-1)-instantons. In this limit the number of isometries is increased to $h^{1,1}(\hat{\mathfrak{Y}}) + 2 = n + 1$, where $4n$ is the real dimension of \mathcal{M}_H . The aim of this paper is to provide a manifestly S-duality invariant description of these two sectors.

2.3 Twistorial description of \mathcal{M}_H

To describe instanton corrections to the classical HM geometry, consistently with supersymmetry, it is indispensable to use the twistor description of QK spaces, very briefly summarized in appendix A. However, before going to the non-perturbative physics, one should understand how the perturbative moduli space is encoded in this formalism. This has been understood in [12, 14], based on the previous results of [17, 43].

As explained in appendix A, the twistor approach allows to encode a QK metric in terms of a set of holomorphic functions $H^{[ij]}$. They describe contact transformations between local Darboux coordinate systems for the complex contact one-form (A.2), which are attached to various patches \mathcal{U}_i of an open covering of the twistor fiber $\mathbb{C}P^1$. To describe the perturbative metric on \mathcal{M}_H , it is sufficient to cover $\mathbb{C}P^1$ by two patches \mathcal{U}_+ , \mathcal{U}_- centered around the north and south poles, $t = 0$ and $t = \infty$, and a third patch⁵ \mathcal{U}_0 which surrounds the equator. The transition functions between these patches are given in terms

⁵The patch \mathcal{U}_0 could in principle be omitted but is very convenient for exposing symplectic invariance.

of the holomorphic prepotential $F(X^\Lambda)$ by

$$H^{[+0]} = F(\xi^\Lambda), \quad H^{[-0]} = \bar{F}(\xi^\Lambda). \quad (2.12)$$

Using (A.4), we find that the Darboux coordinates in the patch \mathcal{U}_0 are given by [12, 43]

$$\begin{aligned} \xi^\Lambda &= \zeta^\Lambda + \frac{1}{2} \tau_2 (t^{-1} z^\Lambda - t \bar{z}^\Lambda), \\ \tilde{\xi}_\Lambda &= \tilde{\zeta}_\Lambda + \frac{1}{2} \tau_2 (t^{-1} F_\Lambda - t \bar{F}_\Lambda), \\ \tilde{\alpha} &= \sigma + \frac{1}{2} \tau_2 (t^{-1} W - t \bar{W}) + \frac{i\chi\mathfrak{y}}{24\pi} \log t, \end{aligned} \quad (2.13)$$

where W denotes the ‘superpotential’

$$W(z) = F_\Lambda(z) \zeta^\Lambda - z^\Lambda \tilde{\zeta}_\Lambda. \quad (2.14)$$

The last, logarithmic term in $\tilde{\alpha}$ encodes the effect of the one-loop g_s -correction and corresponds to an anomalous dimension $c = \chi\mathfrak{y}/(96\pi)$ in the language of appendix A. In (2.13), we have traded the Darboux coordinate α from appendix A for $\tilde{\alpha} \equiv -2\alpha - \tilde{\xi}_\Lambda \xi^\Lambda$, and the integration constants $A^\Lambda, B_\Lambda, B_\alpha$ for

$$\zeta^\Lambda = A^\Lambda, \quad \tilde{\zeta}_\Lambda = B_\Lambda + \text{Re} F_{\Lambda\Sigma} A^\Sigma, \quad \sigma = -2B_\alpha - A^\Lambda B_\Lambda. \quad (2.15)$$

Moreover, the contact potential Φ computed from (A.5) is related to the four- and ten-dimensional string couplings by

$$r = e^\Phi = \frac{\tau_2^2}{16} e^{-\mathcal{K}} + \frac{\chi\mathfrak{y}}{192\pi}, \quad (2.16)$$

which generalizes the first relation in (2.5).

The way to incorporate D-instanton corrections to the above twistor formulation of \mathcal{M}_H was explained in [14, 15], in close analogy with the field theory construction of [25]. We shall refer to this construction as the ‘type IIA construction’, as it is manifestly invariant under symplectic transformations (i.e. monodromies), which are manifest on the type IIA side. Later in this paper, we shall encounter a different ‘type IIB construction’ of the same twistor space, which makes S-duality manifest at the expense of obscuring symplectic transformations.

To explain this construction, one should first recall that, instead of covering the twistor space by open patches surrounded by closed contours, it is possible to consider a set of open contours with associated holomorphic transition functions across them (see appendix A). For a fixed value of the moduli z^a and any state of charge γ with $\Omega(\gamma, z^a) \neq 0$, we then consider the BPS ray

$$\ell_\gamma = \{t \in \mathbb{C}P^1 : Z_\gamma(z^a)/t \in i\mathbb{R}^-\}, \quad (2.17)$$

where $Z_\gamma(z^a)$ is the same central charge function as in (2.10). These rays extend from the north to the south pole and divide the patch \mathcal{U}_0 into angular sectors. Across each BPS ray

ℓ_γ , the Darboux coordinates are required to jump by the complex contact transformation generated by [15]

$$H_\gamma = G_\gamma - \frac{1}{2} q_\Lambda p^\Lambda (G'_\gamma)^2, \quad G_\gamma(\Xi_\gamma) = \frac{\Omega(\gamma)}{(2\pi)^2} \text{Li}_2(\sigma_\gamma e^{-2\pi i \Xi_\gamma}), \quad (2.18)$$

where G'_γ denotes the derivative with respect to $\Xi_\gamma = q_\Lambda \xi_{[\gamma]}^\Lambda - p^\Lambda \tilde{\xi}_\Lambda^{[\gamma]}$, $\Omega(\gamma)$ is the Donaldson-Thomas invariant,⁶ and σ_γ is a certain phase factor depending only on the charge, known as a quadratic refinement. The perhaps unfamiliar formula (2.18) is designed so as to generate jumps of the Darboux coordinates $(\xi^\Lambda, \tilde{\xi}_\Lambda)$ across ℓ_γ given by the usual symplectomorphism [44]

$$\begin{aligned} \Delta \xi^\Lambda &= \frac{1}{2\pi i} \Omega(\gamma) p^\Lambda \log(1 - \sigma_\gamma e^{-2\pi i \Xi_\gamma}), \\ \Delta \tilde{\xi}_\Lambda &= \frac{1}{2\pi i} \Omega(\gamma) q_\Lambda \log(1 - \sigma_\gamma e^{-2\pi i \Xi_\gamma}). \end{aligned} \quad (2.19)$$

The additional term proportional to $(G'_\gamma)^2$ in (2.18) arises upon integrating this symplectomorphism to a generating function $H^{[ij]}$, which depends on the coordinate $\xi_{[i]}^\Lambda$ in one patch and the dual coordinate $\tilde{\xi}_\Lambda^{[j]}$ in the other patch (see appendix A).

Unfortunately, except when all D-instantons are mutually local (i.e. $\langle \gamma, \gamma' \rangle = 0$ for any pair with non-vanishing $\Omega(\gamma), \Omega(\gamma')$), the Darboux coordinates determined by these gluing conditions cannot be expressed in closed form as in (2.13). Instead, they are determined by integral equations (A.4) which upon substitution of (2.18) take the form of a Thermodynamic Bethe Ansatz [25, 45]. Moreover, while the construction outlined above is manifestly covariant under symplectic transformations, it is not manifestly invariant under S-duality. In the presence of D5-brane instantons (i.e. $p^0 \neq 0$), this is of course expected since S-duality mixes D5-branes with NS5-branes, which are not included in the construction above. In the absence of D5-branes however (more precisely, in the large volume limit where both D5-branes and NS5-branes can be ignored), the metric should be invariant under the discrete group $\text{SL}(2, \mathbb{Z})$, yet this is far from obvious from the construction above. This invariance has been shown in the D1-D(-1) sector [34], extending the results of the earlier work [13], and our goal is to extend this construction to D3-D1-D(-1)-instantons. Before doing so however, we return to the simpler case of D1-D(-1)-instantons, improving on the earlier construction in [34].

3 S-duality and D1-D(-1)-instantons

In this section, we revisit the construction of the D1-D(-1)-instanton corrected metric on the HM moduli space in type IIB Calabi-Yau vacua, emphasizing how S-duality is realized in twistor space. In the process we considerably streamline the derivation of the ‘quantum mirror map’ obtained in [34].

⁶The fact that these invariants also depend on the moduli z^a does not spoil the holomorphicity of (2.18) because they are piecewise constant. Across a wall of marginal stability, the Ω 's jump and the BPS rays exchange their order in such a way that the metric on \mathcal{M}_H remains continuous [24, 25].

3.1 S-duality in the classical twistor space

Let us first recall how S-duality is realized in the twistor space \mathcal{Z} of \mathcal{M}_H in the classical, infinite volume limit. This twistor space is described by the Darboux coordinates (2.13), upon dropping the last logarithmic term in $\tilde{\alpha}$ and restricting the prepotential to its large volume limit (2.4). To express them in terms of the type IIB fields, it is sufficient to substitute the classical mirror map (2.5).

An important feature of the twistor space construction is that all isometries of a QK manifold can be lifted to a holomorphic action on \mathcal{Z} . In particular, the $SL(2, \mathbb{R})$ symmetry (2.7) ought to be realized as a holomorphic action on the Darboux coordinates $\xi^\Lambda, \tilde{\xi}_\Lambda$ and α . This indeed turns out to be the case provided the fiber coordinate t transforms as

$$t \mapsto \frac{1 + t_-^{c,d} t}{t_-^{c,d} - t} = -\frac{t_+^{c,d} - t}{1 + t_+^{c,d} t}, \tag{3.1}$$

where $t_\pm^{c,d}$ denote the two roots of the quadratic equation $c\xi^0(t) + d = 0$,

$$t_\pm^{c,d} = \frac{c\tau_1 + d \mp |c\tau + d|}{c\tau_2}, \quad t_+^{c,d} t_-^{c,d} = -1. \tag{3.2}$$

Combining (3.1) with (2.7) and the expressions (2.13) for the Darboux coordinates in terms of the type IIB fields then leads to the following non-linear holomorphic action on the Darboux coordinates in the patch \mathcal{U}_0 [14]⁷

$$\begin{aligned} \xi^0 &\mapsto \frac{a\xi^0 + b}{c\xi^0 + d}, & \xi^a &\mapsto \frac{\xi^a}{c\xi^0 + d}, & \tilde{\xi}_a &\mapsto \tilde{\xi}_a + \frac{c}{2(c\xi^0 + d)} \kappa_{abc} \xi^b \xi^c - c_{2,a} \epsilon(g), \\ \begin{pmatrix} \tilde{\xi}_0 \\ \alpha \end{pmatrix} &\mapsto \begin{pmatrix} d & -c \\ -b & a \end{pmatrix} \begin{pmatrix} \tilde{\xi}_0 \\ \alpha \end{pmatrix} + \frac{1}{6} \kappa_{abc} \xi^a \xi^b \xi^c \begin{pmatrix} c^2 / (c\xi^0 + d) \\ -[c^2(a\xi^0 + b) + 2c] / (c\xi^0 + d)^2 \end{pmatrix}. \end{aligned} \tag{3.3}$$

Under this action, the complex contact one-form (A.2) transforms by an overall holomorphic factor $\mathcal{X}^{[0]} \mapsto \mathcal{X}^{[0]} / (c\xi^0 + d)$, leaving the complex contact structure invariant. Furthermore, the contact potential e^Φ transforms with modular weight $(-\frac{1}{2}, -\frac{1}{2})$, which ensures that the Kähler potential (A.3) varies by a Kähler transformation,

$$e^\Phi \mapsto \frac{e^\Phi}{|c\tau + d|}, \quad K_{\mathcal{Z}} \mapsto K_{\mathcal{Z}} - \log(|c\xi^0 + d|). \tag{3.4}$$

These properties ensure that S-duality is indeed a symmetry of the classical twistor space.

To illuminate the action (3.1) of S-duality on the fiber, it is useful to make a Cayley transformation of the fiber coordinate and define⁸

$$z = \frac{t + i}{t - i}, \quad t = -i \frac{1 + z}{1 - z}. \tag{3.5}$$

⁷In general, since S-duality acts on the twistor fiber, it will map Darboux coordinates in one patch to Darboux coordinates in another patch. The virtue of the patch \mathcal{U}_0 is that it is mapped to itself under S-duality since it contains the points $t = \pm i$, see the remark at the end of this subsection.

⁸We are grateful to J. Manschot for suggesting this redefinition.

Under (3.1), this new coordinate transforms by the compensating $U(1)$ rotation induced by the right action on the coset $U(1)\backslash SL(2, \mathbb{R})$,

$$z \mapsto \frac{c\bar{\tau} + d}{|c\tau + d|} z, \quad (3.6)$$

i.e. z has modular weight $(-\frac{1}{2}, \frac{1}{2})$. In particular, the points $z = 0, \infty$ (corresponding to $t = \pm i$) stay invariant under $SL(2, \mathbb{R})$, whereas the zeros (3.2) of $c\xi^0 + d$ are now given by $z_{\pm}^{c,d} = \mp \sqrt{\frac{c\tau+d}{c\bar{\tau}+d}}$. Rewriting the Darboux coordinate ξ^0 in terms of z ,

$$\xi^0 = \tau_1 + i\tau_2 \frac{z^{-1} + z}{z^{-1} - z}, \quad (3.7)$$

we recognize in ξ^0 and z the analog of the 'canonical' and 'twisting' parameters introduced in [46].

3.2 Type IIB twistorial construction of D1-D(-1)-instanton corrections

As discussed in section 2.2, upon including worldsheet instanton corrections and the one-loop g_s -correction to the metric (2.1), the continuous $SL(2, \mathbb{R})$ action (2.7) is no longer isometric. Nevertheless, it was shown in [13] that invariance under a discrete subgroup $SL(2, \mathbb{Z})$ may be restored by incorporating D1 and D(-1)-instanton corrections. This was achieved in the framework of projective superspace by constructing a modular invariant completion of the hyperkähler potential, a close cousin of our contact potential e^{Φ} defined on the Swann bundle [6] over \mathcal{M}_H . At the same time, these instanton corrections should agree with the type IIA construction presented in section 2.3 restricted to the D1-D(-1) sector, with $p^\Lambda = 0$. This equivalence was demonstrated in [34], where the construction of [13] was translated in the twistorial language, and shown to be related to the type IIA construction by a certain complex contact transformation. Our aim is to revisit the twistorial construction of [34] and expose its invariance under S-duality.

As explained in [34], the projective superspace construction of [13] can be cast in the twistorial language by using a covering of the $\mathbb{C}P^1$ fiber by six patches

$$\mathcal{Z} = \mathcal{U}_+ \cup \mathcal{U}_- \cup \mathcal{U}_{0+} \cup \mathcal{U}_{0-} \cup \mathcal{U}_{\mathbb{R}+} \cup \mathcal{U}_{\mathbb{R}-}. \quad (3.8)$$

Here \mathcal{U}_{\pm} are, as usual, the patches around the north and south poles, $\mathcal{U}_{\mathbb{R}\pm}$ surround the positive and negative real half-axes, and the patches $\mathcal{U}_{0\pm}$ cover the upper and lower half-planes, in particular, containing the S-duality invariant points $t = \pm i$. The transition functions between these patches are taken to be

$$H^{[+0\pm]} = F(\xi^\Lambda), \quad H^{[-0\pm]} = \bar{F}(\xi^\Lambda), \quad H^{[\mathbb{R}+0\pm]} = G_{\text{IIB}}(\xi^\Lambda), \quad H^{[\mathbb{R}-0\pm]} = \bar{G}_{\text{IIB}}(\xi^\Lambda), \quad (3.9)$$

with

$$G_{\text{IIB}}(\xi^\Lambda) = -\frac{i}{(2\pi)^3} \sum_{q_a \gamma^a \in H_2^+(\hat{\mathcal{G}}) \cup \{0\}} n_{q_a}^{(0)} \sum_{\substack{n \in \mathbb{Z} \\ m > 0}} \frac{e^{-2\pi i m q_a \xi^a}}{m^2 (m \xi^0 + n)}, \quad (3.10)$$

where $n_{q_a}^{(0)}$ are the same genus zero Gopakumar-Vafa invariants which govern the worldsheet instantons (2.8).

It is important to note that the transition functions G_{IIB} introducing instanton corrections are independent of $\tilde{\xi}_\Lambda$ and α . This is necessary for the existence of the $n + 1$ commuting isometries corresponding to translations along \tilde{c}_a, \tilde{c}_0 and ψ , or equivalently holomorphic translations along $\tilde{\xi}_\Lambda$ and α . By the moment map construction [47], these $n + 1$ isometries lead to the existence of $n + 1$ $\mathcal{O}(2)$ -valued sections modulo rescalings, which correspond to the Darboux coordinates $(1, \xi^\Lambda)$. Thus the coordinates ξ^Λ are globally well-defined (up to rescalings) and given by their perturbative expressions (2.13) in all patches.

The remaining Darboux coordinates, $\tilde{\xi}_\Lambda$ and α , determined by the transition functions (3.9) were computed explicitly in [34]. In particular, it was shown that it is possible to relate the coordinates $A^\Lambda, B_\Lambda, B_\alpha$ appearing in (A.4) and the type IIB fields in such a way that the Darboux coordinates in the patch $\mathcal{U}_{0\pm}$ transform under the combined action of (2.7) and (3.1) according to the *classical* laws (3.3). This ensures that the twistor space \mathcal{Z} and the original moduli space \mathcal{M}_H carry an isometric action of $\text{SL}(2, \mathbb{Z})$. Moreover, it was also shown that this ‘type IIB’ twistorial construction is related to the usual ‘type IIA’ construction presented in section 2.3 by a set of complex contact transformations. We further discuss and clarify this relation in subsection 3.5.

While these results establish the invariance of the D1-D(-1)-instanton corrected metric under S-duality, it is fair to say that the computation of the quantum corrected mirror map in [34] was rather indirect and unilluminating. Moreover, it is highly desirable to understand how S-duality is realized at the level of transition functions, without having to compute the Darboux coordinates first and then check S-duality invariance. A more conceptual understanding of these two problems is certainly required in order to address D3-brane corrections, since unlike the D1-D(-1) case, the integral equations (A.4) cannot be solved in closed form as soon as D3-instantons are included. In the rest of this section, we disassemble the previous construction into its bare parts and expose the inner workings of S-duality in twistor space.

3.3 The key observations

To understand why the above construction produces a modular invariant twistor space, it is useful to separate the classical and instantonic parts of the construction, and examine how S-duality reshuffles the latter. To this end, let us refine the covering (3.8) and consider (see figure 1)

$$\mathcal{Z} = \mathcal{U}_+ \cup \mathcal{U}_- \cup \mathcal{U}_{0+} \cup \mathcal{U}_{0-} \cup \left(\bigcup'_{m,n} \mathcal{U}_{m,n} \right), \tag{3.11}$$

where m, n run over all pairs of integers⁹ different from $(0, 0)$ and $\mathcal{U}_{m,n}$ is an open set around the point $t_+^{m,n}$ (3.2), which is one of the zeros of $m\xi^0 + n$. The transition functions

⁹This set of patches is highly redundant, since $\mathcal{U}_{km, kn} = \mathcal{U}_{m,n}$ for any positive integer k . One could avoid this redundancy by restricting to coprime integers (m, n) , at the cost of introducing tri-logarithms in the transition functions below, but we prefer to keep it for the sake of simplicity. In particular, note that $\mathcal{U}_{0,k} = \mathcal{U}_+$ and $\mathcal{U}_{0,-k} = \mathcal{U}_-$.




Figure 1. The covering and transition functions incorporating D1-D(-1)-instantons.

between these patches are chosen as

$$H^{[+0^\pm]} = F^{\text{cl}}(\xi^\Lambda), \quad H^{[-0^\pm]} = \bar{F}^{\text{cl}}(\xi^\Lambda), \quad H^{[(m,n)0^\pm]} = G_{m,n}^{\text{D1}}(\xi^\Lambda), \quad (3.12)$$

where $F^{\text{cl}}(X)$ is the tree-level, large volume prepotential (2.4) and

$$G_{m,n}^{\text{D1}}(\xi^\Lambda) = -\frac{i}{(2\pi)^3} \sum_{q_a \gamma^a \in H_2^+(\hat{\mathcal{Q}}) \cup \{0\}} n_{q_a}^{(0)} \begin{cases} \frac{e^{-2\pi i m q_a \xi^a}}{m^2(m\xi^0 + n)}, & m \neq 0, \\ (\xi^0)^2 \frac{e^{2\pi i n q_a \xi^a / \xi^0}}{n^3}, & m = 0. \end{cases} \quad (3.13)$$

Thus, we simply introduced a distinct patch for each term in the sum over (m, n) in (3.10), centered around the point where this term is singular. In addition, we have split the prepotential F into its classical part F^{cl} and its quantum part $F^{\text{w.s.}} = \sum_{n>0} G_{0,n}^{\text{D1}}$, which arises in this construction as the contribution from $m = 0$ in (3.13).

The first two transition functions in (3.12) are the familiar ones arising in the twistorial construction of the tree-level metric, which is known to be S-duality invariant (see section 3.1). Therefore it is sufficient to concentrate on the patches $\mathcal{U}_{m,n}$ and transition functions $G_{m,n}^{\text{D1}}$. The first crucial observation is that the patches $\mathcal{U}_{m,n}$ are mapped into each other under $\text{SL}(2, \mathbb{Z})$ -transformations (3.1) according to

$$\mathcal{U}_{m,n} \mapsto \mathcal{U}_{m',n'}, \quad \begin{pmatrix} m' \\ n' \end{pmatrix} = \begin{pmatrix} a & c \\ b & d \end{pmatrix} \begin{pmatrix} m \\ n \end{pmatrix}, \quad (3.14)$$

whereas the patches \mathcal{U}_{0^\pm} stay invariant.

It is natural to expect that the functions $G_{m,n}^{\text{D1}}$ should similarly be reshuffled under the transformations generated by (3.3). This turns out to be true, up to an important subtlety: the transition functions $G_{m,n}^{\text{D1}}$ transform into each other under $\text{SL}(2, \mathbb{Z})$, up to an

overall factor of $(c\xi^0 + d)^{-1}$ and *up to terms which are regular in the patch $\mathcal{U}_{m',n'}$* ,¹⁰

$$G_{m,n}^{\text{D1}} \mapsto \frac{G_{m',n'}^{\text{D1}}}{c\xi^0 + d} + \text{regular at } t_+^{m',n'}. \quad (3.15)$$

This property is sufficient for S-duality because the regular terms appearing on the r.h.s. of (3.15) can be canceled by a local contact transformation and do not contribute to the contour integrals in eqs. (A.4).

Yet, a further difficulty is the fact the integration kernel $\frac{t'+t}{t'-t} \frac{dt'}{t'}$ appearing in these equations transforms in a complicated fashion, which seems to jeopardize S-duality. Our second key observation is that this kernel can be supplemented with a t -independent term

$$\frac{1}{2} \frac{t'+t}{t'-t} \frac{dt'}{t'} \rightarrow K(t, t') \frac{dt'}{t'} \equiv \frac{1}{2} \left(\frac{t'+t}{t'-t} + \frac{1/t' - t'}{1/t' + t'} \right) \frac{dt'}{t'}, \quad (3.16)$$

in such a way that the resulting kernel becomes $\text{SL}(2, \mathbb{Z})$ -invariant! Indeed, this invariance becomes obvious after expressing (3.16) in terms of the Cayley-rotated coordinate z (3.5),

$$K(t, t') \frac{dt'}{t'} = \frac{(1 + tt')}{(t' - t)(1/t' + t')} \frac{dt'}{t'} = \frac{1}{2} \frac{z' + z}{z' - z} \frac{dz'}{z'}. \quad (3.17)$$

Since the difference between the two kernels is t -independent, it can be absorbed into a redefinition of the constant terms in the t -expansion of Darboux coordinates. Thus, to write them in terms of the new kernel, it is sufficient to redefine $\zeta^\Lambda, \tilde{\zeta}_\Lambda$ and σ and, as we will see in the next subsection, this redefinition is the origin of quantum corrections to the mirror map.

3.4 The quantum mirror map revisited

Now we are ready to demonstrate the explicit invariance under $\text{SL}(2, \mathbb{Z})$ without evaluating explicitly the Penrose type integrals determining the instanton contributions and to derive a simple expression for the mirror map. To this end, we write the integral expressions (A.4) for Darboux coordinates by separating the classical contributions generated by $H^{[\pm 0^\pm]}$ from the quantum corrections coming from $H^{[(m,n)0^\pm]}$. Since ξ^Λ are uncorrected, it is sufficient to consider¹¹

$$\begin{aligned} \tilde{\xi}_\Lambda^{[0^\pm]} &= \tilde{\xi}_\Lambda^{\text{cl}} + \Delta \tilde{\zeta}_\Lambda - \frac{1}{2} \sum'_{m,n} \oint_{C_{m,n}} \frac{dt'}{2\pi i t'} \frac{t'+t}{t'-t} \partial_{\xi^\Lambda} G_{m,n}^{\text{D1}}(t'), \\ \alpha^{[0^\pm]} &= \alpha^{\text{cl}} - \frac{1}{2} \Delta \sigma - \frac{1}{2} \sum'_{m,n} \oint_{C_{m,n}} \frac{dt'}{2\pi i t'} \frac{t'+t}{t'-t} (1 - \xi^\Lambda \partial_{\xi^\Lambda}) G_{m,n}^{\text{D1}}(t'), \end{aligned} \quad (3.18)$$

¹⁰For example, if both m and $m' = dm - cn$ are non zero,

$$\frac{e^{-2\pi i m q_a \xi^a}}{m^2(m\xi^0 + n)} \mapsto \frac{e^{-2\pi i q_a \xi^a} \left(m' - c \frac{m'\xi^0 + n'}{c\xi^0 + d} \right)}{(c\xi^0 + d)(m'\xi^0 + n') \left(m' - c \frac{m'\xi^0 + n'}{c\xi^0 + d} \right)^2} = \frac{e^{-2\pi i m' q_a \xi^a}}{m'^2(c\xi^0 + d)(m'\xi^0 + n')} + O((m'\xi^0 + n')^0).$$

¹¹Here and henceforth the primes on sums over m, n will denote that the value $m = n = 0$ is omitted.

where we denoted by $\tilde{\xi}_\Lambda^{\text{cl}}, \alpha^{\text{cl}}$ the Darboux coordinates (2.13) with the prepotential replaced by its large volume limit $F^{\text{cl}}(X^\Lambda)$ and the logarithmic one-loop correction dropped. Nevertheless, the type IIA fields $\tilde{\zeta}_\Lambda$ and σ entering $\tilde{\xi}_\Lambda^{\text{cl}}$ and α^{cl} are still defined by relations (2.15) with the full prepotential F .¹² This fact is at the origin of two additional contributions in (3.18), given by

$$\begin{aligned}\Delta\tilde{\zeta}_\Lambda &= -\zeta^\Sigma \text{Re} \sum_{n>0} \partial_{z^\Lambda} \partial_{z^\Sigma} G_{0,n}^{\text{D1}}(z^\Lambda), \\ \Delta\sigma &= -\zeta^\Lambda \zeta^\Sigma \text{Re} \sum_{n>0} \partial_{z^\Lambda} \partial_{z^\Sigma} G_{0,n}^{\text{D1}}(z^\Lambda).\end{aligned}\tag{3.19}$$

Next, we perform the kernel replacement (3.16) in (3.18), thus rewriting the Darboux coordinates in the following form

$$\begin{aligned}\tilde{\xi}_\Lambda^{[0^\pm]} &= \left[\tilde{\xi}_\Lambda^{\text{cl}} - \tilde{\zeta}_\Lambda^{\text{inst}} \right] - \sum'_{m,n} \oint_{C_{m,n}} \frac{dt'}{2\pi i t'} K(t, t') \partial_{\xi^\Lambda} G_{m,n}^{\text{D1}}(t'), \\ \alpha^{[0^\pm]} &= \left[\alpha^{\text{cl}} + \frac{1}{2}(\sigma^{\text{inst}} + \tilde{\zeta}_\Lambda^{\text{inst}} \zeta^\Lambda) \right] - \sum'_{m,n} \oint_{C_{m,n}} \frac{dt'}{2\pi i t'} K(t, t') (1 - \xi^\Lambda \partial_{\xi^\Lambda}) G_{m,n}^{\text{D1}}(t'),\end{aligned}\tag{3.20}$$

where we combined all t -independent contributions into

$$\begin{aligned}\tilde{\zeta}_\Lambda^{\text{inst}} &= -\frac{1}{2} \sum'_{m,n} \oint_{C_{m,n}} \frac{dt'}{2\pi i t'} \frac{1/t' - t'}{1/t' + t'} \partial_{\xi^\Lambda} G_{m,n}^{\text{D1}}(t') - \Delta\tilde{\zeta}_\Lambda, \\ \sigma^{\text{inst}} &= -\zeta^\Lambda \tilde{\zeta}_\Lambda^{\text{inst}} + \sum'_{m,n} \oint_{C_{m,n}} \frac{dt'}{2\pi i t'} \frac{1/t' - t'}{1/t' + t'} (1 - \xi^\Lambda \partial_{\xi^\Lambda}) G_{m,n}^{\text{D1}}(t') - \Delta\sigma.\end{aligned}\tag{3.21}$$

Finally, using the property (3.15), it is easy to show that the derivatives of $G_{m,n}^{\text{D1}}$, appearing in (3.20) transform under S-duality as

$$\begin{aligned}\partial_{\xi^a} G_{m,n}^{\text{D1}} &\mapsto \partial_{\xi^a} G_{m',n'}^{\text{D1}} + \text{reg.} \\ \partial_{\xi^0} G_{m,n}^{\text{D1}} &\mapsto d \partial_{\xi^0} G_{m',n'}^{\text{D1}} - c(1 - \xi^\Lambda \partial_{\xi^\Lambda}) G_{m',n'}^{\text{D1}} + \text{reg.} \\ (1 - \xi^\Lambda \partial_{\xi^\Lambda}) G_{m,n}^{\text{D1}} &\mapsto a(1 - \xi^\Lambda \partial_{\xi^\Lambda}) G_{m',n'}^{\text{D1}} - b \partial_{\xi^0} G_{m',n'}^{\text{D1}} + \text{reg.}\end{aligned}\tag{3.22}$$

Using this result and noting that the regular contributions disappear under the contour integrals, the second term in the expression (3.20) for $\tilde{\xi}_a$ is manifestly invariant, whereas the second terms in $(\tilde{\xi}_0, \alpha)$ transform as a doublet under S-duality, consistently with the classical transformation properties (3.3). Thus, to establish that the twistor space carries a holomorphic action of $\text{SL}(2, \mathbb{Z})$, it remains to ensure that the first terms in (3.20) transform as the Darboux coordinates in the classical limit. For this purpose, it suffices to modify

¹²The relations (2.15) are not affected by D-instantons with vanishing p^Λ [14] so that they provide the correct definitions of the type IIA fields in our case. For $p^\Lambda \neq 0$, they receive additional instanton contributions, see eq. (3.10) in [15].

the classical mirror map relations (2.5) into

$$\begin{aligned} \zeta^0 &= \tau_1, & \zeta^a &= -(c^a - \tau_1 b^a), \\ \tilde{\zeta}_a &= \tilde{c}_a + \frac{1}{2} \kappa_{abc} b^b (c^c - \tau_1 b^c) + \tilde{\zeta}_a^{\text{inst}}, & \tilde{\zeta}_0 &= \tilde{c}_0 - \frac{1}{6} \kappa_{abc} b^a b^b (c^c - \tau_1 b^c) + \tilde{\zeta}_0^{\text{inst}}, \\ \sigma &= -2 \left(\psi + \frac{1}{2} \tau_1 \tilde{c}_0 \right) + \tilde{c}_a (c^a - \tau_1 b^a) - \frac{1}{6} \kappa_{abc} b^a c^b (c^c - \tau_1 b^c) + \sigma^{\text{inst}}. \end{aligned} \quad (3.23)$$

The relation between $r = e^\phi$ and τ_2 can be obtained by evaluating the contact potential. In our case, the formula (A.5) gives

$$e^\phi = \frac{\tau_2^2}{2} \mathcal{V} - \frac{\tau_2}{16\pi} \sum'_{m,n} \oint_{C_{m,n}} \frac{dt}{t} (t^{-1} z^\Lambda - t \bar{z}^\Lambda) \partial_{\xi^\Lambda} G_{m,n}^{\text{D1}}. \quad (3.24)$$

Again using (3.22), one may check that the integrand times $\sqrt{\tau_2}$ is invariant under $\text{SL}(2, \mathbb{Z})$ transformations up to a total derivative. As a result, the contact potential (3.24) transforms exactly as required in (3.4).

The formulas (3.21) and (3.24) encode in a nice and compact way the quantum corrections to the mirror map. They can be shown to be consistent with previously known results. Indeed, evaluating the integrals by residues, one can verify that (3.24) coincides with the contact potential found in [13, 34], while the quantum corrections (3.21) reproduce the results in eq. (5.2) [34] up to

$$\delta \tilde{\zeta}_0^{\text{inst}} = \sum_{q_a \geq 0} n_{q_a}^{(0)} \sum'_{m,n} m \delta_{m,n}, \quad \delta \sigma^{\text{inst}} = - \sum_{q_a \geq 0} n_{q_a}^{(0)} \sum'_{m,n} (m\tau_1 + 2n) \delta_{m,n}, \quad (3.25)$$

where

$$\delta_{m,n} = \frac{i n_{q_a}^{(0)}}{16\pi^3} \frac{\tau_2^2}{|m\tau + n|^4} (1 + 2\pi q_a t^a |m\tau + n|) e^{-S_{m,n,q_a}}. \quad (3.26)$$

Nevertheless, these contributions are harmless since the corresponding corrections to the Darboux coordinates, which can be written as

$$\begin{pmatrix} \delta \tilde{\xi}'_0 \\ \delta \alpha' \end{pmatrix} = \sum_{q_a \geq 0} n_{q_a}^{(0)} \sum'_{m,n} \begin{pmatrix} m \\ n \end{pmatrix} \delta_{m,n}, \quad (3.27)$$

transform as a doublet and preserve the transformation properties (3.3). Thus, the additional contributions (3.25) represent an inherent ambiguity of the quantum mirror map and can be dropped at will.

3.5 Equivalence between IIA and IIB constructions

The type IIB construction of the twistor space \mathcal{Z} outlined above is manifestly invariant under S-duality, however its equivalence to the type IIA construction described in section 2.3 is not manifest. In appendix A.2 of [34], it was shown that the type IIA Darboux coordinates differ from their type IIB counterpart by a complex contact transformation

$$\tilde{\xi}_\Lambda^{[i]} \mapsto \tilde{\xi}_\Lambda^{[i]} - \partial_{\xi^\Lambda} H^{[i]}, \quad \alpha^{[i]} \mapsto \alpha^{[i]} - H^{[i]} + \xi^\Lambda \partial_{\xi^\Lambda} H^{[i]}. \quad (3.28)$$

In each of the four quadrants on $\mathbb{C}P^1$ (in the t coordinate), away from the real and imaginary axes, this transformation is generated by

$$H^{[I]} = G_-^{D1}, \quad H^{[II]} = -G_+^{D1}, \quad H^{[III]} = \bar{G}_-^{D1}, \quad H^{[IV]} = -\bar{G}_+^{D1}, \quad (3.29)$$

where we defined

$$G_{\pm}^{D1}(\xi^\Lambda) = \frac{1}{(2\pi)^2} \sum_{\substack{(q_a, q_0) \\ q_0 + q_a b^a > 0, \pm q_a t^a \geq 0}} n_{q_a}^{(0)} \text{Li}_2 \left(e^{-2\pi i q_\Lambda \xi^\Lambda} \right) + \frac{\chi \mathfrak{y}}{96}, \quad (3.30)$$

More succinctly, in all quadrants (3.30) may be written as

$$H = \text{sgn} [\text{Re}(t) \text{Im}(t)] \left(\frac{1}{(2\pi)^2} \sum_{\substack{q_\Lambda \in \mathbb{Z} \\ (q_0 + q_a b^a) \text{Im } t > 0, (t^a q_a) \text{Re}(t) \leq 0}} \text{Li}_2 \left(e^{-2\pi i q_\Lambda \xi^\Lambda} \right) + \frac{\chi \mathfrak{y}}{96} \right). \quad (3.31)$$

To understand the origin of this complex contact transformation, it is useful to recall the structure of the BPS rays ℓ_γ in the D1-D(-1) sector. These rays extend from the north to the south pole along the meridian

$$\arg t = \frac{\pi}{2} + \arg [(q_0 + q_a b^a) + i q_a t^a]. \quad (3.32)$$

The conditions $(q_0 + q_a b^a) \text{Im } t > 0$, $(t^a q_a) \text{Re}(t) < 0$ select precisely those BPS rays which pass through the quadrant where t is located. Thus, the complex contact transformation generated by (3.31) is the product of the elementary contact transformations (2.19) associated with each of the BPS rays encountered in interpolating from a point t in the vicinity of the real axis to the point $t = i$ if $\text{Im } t > 0$, or to the point $t = -i$ if $\text{Im } t < 0$. One can therefore interpret the type IIB Darboux coordinates as the standard Darboux coordinates in the angular sector containing the S-duality invariant points $t = \pm i$, whereas the type IIA Darboux coordinates are the standard Darboux coordinates in the angular sectors containing the (positive or negative) real axis. This is consistent with the fact that the initial step in the computation of [34] was to move all the BPS rays along the positive or negative imaginary axis, before performing the Poisson resummation over the D(-1)-instanton charge q_0 . Had one chosen instead to move all BPS rays along the positive or negative real axis, such a contact transformation would not have arisen.

4 S-duality in twistor space with two isometries

In this section we propose a twistorial construction of a general class of quaternion-Kähler metrics, which consists of non-linear deformations of the c -map metric (2.1) which preserve two translational isometries along the coordinates ψ and \tilde{c}_0 and carry the isometric action (2.7) of $\text{SL}(2, \mathbb{Z})$. This class of metrics includes the D1-D(-1)-instanton corrected hypermultiplet metric, and generalizes the type IIB twistorial construction described in section 3. It is tailored for describing D3-instanton corrections in a manifestly S-duality

invariant fashion, with the coordinates ψ and \tilde{c}_0 playing the role of the axions dual to the D5 and NS5-instantons. However, the identification of the D3-instanton corrected HM metric within this class goes beyond the scope of this work.

Since the continuous isometries along ψ and \tilde{c}_0 lift to translations along the complex Darboux coordinates α and $\tilde{\xi}_0$, the holomorphic transition functions appearing in this construction are allowed to depend on ξ^0, ξ^a and $\tilde{\xi}_a$ in an arbitrary fashion (except for some global constraints following from S-duality). As a result, the Darboux coordinate ξ^0 is (projectively) globally well defined as the moment map of the Killing vector $\partial_{\tilde{c}_0}$, and given in all patches by the same $\mathcal{O}(2)$ section as in (2.13). In contrast, the Darboux coordinates $\xi^a, \tilde{\xi}_\Lambda, \alpha$ are defined only locally.

Let \mathcal{Z} be the $2n + 1$ -complex dimensional contact manifold defined by the infinite covering

$$\mathcal{Z} = \mathcal{U}_+ \cup \mathcal{U}_- \cup \mathcal{U}_0 \cup \left(\bigcup'_{m,n} \mathcal{U}_{m,n} \right), \tag{4.1}$$

and transition functions

$$H^{[+0]} = F^{\text{cl}}(\xi_{[+]}^\Lambda), \quad H^{[-0]} = \bar{F}^{\text{cl}}(\xi_{[-]}^\Lambda), \quad H^{[(m,n)0]} = G_{m,n}(\xi^0, \xi_{[m,n]}^a, \tilde{\xi}_a^{[0]}), \tag{4.2}$$

where $F^{\text{cl}}(X) = -\kappa_{abc} \frac{X^a X^b X^c}{6X^0}$ is an arbitrary cubic prepotential, $a = 1, \dots, n-1$, $\Lambda = 0, 1, \dots, n-1$. Here \mathcal{U}_\pm are the usual patches around the poles of $\mathbb{C}P^1$, $\mathcal{U}_{m,n}$ is a set of patches which are mapped to each other under $\text{SL}(2, \mathbb{Z})$ -transformations, and \mathcal{U}_0 covers the rest. They must be chosen so that \mathcal{U}_0 is mapped to itself under the antipodal map and S-duality, whereas $\mathcal{U}_{m,n}$ are mapped to each other according to

$$\tau[\mathcal{U}_{m,n}] = \mathcal{U}_{-m,-n}, \quad \mathcal{U}_{m,n} \mapsto \mathcal{U}_{m',n'}, \quad \begin{pmatrix} m' \\ n' \end{pmatrix} = \begin{pmatrix} a & c \\ b & d \end{pmatrix} \begin{pmatrix} m \\ n \end{pmatrix}. \tag{4.3}$$

The local holomorphic functions $G_{m,n}$ are assumed to transform under the $\text{SL}(2, \mathbb{Z})$ action (3.3) as

$$G_{m,n} \mapsto \frac{G_{m',n'}}{c\xi^0 + d} - \frac{c}{2} \frac{\kappa_{abc} \partial_{\tilde{\xi}_a^{[0]}} G_{m',n'} \partial_{\tilde{\xi}_b^{[0]}} G_{m',n'}}{(c\xi^0 + d)^2} \left(\xi_{[m',n']}^c - \frac{2}{3} \partial_{\tilde{\xi}_c^{[0]}} G_{m',n'} \right) + \text{reg.} \tag{4.4}$$

where $+\text{reg.}$ denotes equality up to terms which are regular in $\mathcal{U}_{m',n'}$. Then the following statements hold:

1. The Darboux coordinates in the patch \mathcal{U}_0 satisfy the following integral equations:

$$\begin{aligned} \xi^0 &= \zeta^0 + \frac{\tau_2}{2} (t^{-1} - t), \\ \xi_{[0]}^a &= \zeta_{\text{cl}}^a + t^{-1} Y^a - t \bar{Y}^a + \sum'_{m,n} \oint_{C_{m,n}} \frac{dt'}{2\pi i t'} K(t, t') \partial_{\tilde{\xi}_a^{[0]}} G_{m,n}, \\ \tilde{\xi}_\Lambda^{[0]} &= \tilde{\zeta}_\Lambda^{\text{cl}} + t^{-1} F_\Lambda^{\text{cl}}(Y) - t \bar{F}_\Lambda^{\text{cl}}(\bar{Y}) - \sum'_{m,n} \oint_{C_{m,n}} \frac{dt'}{2\pi i t'} K(t, t') \partial_{\tilde{\xi}_{[m,n]}^\Lambda} G_{m,n}, \\ \alpha^{[0]} &= -\frac{1}{2} (\tilde{\sigma} + \zeta^\Lambda \tilde{\zeta}_\Lambda)^{\text{cl}} - (t^{-1} + t) \left(t^{-1} F^{\text{cl}}(Y) + t \bar{F}^{\text{cl}}(\bar{Y}) \right) \end{aligned} \tag{4.5}$$

$$\begin{aligned}
& -\zeta_\Lambda^\Lambda \left(t^{-1} F_\Lambda^{\text{cl}}(Y) - t \bar{F}_\Lambda^{\text{cl}}(\bar{Y}) \right) \\
& - \sum'_{m,n} \oint_{C_{m,n}} \frac{dt'}{2\pi i t'} \left[K(t, t') \left(1 - \xi_{[m,n]}^\Lambda(t') \partial_{\xi_{[m,n]}^\Lambda} \right) G_{m,n} \right. \\
& \qquad \qquad \qquad \left. + \frac{(tt')^{-1} F_a^{\text{cl}}(Y) + t t' \bar{F}_a^{\text{cl}}(\bar{Y})}{1/t' + t'} \partial_{\xi_a^{[0]}} G_{m,n} \right],
\end{aligned}$$

where $C_{m,n}$ are contours on $\mathbb{C}P^1$ surrounding $\mathcal{U}_{m,n}$ counter-clockwise and $(\tau_2, Y^a, \zeta^0, \zeta_{\text{cl}}^a, \tilde{\zeta}_a^{\text{cl}}, \tilde{\sigma}^{\text{cl}})$ are coordinates on \mathcal{M} .

2. Upon identifying these coordinates with the following combinations of the type IIB fields $(\tau_1, \tau_2, t^a, b^a, c^a, \tilde{c}_a, \tilde{c}_0, \psi)$,

$$\begin{aligned}
\zeta^0 &= \tau_1, & \zeta_{\text{cl}}^a &= -(c^a - \tau_1 b^a), \\
\tilde{\zeta}_a^{\text{cl}} &= \tilde{c}_a + \frac{1}{2} \kappa_{abc} b^b (c^c - \tau_1 b^c), & \tilde{\zeta}_0^{\text{cl}} &= \tilde{c}_0 - \frac{1}{6} \kappa_{abc} b^a b^b (c^c - \tau_1 b^c), \\
\tilde{\sigma}^{\text{cl}} &= -2 \left(\psi + \frac{1}{2} \tau_1 \tilde{c}_0 \right) + \tilde{c}_a (c^a - \tau_1 b^a) - \frac{1}{6} \kappa_{abc} b^a c^b (c^c - \tau_1 b^c) + \frac{\tau_2^2}{3} \kappa_{abc} b^a b^b b^c, \\
Y^0 &= \frac{\tau_2}{2}, & Y^a &= \frac{\tau_2}{2} (b^a + i t^a) + \sum'_{m,n} \oint_{C_{m,n}} \frac{dt}{2\pi i t^2} \frac{\partial_{\xi_a^{[0]}} G_{m,n}}{(1/t + t)^2},
\end{aligned} \tag{4.6}$$

the Darboux coordinates (4.5) transform under the $\text{SL}(2, \mathbb{Z})$ transformations (2.7) and (3.1) according to the classical laws (3.3).

3. The same transformation rules also hold for the Darboux coordinates in the patches $\mathcal{U}_{m,n}$, with the understanding that the Darboux coordinates appearing in the r.h.s. are those attached to the patch $\mathcal{U}_{m',n'}$, e.g. $\xi_{[m,n]}^a \mapsto \xi_{[m',n']}^a / (c\xi^0 + d)$.
4. The contact potential is given by

$$\begin{aligned}
e^\Phi &= \frac{2}{3\tau_2} \kappa_{abc} \text{Im } Y^a \text{Im } Y^b \text{Im } Y^c \\
& - \frac{1}{16\pi} \sum'_{m,n} \oint_{C_{m,n}} \frac{dt}{t} \left[(t^{-1} Y^\Lambda - t \bar{Y}^\Lambda) \partial_{\xi_{[m,n]}^\Lambda} G_{m,n} \right. \\
& \qquad \qquad \qquad \left. + \left(t^{-1} F_a^{\text{cl}}(Y) - t \bar{F}_a^{\text{cl}}(\bar{Y}) \right) \partial_{\xi_a^{[0]}} G_{m,n} \right]
\end{aligned} \tag{4.7}$$

and transforms under $\text{SL}(2, \mathbb{Z})$ as in (3.4).

5. As a result, the $4n$ -dimensional quaternion-Kähler manifold \mathcal{M} associated to the twistor space \mathcal{Z} carries an isometric action of $\text{SL}(2, \mathbb{Z})$.
6. Upon freezing the moduli τ_1, τ_2 and ignoring the Darboux coordinates $\alpha, \tilde{\xi}_0$, the remaining ones, ξ^a and $\tilde{\xi}_a$, determine a one-parameter family of $(4n - 4)$ -dimensional hyperkähler metrics \mathcal{M}_τ , with τ taking values in the Poincaré upper half plane, such that \mathcal{M}_τ and $\mathcal{M}_{(a\tau+b)/(c\tau+d)}$ are identified isometrically under (2.7).

Some comments are in order:

- The key transformation law (4.4) expresses the fact that S-duality should commute with the holomorphic transition functions up to local contact transformations, i.e. that the following diagram should be commutative:

$$\begin{array}{ccc}
 \mathcal{U}_0 & \xrightarrow{S} & \mathcal{U}_0 \\
 G_{m,n} \downarrow & & \downarrow G_{m',n'} \\
 \mathcal{U}_{m,n} & \xrightarrow{S} & \mathcal{U}_{m',n'}
 \end{array} \tag{4.8}$$

Here the horizontal arrows correspond to the holomorphic action (3.3) of S-duality on twistor space, which maps the Darboux coordinates in \mathcal{U}_0 (resp., $\mathcal{U}_{m,n}$) to Darboux coordinates in the same patch \mathcal{U}_0 (resp., in the transformed patch $\mathcal{U}_{m',n'}$). The quadratic and cubic terms in (4.4) have the same origin as the quadratic term in (2.18), namely the fact that the generating function of the contact transformation must be expressed in terms of the coordinates ξ^Λ in the original patch and of the coordinates $\tilde{\xi}_\Lambda$ in the final patch. In principle, the condition (4.4) determines all $G_{m,n}$ in terms of $G_{(\text{gcd}(m,n),0)}$, although the non-linearities in (4.4) prevent us from writing a closed expression. In the one-instanton approximation, the quadratic and cubic terms can be omitted, and the transformation law (4.4) simply expresses the invariance of a Čech cocycle in $\mathcal{H}^1(\mathcal{Z}, \mathcal{O}(2))$ under $\text{SL}(2, \mathbb{Z})$. It may be summarized by saying that the formal sum $\sum'_{m,n} G_{m,n}$ transforms as a holomorphic modular form of weight -1 .

- The equations (4.5) are simply the translation of the general integral equations (A.4) for the specific choice of contours and transition functions (4.2), after performing the change of kernel (3.16). In particular the last term in the square bracket in the expression for $\alpha^{[0]}$ originates from the transition functions $H^{[\pm 0]}$ and may be rewritten as $[t^{-1}K(0, t')F_a^{\text{cl}} - tK(\infty, t')\bar{F}_a^{\text{cl}}]\partial_{\tilde{\xi}_a^{[0]}}G_{m,n}$. The kernel substitution does not affect the discontinuity along the contours, and can be reabsorbed by the change of coordinates (4.6).
- If $G_{m,n}$ is independent of $\tilde{\xi}_a^{[0]}$, we recover the construction of the D1-D(-1)-instanton corrected metric in section 3. In that case the patches $\mathcal{U}_{m,n}$ were centered around the zeros of $m\xi^0 + n$, which we denoted by $t_\pm^{m,n}$, and the transition functions are given in (3.13). Their transformation law (3.15) is a particular case of (4.4).
- Instead or in addition to the closed contours $C_{m,n}$ surrounding open patches $\mathcal{U}_{m,n}$, one can include open contours similar to the BPS rays appearing in the construction of section 2.3. In fact, all previous statements hold for an arbitrary set of contours $C_{m,n}$ mapped unto each other by $\text{SL}(2, \mathbb{Z})$. For example, $C_{m,n}$ can be an open contour from $t_+^{m,n}$ to $t_-^{m,n}$. In this case however the structure of patches might be more complicated. The only requirement on the covering is that it should be invariant under the antipodal map and $\text{SL}(2, \mathbb{Z})$ transformations so that a patch \mathcal{U}_i is mapped

to some patch $\mathcal{U}_{i'}$. The Darboux coordinates in each patch are given by the formulas (4.5) and transform under S-duality by (3.3) where the coordinates on the r.h.s. are from the patch $\mathcal{U}_{i'}$. Moreover, if the function $G_{m,n}$ is associated with an open contour, the regular terms in the transformation rule (4.4) are no longer allowed, as they would otherwise not disappear under integration and therefore spoil S-duality.

- Note that the coordinate change (4.6) is almost identical to the classical mirror map (2.5), up to two differences. First, the coordinate $\tilde{\sigma}^{\text{cl}}$ differs from σ in (2.5) by the last term $\frac{\tau^3}{3} \kappa_{abc} b^a b^b b^c$. Although this is a classical contribution, we chose to absorb it into $\tilde{\sigma}^{\text{cl}}$ in order to simplify the expression for α in terms of Y^a . Second and more importantly, Y^a receives instanton corrections away from its classical value $Y^a = \frac{\tau_2}{2} z^a$. This may be viewed as an instanton correction to the mirror map, however note that the notion of mirror map only makes sense for the specific choice of $G_{m,n}$ corresponding to D3-brane instantons. In that case, the full quantum mirror map will require specifying also $\zeta^a, \tilde{\zeta}_\Lambda, \sigma$ in terms of $\zeta_a^{\text{cl}}, \tilde{\zeta}_a^{\text{cl}}, \tilde{\sigma}^{\text{cl}}$.
- The identifications (4.6) are not the only possible ones which lead to manifest $\text{SL}(2, \mathbb{Z})$ invariance. We already met an example of such an ambiguity in (3.25) when studying D1-D(-1)-instanton corrections to the mirror map. Another simple example can be found in appendix B, eq. (B.13), and many other examples can be constructed (see e.g. [35]). Fortunately, all these ambiguities just represent a freedom in the choice of coordinates and do not affect the geometry of the QK manifold.
- The modular invariant family of $4n - 4$ -dimensional HK manifolds \mathcal{M}_τ arises as a rigid limit of the QK space \mathcal{M} , using the same philosophy as in the QK/HK correspondence studied in [24]. Namely, the two isometries $\partial_{\tilde{c}_0}$ and ∂_ψ on \mathcal{M} can be lifted to triholomorphic isometries of the Swann bundle \mathcal{S} (a \mathbb{C}^\times bundle over \mathcal{Z}), with moment maps η^b and $\eta^0 = \xi^0 \eta^b$ in the notations of [12]. The hyperkähler quotient of \mathcal{S} with respect to these two isometries is obtained by enforcing the D-term constraints, i.e. freezing the $\mathcal{O}(2)$ multiplets η^b and η^0 , and quotienting by the action of $\partial_{\tilde{c}_0}$ and ∂_ψ . It produces a family of $(4n - 4)$ -dimensional hyperkähler metrics, parametrized by the moment maps η^b and η^0 . The former can be fixed to an arbitrary non-zero value by a suitable $\text{SU}(2)$ rotation, while the latter yields one complex parameter τ which is extracted from the components of the $\mathcal{O}(2)$ multiplet ξ^0 . For fixed value of τ , the HK metric is then coordinatized by $t^a, b^a, c^a, \tilde{c}_a$, or by the Darboux coordinates $\xi^a, \tilde{\xi}_a$. This class of modular invariant HK metrics should include the Coulomb branch of five-dimensional $\mathcal{N} = 2$ gauge theories compactified on T^2 , where the modular group of T^2 plays the role of S-duality [27] and the monopole strings play the role of D3-instantons.

We emphasize that the statements (1.-6.) hold to all orders in the instanton expansion. It is a tedious, but straightforward exercise to verify them. To this end, one first brings the integral equations (A.4) to the form (4.5) and the contact potential (A.5) to the form (4.7). This can be done by evaluating the integrals with $H^{[\pm 0]}$ and by a suitable identification

of $A^\Lambda, B_\Lambda, B_\alpha$ with $\zeta_{\text{cl}}^\Lambda, \tilde{\zeta}_\Lambda^{\text{cl}}, \tilde{\sigma}^{\text{cl}}$. Then $\text{SL}(2, \mathbb{Z})$ transformations of the Darboux coordinates and the contact potential can be checked by using the transformation properties of the derivatives of $G_{m,n}$, which provide a generalization of (3.22) and can be easily obtained from (4.4),

$$\begin{aligned}
 \partial_{\tilde{\zeta}_a^{[0]}} G_{m,n} &\mapsto \frac{\partial_{\tilde{\zeta}_a^{[0]}} G_{m',n'}}{c\xi^0 + d} + \text{reg.} \\
 \partial_{\xi_{[m,n]}^a} G_{m,n} &\mapsto \partial_{\xi_{[m',n']}^a} G_{m',n'} - \frac{c \kappa_{abc} \partial_{\tilde{\zeta}_b^{[0]}} G_{m',n'}}{c\xi^0 + d} \left(\xi_{[m',n']}^c - \frac{1}{2} \partial_{\tilde{\zeta}_c^{[0]}} G_{m',n'} \right) + \text{reg.} \\
 \partial_{\xi^0} G_{m,n} &\mapsto d \partial_{\xi^0} G_{m',n'} - c \left(1 - \xi_{[m',n']}^\Lambda \partial_{\xi_{[m',n']}^\Lambda} \right) G_{m',n'} \\
 &\quad - \frac{c^2 \kappa_{abc} \partial_{\tilde{\zeta}_a^{[0]}} G_{m',n'}}{2(c\xi^0 + d)} \left(\xi_{[m',n']}^b \xi_{[m',n']}^c - \xi_{[m',n']}^b \partial_{\tilde{\zeta}_c^{[0]}} G_{m',n'} \right. \\
 &\quad \left. + \frac{1}{3} \partial_{\tilde{\zeta}_b^{[0]}} G_{m',n'} \partial_{\tilde{\zeta}_c^{[0]}} G_{m',n'} \right) + \text{reg.}
 \end{aligned} \tag{4.9}$$

In appendix B, we explain in detail how the construction works in the one-instanton approximation, i.e. to first order in $G_{m,n}$.

5 Discussion

In this work we have provided a twistorial construction for a class of quaternion-Kähler metrics which admit two commuting, continuous isometries and an isometric action of $\text{SL}(2, \mathbb{Z})$. These metrics are non-linear deformations of the standard c -map metric derived from a cubic prepotential. They are parametrized by an infinite set of holomorphic functions $G_{m,n}(\xi^\Lambda, \tilde{\xi}_a)$ subject to the transformation rule (4.4) and tailored for describing D3-instanton corrections to the QK metric on the HM moduli space \mathcal{M}_H in type IIB string theory compactified on a Calabi-Yau threefold. As a special case, one recovers the D1-D(-1)-instanton corrected metric on \mathcal{M}_H constructed in [13, 34], whose manifest S-duality invariance is now revealed. Upon taking a rigid limit, the same construction also yields a class of modular invariant hyperkähler metrics (where the modulus τ is now a parameter rather than a coordinate), which should describe the Coulomb branch of five-dimensional $\mathcal{N} = 2$ gauge theories compactified on T^2 , with monopole strings playing the role of D3-instantons.

Clearly, an outstanding question is to determine the set of holomorphic functions $G_{m,n}$ which describes D3-instanton corrections (or similarly, the monopole string instantons) to the real HM moduli space. For this purpose one should find an adequate complex contact transformation which would cast the type IIA construction of the twistor space outlined in section 2.3 into the manifestly S-duality invariant type IIB construction of section 4. This is an important problem, since it would prove that D3-instanton corrections are indeed consistent with S-duality, and it would allow to compute quantum corrections to the mirror map.

An indication that this translation can be done is the fact that, in the one-instanton approximation, the formal sum of transition functions H_γ in (2.18) over all D3-D1-D(-1)

charges is formally identical to the partition function of D4-D2-D0 black holes in type IIA string theory compactified on the same CY threefold $\hat{\mathfrak{Y}}$. The latter is a Jacobi form and admits a natural Poincaré series representation [48–52]. Unfortunately, there are technical difficulties in implementing this idea due to the fact that, for fixed D3-brane charge, the formal sum over D1-instantons leads to an indefinite theta series of signature $(1, b_2(\hat{\mathfrak{Y}}) - 1)$ which is divergent. Moreover, Poincaré series of negative weight are also divergent and need to be regularized, leading to holomorphic or modular anomalies. In [35], we attack this problem from a different angle, and show that the instanton corrections to the type IIA Darboux coordinates, in the large volume limit, are governed by certain Mock theta series. In the analogous problem of the Coulomb branch of $\mathcal{N} = 2$ five dimensional gauge theories, the monopole string instantons are described by the elliptic genus of the $(0, 4)$ superconformal field theory with target space given by the ADHM moduli space of monopoles, which is also expected to be modular invariant [27]. It would be very interesting to see if some of the difficulties of string theory are alleviated in the gauge theory set-up.

Another outstanding problem is to remove the assumption that the metric has two commuting Killing vectors, and to construct a general class of $SL(2, \mathbb{Z})$ invariant QK metrics with no continuous isometries as non-linear perturbations of the c -map metric. Similar to the construction in section 4, this class should be parametrized by functions $G_{m,n}$ which are now allowed to depend on all Darboux coordinates $\xi^\Lambda, \tilde{\xi}_\Lambda, \alpha$. At the linear level, S-duality is again ensured by the condition (4.4) where only the first term on the r.h.s. should be retained. An example of such construction was given in [16], where NS5-brane instanton corrections were inferred by S-duality from the D5-brane instantons and shown to be governed by a set of transition functions transforming with modular weight -1 as in (4.4). However, an extension of this construction beyond the linear level appears to be highly non-trivial.

Acknowledgments

We are grateful to D. Persson and S. Vandoren for valuable discussions, and to J. Manschot for discussions and collaboration on a closely related project.

A Twistorial description of quaternion-Kähler manifolds

A $4n$ -dimensional Riemannian manifold \mathcal{M} is quaternion-Kähler if it has restricted holonomy group $USp(n) \times SU(2) \subset SO(4n)$. The Ricci scalar R is then constant, and the curvature of the $SU(2)$ part of the Levi-Civita connection, rescaled by $1/R$, provides a triplet of quaternionic 2-forms $\vec{\omega}$. While R can take either sign, hypermultiplet moduli spaces in $\mathcal{N} = 2$ supergravity or string theory models have $R < 0$.

A QK manifold \mathcal{M} can be described analytically in terms of its twistor space \mathcal{Z} , the total space of the $\mathbb{C}P^1$ bundle over \mathcal{M} twisted with the *projectivized* $SU(2)$ connection on \mathcal{M} . \mathcal{Z} is a Kähler-Einstein space equipped with a canonical complex contact structure, given by the kernel of the one-form

$$Dt = dt + p_+ - ip_3t + p_-t^2, \tag{A.1}$$

where t is a complex coordinate on $\mathbb{C}P^1$, and p_{\pm}, p_3 is the $SU(2)$ part of the Levi-Civita connection on \mathcal{M} . Note that Dt is defined only projectively, as it rescales under $SU(2)$ rotations. More precisely, it is valued in the $\mathcal{O}(2)$ line bundle on $\mathbb{C}P^1$ [4].

Locally, on a patch of an open covering $\{\mathcal{U}_i\}$ of \mathcal{Z} , one can always find complex Darboux coordinates $(\xi_{[i]}^{\Lambda}, \tilde{\xi}_{\Lambda}^{[i]}, \alpha^{[i]})$ such that the contact one-form (A.1), suitably rescaled by a function $e^{\Phi^{[i]}}$, takes the form

$$\mathcal{X}^{[i]} \equiv 4 e^{\Phi^{[i]}} \frac{Dt}{it} = d\alpha^{[i]} + \tilde{\xi}_{\Lambda}^{[i]} d\xi_{[i]}^{\Lambda}. \quad (\text{A.2})$$

The function $\Phi^{[i]}$, which we refer to as the ‘contact potential’, is holomorphic along the $\mathbb{C}P^1$ fiber and defined up to an additive holomorphic function on \mathcal{U}_i . It provides a Kähler potential for the Kähler-Einstein metric on \mathcal{Z} [12]:

$$K_{\mathcal{Z}}^{[i]} = \log \frac{1+t\bar{t}}{|t|} + \text{Re } \Phi^{[i]}. \quad (\text{A.3})$$

Globally, the complex contact structure on \mathcal{Z} can be specified by a set of generating functions $H^{[ij]}(\xi_{[i]}^{\Lambda}, \tilde{\xi}_{\Lambda}^{[j]}, \alpha^{[j]})$ for complex contact transformations between Darboux coordinates on overlaps $\mathcal{U}_i \cap \mathcal{U}_j$, subject to cocycle and reality conditions [12].

In the case when \mathcal{M} has a quaternionic isometry ∂_{σ} , one may choose the Darboux coordinates such that the Killing vector lifts to the holomorphic action ∂_{α} . As a result, the transition functions $H^{[ij]}$ become independent of the coordinate $\alpha^{[j]}$, and the contact potential $\Phi^{[i]}$ becomes real and constant on $\mathbb{C}P^1$ [14], equal to log-norm of the moment map associated to the isometry. The twistorial construction of such QK manifolds with one quaternionic isometry then becomes isomorphic to the twistorial construction of HK manifolds with one rotational isometry, a relation known as the QK/HK correspondence [24, 53]. In this case the Darboux coordinates are determined by the following system of integral equations:

$$\begin{aligned} \xi_{[i]}^{\Lambda}(t) &= A^{\Lambda} + t^{-1}Y^{\Lambda} - t\bar{Y}^{\Lambda} - \frac{1}{2} \sum_j \int_{C_j} \frac{dt'}{2\pi i t'} \frac{t'+t}{t'-t} \partial_{\tilde{\xi}_{\Lambda}^{[j]}} H^{[ij]}, \\ \tilde{\xi}_{\Lambda}^{[i]}(t) &= B_{\Lambda} + \frac{1}{2} \sum_j \int_{C_j} \frac{dt'}{2\pi i t'} \frac{t'+t}{t'-t} \partial_{\xi_{\Lambda}^{[j]}} H^{[ij]}, \\ \alpha^{[i]}(t) &= B_{\alpha} + \frac{1}{2} \sum_j \int_{C_j} \frac{dt'}{2\pi i t'} \frac{t'+t}{t'-t} \left(H^{[ij]} - \xi_{[i]}^{\Lambda} \partial_{\xi_{\Lambda}^{[i]}} H^{[ij]} \right) + 4ic \log t. \end{aligned} \quad (\text{A.4})$$

Here the complex variables Y^{Λ} , up to an overall phase rotation which can be absorbed into a phase rotation of t , and the real variables $A^{\Lambda}, B_{\Lambda}, B_{\alpha}$ serve as coordinates on \mathcal{M} . It is convenient to fix the phase freedom in Y^{Λ} by requiring $Y^0 \equiv \mathcal{R}$ to be real. Moreover, B_{α} is related to the coordinate σ along the isometric direction, $\partial_{B_{\alpha}} \sim \partial_{\sigma}$. Finally, c is a real constant known as the anomalous dimension [12], which characterizes the singular behavior of the Darboux coordinate α at the north and south poles of $\mathbb{C}P^1$. It plays an important physical role in describing the one-loop correction to the hypermultiplet moduli space metric in type II string compactifications.

The integrals in (A.4) are taken around closed contours C_i surrounding the patches \mathcal{U}_i in the counter-clockwise direction. Nevertheless, the construction is still meaningful if some of the contours are taken to be open. Typically such open contours appear as the degeneration of closed contours in the presence of branch cut singularities in the transition functions $H^{[ij]}$. The holomorphic functions associated with that open contours are then equal to the discontinuity of $H^{[ij]}$ across the branch cut [11]. In particular, this is the case for twistorial constructions of D-brane instantons in type IIA [14, 15] as well as fivebrane instantons in type IIB [16].

Thus, the metric on \mathcal{M} is completely determined by a constant c and a set of open or closed contours in $\mathbb{C}P^1$, together with associated holomorphic functions. The procedure to extract the metric from the solutions of (A.4) was outlined in [12, 14]. An important ingredient in this calculation is the contact potential, which can be computed from the transition functions $H^{[ij]}$ and the solutions of (A.4) using

$$e^\Phi = \frac{1}{16\pi} \sum_j \int_{C_j} \frac{dt}{t} (t^{-1}Y^\Lambda - t\bar{Y}^\Lambda) \partial_{\xi_{[i]}^\Lambda} H^{[ij]} - c. \tag{A.5}$$

B S-duality invariance at linear order

In this appendix we show the consistency of the construction of section 4 in the one-instanton approximation. Our aim is to prove that the Darboux coordinates (4.5) transform under S-duality as in (3.3), provided the coordinates $(\tau_2, Y^a, \zeta^0, \zeta_{\text{cl}}^a, \tilde{\zeta}_a^{\text{cl}}, \tilde{\sigma}^{\text{cl}})$ are related to the type IIB fields $(\tau_1, \tau_2, t^a, b^a, c^a, \tilde{c}_a, \tilde{c}_0, \psi)$ as in (4.6).

We start with the Darboux coordinate $\xi_{[0]}^a$. Under an $SL(2, \mathbb{Z})$ transformation, the kernel $K(t, t')dt'/t'$ is invariant but the derivative $\partial_{\xi_{[0]}^a} G_{m,n}(t')$ transforms with an overall factor of $1/(c\xi^0(t') + d)$ (see (4.9)). Remarking that $t^{-1} + t$ transforms as

$$t^{-1} + t \mapsto \frac{|c\tau + d|}{c\xi^0 + d} (t^{-1} + t), \tag{B.1}$$

we see that the factor $1/(c\xi^0(t') + d)$ can be converted into $1/(c\xi^0(t) + d)$ by inserting an additional factor of $(t^{-1} + t)/(t'^{-1} + t')$ in the integrand. This is the purpose of the field redefinition of Y^a in (4.6). Indeed, by expressing Y^a in terms of $z^a = b^a + it^a$ in the second line of (4.5) we arrive at

$$\xi_{[0]}^a = \zeta_{\text{cl}}^a + \frac{\tau_2}{2} (t^{-1}z^a - t\bar{z}^a) + \delta\xi^a, \tag{B.2}$$

where

$$\delta\xi^a = \sum'_{m,n} \oint_{C_{m,n}} \frac{dt'}{2\pi i t'} \frac{1/t + t}{1/t' + t'} K(t, t') \partial_{\xi_a} G_{m,n}. \tag{B.3}$$

Both terms then manifestly transform as in (3.3), under the assumption that ζ_{cl}^a is related to c^a as in (4.6).

To investigate the other Darboux coordinates in (4.5), we first rewrite them in terms of the type IIB Kähler moduli z^a as

$$\begin{aligned}
 \tilde{\xi}_a^{[0]} &= \tilde{\zeta}_a^{\text{cl}} - \frac{\tau_2}{4} \kappa_{abc} \left(t^{-1} z^b z^c - t \bar{z}^b \bar{z}^c \right) + \delta \tilde{\xi}_a, \\
 \delta \tilde{\xi}_a &= - \sum'_{m,n} \oint_{C_{m,n}} \frac{dt'}{2\pi i t'} \left(K(t, t') \partial_{\xi^a} G_{m,n} + \kappa_{abc} \frac{(tt')^{-1} z^b - tt' \bar{z}^b}{(1/t' + t')^2} \partial_{\tilde{\xi}_c} G_{m,n} \right), \\
 \tilde{\xi}_0^{[0]} &= \tilde{\zeta}_0^{\text{cl}} + \frac{\tau_2}{12} \kappa_{abc} \left(t^{-1} z^a z^b z^c - t \bar{z}^a \bar{z}^b \bar{z}^c \right) + \delta \tilde{\xi}_0, \\
 \delta \tilde{\xi}_0 &= - \sum'_{m,n} \oint_{C_{m,n}} \frac{dt'}{2\pi i t'} \left(K(t, t') \partial_{\xi^0} G_{m,n} - \kappa_{abc} \frac{(tt')^{-1} z^a z^b - tt' \bar{z}^a \bar{z}^b}{2(1/t' + t')^2} \partial_{\tilde{\xi}_c} G_{m,n} \right), \\
 \alpha^{[0]} &= - \frac{1}{2} (\sigma + \zeta^\Lambda \tilde{\zeta}_\Lambda)_{\text{cl}} + \frac{\tau_2^2}{8} \left(\bar{z}^\Lambda F_\Lambda^{\text{cl}}(z) + z^\Lambda \bar{F}_\Lambda^{\text{cl}}(\bar{z}) \right) \\
 &\quad - \frac{\tau_2^2}{4} \left(t^{-2} F^{\text{cl}}(z) + t^2 \bar{F}^{\text{cl}}(\bar{z}) \right) - \frac{\tau_2 \zeta^\Lambda}{2} \left(t^{-1} F_\Lambda^{\text{cl}}(z) - t \bar{F}_\Lambda^{\text{cl}}(\bar{z}) \right) + \delta \alpha, \\
 \delta \alpha &= - \sum'_{m,n} \oint_{C_{m,n}} \frac{dt'}{2\pi i t'} \left(K(t, t') (1 - \xi^\Lambda(t') \partial_{\xi^\Lambda}) G_{m,n} - \kappa_{abc} \frac{V^{ab}(t, t')}{(1/t' + t')^2} \partial_{\tilde{\xi}_c} G_{m,n} \right),
 \end{aligned} \tag{B.4}$$

where, in the last expression for $\delta \alpha$,

$$\begin{aligned}
 V^{ab}(t, t') &= -c^a \left(b^b \frac{1 - (tt')^2}{tt'} + i t^b \frac{1 + (tt')^2}{tt'} \right) + \frac{\tau_1}{2} \left(b^a b^b + t^a t^b \right) \frac{1 - (tt')^2}{tt'} \\
 &\quad + \frac{\tau_2}{4} (t^{-1} + t + t'^{-1} + t') \left((b^a b^b - t^a t^b) \frac{1 + (tt')^2}{tt'} + 2i b^a t^b \frac{1 - (tt')^2}{tt'} \right).
 \end{aligned} \tag{B.5}$$

It may be checked by a direct but tedious calculation that each of these Darboux coordinates transforms as in (3.3), provided $\tilde{\zeta}_\Lambda^{\text{cl}}, \sigma_{\text{cl}}$ are related to c_a, c_0, ψ as in (4.6).

This computation is however best performed by introducing first the following linear combinations of the instanton contributions

$$\begin{aligned}
 \hat{\delta} \tilde{\xi}_a &= \delta \tilde{\xi}_a + \kappa_{abc} \left(b^b + i t^b \frac{1/t - t}{1/t + t} \right) \delta \xi^c, \\
 \hat{\delta}_+ \alpha &= \delta \alpha + \tau \delta \tilde{\xi}_0 - \frac{1}{2} \kappa_{abc} \left(b^a + i t^a \frac{1/t - t}{1/t + t} \right) \left(\xi^b - c^b + \tau b^b - \frac{2t^b (\xi^0 - \tau)}{1/t + t} \right) \delta \xi^c, \\
 \hat{\delta}_- \alpha &= \delta \alpha + \bar{\tau} \delta \tilde{\xi}_0 - \frac{1}{2} \kappa_{abc} \left(b^a + i t^a \frac{1/t - t}{1/t + t} \right) \left(\xi^b - c^b + \bar{\tau} b^b + \frac{2t^b (\xi^0 - \bar{\tau})}{1/t + t} \right) \delta \xi^c.
 \end{aligned} \tag{B.6}$$

These combinations have been designed such that the complicated transformation rules of $\delta \tilde{\xi}_\Lambda, \delta \alpha$, obtained by linearizing (3.3), translate into the simple properties

$$\hat{\delta} \tilde{\xi}_a \mapsto \hat{\delta} \xi_a, \quad \hat{\delta}_+ \alpha \mapsto \frac{\hat{\delta}_+ \alpha}{c\tau + d}, \quad \hat{\delta}_- \alpha \mapsto \frac{\hat{\delta}_- \alpha}{c\bar{\tau} + d}, \tag{B.7}$$

which we now need to prove for the instanton contributions given by (B.4). To this end, we note that the result (4.9) implies that $-\partial_{\tilde{\xi}_a} G, \partial_{\xi^a} G, (1 - \xi^\Lambda \partial_{\xi^\Lambda}) G$ transform in the same way

as $\delta\xi^a, \delta\tilde{\xi}_\Lambda, \delta\alpha$. Due to this, from these derivatives one can form similar linear combinations

$$\begin{aligned} \hat{\partial}_{\xi^a} G &= \partial_{\xi^a} G - \kappa_{abc} \left(b^b + it^b \frac{1/t - t}{1/t + t} \right) \partial_{\xi_c} G, \\ \hat{G}_+ &= (1 - \xi^\Lambda \partial_{\xi^\Lambda} + \tau \partial_{\xi^0}) G + \frac{1}{2} \kappa_{abc} \left(b^a + it^a \frac{1/t - t}{1/t + t} \right) \left(\xi^b - c^b + \tau b^b - \frac{2t^b(\xi^0 - \tau)}{1/t + t} \right) \partial_{\xi_c} G, \\ \hat{G}_- &= (1 - \xi^\Lambda \partial_{\xi^\Lambda} + \bar{\tau} \partial_{\xi^0}) G + \frac{1}{2} \kappa_{abc} \left(b^a + it^a \frac{1/t - t}{1/t + t} \right) \left(\xi^b - c^b + \bar{\tau} b^b + \frac{2t^b(\xi^0 - \bar{\tau})}{1/t + t} \right) \partial_{\xi_c} G, \end{aligned} \tag{B.8}$$

which satisfy the same transformation laws as in (B.7),

$$\hat{\partial}_{\xi^a} G \mapsto \hat{\partial}_{\xi^a} G, \quad \hat{G}_+ \mapsto \frac{\hat{G}_+}{c\tau + d}, \quad \hat{G}_- \mapsto \frac{\hat{G}_-}{c\bar{\tau} + d}. \tag{B.9}$$

Using (B.4) to express (B.6) in terms of (B.8), one finds

$$\begin{aligned} \hat{\delta}_{\tilde{\xi}^a} &= -\sum'_{m,n} \oint_{C_{m,n}} \frac{dt'}{2\pi it'} \left[K(t, t') \hat{\partial}_{\xi^a} G - 2i \frac{\kappa_{abc} t^b \partial_{\xi_c} G_{m,n}}{(1/t' + t')^2} \right], \\ \hat{\delta}_+ \alpha &= -\sum'_{m,n} \oint_{C_{m,n}} \frac{dt'}{2\pi it'} \left[K(t, t') \hat{G}_+ + 2i \kappa_{abc} \left(\frac{(1 + tt') \tau_2 t^a}{(t - i)(t' - i)} - (c^a - \tau b^a) \right) \frac{t^b \partial_{\xi_c} G_{m,n}}{(1/t' + t')^2} \right], \\ \hat{\delta}_- \alpha &= -\sum'_{m,n} \oint_{C_{m,n}} \frac{dt'}{2\pi it'} \left[K(t, t') \hat{G}_- - 2i \kappa_{abc} \left(\frac{(1 + tt') \tau_2 t^a}{(t + i)(t' + i)} + (c^a - \bar{\tau} b^a) \right) \frac{t^b \partial_{\xi_c} G_{m,n}}{(1/t' + t')^2} \right]. \end{aligned} \tag{B.10}$$

In each of these expressions, the first term in the square bracket manifestly transforms as in (B.7). To see that this is also true for the remaining terms, we rewrite them in terms of the Cayley-rotated coordinate z as

$$\begin{aligned} \hat{\delta}^{(2)} \tilde{\xi}^a &= -\kappa_{abc} t^b \sum'_{m,n} \oint_{C_{m,n}} \frac{dz'}{8\pi^2 z'^2} (1 - z'^2) \partial_{\tilde{\xi}_c^{[0]}} G_{m,n}, \\ \hat{\delta}_+^{(2)} \alpha &= -\kappa_{abc} t^a \sum'_{m,n} \oint_{C_{m,n}} \frac{dz'}{8\pi z'^2} \left(c^b - \tau b^b - \frac{1}{2} (z + z') \tau_2 t^b \right) (1 - z'^2) \partial_{\tilde{\xi}_c^{[0]}} G_{m,n}, \\ \hat{\delta}_-^{(2)} \alpha &= \kappa_{abc} t^a \sum'_{m,n} \oint_{C_{m,n}} \frac{dz'}{8\pi z'^2} \left(c^b - \bar{\tau} b^b + \frac{1}{2} \frac{z + z'}{zz'} \tau_2 t^b \right) (1 - z'^2) \partial_{\tilde{\xi}_c^{[0]}} G_{m,n}. \end{aligned} \tag{B.11}$$

Given that, under S-duality,

$$(1 - z^2) \partial_{\tilde{\xi}_a} G \mapsto \frac{(1 - z^2) \partial_{\tilde{\xi}_a} G}{c\tau + d}, \tag{B.12}$$

our claim trivially follows. It is interesting to note that the contribution $\hat{\delta}^{(2)} \tilde{\xi}_a$ can be removed by adjusting the mirror map (4.6) by a further shift of $\tilde{\zeta}_a$ by

$$\Delta \tilde{\zeta}_a = -\kappa_{abc} t^b \sum'_{m,n} \oint_{C_{m,n}} \frac{dt}{\pi t} \frac{\partial_{\tilde{\xi}_c^{[0]}} G_{m,n}}{(1/t + t)^2}. \tag{B.13}$$

However, such a correction is modular invariant by itself and thus represents an inherent ambiguity in the definition of the mirror-map.

Finally, we turn to the contact potential (4.7). In the one-instanton approximation, taking into account the correction to Y^a coming from (4.6), we find

$$e^\Phi = \frac{\tau_2^2}{2} \mathcal{V}(t^a) - \frac{1}{16\pi} \sum'_{m,n} \oint_{C_{m,n}} \frac{dt}{t} A, \tag{B.14}$$

where

$$A = (t^{-1}Y^\Lambda - t\bar{Y}^\Lambda) \partial_{\xi^\Lambda} G_{m,n} + (t^{-1}F_a(Y) - t\bar{F}_a(\bar{Y})) \partial_{\bar{\xi}_a} G_{m,n} + 2\tau_2 \kappa_{abc} t^a t^b \frac{t^{-1} - t}{(t^{-1} + t)^2} \partial_{\bar{\xi}_c} G_{m,n}. \tag{B.15}$$

Using the transformation properties

$$\begin{aligned} \frac{dt}{t} &\mapsto \frac{|c\tau + d|}{c\xi^0 + d} \frac{dt}{t}, & \frac{t^{-1} - t}{(t^{-1} + t)^2} &\mapsto \frac{(c\xi^0 + d)^2}{|c\tau + d|^2} \frac{t^{-1} - t}{(t^{-1} + t)^2} - \frac{c\tau_2}{2} \frac{c\xi^0 + d}{|c\tau + d|^2}, \\ A &\mapsto \frac{(c\xi^0 + d)}{|c\tau + d|^2} A + \frac{c(c\xi^0 + d)}{|c\tau + d|^2} t \partial_t \left(\frac{t^{-1} + t}{c\xi^0 + d} G_{m',n'} \right) + \text{reg.} \end{aligned} \tag{B.16}$$

we see that the integrand transforms by a total derivative, and therefore that the contact potential transforms as in (3.4).

Open Access. This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] B. de Wit, P. Lauwers and A. Van Proeyen, *Lagrangians of $N = 2$ supergravity-matter systems*, *Nucl. Phys. B* **255** (1985) 569 [INSPIRE].
- [2] J. Bagger and E. Witten, *Matter couplings in $N = 2$ supergravity*, *Nucl. Phys. B* **222** (1983) 1 [INSPIRE].
- [3] S. Alexandrov, *Twistor approach to string compactifications: a review*, [arXiv:1111.2892](https://arxiv.org/abs/1111.2892) [INSPIRE].
- [4] S.M. Salamon, *Quaternionic Kähler manifolds*, *Invent. Math.* **67** (1982) 143.
- [5] C. LeBrun, *Quaternionic-Kähler manifolds and conformal geometry*, *Math. Ann.* **284** (1989) 353.
- [6] A. Swann, *Hyper-Kähler and quaternionic Kähler geometry*, *Math. Ann.* **289** (1991) 421.
- [7] A. Karlhede, U. Lindström and M. Roček, *Selfinteracting tensor multiplets in $N = 2$ superspace*, *Phys. Lett. B* **147** (1984) 297 [INSPIRE].
- [8] N.J. Hitchin, A. Karlhede, U. Lindström and M. Roček, *Hyper-Kähler metrics and supersymmetry*, *Commun. Math. Phys.* **108** (1987) 535 [INSPIRE].
- [9] B. de Wit, M. Roček and S. Vandoren, *Hypermultiplets, hyper-Kähler cones and quaternion Kähler geometry*, *JHEP* **02** (2001) 039 [[hep-th/0101161](https://arxiv.org/abs/hep-th/0101161)] [INSPIRE].

- [10] U. Lindström and M. Roček, *Properties of hyper-Kähler manifolds and their twistor spaces*, *Commun. Math. Phys.* **293** (2010) 257 [[arXiv:0807.1366](#)] [[INSPIRE](#)].
- [11] S. Alexandrov, B. Pioline, F. Saueressig and S. Vandoren, *Linear perturbations of hyper-Kähler metrics*, *Lett. Math. Phys.* **87** (2009) 225 [[arXiv:0806.4620](#)] [[INSPIRE](#)].
- [12] S. Alexandrov, B. Pioline, F. Saueressig and S. Vandoren, *Linear perturbations of quaternionic metrics*, *Commun. Math. Phys.* **296** (2010) 353 [[arXiv:0810.1675](#)] [[INSPIRE](#)].
- [13] D. Robles-Llana, M. Roček, F. Saueressig, U. Theis and S. Vandoren, *Nonperturbative corrections to 4D string theory effective actions from $SL(2, \mathbb{Z})$ duality and supersymmetry*, *Phys. Rev. Lett.* **98** (2007) 211602 [[hep-th/0612027](#)] [[INSPIRE](#)].
- [14] S. Alexandrov, B. Pioline, F. Saueressig and S. Vandoren, *D-instantons and twistors*, *JHEP* **03** (2009) 044 [[arXiv:0812.4219](#)] [[INSPIRE](#)].
- [15] S. Alexandrov, *D-instantons and twistors: some exact results*, *J. Phys. A* **42** (2009) 335402 [[arXiv:0902.2761](#)] [[INSPIRE](#)].
- [16] S. Alexandrov, D. Persson and B. Pioline, *Fivebrane instantons, topological wave functions and hypermultiplet moduli spaces*, *JHEP* **03** (2011) 111 [[arXiv:1010.5792](#)] [[INSPIRE](#)].
- [17] M. Roček, C. Vafa and S. Vandoren, *Hypermultiplets and topological strings*, *JHEP* **02** (2006) 062 [[hep-th/0512206](#)] [[INSPIRE](#)].
- [18] S. Alexandrov, F. Saueressig and S. Vandoren, *Membrane and fivebrane instantons from quaternionic geometry*, *JHEP* **09** (2006) 040 [[hep-th/0606259](#)] [[INSPIRE](#)].
- [19] D. Robles-Llana, F. Saueressig, U. Theis and S. Vandoren, *Membrane instantons from mirror symmetry*, *Commun. Num. Theor. Phys.* **1** (2007) 681 [[arXiv:0707.0838](#)] [[INSPIRE](#)].
- [20] B. Pioline and D. Persson, *The automorphic NS5-brane*, *Commun. Num. Theor. Phys.* **3** (2009) 697 [[arXiv:0902.3274](#)] [[INSPIRE](#)].
- [21] L. Bao, A. Kleinschmidt, B.E. Nilsson, D. Persson and B. Pioline, *Instanton corrections to the universal hypermultiplet and automorphic forms on $SU(2, 1)$* , *Commun. Num. Theor. Phys.* **4** (2010) 187 [[arXiv:0909.4299](#)] [[INSPIRE](#)].
- [22] B. Pioline and S. Vandoren, *Large D-instanton effects in string theory*, *JHEP* **07** (2009) 008 [[arXiv:0904.2303](#)] [[INSPIRE](#)].
- [23] S. Alexandrov, D. Persson and B. Pioline, *On the topology of the hypermultiplet moduli space in type-II/CY string vacua*, *Phys. Rev. D* **83** (2011) 026001 [[arXiv:1009.3026](#)] [[INSPIRE](#)].
- [24] S. Alexandrov, D. Persson and B. Pioline, *Wall-crossing, Rogers dilogarithm and the QK/HK correspondence*, *JHEP* **12** (2011) 027 [[arXiv:1110.0466](#)] [[INSPIRE](#)].
- [25] D. Gaiotto, G.W. Moore and A. Neitzke, *Four-dimensional wall-crossing via three-dimensional field theory*, *Commun. Math. Phys.* **299** (2010) 163 [[arXiv:0807.4723](#)] [[INSPIRE](#)].
- [26] D. Gaiotto, G.W. Moore and A. Neitzke, *Framed BPS states*, [arXiv:1006.0146](#) [[INSPIRE](#)].
- [27] B. Haghighat and S. Vandoren, *Five-dimensional gauge theory and compactification on a torus*, *JHEP* **09** (2011) 060 [[arXiv:1107.2847](#)] [[INSPIRE](#)].
- [28] A. Neitzke, *On a hyperholomorphic line bundle over the Coulomb branch*, [arXiv:1110.1619](#) [[INSPIRE](#)].

- [29] I. Antoniadis, S. Ferrara, R. Minasian and K. Narain, R^4 couplings in M and type-II theories on Calabi-Yau spaces, *Nucl. Phys. B* **507** (1997) 571 [[hep-th/9707013](#)] [[INSPIRE](#)].
- [30] H. Gunther, C. Herrmann and J. Louis, Quantum corrections in the hypermultiplet moduli space, *Fortsch. Phys.* **48** (2000) 119 [[hep-th/9901137](#)] [[INSPIRE](#)].
- [31] I. Antoniadis, R. Minasian, S. Theisen and P. Vanhove, String loop corrections to the universal hypermultiplet, *Class. Quant. Grav.* **20** (2003) 5079 [[hep-th/0307268](#)] [[INSPIRE](#)].
- [32] S. Cecotti, S. Ferrara and L. Girardello, Geometry of type II superstrings and the moduli of superconformal field theories, *Int. J. Mod. Phys. A* **4** (1989) 2475 [[INSPIRE](#)].
- [33] S. Ferrara and S. Sabharwal, Quaternionic manifolds for type II superstring vacua of Calabi-Yau spaces, *Nucl. Phys. B* **332** (1990) 317 [[INSPIRE](#)].
- [34] S. Alexandrov and F. Saueressig, Quantum mirror symmetry and twistors, *JHEP* **09** (2009) 108 [[arXiv:0906.3743](#)] [[INSPIRE](#)].
- [35] S. Alexandrov, J. Manschot and B. Pioline, $D3$ -instantons, mock theta series and twistors, [arXiv:1207.1109](#) [[INSPIRE](#)].
- [36] R. Bohm, H. Gunther, C. Herrmann and J. Louis, Compactification of type IIB string theory on Calabi-Yau threefolds, *Nucl. Phys. B* **569** (2000) 229 [[hep-th/9908007](#)] [[INSPIRE](#)].
- [37] P. Candelas, X.C. De La Ossa, P.S. Green and L. Parkes, A pair of Calabi-Yau manifolds as an exactly soluble superconformal theory, *Nucl. Phys. B* **359** (1991) 21 [[INSPIRE](#)].
- [38] S. Hosono, A. Klemm, S. Theisen and S.-T. Yau, Mirror symmetry, mirror map and applications to Calabi-Yau hypersurfaces, *Commun. Math. Phys.* **167** (1995) 301 [[hep-th/9308122](#)] [[INSPIRE](#)].
- [39] D. Robles-Llana, F. Saueressig and S. Vandoren, String loop corrected hypermultiplet moduli spaces, *JHEP* **03** (2006) 081 [[hep-th/0602164](#)] [[INSPIRE](#)].
- [40] S. Alexandrov, Quantum covariant c -map, *JHEP* **05** (2007) 094 [[hep-th/0702203](#)] [[INSPIRE](#)].
- [41] K. Becker, M. Becker and A. Strominger, Five-branes, membranes and nonperturbative string theory, *Nucl. Phys. B* **456** (1995) 130 [[hep-th/9507158](#)] [[INSPIRE](#)].
- [42] M.B. Green and M. Gutperle, Effects of D instantons, *Nucl. Phys. B* **498** (1997) 195 [[hep-th/9701093](#)] [[INSPIRE](#)].
- [43] A. Neitzke, B. Pioline and S. Vandoren, Twistors and black holes, *JHEP* **04** (2007) 038 [[hep-th/0701214](#)] [[INSPIRE](#)].
- [44] M. Kontsevich and Y. Soibelman, Stability structures, motivic Donaldson-Thomas invariants and cluster transformations, [arXiv:0811.2435](#).
- [45] S. Alexandrov and P. Roche, TBA for non-perturbative moduli spaces, *JHEP* **06** (2010) 066 [[arXiv:1003.3964](#)] [[INSPIRE](#)].
- [46] A. Kapustin and E. Witten, Electric-magnetic duality and the geometric Langlands program, *Commun. Num. Theor. Phys.* **1** (2007) 1 [[hep-th/0604151](#)] [[INSPIRE](#)].
- [47] K. Galicki, A generalization of the momentum mapping construction for quaternionic Kähler manifolds, *Commun. Math. Phys.* **108** (1987) 117 [[INSPIRE](#)].
- [48] J.M. Maldacena, A. Strominger and E. Witten, Black hole entropy in M -theory, *JHEP* **12** (1997) 002 [[hep-th/9711053](#)] [[INSPIRE](#)].

- [49] D. Gaiotto, A. Strominger and X. Yin, *From AdS_3/CFT_2 to black holes/topological strings*, *JHEP* **09** (2007) 050 [[hep-th/0602046](#)] [[INSPIRE](#)].
- [50] D. Gaiotto, A. Strominger and X. Yin, *The M5-brane elliptic genus: modularity and BPS states*, *JHEP* **08** (2007) 070 [[hep-th/0607010](#)] [[INSPIRE](#)].
- [51] J. de Boer, M.C. Cheng, R. Dijkgraaf, J. Manschot and E. Verlinde, *A Farey tail for attractor black holes*, *JHEP* **11** (2006) 024 [[hep-th/0608059](#)] [[INSPIRE](#)].
- [52] F. Denef and G.W. Moore, *Split states, entropy enigmas, holes and halos*, *JHEP* **11** (2011) 129 [[hep-th/0702146](#)] [[INSPIRE](#)].
- [53] A. Haydys, *Hyper-Kähler and quaternionic Kähler manifolds with S^1 -symmetries*, *J. Geom. Phys.* **58** (2008) 293.