

Three new triterpenoids from *Rubia schumanniana*

Bin KUANG,^{a,b} Jing HAN,^{a,b} Guang-Zhi ZENG,^a Xiao-Qiang CHEN,^a Wen-Jun HE,^a and Ning-Hua TAN^{a,*}^aState Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650201, China^bGraduate University of Chinese Academy of Sciences, Beijing 100049, China

Received 15 May 2012; Accepted 25 June 2012

© The Author(s) 2012. This article is published with open access at Springerlink.com

Abstract: Three new triterpenoids, 3 β -hydroxy-urs-30-*p*-Z-hydroxycinnamoyl-12-en-28-oic-acid (**1**), 3 β -hydroxy-olean-30-*p*-E-hydroxycinnamoyl-12-en-28-oic-acid (**2**) and 3 β ,6 α -dihydroxy-urs-14-en-12-one (**3**), together with seven known triterpenoids, were isolated from the roots of *Rubia schumanniana*. Their structures were established by means of spectroscopic analysis. All compounds were evaluated for cytotoxic activity, and compounds **2–6** showed cytotoxicity with the IC₅₀ values of 10.75–18.87 μ g/mL.


Keywords: *Rubia schumanniana*, triterpenoid, cytotoxicity

Introduction

Rubia schumanniana, an endemic species, is mainly distributed in southwest China. As one of the substitutes of traditional Chinese medicine *R. cordifolia*, its roots have been used for the treatment of tuberculosis, rheumatism, contusion, febrility and menoxenia. Previous studies on this plant have resulted in the isolation of seven quinones and β -sitosterol.^{1,2} As part of our continuing research on chemical constituents of medicinal plants from the genus *Rubia*, a systematic phytochemical investigation of the roots of *R. schumanniana* was carried out, which led to the isolation of three new triterpenoids (**1–3**), along with seven known triterpenoids, zanic acid (**4**),³ maslinic acid (**5**),⁴ ursolic acid (**6**),⁵ rubifolic acid (**7**),⁶ oleanolic acid (**8**),⁷ karachic acid (**9**),⁸ and rubiarbonol K (**10**).⁹ All compounds were evaluated for cytotoxicity against three human cancer cell lines (Hela, BGC-823, A549). Herein, we report the isolation, structural determination, and cytotoxic activities of these compounds.

Results and Discussion

Compound **1** was obtained as a white powder with a positive specific rotation ($[\alpha]_D^{25} + 6.5$). Its molecular formula, C₃₉H₅₄O₆, was deduced by HRESIMS (m/z 617.3856 [M – H][–]), indicating 13 degrees of unsaturation. The IR spectrum showed absorption bands for hydroxyl (3426 cm^{–1}), carbonyl (1689 cm^{–1}) and olefinic (1632 cm^{–1}) groups. The ¹³C NMR spectrum of **1** (Table 1) exhibited 39 carbons, including one trisubstituted double bond (δ_C 126.7, 139.3), one carboxyl (δ_C 180.3) and one *p*-hydroxycinnamoyl group, six methyls, ten



methylenes (one oxygenated), six methines (one oxygenated), and five quaternary carbons. Comparison of the NMR data of **1** with those of zanic acid (**4**) revealed that both compounds are ursane-type triterpenoids. The only difference between them was that the coupling constant of the disubstituted double bond in *p*-hydroxycinnamoyl group is 13.0 Hz in **1** while 16.0 Hz in **4**. The HMBC correlations of H-30 with the ester carbonyl carbon, C-19, C-20, and C-21 enabled the *p*-Z-hydroxycinnamoyl group to be placed at C-30 (Figure 1). The relative configuration of **1** was deduced from the analysis of its ROESY spectrum (Figure 2). The observed NOE correlations of H-3/H-5 and Me-23, H-5/H-9, and H-9/Me-27 indicated that H-3, H-5, H-9, Me-23 and Me-27 are cofacial and assigned as α -oriented. In turn the cross-peaks of Me-25/Me-24 and Me-26, and H-20/H-18 and Me-29 indicated the β -oriented of H-18, H-20, Me-24, Me-25, Me-26 and Me-29. From the above evidences, the structure of **1** was established as 3 β -hydroxy-urs-30-*p*-Z-hydroxycinnamoyl-12-en-28-oic-acid.

Compound **2** exhibited the same molecular formula C₃₉H₅₄O₆ as **1**, as established by HREIMS at m/z 618.3890 [M]⁺. The NMR data of **2** (Table 1) were similar to those of oleanolic acid (**8**) except for the presence of one *p*-E-hydroxycinnamoyl group in the downfield region of **2** and the

*To whom correspondence should be addressed. E-mail: nhtan@mail.kib.ac.cn


Table 1. ^1H NMR and ^{13}C NMR data for compounds 1–3 in pyridine- d_5

pos.	1		2		3	
	$\delta_{\text{H}}^{\text{d}}$ (J in Hz)	$\delta_{\text{C}}^{\text{a}}$, type	$\delta_{\text{H}}^{\text{d}}$ (J in Hz)	$\delta_{\text{C}}^{\text{b}}$, type	$\delta_{\text{H}}^{\text{c}}$ (J in Hz)	$\delta_{\text{C}}^{\text{a}}$, type
1a	0.98, overlap	39.5, CH ₂	1.01, overlap	39.0, CH ₂	1.09, overlap	38.4, CH ₂
1b	1.56, overlap		1.54, m			
2a	1.85, overlap	28.6, CH ₂	1.85, m	28.1, CH ₂	1.92, m	28.2, CH ₂
2b	2.34, m		2.18, m			
3	3.48, dd (10.0, 6.0)	78.6, CH	3.46, m	78.1, CH	3.57, m	78.8, CH
4		39.9, C		39.4, C		40.6, C
5	0.86, overlap	56.3, CH	0.87, overlap	55.8, CH	1.34, d (10.6)	61.4, CH
6a	1.37, overlap	19.3, CH ₂	1.37, m	18.8, CH ₂	4.43, dt (10.6, 3.6)	68.0, CH
6b	1.56, overlap		1.56, overlap			
7a	1.37, overlap	34.0, CH ₂	1.32, overlap	33.3, CH ₂	1.96, overlap	53.1, CH ₂
7b	1.56, overlap		1.49, m		2.58, dd (12.0, 3.6)	
8		40.4, C		39.8, C		42.1, C
9	1.64, overlap	48.5, CH	1.67, overlap	48.1, CH	2.25, overlap	51.2, CH
10		37.8, C		37.4, C		40.1, C
11a	1.97, overlap	24.1, CH ₂	1.94, m	23.8, CH ₂	2.48, dd (16.8, 11.2)	38.2, CH ₂
11b					2.80, dd (16.8, 9.2)	
12	5.51, br. s	126.7, CH	5.56, br. s	123.1, CH		215.8, C
13		139.3, C		139.9, C		54.2, C
14		42.9, C		42.2, C		157.2, C
15	2.33, m	29.1, CH ₂	2.20, m	28.3, CH ₂	5.75, dd (8.0, 2.0)	119.1, CH
16a	2.05, overlap	25.3, CH ₂	2.01, m	23.9, CH ₂	1.52, overlap	38.1, CH ₂
16b	2.14, m		2.18, m		2.13, d (14.8)	
17		48.3, C		47.0, C		35.5, C
18	2.69, d (11.0)	53.8, CH	3.40, m	41.1, CH	2.25, overlap	49.3, CH
19	1.85, overlap	34.9, CH	2.06, m	41.1, CH ₂	1.43, overlap	37.6, CH
20	1.42, overlap	44.4, CH		35.2, C	1.30, overlap	37.4, CH
21a	1.62, overlap	26.1, CH ₂	1.37, m	29.2, CH ₂	1.05, overlap	29.3, CH ₂
21b	1.85, overlap		1.69, m		1.60, m	
22a	1.97, overlap	37.4, CH ₂	1.90, m	32.3, CH ₂	1.25, overlap	38.7, CH ₂
22b	2.05, overlap		2.09, m			
23	1.26, s	29.3, CH ₃	1.25, s	28.8, CH ₃	2.02, s	32.4, CH ₃
24	1.04, s	17.1, CH ₃	1.05, s	16.6, CH ₃	1.48, s	17.1, CH ₃
25	0.90, s	16.2, CH ₃	0.91, s	15.6, CH ₃	1.03, s	17.1, CH ₃
26	1.07, s	17.9, CH ₃	1.05, s	17.5, CH ₃	1.07, s	25.5, CH ₃
27	1.22, s	24.4, CH ₃	1.32, s	26.2, CH ₃	1.43, s	21.1, CH ₃
28		180.3, C		179.9, C	0.88, s	34.0, CH ₃
29	1.07, overlap	17.6, CH ₃	1.19, s	19.5, CH ₃	1.14, d (6.4)	25.1, CH ₃
30a	4.26, dd (11.0, 7.5)	68.2, CH ₂	4.14, d (10.5)	74.9, CH ₂	0.97, d (6.4)	22.8, CH ₃
30b	4.49, dd (11.0, 3.0)		4.23, d (10.5)			
1'		167.8, C		167.7, C		
2'	6.07, d (13.0)	116.9, CH	6.77, d (16.0)	115.3, CH		
3'	7.01, d (13.0)	144.6, CH	8.07, d (16.0)	144.6, CH		
4'		127.1, C		126.3, C		
5'	8.10, d (8.5)	134.1, CH	7.68, d (8.0)	130.8, CH		
6'	7.22, overlap	116.5, CH	7.21, overlap	116.9, CH		
7'		161.1, C		161.7, C		
8'	7.22, overlap	116.5, CH	7.21, overlap	116.9, CH		
9'	8.10, d (8.5)	134.1, CH	7.68, d (8.0)	130.8, CH		

^aDate were measured at 100 MHz; ^bDate were measured at 125 MHz; ^cDate were measured at 400 MHz; ^dDate were measured at 500 MHz.

replacement of one methyl in **8** by one hydroxymethyl group (δ_{C} 74.9). HMBC correlations of H-30 with the ester carbonyl carbon (δ_{C} 167.7), C-19, C-20, C-21, and Me-29 indicated that the *p*-*E*-hydroxycinnamoyl group located at C-30 (Figure 1). The observed NOE correlations (Figure 2) of H-3/H-5 and Me-23, H-5/H-9, and H-9/Me-27 indicated that H-3, H-5, H-9, Me-23 and Me-27 are cofacial and assigned as α -oriented. In turn the cross-peaks of Me-25/Me-24 and Me-26, and H-18/Me-29 indicated the β -oriented of H-18, Me-24, Me-25, Me-26 and Me-29. Thus, the structure of **2** was assigned as 3β -hydroxy-olean-30-*p*-*E*-hydroxycinnamoyl-12-en-28-oic-acid.

Compound **3** gave the molecular formula $\text{C}_{30}\text{H}_{48}\text{O}_3$, based on HRESIMS (m/z 479.3504 $[\text{M} + \text{Na}]^+$), requiring seven degrees of unsaturation. The IR spectrum showed absorption bands for hydroxyl (3442 cm^{-1}), carbonyl (1705 cm^{-1}) and olefinic (1640 cm^{-1}) groups. The ^{13}C NMR spectrum data (Table 1) showed the presence of 30 carbon signals due to one trisubstituted double bond (δ_{C} 119.1, 157.2), one ketone carbon (δ_{C} 215.8), eight methyls, seven methylenes, seven

**Figure 1.** Key ^1H - ^1H COSY and HMBC correlations of 1–3

methines (two oxygenated), and five quaternary carbons. Comparison of the NMR data of **3** with those of ursolic acid suggested that their structures are closely related.¹⁰ The main differences were that one characteristic trisubstituted double bond at C-12/C-13 in conventional ursane-type triterpenoids was absent in **3**, while one different trisubstituted double bond, one carbonyl group, and one additional hydroxy group were present. The double bond was placed between C-14 and C-15, as determined by HMBC correlations (Figure 1) of H-15 (δ_{H} 5.75) with C-17 (δ_{C} 35.5) and of Me-26 and Me-27 with C-14 (δ_{C} 157.2). The location of the ketone carbon (δ_{C} 215.8) at C-12 was elucidated by HMBC correlations of H-9, H-11 and Me-27 with C-12. In addition, the position of the additional hydroxy group at C-6 was deduced by correlations of H-6 with H-5 in the ^1H - ^1H COSY spectrum combined with HMBC correlations of H-5 with C-6 (Figure 1). Thus, the planar structure of **3** was established. The α -orientations of H-3, H-5, H-9, and Me-23 were established by NOE correlations of H-3/H-5 and Me-23 and H-5/H-9, and the β -orientations of H-6, H-18, H-20, Me-24, Me-25, Me-26, Me-27, Me-28, Me-29 were deduced by NOE correlations of H-6/Me-24 and Me-25, H-7 β /H-6 and Me-26, H-18/Me-27, Me-28 and Me-29, and Me-29/H-20 (Figure 2). Accordingly, the structure of **3** was elucidated as 3 β , 6 α -dihydroxy-urs-14-en-12-one.




Figure 2. Key ROESY correlations of **1–3**

All compounds were evaluated for cytotoxicity against three human cancer cell lines, HeLa (human cervical carcinoma), BGC-823 (human stomach adenocarcinoma), and A549 (human lung adenocarcinoma), and results indicated that compounds **2–6** showed cytotoxicity with the IC_{50} values of 10.75–18.87 $\mu\text{g}/\text{mL}$ (Table 2).

Experimental Section

General Experimental Procedures. Optical rotations were measured with a Horiba SEPA-300 polarimeter. IR spectra were obtained by a Bruker FT-IR Tensor 27 spectrophotometer using KBr pellets. UV spectra were obtained using a Shimadzu UV-2401A spectrophotometer. 1D and 2D NMR spectra were recorded on Bruker AX-400, DRX-500, or AV-600 spectrometers with TMS as an internal standard. Chemical shifts (δ) were expressed in ppm with reference to solvent signals. HREIMS were recorded on a Waters Auto Premier

Table 2. Cytotoxicity of **1–10** against cancer cell lines^a with IC_{50} ($\mu\text{g}/\text{mL}$)

compound	HeLa	A549	BGC-823
1	> 20	> 20	> 20
2	16.18	> 20	> 20
3	> 20	15.74	> 20
4	13.53	> 20	14.18
5	12.39	18.87	10.75
6	11.80	> 20	11.89
7	> 20	> 20	> 20
8	> 20	> 20	> 20
9	> 20	> 20	> 20
10	> 20	> 20	> 20
Taxol ^b	0.38	0.02	0.01

^aCell lines: HeLa human cervical carcinoma; BGC-823 human stomach adenocarcinoma; A549 human lung adenocarcinoma.

^bpositive control.

P776 spectrometer. HRESIMS were recorded on an API QSTER time-of-flight spectrometer. Analytical or Semipreparative HPLC was performed on an Agilent 1100 liquid chromatograph with a Zorbax Eclipse-C₁₈ (4.6 mm \times 150 mm; 9.4 mm \times 250 mm) column. Column chromatographies were performed using silica gel (200–300 mesh, Qingdao Yu-Ming-Yuan Chemical Co. Ltd., Qingdao, China), Sephadex LH-20 (Pharmacia Fine Chemical Co., Uppsala, Sweden), and Lichroprep RP-18 gel (40–63 μm , Merck, Darmstadt, Germany). Fractions were monitored by TLC (GF 254, Qingdao Yu-Ming-Yuan Chemical Co. Ltd., Qingdao, China), and spots were visualized by heating silica gel plates sprayed with 10% H_2SO_4 in EtOH.

Plant Material. The roots of *R. schumanniana* were purchased in August 2009 from the Yunnan Lv-Sheng Pharmaceutical Co. Ltd., Kunming, China. The material was identified by Prof. Xi-Wen Li of Kunming Institute of Botany. A voucher specimen (KUN0328859) was deposited at the Herbarium of Kunming Institute of Botany, Chinese Academy of Sciences.

Extraction and Isolation. The dried and powdered roots of *R. schumanniana* (50 kg) were extracted with 70% aqueous MeOH (40 L \times 3) for 12 hours at room temperature. After removal of the solvent under reduced pressure, the MeOH extract (18.6 kg) was suspended in H_2O and partitioned successively with EtOAc and *n*-BuOH to give an EtOAc-soluble portion (3.7 kg) and a *n*-BuOH-soluble portion (4.2 kg). The EtOAc part was chromatographed on silica gel column eluting with chloroform-methanol (1:0, 95:5, 9:1, 8:2, 7:3, and 0:1) to afford fractions I–IV. Fraction I (9:1, 163 g) was further chromatographed on silica gel using a petroleum ether-acetone gradient (10:1 to 0:1) as the eluent to yield 6 subfractions, I-1–I-6. Subfraction I-1 was chromatographed with RP-18, and then separated by semi-preparative HPLC ($\text{CH}_3\text{CN}:\text{H}_2\text{O}$, 80:20) to yield **1** (12 mg) and **2** (2.6 mg). **4** (23 mg) was purified from subfractions I-2 by repeated chromatography with silica gel. Subfraction I-4 was chromatographed on silica gel using a chloroform-methanol gradient (50:1 to 10:1) as the eluent, and then purified over Sephadex LH-20 eluted with chloroform-methanol (1:1), then by semi-preparative HPLC ($\text{CH}_3\text{CN}:\text{H}_2\text{O}$, 60:40 and 73:27) to yield **5** (22 mg), **6** (8 mg), **8** (35 mg) and **3** (7 mg), respectively. Subfraction I-6 was chromatographed on silica gel using a

chloroform-acetone gradient (50:1 to 0:1) as the eluent, and then purified by semi-preparative HPLC (CH₃CN:H₂O, 65:35) to yield **7** (15 mg), **9** (10.2 mg) and **10** (6 mg).

3 β -Hydroxy-urs-30-*p*-Z-hydroxycinnamoyl-12-en-28-oic-acid (1): white powder; $[\alpha]_D^{16} + 6.5$ (*c* 0.08, MeOH:CHCl₃ = 1:1); UV (MeOH) λ_{\max} (log ϵ) 202 (4.16), 312 (4.14) nm; IR (KBr) ν_{\max} 3426, 2965, 2937, 2873, 1689, 1632, 1606, 1514, 1456, 1377, 1311, 1277, 1258, 1202, 1184, 1029, 997, 833, 519 cm⁻¹; ¹H and ¹³C NMR data, see Table 1; negative ESIMS *m/z* 617 [M – H]⁻; negative HRESIMS *m/z* 617.3856 [M – H]⁻ (calcd for C₃₉H₅₃O₆, 617.3842).

3 β -Hydroxy-olean-30-*p*-E-hydroxycinnamoyl-12-en-28-oic-acid (2): white amorphous powder; $[\alpha]_D^{16} + 13.1$ (*c* 0.13, MeOH); UV (MeOH) λ_{\max} (log ϵ) 202 (4.16), 227 (4.01), 313 (4.22) nm; IR (KBr) ν_{\max} 3429, 2938, 2875, 1692, 1632, 1606, 1515, 1456, 1387, 1167, 1027, 996, 833, 519 cm⁻¹; ¹H and ¹³C NMR data, see Table 1; positive EIMS *m/z* 618 [M]⁺; positive HRESIMS *m/z* 618.3890 [M]⁺ (calcd for C₃₉H₅₄O₆, 618.3920).

3 β , 6 α -Dihydroxy-urs-14-en-12-one (3): white amorphous powder; $[\alpha]_D^{20} - 10.7$ (*c* 0.10, MeOH); UV (MeOH) λ_{\max} (log ϵ) 201 (3.41) nm; IR (KBr) ν_{\max} 3442, 2927, 2866, 1705, 1640, 1462, 1382, 1140, 1036, 987 cm⁻¹; ¹H and ¹³C NMR data, see Table 1; positive ESIMS *m/z* 479 [M + Na]⁺; positive HRESIMS *m/z* 479.3504 [M + Na]⁺ (calcd for C₃₀H₄₈O₃Na, 479.3501).

Cytotoxicity Assay. The cytotoxicity of all compounds against Hela, A549, and BGC-823 cancer cell lines was measured using the sulforhodamine B (SRB) assay. Taxol was used as positive control. Cells were plated in 96-well culture plates for 24h before treated with serial dilutions of all compounds. After being incubated for 48 h, cells were fixed with 25 μ L of ice-cold 50% trichloroacetic acid and incubated at 4 °C for 1 h. After washing with distilled water and air-drying, the plate was stained for 15 min with 100 μ L of 0.4% SRB (Sigma) in 1% glacial acetic acid. The plates were washed with 1% acetic acid and air-dried. For reading the

plate, the protein-bound dye was dissolved in 100 μ L of 10 mM Tris base. The absorbance was measured at 560 nm. All tests were performed in triplicate, and results are expressed as IC₅₀ values.

Electronic Supplementary Material

Supplementary material is available in the online version of this article at <http://dx.doi.org/10.1007/s13659-012-0038-8> and is accessible for authorized users.

Acknowledgments

This work was supported by the National Natural Science Foundation of China (U1032602, 91013002, 30725048), the National New Drug Innovation Major Project of China (2011ZX09307-002-02), and the National Basic Research Program of China (2009 CB522300).

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] Liu, Y. L.; Bai, Y. L. *Acta Pharm. Sin.* **1985**, *20*, 53–58.
- [2] Liu, Y. L.; Chen, B. Z.; Bai, Y. L.; Duddeck, H.; Hiegemann, M. *Phytochemistry* **1991**, *30*, 947–949.
- [3] Siddiqui, B. S.; Firdous; Begum, S. *Phytochemistry* **1999**, *52*, 1111–1115.
- [4] Wu, Y. X.; Zhang, W.; Li, J. C.; Yang, L. J.; Liu, N. *Chin. Tradit. Herb. Drugs* **2011**, *42*, 2402–2406.
- [5] Zhu, C. C.; Gao, L.; Zhao, Z. X.; Lin, C. Z. *Acta Pharm. Sin.* **2012**, *47*, 77–83.
- [6] Talapatra, S. K.; Sarkar, A. C.; Talapatra, B. *Phytochemistry* **1981**, *20*, 1923–1927.
- [7] Sun, W. *Chin. Tradit. Herb. Drugs* **2012**, *43*, 23–26.
- [8] Khan, M. A.; Atta-ur-Rahman. *Phytochemistry* **1975**, *14*, 789–791.
- [9] Zou, C.; Hao, X. J.; Chen, C. X.; Zhou, J. *Acta Bot. Yunnan.* **1992**, *14*, 114.
- [10] Seebacher, W.; Simic, N.; Weis, R.; Saf, R.; Kunert, O. *Magn. Reson. Chem.* **2003**, *41*, 636–638.