

RESEARCH ARTICLE

Open Access

Antiprotozoal and antimycobacterial activities of *Persea americana* seeds

Adelina Jiménez-Arellanes^{1*}, Julieta Luna-Herrera², Ricardo Ruiz-Nicolás², Jorge Cornejo-Garrido¹, Amparo Tapia³ and Lilián Yépez-Mulia³

Abstract

Background: *Persea americana* seeds are widely used in traditional Mexican medicine to treat rheumatism, asthma, infectious processes as well as diarrhea and dysentery caused by intestinal parasites.

Methods: The chloroformic and ethanolic extracts of *P. americana* seeds were prepared by maceration and their amoebicidal, giardicidal and trichomonocidal activity was evaluated. These extracts were also tested against *Mycobacterium tuberculosis* H37Rv, four mono-resistant and two multidrug resistant strains of *M. tuberculosis* as well as five non tuberculosis mycobacterium strains by MABA assay.

Results: The chloroformic and ethanolic extracts of *P. americana* seeds showed significant activity against *E. histolytica*, *G. lamblia* and *T. vaginalis* (IC₅₀ <0.634 µg/ml). The chloroformic extract inhibited the growth of *M. tuberculosis* H37Rv, *M. tuberculosis* MDR SIN 4 isolate, three *M. tuberculosis* H37Rv mono-resistant reference strains and four non tuberculosis mycobacteria (*M. fortuitum*, *M. avium*, *M. smegmatis* and *M. abscessus*) showing MIC values ≤50 µg/ml. Contrariwise, the ethanolic extract affected only the growth of two mono-resistant strains of *M. tuberculosis* H37Rv and *M. smegmatis* (MIC ≤50 µg/ml).

Conclusions: The CHCl₃ and EtOH seed extracts from *P. americana* showed amoebicidal and giardicidal activity. Importantly, the CHCl₃ extract inhibited the growth of a MDR *M. tuberculosis* isolate and three out of four mono-resistant reference strains of *M. tuberculosis* H37Rv, showing a MIC = 50 µg/ml. This extract was also active against the NTM strains, *M. fortuitum*, *M. avium*, *M. smegmatis* and *M. abscessus*, with MIC values <50 µg/ml.

Keywords: Medicinal plant, Seeds, Antimycobacterial, Antiprotozoal, *Persea americana*

Background

Persea americana Mill. (Lauraceae) is an edible fruit commonly known as *aguacate* (avocado) that grows throughout the tropics. The seeds (crude or toasted) are employed in traditional Mexican medicine to treat skin rashes, diarrhea, and dysentery caused by helminths and amoebas, for the cure of infectious processes caused by fungi and bacteria, as well as for the treatment of asthma, high blood pressure, and rheumatism [1-5]. The seeds of *P. americana* used alone or mixed with other species, such as *Psidium guajava*, *Mentha piperita* or *Ocimum basilicum*, are mainly employed for the treatment of diarrhea [4].

The presence of fatty acids (linoleic, oleic, palmitic, stearic, linolenic, capric and myristic acids), polyphenols (catechin, isocatechin, procyanidin, flavonoids, tannins and proanthocyanidin monomers), saponins, glucosides (D-perseitol, D-α-manoheptitol, D-monoheptulose, perseitol), sterols (β-sitosterol, campesterol, stigmaterol, cholesterol), the amino acid carnitine and two glucosides of abscisic acid has been described for *P. americana* seeds [4-8]. High concentrations of catechins, procyanidins and hydroxycinnamic acid have recently been determined in 100% ethyl acetate (EtOAc), in 70% acetone and 70% methanol (MeOH) extracts obtained from *P. americana* peel and seeds, while the pulp extract was rich in hydroxybenzoic acid, hydroxycinnamic acid and procyanidins [9].

Interestingly, the hypolipemic effect of the MeOH extract obtained from *P. americana* seeds has been demonstrated in male rats with induced hypercholesterolemia

* Correspondence: adelinajim08@prodigy.net.mx

¹Unidad de Investigación Médica en Farmacología (UIM), Hospital de Especialidades, CMNS XXI, Instituto Mexicano del Seguro Social (IMSS), Av. Cuauhtémoc 330, Col. Doctores, 06720, México, DF, Mexico
Full list of author information is available at the end of the article

[10,11]. This extract reduced total cholesterol levels, triglycerides and Low density lipoprotein (LDL); on the other hand, increased the levels of High density lipoprotein (HDL). The same effect was described for the aqueous extract, which also reduced blood pressure both in normal rats and those with high blood pressure; in addition, it exerted a hypoglycemic effect on rats and rabbits with diabetes [12-15]. The aqueous extract showed a median Lethal dose (LD₅₀) = 10 g/kg in rats when it was administered orally. Importantly, it did not alter the hematological parameters nor the levels of Alanine aminotransferase (ALT), Aspartate aminotransferase (AST), albumin, and creatinine in male and female rats were treated for 28 days [16].

The hexanic and MeOH seed extracts of *P. americana* have been described to have a Minimum inhibitory concentration (MIC) of <1.25 µg/ml against *Candida* ssp., *Cryptococcus neoformans* and *Malassezia pachydermatis*. These extracts were also active against *Artemia salina*, with Lethal concentration (LC₅₀) values of 2.37 and 24.13 mg/ml, respectively. They were also active against *Aedes aegypti* larvae with LC₅₀ values of 16.7 and 8.9 mg/ml, respectively [17]. On the other hand, the MeOH extract from *P. americana* leaves inhibited completely the growth of *M. tuberculosis* H37Ra (MIC = 125 µg/ml) and H37Rv (MIC = 62.5 µg/ml); furthermore, the hexane fraction inhibited the growth of both mycobacteria with MIC = 31.2 µg/ml [18]. In addition, the EtOH extract was active against both Gram-positive and -negative bacteria (with the exception of *Staphylococcus epidermis* and *Escherichia coli*) with MIC = 500 µg/ml [19]. Regarding the bacterial activity of *P. americana* (var Hass and Fuerte), the acetone seed extract exhibited moderate activity against *Bacillus cereus*, *Staphylococcus aureus* and *Listeria monocytogenes* [9]. The trypanomicidal activity of the MeOH extract from *P. americana* seeds has been also tested [20]. It showed moderate activity when was evaluated at the concentration range of 250–500 µg/ml. In the case of the aqueous seed extract, it had a slight anti-*Giardia duodenalis* (syn *G. lamblia*) activity, inducing 23% of mortality at 4 mg/ml [21].

An important antioxidant activity (AOA) of the MeOH extract of *P. americana* seeds and leaves has been described by different methods [8,22-24]. Besides, AOA has been reported in the 100% EtOAc, 70% acetone and 70% MeOH of the peel, pulp and seed extract [9].

Up to now, the activity of the chloroformic (CHCl₃) and ethanol (EtOH) extracts obtained from *P. americana* seeds against anaerobic protozoan and *M. tuberculosis* H37Rv strains with different level of drug resistance has not yet been evaluated. Therefore, herein, the activity of both extracts was tested against the anaerobic protozoa *Giardia lamblia*, *Entamoeba histolytica* and *Trichomonas*

vaginalis. In addition, their antimycobacterial activity was evaluated against four mono-resistant reference strains of *M. tuberculosis* H37Rv, two MDR *M. tuberculosis* clinical isolates and five non-tuberculosis mycobacterium (NTM).

Methods

Plant material

P. americana seeds were obtained from the town of Ario de Rosales in the state of Michoacan, Mexico in August 2009. This material was then dried at room temperature and under conditions of darkness; the material was then ground. The plant was botanically identified by Abigail Aguilar, M.Sc., and a voucher specimen was deposited at the Herbarium of the Instituto Mexicano del Seguro Social, Mexico (IMSSM) with code number 14256.

Preparation of extracts

The dry and powdered plant material (1.364 kg) was macerated three times with CHCl₃ analytical reagent - AR- (J.T. Baker) at room temperature for 7 days. The extract was filtered and concentrated to dryness under low pressure at 40°C. The plant material was later macerated with EtOH AR grade (J.T. Baker) three times for 7 days and the solvent was eliminated under reduced pressure until the solvent-free extract was obtained.

Antiprotozoal activity evaluation

For this assay, the *E. histolytica* strain HM1-IMSS and the *T. vaginalis* GT9 strain were cultured in a TYI-S-33 modified medium supplemented with 10% calf serum; *G. lamblia* strain IMSS:0989:1 was maintained in a TYI-S-33 medium supplemented with 10% calf serum and bovine bile. *In vitro* susceptibility assays were performed according to the method previously described [25,26]. Briefly, 5 × 10⁴ trophozoites of *G. lamblia* were incubated for 48 h at 37°C with increasing concentrations of the EtOH and CHCl₃ extracts of *P. americana* seeds using Dimethyl sulfoxide (DMSO) as a suitable solvent. After incubation, *G. lamblia* trophozoites were washed and subcultured for an additional 48 h in a fresh medium alone. For *E. histolytica* and *T. vaginalis*, 6 × 10³ trophozoites were incubated for 72 h at 37°C with increasing concentrations of the samples tested. Metronidazole was included as a positive control; parasites without treatment but with the highest DMSO concentration used for sample dilutions were included as a negative control. *G. lamblia*, *T. vaginalis* and *E. histolytica* trophozoites were counted and the 50% Inhibitory concentration (IC₅₀) was calculated by Probit analysis. The experiments were carried out in triplicate and repeated at least twice.

Antimycobacterial evaluation

For this assay, 12 Mycobacterium strains were employed: *M. tuberculosis* H37Rv (ATCC 27294, a strain sensitive to streptomycin (STR), isoniazid (INH), rifampicin (RIF), etambutol (EMB), or pyrazinamide; *M. tuberculosis* SIN 4 (a MDR clinical isolate with resistance to first-line drugs); *M. tuberculosis* MMDO (an MDR clinical isolate with resistance to INH and EMB); four mono-resistant strains of *M. tuberculosis* H37Rv (INH-R, ATCC 35822; STR-R, ATCC 35820; RIF-R, ATCC 35838, and EMB-R, ATCC 35837) and four NTM clinical isolates (*M. fortuitum*, *M. chelonae*, *M. abscessus* and *M. avium*) and *M. smegmatis* (ATCC 35798). The Mycobacterium strains were grown and maintained in Middlebrook 7H9 broth supplemented with 10% OADC enrichment (Becton Dickenson, USA) at 37°C until a logarithmic growth phase was achieved. At the moment of evaluation, *M. tuberculosis* strains and NTM were diluted 1:20 and 1:50, respectively in the 7H9 medium.

The antimycobacterial activity was carried out by means of the Microplate alamar blue assay (MABA) previously described [26,27]. EtOH and CHCl₃ extracts (10 mg) were solubilized in 500 µL of DMSO and from these stock solutions, several dilutions were prepared to achieve concentrations between 200 and 3.13 µg/ml. The maximum DMSO concentration used in the MABA assay does not affect mycobacterial growth, as it was previously reported [28]. Extracts that presented MIC <100 µg/ml were considered to have good antimycobacterial activity. Rifampicin and isoniazid at 0.06 µg/mL were included as a positive control for the H37Rv strain, for the MDR clinical isolates and for NTM; the same drugs were used but at concentrations of 100 and 3.1 µg/ml, respectively.

Results and discussion

By means of the maceration process, 88.7 g of the CHCl₃ extract and 77.2 g of the EtOH extract from *P. americana* seeds were obtained with an average yield of 6% with respect to the plant material's dry weight. A preliminary phytochemical analysis by Thin layer chromatography (TLC) of the CHCl₃ extract led to the detection of the presence of β-sitosterol, phytol and palmitic acid. On the other hand, catechin and epicatechin were detected in the EtOH extract by TLC. All compounds were identified by comparison of the Retention factor (R_f) with their commercial reference. A previous study by Rodríguez-Carpena et al. [9] reported a high concentration of catechin and epicatechin in 100% EtOAc, 70% acetone and 70% MeOH extracts obtained from *P. americana* seeds and peel.

The EtOH and CHCl₃ extracts displayed significant antiprotozoal activity. For *G. lamblia*, *E. histolytica* and *T. vaginalis*, the CHCl₃ extract showed values of IC₅₀ = 0.634, 0.417 and 0.524 µg/ml, respectively. On the other hand, the EtOH extract exhibited IC₅₀

Table 1 Antiprotozoal activity of the CHCl₃ and EtOH extracts from *Persea americana* seeds

Sample	IC ₅₀ (µg/ml)		
	<i>G. lamblia</i>	<i>E. histolytica</i>	<i>T. vaginalis</i>
CHCl ₃ extract	0.634	0.417	0.524
EtOH extract	0.486	0.386	0.533
Metronidazole	0.210	0.06	0.037

IC₅₀: medium inhibitory concentration.

values of 0.486, 0.386 and 0.533 µg/ml against *G. lamblia*, *E. histolytica* and *T. vaginalis*, respectively (Table 1). Although the giardicidal activity of the aqueous seed extract has been previously evaluated [21], the authors used the 3-(4,5-dimetiltiazol-2-ilo)-2,5-difeniltetrazol (MTT) colorimetric method; therefore our data can not be compared.

It is interesting to note that, the reference drug, metronidazole (IC₅₀ = 0.210 µg/ml) showed only three and two times higher anti-*Giardia* activity than the CHCl₃ and EtOH extracts. However, in the case of *E. histolytica*, the CHCl₃ and EtOH extracts were seven and six times less potent than metronidazole (IC₅₀ = 0.060 µg/ml). For

Table 2 Antimycobacterial effect of the CHCl₃ and EtOH extracts from *Persea americana* seeds

<i>Mycobacterium tuberculosis</i>	Drug-resistance pattern	MIC (µg/ml)	
		CHCl ₃ extract	EtOH extract
H37Rv	INH-, RIF-, STR-, and EMB- susceptible	50	>100
Clinical isolates			
SIN4	STR, INH, RIF, EMB, RFB, ETH, and OFX	50	>100
MMDO	INH, EMB	100	>100
Mono-resistant			
RIF-R	RIF	>100	>100
INH-R	INH	50	>100
STR-R	STR	50	50
EMB-R	EMB	50	50
NTB <i>Mycobacterium</i>			
<i>M. fortuitum</i>		50	100
<i>M. avium</i>		25	100
<i>M. chelonae</i>		100	100
<i>M. smegmatis</i>		12.5	25
<i>M. abscessus</i>		25	100

SIN 4 MDR clinical isolate of *M. tuberculosis* resistant to STR: Streptomycin, INH: Isoniazid, RIF: Rifampicin, EMB: Ethambutol, RFB: Rifabutin, ETH: Ethionamide and OFX: Ofloxacin. MMDO MDR clinical isolates of *M. tuberculosis* resistant to INH and EMB; the resistance pattern was determined by Microdilution alamar blue assay (MABA). NTB: non-tuberculosis mycobacteria; MIC: Minimum inhibitory concentration. Data are means of three determinations.

T. vaginalis, these extracts showed weak activity, being 16 (CHCl₃) and 17 (EtOH) times less potent than metronidazole (CI₅₀ = 0.037 µg/ml).

In traditional Mexican medicine, *P. americana* seeds are used to treat diarrhea [1-4]. It is shown here that *P. americana* EtOH and CHCl₃ seed extracts are indeed responsible for the activity against two anaerobic parasites that cause diarrhea: *G. lamblia*, and *E. histolytica*. The *in vitro* and *in vivo* giardicidal activity of epicatechin isolated from plants such as *Rubus coriifolius* and *Heliantemum glomeratus* has been reported [29,30]. In this regard, epicatechin was detected in *P. americana* EtOH seed extract by TLC analysis; hence, this compound may be responsible for the giardicidal activity observed in this study. Further studies are required to define the active compound(s) responsible for the antiprotozoal activity of the CHCl₃ extract.

It is worth considering the fact that the EtOH and CHCl₃ extracts constitute potential sources of compounds that can be employed as prototype molecules for the development of novel antiprotozoal agents as an alternative treatment of clinical isolates with metronidazole resistance.

The results of the antimycobacterial activity of the EtOH and CHCl₃ extracts determined by MABA assay are presented in Table 2. It is important to note that the CHCl₃ extract inhibited the growth of *M. tuberculosis* H37Rv, MDR *M. tuberculosis* SIN 4 and three out of four mono-resistant reference strains of *M. tuberculosis* H37Rv (INH-R, STR-R, and EMB-R), showing a MIC = 50 µg/ml. This extract was also active against the NTM: *M. fortuitum*, *M. avium*, *M. smegmatis* and *M. abscessus* with MIC values < 50 µg/ml. However, the EtOH extract affected only the growth of *M. smegmatis* (MIC = 25 µg/ml) and the mono-resistant strains of *M. tuberculosis* H37Rv STR-R and EMB-R (MIC = 50 µg/ml).

To date, the hexanic and MeOH extracts obtained from the stems and leaves of *P. americana* have been reported to inhibit the growth of *M. tuberculosis* H37Rv and *M. tuberculosis* H37Ra strains [18,31]. The reported antimycobacterial activity of the MeOH extracts was attributed to the presence of lignans [31] and the MIC was slightly lower than the CHCl₃ extract tested in this study.

Since the presence of MDR and extensively drug-resistant (XDR) *M. tuberculosis* cases is increasing rapidly and current chemotherapy is prolonged, poorly effective, expensive and is accompanied by severe side effects [26]; it is necessary to have recourse to an alternative treatment against these strains or that can even aid in and/or shorten the currently available that could have a different mechanism of action. Therefore, the high activity exhibited by the *P. americana* CHCl₃ seed extract against mono-resistant strains of *M. tuberculosis* H37Rv, MDR clinical isolates and NTM that is described in this study, it is of great interest.

From the clinical point of view, NTM are becoming relevant, because of the so-called mycobacteriosis and are currently recognized as important pathogens associated with both immune-deficient and immune-competent patients. The mycobacteria tested in this study are representative of the most common NTM isolated from pulmonary cases (*M. abscessus* and *M. avium*) or associated with soft tissue infections (*M. fortuitum* and *M. chelonae*) [32]. Although *M. smegmatis* is a poor pathogenic bacterium, it was included in the NTM group because it is widely employed in the determination of the antimycobacterial activity of new compounds [32,33]. In the case of NTM, the majority is naturally resistant to some of the first-line anti-TB drugs such as INH and RIF; thus, effective drugs against NTM are scarce than those for TB, emphasizing the urgency of finding novel active compounds that could be used in the treatment of the NTM group. Based on our results, *P. americana* seeds may be a source for potential moieties (molecules) against NTM. We are currently conducting the isolation and identification of the active compounds responsible for the antimycobacterial activity observed with the CHCl₃ extract.

Conclusions

Herein, to the best of our knowledge, the activity of the CHCl₃ and EtOH seed extracts from *P. americana* against two intestinal parasites that cause diarrhea: *E. histolytica*, and *G. lamblia* has been demonstrated. In addition, based on our results, CHCl₃ seed extract may be a source for potential moieties (molecules) against *M. tuberculosis* drug-resistant species as well as NTM.

Further studies are required for the identification of the active compounds responsible for the antiprotozoal and antimycobacterial activity observed with the EtOH and CHCl₃ seed extracts from *P. americana*.

Abbreviations

ALT: Alanine aminotransferase; AST: Aspartate aminotransferase; AOA: Antioxidant activity; CHCl₃: Chloroformic; AR: Analytical reagent; DMSO: Dimethyl sulfoxide; EtOAc: Ethyl acetate; EtOH: Ethanol; MeOH: Methanol; MTT: 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazol; IMSSM: Instituto Mexicano del Seguro Social Mexico; *P. americana*: *Persea americana*; HDL: High density lipoprotein; LDL: Low density lipoprotein; R_f: Retention factor; TLC: Thin layer chromatography; EMB: Etambutol; INH: Isoniazid; STR: Streptomycin; RIF: Rifampicin; LC₅₀: Lethal concentration; LD₅₀: Lethal dose; MIC: Minimum inhibitory concentration; IC₅₀: Medium Inhibitory concentration; MABA: Microplate alamar blue assay; MDR: Multidrug-resistant; XDR: Extensively drug-resistant; *M. fortuitum*: *Mycobacterium fortuitum*; *M. avium*: *Mycobacterium avium*; *M. smegmatis*: *Mycobacterium smegmatis*; *M. abscessus*: *Mycobacterium abscessus*; *M. tuberculosis*: *Mycobacterium tuberculosis*; NTM: non-tuberculosis mycobacterium; *E. histolytica*: *Entamoeba histolytica*; *G. lamblia*: *Giardia lamblia*; *T. vaginalis*: *Trichomonas vaginalis*.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

AJ-A designed and coordinated the study, prepared the CHCl₃ and EtOH extracts and carried out their phytochemical analysis and wrote the

manuscript. C-G collected the *P. americana* seeds and contributed to the preparing the CHCl₃ and EtOH extracts. JL-H and RR-N evaluated the antimycobacterial activity from CHCl₃ and EtOH extracts. The antiprotozoal activity from both extracts was determined by AT and LY-M, who also contributed to the manuscript preparation. All authors have read and approved the final manuscript.

Acknowledgments

Part of this study was developed with the financial support of CONACyT-México project 48339 and FIS/IMSS/PROT/G12/1126.

Author details

¹Unidad de Investigación Médica en Farmacología (UIM), Hospital de Especialidades, CMNS XXI, Instituto Mexicano del Seguro Social (IMSS), Av. Cuauhtémoc 330, Col. Doctores, 06720, México, DF, Mexico. ²Departamento de Inmunología, Escuela Nacional de Ciencias Biológicas, Instituto Politécnico Nacional, Plan de Ayala s/n, Col. Casco de Santo Tomás, 11340, México, DF, Mexico. ³UIM en Enfermedades Infecciosas y Parasitarias, Hospital de Pediatría, 2º piso, CMNS XXI, IMSS, Av. Cuauhtémoc 330, Col. Doctores, 06720, México, DF, Mexico.

Received: 4 December 2012 Accepted: 8 May 2013

Published: 16 May 2013

References

1. Aguilar A, Aguilar A: *Herbario Medicinal del IMSS*. 1st edition. México: Editorial Redact, S.A; 1994.
2. Argueta A, Cano L, Rodarte M: *Atlas de las Plantas de la Medicina Tradicional Mexicana*, Vols. 2, 3. 1st edition. México, DF: Editorial Instituto Nacional Indigenista; 1994.
3. Moreno-Urbe V: *Herbolario y tradición en la región de Xico*. 1st edition. Mexico: Veracruz, Diseño Editorial; 2008.
4. Osuna-Torres L, Tapia-Pérez ME, Aguilar-Contreras A: *Plantas medicinales de la medicina tradicional mexicana para tratar afecciones gastrointestinales*. 1st edition. España: Editorial Universidad de Barcelona; 2005.
5. Del Refugio Ramos M, Jerz G, Villanueva S, López-Dellamary F, Waibel R, Winterhalter P: **Two glucosylated abscisic acid derivatives from avocado seeds (*Persea americana* Mill. Lauraceae cv. Hass).** *Phytochem* 2004, **65**(7):955–962.
6. Nwaogu LA, Alisi CS, Ojako OA: **Studies on the nutritional and phytochemical properties of *Persea americana* seed.** *Bio-Research* 2008, **6**(1):320–322.
7. Takenaga F, Matsuyama K, Abe S, Torii Y, Itoh S: **Lipid and fatty acid composition of mesocarp and seed of avocado fruits harvested at northern range in Japan.** *J Oleo Sci* 2008, **57**(11):591–597.
8. Wang W, Bostic TR, Gu L: **Antioxidant capacities, procyanidins and pigments in avocados of different strains and cultivars.** *Food Chem* 2010, **122**(4):1193–1198.
9. Rodríguez-Carpena JG, Morcuende D, Andrade MJ, Kylli P, Estévez M: **Avocado (*Persea americana* Mill.) phenolics, *in vitro* antioxidant and antimicrobial activities, and inhibition of lipid and protein oxidation in porcine patties.** *J Agric Food Chem* 2011, **59**(10):5625–5635.
10. Asaolu MF, Asaolu SS, Oyeyemi AO, Aluko BT: **Hypolipemic effects of methanolic extract of *Persea americana* seeds in hypercholesterolemic rats.** *J Med Medical Sciences* 2010, **1**(4):126–128.
11. Imafidon KE, Amaechina FC: **Effects of aqueous seed extract of *Persea americana* Mill. (avocado) on blood pressure and lipid profile in hypertensive rats.** *Adv Biol Res* 2010, **4**(2):116–121.
12. Ogochukwu NA, Ozolua RI, Okpo SO: **Effect of the aqueous seed extract of *Persea americana* Mill. (Lauraceae) on the blood pressure of Sprague Dawley rats.** *Afric J Pharm Pharmacol* 2009, **3**(10):485–490.
13. Edem D, Ekanem I, Ebong P: **Effect of aqueous extracts of alligator pear seed (*Persea americana* Mill.) on blood glucose and histopathology of pancreas in alloxan-induced diabetic rats.** *Pak J Pharm Sci* 2009, **22**(3):272–276.
14. Koffi N, Kouakou EA, Dodiomon S: **Effect of aqueous extract of *Persea americana* seeds on the glycemia of diabetic rabbits.** *Europ J Sci Res* 2009, **26**(3):376–385.
15. Okonta M, Okonta L, Aguwa CN: **Blood glucose lowering activities of seeds of *Persea americana* on alloxan induced diabetic rats.** *Nig J Nat Prod Med* 2007, **11**:26–28.
16. Ozolua RI, Anaka ON, Okpo SO, Idogun SE: **Acute and sub-acute toxicological assessment of the aqueous seed extract of *Persea americana* Mill (Lauraceae) in rats.** *Afr J Tradit Complement Altern Med* 2009, **6**(4):573–578.
17. Giffoni LJ, Salles EH, Aguiar R, Nogueira RS, Costa JJ, Medeiros SL, De Moraes S, Gadelha MF: **Chemical composition, toxicity and larvicity and antifungal activities of *Persea americana* (avocado) seed extracts.** *Rev Soc Bras Med Trop* 2009, **42**:110–113.
18. Gomez-Flores R, Arzate-Quintana C, Quintanilla-Licea R, Tamez-Guerra P, Tamez-Guerra R, Monreal-Cuevas E, Rodríguez-Padilla C: **Antimicrobial activity of *Persea americana* Mill (Lauraceae) (Avocado) and *Gymnosperma glutinosum* (Spreng.) Less (Asteraceae) leaf extracts and activities fractions against *Mycobacterium tuberculosis*.** *Amer-Eur J Scient Res* 2008, **3**(2):188–194.
19. Raymond Chia TW, Dykes GA: **Antimicrobial activity of crude epicarp and seed extracts from mature avocado fruit (*Persea americana*) of three cultivars.** *Pharm Biol* 2011, **48**(7):753–756.
20. Abe F, Nagafuji S, Okawa M, Kinjo J, Akahane H, Ogura T, Martínez-Alfaro MA, Reyes-Chilpa R: **Trypanocidal constituents in plants. 5. Evaluation of some Mexican plants for their trypanocidal activity and active constituents in the seed of *Persea americana*.** *Biol Pharm Bull* 2005, **28**(7):1314–1317.
21. Ponce-Macotela M, Navarro-Alegria I, Martínez-Gordillo MN, Álvarez-Chacón R: **Efecto antiangiardiaco *in vitro* de 14 extractos de plantas.** *Rev Invest Clin* 1994, **46**:343–3474.
22. Matsusaka Y, Kawabata J, Takanori K: **Antioxidative constituents in avocado (*Persea americana* Mill.) seeds.** *J Jap Soc Food Sci Technol* 2003, **50**(11):550–552.
23. Yean-Yean S, Barlow PJ: **Antioxidant activity and phenolic content of selected fruit seeds.** *Food Chem* 2004, **88**(3):411–417.
24. Asaolu MF, Asaolu SS, Fakunle JB, Emman-Okon BO, Ajayi EO, Togun RA: **Evaluation of *in vitro* antioxidant activities of methanol extracts of *Persea americana* and *Cnidiosculus aconitifolius*.** *Pak J Nutr* 2010, **9**:1074–1077.
25. Cedillo-Rivera R, Chávez B, González-Robles A, Tapia A, Yépez-Mulia L: ***In vitro* effect of nitazoxanide against *Entamoeba histolytica*, *Giardia intestinalis* and *Trichomonas vaginalis* trophozoites.** *J Eukaryot Microbiol* 2002, **49**(3):201–208.
26. Jiménez-Arellanes A, León-Díaz R, Meckes M, Tapia A, Molina-Salinas GM, Luna-Herrera J, Yépez-Mulia L: **Antiprotozoal and antimycobacterial activities of pure compounds from *Aristolochia elegans* rhizomes.** *Evid Based Complement Alternat Med* 2012, **201**(2):593403. doi:10.1155/2012/593403.
27. León-Díaz R, Meckes M, Said-Fernández S, Molina-Salinas GM, Vargas-Villareal J, Torres J, Luna-Herrera J, Jiménez-Arellanes A: **Antimycobacterial neolignans isolated from *Aristolochia taliscana*.** *Mem Inst Oswaldo Cruz* 2010, **105**(1):45–51.
28. Molina-Salinas GM, Ramos-Guerra MC, Vargas-Villareal J, Mata-Cárdenas BD, Becerril-Montes P, Said-Fernández S: **Bactericidal activity of organic extracts from *Fluorensia cernua* DC against strains of *Mycobacterium tuberculosis*.** *Arch Med Res* 2006, **37**(1):45–49.
29. Alanis AD, Calzada F, Cedillo-Rivera R, Meckes M: **Antiprotozoal activity of the constituents of *Rubus coriifolius*.** *Phytother Res* 2003, **17**(6):681–682.
30. Barbosa E, Calzada F, Campos R: ***In vivo* antiangiardiaco activity of three flavonoids isolated of some medicinal plants used in Mexican traditional medicine for the treatment of diarrhea.** *J Ethnopharmacol* 2007, **109**(3):552–554.
31. Baquero SE, Benavides J, Sepúlveda L, Quiñones W, Torres F, Cardona D, Archbold R, Guzmán JD, Cuca LE, Franzblau S, Echeverri F: **Actividad antimicobacteriana de algunas plantas de la flora colombiana.** *Sci Techn* 2007, **XIII**:133–136.
32. Cassidy PM, Hedberg K, Saulson A, McNelly E, Winthrop KL: **Nontuberculous mycobacterial disease prevalence and risk factors: a changing epidemiology.** *Clin Infect Dis* 2009, **49**(12):e124–e129.
33. Pauli GF, Case RJ, Inui T, Wang Y, Cho S, Fischer NH, Franzblau SG: **New perspectives on natural products in TB drug research.** *Life Sci* 2005, **78**(5):485–494.

doi:10.1186/1472-6882-13-109

Cite this article as: Jiménez-Arellanes et al.: Antiprotozoal and antimycobacterial activities of *Persea americana* seeds. *BMC Complementary and Alternative Medicine* 2013 **13**:109.