

Hindawi Publishing Corporation
 Advances in Difference Equations
 Volume 2008, Article ID 143723, 6 pages
 doi:10.1155/2008/143723

Research Article

The Periodic Character of the Difference Equation

$$x_{n+1} = f(x_{n-l+1}, x_{n-2k+1})$$

Taixiang Sun¹ and Hongjian Xi²

¹Department of Mathematics, College of Mathematics and Information Science, Guangxi University, Nanning 530004, Guangxi, China

²Department of Mathematics, Guangxi College of Finance and Economics, Nanning 530003, Guangxi, China

Correspondence should be addressed to Taixiang Sun, stx1963@163.com

Received 3 February 2007; Revised 18 September 2007; Accepted 27 November 2007

Recommended by H. Bevan Thompson

In this paper, we consider the nonlinear difference equation $x_{n+1} = f(x_{n-l+1}, x_{n-2k+1})$, $n = 0, 1, \dots$, where $k, l \in \{1, 2, \dots\}$ with $2k \neq l$ and $\gcd(2k, l) = 1$ and the initial values $x_{-\alpha}, x_{-\alpha+1}, \dots, x_0 \in (0, +\infty)$ with $\alpha = \max\{l-1, 2k-1\}$. We give sufficient conditions under which every positive solution of this equation converges to a (not necessarily prime) 2-periodic solution, which extends and includes corresponding results obtained in the recent literature.

Copyright © 2008 T. Sun and H. Xi. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

In this paper, we consider a nonlinear difference equation and deal with the question of whether every positive solution of this equation converges to a periodic solution. Recently, there has been a lot of interest in studying the global attractivity, the boundedness character, and the periodic nature of nonlinear difference equations (e.g., see [1, 2]). In [3], Grove et al. considered the following difference equation:

$$x_{n+1} = \frac{p + x_{n-(2m+1)}}{1 + x_{n-2r}}, \quad n = 0, 1, \dots, \quad (E1)$$

where $p \in (0, +\infty)$ and the initial values $x_{-\alpha}, x_{-\alpha+1}, \dots, x_0 \in (0, +\infty)$ with $\alpha = \max\{2r, 2m+1\}$, and proved that every positive solution of (E1) converges to (not necessarily prime) a 2s-periodic solution with $s = \gcd(m+1, 2r+1)$. In [4], Stević investigated the periodic character of positive solutions of the following difference equation:

$$x_{n+1} = 1 + \frac{x_{n-2s+1}}{x_{n-(2r+1)s+1}}, \quad n = 0, 1, \dots, \quad (E2)$$

and proved that every positive solution of (E2) converges to (not necessarily prime) a 2s-periodic solution, which generalized the main result of [5]. Furthermore, Stević [6] studied the periodic character of positive solutions of the following difference equation:

$$x_n = 1 + \frac{\sum_{i=1}^k \alpha_i x_{n-p_i}}{\sum_{j=1}^m \beta_j x_{n-q_j}}, \quad n = 1, 2, \dots, \quad (\text{E3})$$

where $\alpha_i, i \in \{1, \dots, k\}$, and $\beta_j, j \in \{1, \dots, m\}$, are positive numbers such that $\sum_{i=1}^k \alpha_i = \sum_{j=1}^m \beta_j = 1$, and $p_i, i \in \{1, \dots, k\}$, and $q_j, j \in \{1, \dots, m\}$, are natural numbers such that $p_1 < p_2 < \dots < p_k$ and $q_1 < q_2 < \dots < q_m$. For closely related results, see [7, 8].

In this paper, we consider the more general equation

$$x_{n+1} = f(x_{n-l+1}, x_{n-2k+1}), \quad n = 0, 1, 2, \dots, \quad (\text{1.1})$$

where $k, l \in \{1, 2, \dots\}$ with $2k \neq l$ and $\gcd(2k, l) = 1$, the initial values $x_{-\alpha}, x_{-\alpha+1}, \dots, x_0 \in (0, +\infty)$ with $\alpha = \max\{l-1, 2k-1\}$, and f satisfies the following hypotheses:

(H₁) $f \in C(E \times E, (0, +\infty))$ with $a = \inf_{(u,v) \in E \times E} f(u, v) \in E$, where $E \in \{(0, +\infty), [0, +\infty)\}$;

(H₂) $f(u, v)$ is decreasing in u and increasing in v ;

(H₃) there exists a decreasing function $g \in C((a, +\infty), (a, +\infty))$ such that

(i) for any $x > a$, $g(g(x)) = x$ and $x = f(g(x), x)$;

(ii) $\lim_{x \rightarrow a^+} g(x) = +\infty$ and $\lim_{x \rightarrow +\infty} g(x) = a$.

The main result of this paper is the following theorem.

Theorem 1.1. *Every positive solution of (1.1) converges to (not necessarily prime) a 2-periodic solution.*

2. Proof of Theorem 1.1

In this section, we will prove Theorem 1.1. Without loss of generality, we may assume $l < 2k$ (the proof for the case $l > 2k$ is similar); then

$$\{l, 2l, 3l, \dots, 2kl\} = \{0, 1, 2, \dots, 2k-1\} \pmod{2k}. \quad (\text{2.1})$$

Lemma 2.1. *Let $\{x_n\}_{n=-\alpha}^{\infty}$ be a positive solution of (1.1). Then there exists a real number $L \in (a, +\infty)$ such that $L \leq x_n \leq g(L)$ for all $n \geq 1$. Furthermore, let $\limsup x_n = M$ and $\liminf x_n = m$, then $M = g(m)$ and $m = g(M)$.*

Proof. By (H₁) and (H₂), we have

$$x_i = f(x_{i-l}, x_{i-2k}) > f(x_{i-l} + 1, x_{i-2k}) \geq a \quad \text{for every } 1 \leq i \leq \alpha + 1. \quad (\text{2.2})$$

Then there exists $L \in (a, +\infty)$ with $L < g(L)$ such that

$$L \leq x_i \leq g(L) \quad \text{for every } 1 \leq i \leq \alpha + 1. \quad (\text{2.3})$$

It follows from (2.3) and (H₃) that

$$g(L) = f(L, g(L)) \geq x_{\alpha+2} = f(x_{\alpha+2-l}, x_{\alpha+2-2k}) \geq f(g(L), L) = L. \quad (2.4)$$

Inductively, it follows that $L \leq x_n \leq g(L)$ for all $n \geq 1$.

Let $\limsup x_n = M$ and $\liminf x_n = m$, then there exist $A, B, C, D \in [m, M]$ and sequences $t_n \geq 1$ and $r_n \geq 1$ such that

$$\begin{aligned} \lim_{n \rightarrow \infty} x_{t_n} &= M, & \lim_{n \rightarrow \infty} x_{t_n-l} &= A, & \lim_{n \rightarrow \infty} x_{t_n-2k} &= B, \\ \lim_{n \rightarrow \infty} x_{r_n} &= m, & \lim_{n \rightarrow \infty} x_{r_n-l} &= C, & \lim_{n \rightarrow \infty} x_{r_n-2k} &= D. \end{aligned} \quad (2.5)$$

Thus by (1.1), (H₂), and (H₃), we have

$$\begin{aligned} f(g(M), M) &= M = f(A, B) \leq f(m, M), \\ f(g(m), m) &= m = f(C, D) \geq f(M, m), \end{aligned} \quad (2.6)$$

from which it follows that $g(M) \geq m$ and $g(m) \leq M$. Since g is decreasing, it follows that

$$m = g(g(m)) \geq g(M), \quad M = g(g(M)) \leq g(m). \quad (2.7)$$

Therefore, $M = g(m)$ and $m = g(M)$. The proof is complete. \square

Proof of Theorem 1.1. Let $\{x_n\}_{n=-\alpha}^{\infty}$ be a positive solution of (1.1) with the initial conditions $x_0, x_{-1}, \dots, x_{-\alpha} \in (0, +\infty)$. It follows from Lemma 2.1 that

$$a < \liminf x_n = m = g(M) \leq \limsup x_n = M < +\infty. \quad (2.8)$$

Obviously, every sequence

$$L, g(L), L, g(L), \dots \quad (2.9)$$

is a 2-periodic (not necessarily prime) solution of (1.1), where $L \in \{M, m\}$.

By taking a subsequence, we may assume that there exists a sequence $t_n \geq 2kl + 1$ such that

$$\begin{aligned} \lim_{n \rightarrow \infty} x_{t_n} &= M, \\ \lim_{n \rightarrow \infty} x_{t_n-j} &= A_j \in [g(M), M] \quad \text{for } j \in \{1, 2, \dots, 2kl\}. \end{aligned} \quad (2.10)$$

According to (1.1), (2.10), and (H₃), we obtain

$$f(g(M), M) = M = f(A_l, A_{2k}) \leq f(g(M), M), \quad (2.11)$$

from which it follows that

$$A_l = g(M), \quad A_{2k} = M. \quad (2.12)$$

In a similar fashion, we can obtain

$$\begin{aligned} f(g(M), M) &= M = A_{2k} = f(A_{2k+l}, A_{4k}) \leq f(g(M), M), \\ f(M, g(M)) &= g(M) = A_l = f(A_{2l}, A_{l+2k}) \geq f(M, g(M)), \end{aligned} \quad (2.13)$$

from which it follows that

$$A_{4k} = A_{2k} = A_{2l} = M, \quad A_{2k+l} = A_l = g(M). \quad (2.14)$$

Inductively, we have

$$\begin{aligned} A_{j2k} &= M \quad \text{for } j \in \{1, 2, \dots, l\}, \\ A_{jl} &= g(M) \quad \text{for } j \in \{1, 3, \dots, 2k-1\}, \\ A_{jl} &= M \quad \text{for } j \in \{0, 2, \dots, 2k\}, \\ A_{j+l2k} &= A_{jl} \quad \text{for } j \in \{0, 1, \dots, 2k\}, \quad r \in \{0, 1, \dots, l\}, \quad jl + r2k \leq 2kl. \end{aligned} \quad (2.15)$$

For every $r \in \{0, 1, 2, 3, \dots, 2k-1\}$, there exist $j_r \in \{0, 1, 2, 3, \dots, 2k-1\}$ and $p_r \in \{0, 1, \dots, l-1\}$ such that $j_r l = 2kp_r + r$, from which, with (2.15), it follows that

$$A_{2k(l-1)+r} = A_{j_r l} = \begin{cases} M & \text{for } r \in \{0, 2, 4, \dots, 2k-2\}, \\ g(M) & \text{for } r \in \{1, 3, \dots, 2k-1\}, \end{cases} \quad (2.16)$$

$$\begin{aligned} \lim_{n \rightarrow \infty} x_{t_n - 2k(l-1)-j} &= M \quad \text{for } j \in \{0, 2, \dots, 2k\}, \\ \lim_{n \rightarrow \infty} x_{t_n - 2k(l-1)-j} &= g(M) \quad \text{for } j \in \{1, 3, \dots, 2k-1\}. \end{aligned} \quad (2.17)$$

In view of (2.17), for any $0 < \varepsilon < M - a$, there exists some $t_\beta \geq 4kl$ such that

$$\begin{aligned} M - \varepsilon &< x_{t_\beta - 2k(l-1)-j} < M + \varepsilon \quad \text{if } j \in \{0, 2, \dots, 2k\}, \\ g(M + \varepsilon) &< x_{t_\beta - 2k(l-1)-j} < g(M - \varepsilon) \quad \text{if } j \in \{1, 3, \dots, 2k-1\}. \end{aligned} \quad (2.18)$$

By (1.1) and (2.18), we have

$$x_{t_\beta - 2k(l-1)+1} = f(x_{t_\beta - 2k(l-1)-l+1}, x_{t_\beta - 2kl+1}) < f(M - \varepsilon, g(M - \varepsilon)) = g(M - \varepsilon). \quad (2.19)$$

Also (1.1), (2.18), and (2.19) imply that

$$x_{t_\beta - 2k(l-1)+2} = f(x_{t_\beta - 2k(l-1)-l+2}, x_{t_\beta - 2kl+2}) > f(g(M - \varepsilon), M - \varepsilon) = M - \varepsilon. \quad (2.20)$$

Inductively, it follows that

$$\begin{aligned} x_{t_\beta - 2k(l-1)+2n} &> M - \varepsilon \quad \forall n \geq 0, \\ x_{t_\beta - 2k(l-1)+2n+1} &< g(M - \varepsilon) \quad \forall n \geq 0. \end{aligned} \quad (2.21)$$

Therefore,

$$\lim_{n \rightarrow \infty} x_{2n} = M, \quad \lim_{n \rightarrow \infty} x_{2n+1} = g(M) \quad (2.22)$$

or

$$\lim_{n \rightarrow \infty} x_{2n} = g(M), \quad \lim_{n \rightarrow \infty} x_{2n+1} = M. \quad (2.23)$$

The proof is complete. \square

Remark 2.2. (1) The proofs of Lemma 2.1 and Theorem 1.1 draw on ideas from the proofs of Theorems 2.1 and 2.2 in [6].

(2) Consider the nonlinear difference equation

$$x_{n+1} = f(x_{n-ls+1}, x_{n-2ks+1}), \quad n = 0, 1, \dots, \quad (2.24)$$

where $s, k, l \in \{1, 2, \dots\}$ with $2k \neq l$ and $\gcd(2k, l) = 1$, the initial values $x_{-\alpha}, x_{-\alpha+1}, \dots, x_0 \in (0, +\infty)$ with $\alpha = \max\{ls - 1, 2ks - 1\}$, and f satisfies (H₁)–(H₃). Let $y_{n+1}^i = x_{ns+i+1}$ for every $0 \leq i \leq s - 1$ and $n = 0, 1, 2, \dots$, then (2.24) reduces to the equation

$$y_{n+1}^i = f(y_{n-l+1}^i, y_{n-2k+1}^i), \quad 0 \leq i \leq s - 1, \quad n = 0, 1, 2, \dots \quad (2.25)$$

It follows from Theorem 1.1 that for any $0 \leq i \leq s - 1$, every positive solution of the equation $y_{n+1}^i = f(y_{n-l+1}^i, y_{n-2k+1}^i)$ converges to (not necessarily prime) a 2-periodic solution. Thus every positive solution of (2.24) converges to (not necessarily prime) a $2s$ -periodic solution.

3. Examples

To illustrate the applicability of Theorem 1.1, we present the following examples.

Example 3.1. Consider the equation

$$x_{n+1} = \frac{p + \sum_{i=1}^{m+1} x_{n-2k+1}^i}{\sum_{i=0}^m x_{n-2k+1}^i + x_{n-l+1}}, \quad n = 0, 1, \dots, \quad (3.1)$$

where $m, k, l \in \{1, 2, \dots\}$ with $2k \neq l$ and $\gcd(2k, l) = 1$ and the initial values $x_{-\alpha}, x_{-\alpha+1}, \dots, x_0 \in (0, +\infty)$ with $\alpha = \max\{l - 1, 2k - 1\}$, $0 < p \leq 1$. Let $E = [0, +\infty)$ and

$$f(x, y) = \frac{p + \sum_{i=1}^{m+1} y^i}{\sum_{i=0}^m y^i + x} \quad (x \geq 0, y \geq 0), \quad g(x) = \frac{p}{x} \quad (x > 0). \quad (3.2)$$

It is easy to verify that (H₁)–(H₃) hold for (3.1). It follows from Theorem 1.1 that every solution of (3.1) converges to (not necessarily prime) a 2-periodic solution.

Example 3.2. Consider the equation

$$x_{n+1} = 1 + \frac{x_{n-2k+1}^{m+1}}{\sum_{i=1}^m x_{n-2k+1}^i + x_{n-l+1}}, \quad n = 0, 1, \dots, \quad (3.3)$$

where $m, k, l \in \{1, 2, \dots\}$ with $2k \neq l$ and $\gcd(2k, l) = 1$ and the initial values $x_{-\alpha}, x_{-\alpha+1}, \dots, x_0 \in (0, +\infty)$ with $\alpha = \max\{l-1, 2k-1\}$. Let $E = (0, +\infty)$ and

$$f(x, y) = 1 + \frac{y^{m+1}}{\sum_{i=1}^m y^i + x} \quad (x > 0, y > 0), \quad g(x) = \frac{x}{x-1} \quad (x > 1). \quad (3.4)$$

It is easy to verify that (H₁)–(H₃) hold for (3.3). It follows from Theorem 1.1 that every solution of (3.3) converges to (not necessarily prime) a 2-periodic solution.

Acknowledgments

The authors would like to thank the referees for some valuable and constructive comments and suggestions. The project is supported by NNSF of China (10461001) and NSF of Guangxi (0640205, 0728002).

References

- [1] S. Stevo, "The recursive sequence $x_{n+1} = g(x_n, x_{n-1})/(A+x_n)$," *Applied Mathematics Letters*, vol. 15, no. 3, pp. 305–308, 2002.
- [2] R. DeVault, C. Kent, and W. Kosmala, "On the recursive sequence $x_{n+1} = p + x_{n-k}/x_n$," *Journal of Difference Equations and Applications*, vol. 9, no. 8, pp. 721–730, 2003.
- [3] E. A. Grove, G. Ladas, and M. Predescu, "On the periodic character of the difference equation $x_{n+1} = (px_{n-2l} + x_{n-(2k+1)})/(1 + x_{n-2l})$," *Mathematical Sciences Research Journal*, vol. 6, no. 5, pp. 221–233, 2002.
- [4] S. Stević, "A note on periodic character of a difference equation," *Journal of Difference Equations and Applications*, vol. 10, no. 10, pp. 929–932, 2004.
- [5] W. T. Patula and H. D. Voulov, "On the oscillation and periodic character of a third order rational difference equation," *Proceedings of the American Mathematical Society*, vol. 131, no. 3, pp. 905–909, 2003.
- [6] S. Stević, "On the recursive sequence $x_n = 1 + \sum_{i=1}^k \alpha_i x_{n-p_i} / \sum_{j=1}^m \beta_j x_{n-q_j}$," *Discrete Dynamics in Nature and Society*, vol. 2007, Article ID 39404, 7 pages, 2007.
- [7] T. Sun and H. Xi, "The periodic character of positive solutions of the difference equation $x_{n+1} = f(x_n, x_{n-k})$," *Computers & Mathematics with Applications*, vol. 51, no. 9-10, pp. 1431–1436, 2006.
- [8] K. S. Berenhaut, J. D. Foley, and S. Stević, "The global attractivity of the rational difference equation $y_n = 1 + y_{n-k}/y_{n-m}$," *Proceedings of the American Mathematical Society*, vol. 135, no. 4, pp. 1133–1140, 2007.