

Single meson contributions to the muon's anomalous magnetic moment

Vladyslav Pauk^{1,2,3,a}, Marc Vanderhaeghen^{1,2}¹ Institut für Kernphysik, Johannes Gutenberg-Universität, Mainz, Germany² PRISMA Cluster of Excellence, Johannes Gutenberg-Universität, Mainz, Germany³ Department of Physics, Taras Shevchenko National University of Kyiv, Kyiv, UkraineReceived: 12 May 2014 / Accepted: 25 July 2014 / Published online: 9 August 2014
© The Author(s) 2014. This article is published with open access at Springerlink.com

Abstract We develop the formalism to provide an improved estimate for the hadronic light-by-light correction to the muon's anomalous magnetic moment a_μ , by considering single meson contributions beyond the leading pseudoscalar mesons. We incorporate available experimental input as well as constraints from light-by-light scattering sum rules to estimate the effects of axial-vector, scalar, and tensor mesons. We give numerical evaluations for the hadronic light-by-light contribution of these states to a_μ . The presented formalism allows one to further improve on these estimates, once new data for such meson states will become available.

1 Introduction

The anomalous magnetic moment of the muon $a_\mu = (g-2)/2$ is one of the most precisely measured quantities in particle physics. It has been playing a vital role in testing the framework of quantum field theory since its development more than half a century ago, as well as in searching for new physics beyond the Standard Model (SM) of particles and interactions, for a comprehensive review, see [1,2] and references therein. On the experimental side, the present world average for a_μ is [3,4]

$$a_\mu(\text{exp}) = (116,592,089 \pm 63) \times 10^{-11}, \quad (1)$$

which corresponds to a relative precision of 0.54 parts per million. From the theoretical point of view, in the SM a_μ is defined by electromagnetic (QED), electroweak, and hadronic contributions. The dominant QED contribution, which at present has been calculated including all terms up to fifth-order in the fine structure constant [5,6], is known

to an impressive theoretical precision corresponding with $\delta a_\mu(\text{QED}) = 8 \times 10^{-13}$. The much smaller electroweak contribution, which has been calculated up to two-loop order [7–10], is also known with good accuracy; its inaccuracy $\delta a_\mu(\text{weak}) = 2 \times 10^{-11}$, which is more than a factor of 30 smaller than the present experimental precision. Within the SM, the largest source of uncertainty is given by the hadronic contribution, which contains two parts, the hadronic vacuum polarization (HVP) together with the hadronic light-by-light scattering (HLbL); see Fig. 1. The HVP has been estimated based on data for $e^+e^- \rightarrow \text{hadrons}$, data for $e^+e^- \rightarrow \gamma + \text{hadrons}$, as well as τ decays, by several groups [11–21]. A recent evaluation of the leading order HVP has found [21]:

$$a_\mu(\text{l.o. HVP}) = (6,886.0 \pm 42.4) \times 10^{-11}. \quad (2)$$

The next-to-leading order HVP has been estimated as [19]

$$a_\mu(\text{n.l.o. HVP}) = (-98.4 \pm 0.7) \times 10^{-11}. \quad (3)$$

The HLbL, although much smaller in size than the HVP, has a comparable theoretical uncertainty. It has been estimated by different groups to be

$$a_\mu(\text{HLbL}) = (116 \pm 39) \times 10^{-11} \quad \text{Ref. [1, 2]}, \quad (4)$$

$$a_\mu(\text{HLbL}) = (105 \pm 26) \times 10^{-11} \quad \text{Ref. [22]}. \quad (5)$$

When comparing theory with experiment for a_μ , the difference has recently been evaluated as [21]

$$a_\mu(\text{exp}) - a_\mu(\text{theory}) = (312.5 \pm 57.6(\text{theory}) \pm 63(\text{exp})) \times 10^{-11}, \quad (6)$$

which corresponds with a 3.7σ discrepancy. The different analyses for the l.o. HVP and HLbL contributions give results which all agree within 1σ [23].

^a e-mail: pauk@uni-mainz.de




Fig. 1 The hadronic contributions to the anomalous magnetic moment of the muon. *Left panel* hadronic vacuum polarization (HVP). *Right panel* hadronic light-by-light contribution (HLbL). The *gray blobs* denote hadronic intermediate states

In order to conclude whether this discrepancy is a sign of new physics beyond the standard model, new experiments are planned in the near future both at Fermilab [24] as well as at J-PARC [25] to further improve on the precision. The Fermilab experiment aims to reduce the experimental uncertainty by a factor 4 to $\delta a_\mu \approx 16 \times 10^{-11}$. Such improvement also calls to improve on the theoretical accuracy by at least a factor of 2 in order to obtain a definitive test for the presently observed discrepancy. As the theoretical uncertainty is totally dominated by the knowledge of the HVP, Eq. (2), and the HLbL, Eq. (4), the main effort on the theoretical side will be to improve on both estimates. For the HVP, new data from ongoing experiments at Novosibirsk and BES-III will provide valuable experimental input to further constrain this contribution. It was estimated in Ref. [23] that such data will allow us to reduce the uncertainty in the HVP by around a factor of two, down to $\delta a_\mu(\text{i.o. HVP}) = 26 \times 10^{-11}$. For the HLbL scattering, new data are expected from KLOE-2 for the $\gamma^* \gamma \rightarrow \pi^0$ transition form factor at very low photon virtualities, and from BES-III for the reactions $\gamma^* \gamma \rightarrow X$, where $X = \pi^0, \eta, \eta', 2\pi$. Such data do require a theoretical analysis in order to further constrain the HLbL evaluation.

The aim of the present work is to provide an improved estimate for the HLbL contribution, by considering single meson contributions beyond the leading pseudoscalar mesons (π^0, η, η'), which have been evaluated in the pioneering work of Ref. [26]. We will incorporate available experimental input as well as constraints from light-by-light scattering sum rules [27, 28] to estimate the effects of axial-vector, scalar, and tensor mesons to the HLbL contribution. The framework which will be presented will also allow one to further improve on the estimate, once new data, in particular from BES-III, for such meson states will become available.

2 Meson pole contributions to the hadronic light-by-light scattering

The HLbL contribution to the muon’s anomalous magnetic moment is a $\mathcal{O}(\alpha^3)$ correction to the Pauli form factor due

to the second diagram in Fig. 1. It may be isolated from the vertex matrix element

$$\langle \mu^-(p') | j_{em}^\mu(0) | \mu^-(p) \rangle = (-ie) \bar{u}(p') \Gamma^\mu(p', p) u(p), \quad (7)$$

where p (p') denote the initial (final) muon momenta, when considering the classical limit ($p' - p \equiv k \rightarrow 0$) by using a projection operator technique [29]. This amounts to the two-loop integral representation

$$a_\mu^{LbL} = \lim_{k \rightarrow 0} i e^6 \int \frac{d^4 q_1}{(2\pi)^4} \int \frac{d^4 q_2}{(2\pi)^4} \frac{1}{q_1^2 q_2^2 (k - q_1 - q_2)^2} \times \frac{1}{(p + q_1)^2 - m^2} \frac{1}{(p' - q_2)^2 - m^2} \times T^{\mu\nu\lambda\sigma}(q_1, k - q_1 - q_2, q_2) \Pi_{\mu\nu\lambda\sigma}(q_1, k - q_1 - q_2, q_2), \quad (8)$$

with the projector

$$\Lambda_\mu(p', p) = \frac{m^2}{k^2(4m^2 - k^2)} \left[\gamma_\mu + \frac{k^2 + 2m^2}{m(k^2 - 4m^2)} (p' + p)_\mu \right], \quad (9)$$

where m denotes the muon mass. Furthermore, in Eq. (8) the leptonic tensor T is given by

$$T^{\mu\nu\lambda\sigma}(q_1, k - q_1 - q_2, q_2) = \text{Tr} \left[(\not{p} + m) \Lambda^\sigma(p', p) (\not{p}' + m) \gamma^\lambda (\not{p}' - \not{q}_2 + m) \times \gamma^\nu (\not{p} + \not{q}_1 + m) \gamma^\mu \right]. \quad (10)$$

The tensor $\Pi_{\mu\nu\lambda\sigma}(q_1, q_2, q_3)$ denotes the light-by-light vacuum polarization tensor. In this work, we will consider the contributions of a single meson with an arbitrary spin to Π , which have the general form:

$$(ie)^4 \Pi_{\mu\nu\lambda\sigma}(q_1, k - q_1 - q_2, q_2) = \mathcal{M}_{\mu\nu, \{\alpha\}}(q_1, k - q_1 - q_2) \frac{i P^{\{\alpha\}, \{\beta\}}(k - q_2)}{(k - q_2)^2 - M^2} \times \mathcal{M}_{\lambda\sigma, \{\beta\}}(q_2, -k) + \mathcal{M}_{\mu\sigma, \{\alpha\}}(q_1, -k) \frac{i P^{\{\alpha\}, \{\beta\}}(k - q_1)}{(k - q_1)^2 - M^2} \times \mathcal{M}_{\nu\lambda, \{\beta\}}(k - q_1 - q_2, q_2) + \mathcal{M}_{\mu\lambda, \{\alpha\}}(q_1, q_2) \frac{i P^{\{\alpha\}, \{\beta\}}(q_1 + q_2)}{(q_1 + q_2)^2 - M^2} \times \mathcal{M}_{\nu\sigma, \{\beta\}}(k - q_1 - q_2, -k), \quad (11)$$

where the three terms correspond with the three topologies shown in Fig. 2. In Eq. (11), the Lorentz amplitude $\mathcal{M}_{\mu\nu, \{\alpha\}}(q_1, q_2)$ describes the transition from the initial state of two virtual photons with momenta q_1 and q_2 to a C -even (J^{PC}) meson with mass M . Depending on the spin J of the meson, the amplitude $\mathcal{M}_{\mu\nu, \{\alpha\}}$ has different Lorentz structures: for the case of a pseudoscalar ($J^{PC} = 0^{-+}$) and a scalar ($J^{PC} = 0^{++}$) meson the amplitude is a rank two tensor, for the case of an axial-vector ($J^{PC} = 1^{+-}$) and a tensor




Fig. 2 The single meson contributions to the hadronic light-by-light scattering

($J^{PC} = 2^{++}$) meson it is a rank three tensor and a rank four tensor, respectively. The projector P for spin J entering the meson propagator is defined by the spin sum

$$P^{\{\alpha\},\{\beta\}}(p) = \sum_{\sigma=-J}^J \varepsilon_{\sigma}^{\{\alpha\}}(p)\varepsilon_{\sigma}^{\{\beta\}*}(p), \tag{12}$$

where the $\varepsilon_{\sigma}^{\{\alpha\}}$ denote the corresponding polarization tensors. In the following, we give the specific expressions which we use for the transition amplitudes and the polarization sums in our calculation. The transition amplitudes are defined in such a way that the non-perturbative physics is contained in the meson transition Form Factors (FFs). It is important to note that these FFs depend on three invariants in the general case [1,2]. However, mainly due to the absence of reliable information about the off-shell dependence on the virtuality of the exchanged meson we will assume, for the following estimates, the pole-dominance approximation for the FFs:

$$F(q_1^2, q_2^2, (q_1+q_2)^2) = F(q_1^2, q_2^2, M^2) \equiv F_{\mathcal{M}\gamma^*\gamma^*}(q_1^2, q_2^2), \tag{13}$$

where q_1^2, q_2^2 denote the two photon virtualities, and $(q_1 + q_2)^2$ denotes the meson virtuality.

For a *pseudoscalar meson* (\mathcal{P}), the transition amplitude is defined by

$$\mathcal{M}_{\mu\nu}^{(\mathcal{P})}(q_1, q_2) = -ie^2\varepsilon_{\mu\nu\alpha\beta}q_1^\alpha q_2^\beta F_{\mathcal{P}\gamma^*\gamma^*}(q_1^2, q_2^2). \tag{14}$$

The spin projection operator for $J = 0$ has a trivial form:

$$P(q) = 1. \tag{15}$$

A *scalar meson* (\mathcal{S}) may be produced either by two transverse or by two longitudinal photons [30,31]. As the main contribution to the magnetic moment comes from the region of small photon virtualities, the contribution of the transverse amplitude is dominating. Furthermore, there is no empirical information on the structure of the longitudinal FFs at present. Thus in this work, we will only consider the transverse part of the scalar meson production amplitude which is described by

$$\mathcal{M}_{\mu\nu}^{(\mathcal{S})}(q_1, q_2) = -e^2\frac{(q_1\cdot q_2)}{M} R_{\mu\nu}(q_1, q_2) F_{\mathcal{S}\gamma^*\gamma^*}(q_1^2, q_2^2), \tag{16}$$

where the symmetric transverse tensor $R^{\mu\nu}$ is defined by

$$R^{\mu\nu}(q_1, q_2) \equiv -g^{\mu\nu} + \frac{1}{X} \{ (q_1 \cdot q_2) (q_1^\mu q_2^\nu + q_2^\mu q_1^\nu) - q_1^2 q_2^\mu q_2^\nu - q_2^2 q_1^\mu q_1^\nu \}, \tag{17}$$

and $X \equiv (q_1 \cdot q_2)^2 - q_1^2 q_2^2$.

Although the production of an *axial-vector meson* (\mathcal{A}) by two real photons is forbidden by the Landau–Yang theorem [32,33], an axial-vector meson can be produced in two-photon processes when one or both photons are virtual. Existing phenomenological analyses have used the expression for the transition amplitude derived from a non-relativistic quark model calculation [34–37]:

$$\begin{aligned} \mathcal{M}_{\mu\nu\alpha}^{(\mathcal{A})}(q_1, q_2) &= ie^2\varepsilon_{\rho\nu\tau\alpha} \left[(q_1^2 g_\mu^\rho - q_1^\rho q_{1\mu}) q_2^\tau - (q_2^2 g_\mu^\rho - q_2^\rho q_{2\mu}) q_1^\tau \right] \\ &\times \frac{1}{M^2} F_{\mathcal{A}\gamma^*\gamma^*}(q_1^2, q_2^2). \end{aligned} \tag{18}$$

Note that a general discussion of the $\mathcal{A}\gamma^*\gamma^*$ vertex, has to allow for three independent Lorentz structures [30,31]. However, as no phenomenological information is available at present to disentangle the three helicity structures, we will use the simplified vertex of Eq. (18) in the present work. According to Eq. (12) the projection operator for spin $J = 1$ is

$$P^{\alpha\beta}(p) = g^{\alpha\beta} - \frac{p^\alpha p^\beta}{M^2}. \tag{19}$$

In this work, we also consider the two-photon production of *tensor mesons* (\mathcal{T}). For the light quark tensor mesons, the experimental analyses of decay angular distributions for $\gamma\gamma$ cross sections to $\pi^+\pi^-$, $\pi^0\pi^0$, $\eta\pi^0$, and K^+K^- channels have shown [38] that the $J = 2$ mesons are produced predominantly (around 95 % or more) in a state of helicity $\Lambda = 2$. In addition, it has been shown in Ref. [39] that when saturating sum rules for $\gamma\gamma$ scattering by lowest lying mesons, the tensor meson decay coupling into two photons with total helicity $\Lambda = 2$ is much larger than the decay into two photons with total helicity $\Lambda = 0$. We will therefore assume in all of the following analyses that the hadronic light-by-light amplitude for tensor states is dominated by the helicity $\Lambda = 2$ exchange. Therefore, the two-photon decay rate is $\Gamma_{\gamma\gamma}(\mathcal{T}(\Lambda = 2)) \approx \Gamma_{\gamma\gamma}(\mathcal{T})$, and we will safely neglect the contribution of the remaining four helicity amplitudes. The relevant part of the Lorentz amplitude for the dominant process $\gamma^*\gamma^* \rightarrow \mathcal{T}(\Lambda = 2)$ can be parameterized as [28]

$$\begin{aligned} \mathcal{M}_{\mu\nu\alpha\beta}^{(\mathcal{T})}(q_1, q_2) &\equiv e^2 \frac{(q_1 \cdot q_2)}{M} M_{\mu\nu\alpha\beta}(q_1, q_2) \\ &\times F_{\mathcal{T}\gamma^*\gamma^*}(q_1^2, q_2^2), \end{aligned} \tag{20}$$

with

$$M_{\mu\nu\alpha\beta} = \left\{ R_{\mu\alpha}(q_1, q_2)R_{\nu\beta}(q_1, q_2) + \frac{1}{8(q_1 + q_2)^2 [(q_1 \cdot q_2)^2 - q_1^2 q_2^2]} R_{\mu\nu}(q_1, q_2) \times \left[(q_1 + q_2)^2 (q_1 - q_2)_\alpha - (q_1^2 - q_2^2)(q_1 + q_2)_\alpha \right] \times \left[(q_1 + q_2)^2 (q_1 - q_2)_\beta - (q_1^2 - q_2^2)(q_1 + q_2)_\beta \right] \right\}. \tag{21}$$

The projection operator for $J = 2$ has the form

$$P_{\alpha\beta,\gamma\delta} = \frac{1}{2} (K_{\alpha\gamma}K_{\beta\delta} + K_{\alpha\delta}K_{\beta\gamma}) - \frac{1}{3} K_{\alpha\beta}K_{\gamma\delta}, \tag{22}$$

with $K_{\alpha\beta} \equiv -g_{\alpha\beta} + p_\alpha p_\beta / p^2$.

By an appropriate change of variables one can show that the first and second diagrams of Fig. 2 give equal contributions to Eq. (8). Thus, the two-loop integral for a_μ may be defined as a sum of two terms:

$$a_\mu^{LbL} = \lim_{k \rightarrow 0} -e^6 \int \frac{d^4 q_1}{(2\pi)^4} \int \frac{d^4 q_2}{(2\pi)^4} \frac{1}{q_1^2 q_2^2 (k - q_1 - q_2)^2} \times \frac{1}{[(p + q_1)^2 - m^2][(p + k - q_2)^2 - m^2]} \times \left[\frac{F_{\mathcal{M}\gamma^*\gamma^*}(q_1^2, (k - q_1 - q_2)^2) F_{\mathcal{M}\gamma^*\gamma^*}(k^2, q_2^2)}{(k - q_2)^2 - M^2} T_1 + \frac{F_{\mathcal{M}\gamma^*\gamma^*}(q_1^2, q_2^2) F_{\mathcal{M}\gamma^*\gamma^*}((k - q_1 - q_2)^2, k^2)}{(q_1 + q_2)^2 - M^2} T_2 \right], \tag{23}$$

where $T_{1,2} = T_{1,2}(q_1, k - q_1 - q_2, q_2)$. Here, T_1 is given by the contraction of the first two terms in Eq. (11) with the tensor $T^{\mu\nu\lambda\sigma}$; whereas T_2 is defined by the contraction of $T^{\mu\nu\lambda\sigma}$ with the third term. Computing the Dirac traces (for which we used the computer algebraic system FORM [40]), we find that T_1 and T_2 contain a set of structures of three types:

$$(q_1 \cdot k)^i (q_2 \cdot k)^j, \quad \frac{(q_1 \cdot k)^i (q_2 \cdot k)^j}{(q_2 \cdot k)^2 - q_2^2 k^2}, \quad \frac{(q_1 \cdot k)^i (q_2 \cdot k)^j}{[(q_2 \cdot k)^2 - q_2^2 k^2]^2}.$$

Before taking the limit $k \rightarrow 0$, we first need to eliminate the dependence on the spatial direction of k . Since the trace under consideration projects to a scalar, we may average the $\Omega(\hat{\mathbf{k}})$ dependence over all spatial directions without changing the result for a_μ :

$$\int \frac{d\Omega(\hat{\mathbf{k}})}{4\pi} a_\mu(\mathbf{k}) = a_\mu. \tag{24}$$

After taking the limit $k \rightarrow 0$ explicitly, we integrate the angular dependence on $\Omega(\hat{\mathbf{q}}_1)$ and $\Omega(\hat{\mathbf{q}}_2)$. For the angular integrations, we use the technique based on the properties of

the Legendre polynomials. Given a particular parametrization of the FFs, the angular integrals may be performed analytically. We will give more technical details in a forthcoming publication. In this work, we will use both monopole (M) and dipole (D) parameterizations of the form

$$\frac{F_{\mathcal{M}\gamma^*\gamma^*}^M(q_1^2, q_2^2)}{F_{\mathcal{M}\gamma^*\gamma^*}^M(0, 0)} = \frac{1}{(1 - q_1^2/\Lambda_M^2)} \frac{1}{(1 - q_2^2/\Lambda_M^2)}, \tag{25}$$

$$\frac{F_{\mathcal{M}\gamma^*\gamma^*}^D(q_1^2, q_2^2)}{F_{\mathcal{M}\gamma^*\gamma^*}^D(0, 0)} = \frac{1}{(1 - q_1^2/\Lambda_D^2)^2} \frac{1}{(1 - q_2^2/\Lambda_D^2)^2}, \tag{26}$$

where Λ_M (Λ_D) are the monopole (dipole) mass parameters, respectively, which are to be determined from phenomenology.

After working out the angular integrations analytically, we perform the remaining integrals numerically. For convenience, we perform a Wick rotation for the energy component of the four-momenta q_1 and q_2 and carry out the numerical integration in polar coordinates. In particular, we make the change of variables

$$Q_i^0 = Q_i \cos \psi_i, \quad |\mathbf{Q}_i| = Q_i \sin \psi_i, \tag{27}$$

where $Q_i^2 \equiv -q_i^2$. The resulting four-dimensional integral for the case of a monopole FF takes e.g. the form

$$a_\mu^{LbL} = -\frac{4\alpha^3}{\pi^3} (2J + 1) |F_{\mathcal{M}\gamma^*\gamma^*}(0, 0)|^2 \times \int_0^\infty dQ_1 \int_0^\pi d\psi_1 \int_0^\infty dQ_2 \int_0^\pi d\psi_2 \frac{1}{Q_1^2/\Lambda^2 + 1} \frac{1}{Q_2^2/\Lambda^2 + 1} \times \frac{1}{Q_2^2 + M^2} \frac{\sin^2 \psi_1 \sin^2 \psi_2}{Q_1 + 2im \cos \psi_1} \left[2 \frac{\tilde{T}_1(Q_1, Q_2, \psi_1, \psi_2)}{Q_2 - 2im \cos \psi_2} + Q_2 \tilde{T}_2(Q_1, Q_2, \psi_1, \psi_2) \right], \tag{28}$$

where the factor $[(Q_1 + Q_2)^2 + \Lambda^2]^{-1}$ is absorbed in the expressions for \tilde{T}_1 and \tilde{T}_2 and in the third diagram a factor $[(Q_1 + Q_2)^2 + 2im(Q_1 \cos \psi_1 + Q_2 \cos \psi_2)]^{-1}$ has been absorbed into \tilde{T}_2 .

3 Result and discussion

To test our formalism, we have firstly applied it to the case of pseudoscalar meson exchanges. This case had been worked out in Ref. [26] using the Gegenbauer polynomial technique, where for pole parametrizations of the FFs the HLbL contribution to a_μ had been given by a two-dimensional numerical integral over Q_1 and Q_2 . We checked that using e.g. a monopole FF, the result obtained from Eq. (28) is in exact agreement with the result of [26]. Due to the more complicated vertex structure for axial-vector, scalar and tensor

mesons, the Gegenbauer polynomial technique cannot easily be extended, which is why we resort to the four-dimensional expression of Eq. (28). Using this formalism, we subsequently discuss our estimates for the HLbL contribution to a_μ due to axial-vector, scalar and tensor mesons.

3.1 Axial-vector mesons

For an axial-vector meson, it is conventional to define an equivalent two-photon decay width to describe its decay into one quasi-real longitudinal photon (with virtuality Q_1^2) and a transverse (real) photon as [31,36,37]

$$\tilde{\Gamma}_{\gamma\gamma} \equiv \lim_{Q_1^2 \rightarrow 0} \frac{M^2}{Q_1^2} \frac{1}{2} \Gamma(\mathcal{A} \rightarrow \gamma_L^* \gamma_T), \tag{29}$$

which allows us to express the FF normalization entering the $\mathcal{A}\gamma^*\gamma^*$ vertex of Eq. (18) as [28]

$$[F_{\mathcal{A}\gamma^*\gamma^*}(0, 0)]^2 = \frac{3}{M} \frac{4}{\pi \alpha^2} \tilde{\Gamma}_{\gamma\gamma}. \tag{30}$$

Phenomenologically, the two-photon production cross sections have been measured for the two lowest lying axial-vector mesons: $f_1(1,285)$ and $f_1(1,420)$. The most recent measurements were performed by the L3 Collaboration [36,37]. In those works, the non-relativistic quark model expression of Eq. (18) in terms of a single FF $F_{\mathcal{A}\gamma^*\gamma^*}$ has been assumed, and the resulting FF has been parameterized by a dipole form as in Eq. (26). Table 1 shows the present experimental status of the equivalent 2γ decay widths for $f_1(1,285)$, and $f_1(1,420)$, as well as the phenomenological values for the dipole mass parameters Λ_D .

Using these values, we can calculate the HLbL contributions of $f_1(1,285)$ and $f_1(1,420)$ to a_μ , which are shown in Table 1. Both contributions sum up to a value of 6.4×10^{-11} , which is roughly one order of magnitude smaller than the dominant HLbL contribution to a_μ due to the π^0 [26]. We like to emphasize that our estimate for the two dominant axial-vector meson contributions is based on available experimental information. In this way, we are also able to provide an error estimate, which derives from the experimental uncertainties in the equivalent 2γ decay widths and from the FF parameterization.

Table 1 Present values [42] of the $f_1(1,285)$ meson and $f_1(1,420)$ meson masses M , their equivalent 2γ decay widths $\tilde{\Gamma}_{\gamma\gamma}$, defined according to Eq. (29), as well as their dipole masses Λ_D entering the FF of Eq. (18). For $\tilde{\Gamma}_{\gamma\gamma}$, we use the experimental results from the L3 Collaboration: $f_1(1,285)$ from [36], $f_1(1,420)$ from [37]

	M (MeV)	$\tilde{\Gamma}_{\gamma\gamma}$ (keV)	Λ_D (MeV)	$a_\mu(10^{-11})$
$f_1(1,285)$	$1,281.8 \pm 0.6$	3.5 ± 0.8	$1,040 \pm 78$	5.0 ± 2.0
$f_1(1,420)$	$1,426.4 \pm 0.9$	3.2 ± 0.9	926 ± 78	1.4 ± 0.7
Sum				6.4 ± 2.0




Fig. 3 The density ρ_a as defined in Eq. (31), in units $10^{-10} \text{ GeV}^{-2}$, for the axial-vector meson $f_1(1,285)$. Left panel corresponds with left two diagrams of Fig. 2, right panel corresponds with right diagram of Fig. 2

In order to have a better understanding which region of virtualities in the axial-vector meson FFs is contributing mostly to this result, it is instructive to define a density function ρ_a as

$$a_\mu^{\text{LbL}} = \int_0^\infty dQ_1 \int_0^\infty dQ_2 \rho_a(Q_1, Q_2). \tag{31}$$

We show the dependence of ρ_a on the photon virtualities Q_1 and Q_2 , which enter the HLbL scattering diagram, for the axial-vector meson $f_1(1,285)$ in Fig. 3. One notices that the dominant contribution arises from the region around $Q_1 \approx Q_2 \approx 0.5 \text{ GeV}$. One also sees that the contribution beyond $Q_{1,2} \geq 1.5 \text{ GeV}$ becomes negligible.

3.2 Scalar mesons

We next proceed to the estimate for scalar mesons (S). The normalization of the FF corresponding with two transverse photons is related to the two-photon decay width of the scalar meson as [28]

Table 2 Scalar meson pole contribution to a_μ based on the present PDG values [42] of the scalar meson 2γ decay widths $\Gamma_{\gamma\gamma}$

	$\Gamma_{\gamma\gamma}$ (keV)	$a_\mu [10^{-11}] (\Lambda_M = 1 \text{ GeV})$	$a_\mu [10^{-11}] (\Lambda_M = 2 \text{ GeV})$
$f_0(980)$	0.29 ± 0.07	-0.19 ± 0.05	-0.61 ± 0.15
$f'_0(1,370)$	3.8 ± 1.5	-0.54 ± 0.21	-1.84 ± 0.73
$a_0(980)$	0.3 ± 0.1	-0.20 ± 0.07	-0.63 ± 0.21
Sum		-0.9 ± 0.2	-3.1 ± 0.8

$$[F_{S\gamma^*\gamma^*}(0, 0)]^2 = \frac{1}{M} \frac{4}{\pi\alpha^2} \Gamma_{\gamma\gamma}. \tag{32}$$

When going to virtual photons, unfortunately no empirical information is available at present for the $S\gamma^*\gamma^*$ transition FFs. We will assume a simple monopole behavior of the FF. The monopole mass Λ_M is considered as a free parameter, which we will vary in the expected hadronic range $\Lambda_M = 1 - 2 \text{ GeV}$, in order to obtain the numerical estimates for a_μ . We show our results for the HLbL contribution to a_μ due to the leading scalar mesons $f_0, f'_0,$ and a_0 in Table 2. We find a negative contribution of the scalar mesons to a_μ which is in the range -1 to -3 (in units 10^{-11}), when varying Λ_M in the range 1 to 2 GeV.

Note that in this work we do not include the contribution from the low-lying and very broad $f_0(500)$ state, which requires a full treatment of the HLbL contribution to a_μ due to 2π intermediate states [41]. Such a treatment goes beyond the pole model framework described here.

3.3 Tensor mesons

In this work, we also estimate the HLbL contribution to a_μ due to tensor mesons (\mathcal{T}). The dominant tensor mesons produced in two-photon fusion processes are given by $f_2(1,270), a_2(1,320), f_2(1,565),$ and $a_2(1,700)$; see Table 3. As described above, we will assume in our analysis that the tensor meson is only produced in a state of helicity 2. This allows one to express the normalization of the dominant (helicity-2) FF entering the $\mathcal{T}\gamma^*\gamma^*$ vertex as [28]

$$[F_{\mathcal{T}\gamma^*\gamma^*}(0, 0)]^2 = \frac{5}{M} \frac{4}{\pi\alpha^2} \Gamma_{\gamma\gamma}. \tag{33}$$

At the present moment there is unfortunately no direct experimental information about the Q^2 dependence of the tensor meson transition FFs. One can, however, resort to other phenomenological information based on exact forward sum rules for the light-by-light scattering. For $\gamma^*\gamma \rightarrow X$ fusion cross sections, with one real photon (γ) and one virtual photon (γ^*), three super convergence sum rules were derived in Refs. [27,28]. Applied to the $\gamma^*\gamma$ production of mesons, this leads to intricate relations between transition FFs of pseudoscalar, axial and tensor mesons. In order to saturate these sum rules, one obtains approximate expressions for the dominant tensor meson transition FFs, given the knowledge of the

Table 3 Tensor meson pole contribution to a_μ based on the present PDG values [42] of the tensor meson masses M and their 2γ decay widths $\Gamma_{\gamma\gamma}$

	M (MeV)	$\Gamma_{\gamma\gamma}$ (keV)	$a_\mu (\Lambda_D = 1.5 \text{ GeV}) [10^{-11}]$
$f_2(1,270)$	$1,275.1 \pm 1.2$	3.03 ± 0.35	0.79 ± 0.09
$f_2(1,565)$	$1,562 \pm 13$	0.70 ± 0.14	0.07 ± 0.01
$a_2(1,320)$	$1,318.3 \pm 0.6$	1.00 ± 0.06	0.22 ± 0.01
$a_2(1,700)$	$1,732 \pm 16$	0.30 ± 0.05	0.02 ± 0.003
Sum			1.1 ± 0.1

transition FFs for the pseudoscalar mesons. In particular, it was shown in Ref. [28] that the $\eta,$ and η' transition FFs constrain the transition FF for $f_2(1,270)$ and the π^0 transition FF constrains the corresponding transition FF for the $a_2(1,320)$ state. We found that these relations can approximately be expressed by choosing a dipole form for the tensor meson transition FF with dipole mass parameter $\Lambda_D = 1.5 \text{ GeV}$. We use this estimate in calculating the HLbL contribution to a_μ due to tensor mesons, which is shown in Table 3. We see that the four dominant tensor meson contributions add up to a contribution to a_μ of around 1 (in units 10^{-11}).

3.4 Comparison with previous works

Our results can be compared with previous estimates for axial-vector and scalar mesons, which are shown in Table 4. For tensor mesons, our results are the first estimates.

The previous estimates for axial-vector mesons differ quite a lot. The BPP estimate [43,44] is based on an extended Nambu–Jona-Lasinio model in which both a $1/N_c$ and chiral counting was used. The HKS estimate [45–47] for axial-vector meson FFs was based on a hidden local gauge symmetry model. The MV estimate [48], which was also adopted in JN [1,2] is an order of magnitude larger than the BPP and HKS estimates, and around a factor 3 larger than our estimate. The large value of Ref. [48] was obtained because a constant FF was used at the external vertex to reproduce the QCD short-distance constraints. Although such short-distance constraints are surely important for the large Q^2 behavior of the FFs, one can see from Fig. 3 that using the empirical information for the $f_1(1,285)$ transition FF, the region which dominates the HLbL contribution to a_μ is for

Table 4 HLbL contribution to a_μ (in units 10^{-11}) due to axial-vector, scalar, and tensor mesons obtained in this work, compared with various previous estimates. For our scalar meson estimate, we have quoted the value corresponding with $\Lambda_{\text{mon}} = 2 \text{ GeV}$

	Axial-vectors	Scalars	Tensors
BPP [43,44]	2.5 ± 1.0	-7 ± 2	–
HKS [45–47]	1.7 ± 1.7	–	–
MV [48]	22 ± 5	–	–
PdRV [22]	15 ± 10	-7 ± 7	–
N/JN [1,2]	22 ± 5	-7 ± 2	–
This work	6.4 ± 2.0	-3.1 ± 0.8	1.1 ± 0.1

virtualities around and below 1 GeV^2 . Besides, in case of the two-photon production of axial-vector mesons, the helicity amplitudes need to vanish in the limit of both real photons due to the Landau–Yang theorem, which is not incorporated in the QCD short-distance constraints. It has furthermore been argued by PdRV [22] that the errors in the BPP and HKS estimates were underestimated, and an intermediate estimate with larger error has been suggested, which is in agreement within 1σ with our estimate.

For the scalar mesons, BPP has performed an estimate, which was adopted by N/JN and PdRV (by increasing the error bar to 100 %). Compared with the result of BPP, our result also has the negative sign but is around a factor of 2 smaller in magnitude. Given that there is no empirical information at all on the scalar meson transition FFs, future data from BES-III would be mostly welcome here to better constrain this contribution.

For tensor mesons, no quantitative estimates had been given so far. In Ref. [22], it has been noted that if the tensor meson decays predominantly into two photons with total helicity $\Lambda = 0$, its contributions to a_μ should be negative, similar to the case for scalar mesons. In our analysis, we follow, however, the phenomenological indication that at low virtualities, the tensor meson decays predominantly into two photons with total helicity $\Lambda = 2$ [38]. For the latter decay, we find a positive contribution to a_μ .

4 Conclusions

In this letter we have presented the formalism to calculate the HLbL contribution to the muon’s anomalous magnetic moment a_μ due to axial-vector, scalar and tensor meson poles. In this way, we have extended the framework of Ref. [26], where the leading HLbL contribution due to pseudoscalar mesons was evaluated. To allow for the different Lorentz structures of the $\gamma^*\gamma^* \rightarrow$ meson vertex, we have performed a combined analytical and numerical technique, where the angular integrals over the virtual photon momenta

were performed analytically using the Legendre polynomial technique, and where the resulting four-dimensional integral for a_μ was performed numerically. We validated our method by reproducing the known result for pseudoscalar mesons. To estimate the HLbL contribution to a_μ from axial-vector, scalar and tensor mesons, we incorporated available experimental input as well as constraints from light-by-light scattering sum rules. For those mesons which have the largest known couplings to two virtual photons, we obtained as estimates

$$a_\mu(f_1, f'_1) = (6.4 \pm 2.0) \times 10^{-11},$$

$$\alpha_\mu(f_0, f'_0, a_0) = [(-0.9 \pm 0.2) \text{ to } (-3.1 \pm 0.8)] \times 10^{-11},$$

$$a_\mu(f_2, f'_2, a_2, a'_2) = (1.1 \pm 0.1) \times 10^{-11}.$$

The size of such contributions is about an order of magnitude smaller than the dominant π^0 HLbL contribution. Given a new muon $g - 2$ experiment at Fermilab, which aims to reduce the experimental uncertainty to $\delta a_\mu \approx 16 \times 10^{-11}$, it is, however, crucial to further constrain the theoretical uncertainty due to the HLbL contribution. In this respect, it would be particularly helpful to have measurements of $\gamma^*\gamma^* \rightarrow$ meson transition form factors with one and two virtual photons for axial-vector, scalar, and tensor states. As such information will become available, in particular from future measurements from BES-III, the formalism here developed can be used to further improve on the estimate of the HLbL contribution to a_μ .

Acknowledgments We like to thank A. Denig and P. Masjuan for useful discussions. This work was supported in part by the Deutsche Forschungsgemeinschaft DFG in part through the Collaborative Research Center “The Low-Energy Frontier of the Standard Model” (SFB 1044), in part through the graduate school Graduate School “Symmetry Breaking in Fundamental Interactions” (GRK 1581), and in part through the Cluster of Excellence “Precision Physics, Fundamental Interactions and Structure of Matter” (PRISMA).

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

Funded by SCOAP³ / License Version CC BY 4.0.

References

1. F. Jegerlehner, A. Nyffeler, Phys. Rept. **477**, 1 (2009)
2. F. Jegerlehner, Springer Tracts Mod. Phys. **226**, 1 (2008)
3. G.W. Bennett et al. [Muon $g-2$ Collaboration], Phys. Rev. Lett. **92**, 161802 (2004)
4. G.W. Bennett et al. [Muon $G-2$ Collaboration], Phys. Rev. D **73**, 072003 (2006)
5. T. Aoyama, M. Hayakawa, T. Kinoshita, M. Nio, Phys. Rev. Lett. **109**, 111808 (2012)
6. T. Aoyama, M. Hayakawa, T. Kinoshita, M. Nio, Phys. Rev. D **85**, 093013 (2012)
7. K. Fujikawa, B.W. Lee, A.I. Sanda, Phys. Rev. D **6**, 2923 (1972)

8. A. Czarnecki, B. Krause, W.J. Marciano, Phys. Rev. Lett. **76**, 3267 (1996)
9. M. Knecht, S. Peris, M. Perrottet, E. De Rafael, JHEP **0211**, 003 (2002)
10. A. Czarnecki, W.J. Marciano, A. Vainshtein, Phys. Rev. D **67**, 073006 (2003) [Erratum-ibid. D **73**, 119901 (2006)]
11. M. Davier, S. Eidelman, A. Höcker, Z. Zhang, Eur. Phys. J. C **27**, 497 (2003)
12. M. Benayoun et al., Eur. Phys. J. C **55**, 199 (2008)
13. M. Benayoun, P. David, L. DelBuono, O. Leitner, Eur. Phys. J. C **65**, 211 (2010)
14. M. Benayoun, P. David, L. DelBuono, O. Leitner, Eur. Phys. J. C **68**, 355 (2010)
15. M. Davier et al., Eur. Phys. J. C **66**, 1 (2010)
16. M. Davier et al., Eur. Phys. J. C **66**, 127 (2010)
17. M. Davier, A. Höcker, B. Malaescu, Z. Zhang, Eur. Phys. J. C **71**, 1515 (2011) [Erratum-ibid. C **72**, 1874 (2012)]
18. F. Jegerlehner, R. Szafron, Eur. Phys. J. C **71**, 1632 (2011)
19. K. Hagiwara, R. Liao, A.D. Martin, D. Nomura, T. Teubner, J. Phys. G **38**, 085003 (2011)
20. M. Benayoun, P. David, L. DelBuono, F. Jegerlehner, Eur. Phys. J. C **73**, 2453 (2013)
21. F. Jegerlehner, Acta Phys. Polon. B **44**(11), 2257 (2013)
22. J. Prades, E. de Rafael, A. Vainshtein. [arXiv:0901.0306](https://arxiv.org/abs/0901.0306) [hep-ph]
23. T. Blum, A. Denig, I. Logashenko, E. de Rafael, B. L. Roberts, T. Teubner, G. Venanzoni. [arXiv:1311.2198](https://arxiv.org/abs/1311.2198) [hep-ph]
24. B. Lee Roberts [Fermilab P989 Collaboration], Nucl. Phys. Proc. Suppl. **218**, 237 (2011)
25. H. Iinuma [J-PARC New g-2/EDM experiment Collaboration], J. Phys. Conf. Ser. **295**, 012032 (2011)
26. M. Knecht, A. Nyffeler, Phys. Rev. D **65**, 073034 (2002)
27. V. Pascalutsa, M. Vanderhaeghen, Phys. Rev. Lett. **105**, 201603 (2010)
28. V. Pascalutsa, V. Pauk, M. Vanderhaeghen, Phys. Rev. D **85**, 116001 (2012)
29. S.J. Brodsky, J.D. Sullivan, Phys. Rev. **156**, 1644 (1967)
30. M. Poppe, Int. J. Mod. Phys. A **1**, 545 (1986)
31. G.A. Schuler, F.A. Berends, R. van Gulik, Nucl. Phys. B **523**, 423 (1998)
32. L.D. Landau, Dokl. Akad. Nauk Ser. Fiz. **60**, 207 (1948)
33. C.-N. Yang, Phys. Rev. **77**, 242 (1950)
34. R.N. Cahn, Phys. Rev. D **35**, 3342 (1987)
35. R.N. Cahn, Phys. Rev. D **37**, 833 (1988)
36. P. Achard et al. [L3 Collaboration], Phys. Lett. B **526**, 269 (2002)
37. P. Achard et al. [L3 Collaboration], JHEP **0703**, 018 (2007)
38. M.R. Pennington, T. Mori, S. Uehara, Y. Watanabe, Eur. Phys. J. C **56**, 1 (2008)
39. P. Grassberger, R. Kogerler, Nucl. Phys. B **106**, 451 (1976)
40. J. Kuipers, T. Ueda, J.A.M. Vermaseren, J. Vollinga, Comput. Phys. Commun. **184**, 1453 (2013)
41. G. Colangelo, M. Hoferichter, M. Procura, P. Stoffer. [arXiv:1402.7081](https://arxiv.org/abs/1402.7081) [hep-ph]
42. K. Nakamura et al. [Particle Data Group Collaboration], J. Phys. G **37**, 075021 (2010)
43. J. Bijnens, E. Pallante, J. Prades, Phys. Rev. Lett. **75**, 1447 (1995) [Erratum-ibid. **75**, 3781 (1995)]
44. J. Bijnens, E. Pallante, J. Prades, Nucl. Phys. B **474**, 379 (1996) [Erratum-ibid. **626**, 410 (2002)]
45. M. Hayakawa, T. Kinoshita, A.I. Sanda, Phys. Rev. Lett. **75**, 790 (1995)
46. M. Hayakawa, T. Kinoshita, A.I. Sanda, Phys. Rev. D **54**, 3137 (1996)
47. M. Hayakawa, T. Kinoshita, Phys. Rev. D **57**, 465 (1998) [Erratum-ibid. D **66**, 019902 (2002)]
48. K. Melnikov, A. Vainshtein, Phys. Rev. D **70**, 113006 (2004)