



Factors associated with post-stroke depression and fatigue: lesion location and coping styles

Changjuan Wei¹ · Fang Zhang¹ · Li Chen¹ · Xiaofeng Ma¹ · Nan Zhang¹ · Junwei Hao¹

Received: 28 July 2015 / Revised: 24 October 2015 / Accepted: 26 October 2015 / Published online: 14 November 2015
© The Author(s) 2015. This article is published with open access at Springerlink.com

Abstract Post-stroke depression (PSD) and post-stroke fatigue (PSF) are frequent and persistent problems among stroke survivors. Therefore, awareness of signs and symptoms of PSD and PSF is important for their treatment and recovery from stroke. Additionally, since sudden serious illness can result in disequilibrium, early institution of a coping process is essential to restoring stability. The brain damage of stroke leaves patients with unique physical and mental dysfunctions for which coping maybe a key resource while rebuilding lives. We evaluated 368 consecutive patients with acute ischemic stroke for post-stroke emotional disorders at admission and 3 months later. PSD was evaluated by using the Beck Depression Inventory, and PSF was scored with the Fatigue Severity Scale. The Social Support Rating Scale and Medical Coping Modes Questionnaire were also used as measurement tools. Locations of lesions were based on MRI. Those scans revealed infarcts located in the basal ganglia, corona radiate and internal capsule and constituted the independent factors associated with PSF 3 months after stroke occurrence. Conversely, PSD was not related to lesion location. Acceptance-resignation related to PSD and PSF both at admission and 3 months after stroke. Avoidance was the independent factor most closely related to PSD, whereas confrontation was the independent factor best related to PSF at 3 months after stroke onset.

Keywords Post-stroke depression · Post-stroke fatigue · Coping styles · Lesion location

Introduction

Each year, of the approximately 15 million human stroke victims worldwide, at least five million die, and one-third remain disabled [1]. Overwhelming feelings of fatigue, that is, tiredness and lack of energy, is the chief complaint in more than half of the 10 million survivors [2]. Fatigue as a common post-stroke emotional disturbance always impairs patients' ability to regain lost functions [3] and leads to negative long-term outcomes. Similarly, post-stroke depression (PSD) mainly manifested as sadness, reduction of interest and pleasure, and multiple psychological and vegetative symptoms [4] afflicts 20–65 % of these patients, depending on the population studied, the assessment measures, and the definition of depression applied [5]. As the most common psychological sequel of stroke, patients who had PSD experienced the least benefit from rehabilitation, the poorest quality of life and a substantially increased risk of suicide [6].

Several cross-sectional papers have linked fatigue with depression after stroke [7]. In a Swedish study, 49 % of patients with fatigue 1 year after stroke were diagnosed with depression compared with 39 % in the total sample [8]. Similarly, in a Korean study, 34 % of patients diagnosed as depressed were among those with fatigue approximately 15 months after stroke [7]. Fatigue was considered an essential component of PSD [9], and depression-related symptoms such as insomnia or appetite loss resulted in fatigue [10]. In contrast, PSF accompanying PSD is often relieved when the depression is adequately treated [11]. An association between these two syndromes

✉ Junwei Hao
hjw@tjmu.edu.cn

¹ Department of Neurology and Tianjin Neurological Institute, Tianjin Medical University General Hospital, Tianjin 300052, China

at a particular time point does not necessarily mean that one causes the other or exclude the possibility that a third factor causes both. Nevertheless, among all these attempts to make sense of PSD and PSF, identifying predictors and early signs is crucial for taking preventive measures, promoting early diagnosis, implementing early and adequate treatment, and improving quality of life, both for patients with stroke and for their caregivers.

Some studies invoke more physiologic than behavioral causes for PSD and PSF by suggesting that the extent of stroke-induced functional residual impairment is the major risk factor. Others suggest a possible biological relationship between the occurrence of emotional disturbances and structural brain damage [5]. However, the effect of lesion location has remained the most controversial purported cause of PSD and PSF. With, social support and coping styles also prominent considerations. Coping refers to the person's cognitive and behavioral efforts to manage (reduce, minimize, master, or tolerate) the internal and external demands of his/her environment that seem too taxing or exceed personal resources [12]. Moos and Tsu [13] proposed that sudden serious illness often results in disequilibrium, which triggers adaptive processes to restore equilibrium. The coping process, initiated to restore equilibrium (adaptation), includes cognitive appraisal of the illness, identification of adaptive tasks and coping skills. Contextual factors (background, patient illness and social–environmental variables) shape these coping factors. Emotional disturbance and physical health are designated as adaptation outcomes. Since stroke is considered a special physical illness with neurological dysfunction and brain damage, coping may be the key psychological resource involved in rebuilding the lives of patients. Studies assessing coping strategies in patients after stroke have produced inconsistent results [14]. In this context, we evaluated PSD and PSF at different stages after cerebral infarction and attempted to correlate patients' symptoms with lesion location, coping styles and other variables. The goal of this research is to provide new insight into the rehabilitation and management of stroke.

Subjects and methods

Participants

This descriptive, cohort study was conducted with stroke patients upon admission and at 3 months afterward. Our study was approved by the ethics committees of Tianjin Medical University General Hospital. All participants provided written informed consent. We evaluated 989 consecutive patients who were admitted to Tianjin Medical University General Hospital (Tianjin, China) with a

diagnosis of acute stroke, between Jan 2012 and Jun 2014. The diagnosis was confirmed based on CT or MRI findings within 7 days after stroke onset. Exclusion criteria applied to patients with hemorrhagic stroke ($n = 83$); those with unusual causes such as dissection, venous infarction, or moyamoya disease ($n = 77$); those with transient ischemic attack without progression to stroke ($n = 120$); those with communication problems (decreased consciousness, confusion, aphasia, dementia, or dysarthria) severe enough to preclude a reliable interview ($n = 142$); and those with very severe neurologic or medical conditions (such as metastatic cancer) ($n = 54$). We also excluded patients who scored ≤ 23 on the Mini-Mental State Examination ($n = 31$), patients diagnosed with depression or other psychiatric illnesses or treated with selective serotonin reuptake inhibitors before the onset of stroke ($n = 16$), patients who lived alone (because no information was available from relatives) ($n = 9$), patients who did not undergo a complete follow-up ($n = 37$), and patients who did not sign an informed consent ($n = 52$). Finally, 368 patients were enrolled into the study.

Before the project began officially, all interviewers/raters completed and passed a consistency-training examination. Two researchers (L.C. and F. Z.) recorded all clinical and socio-demographic information. The neuropsychological evaluations were recorded by one of the authors (C. W.) and supervised by a psychiatrist (L. L.). An experienced stroke neurologist (N. Z.) documented patients' neurologic findings during the 3-month follow-up, and one of the authors with experience in neuroimaging (F. Z.) analyzed MRI and Fazeka's scale scores. Questions arising during subsequent interviews were brought to a research team meeting for group consensus on the appropriate answer.

Testing instruments

Depression was assessed with the Beck Depression Inventory (BDI) [15], which was self-recorded. Fatigue was measured on the fatigue severity scale [16, 17]. To calculate cognitive functioning we applied the Mini-Mental State Examination (MMSE) [18], a measure of global cognitive decline that encompasses basic cognitive domains. The Social Support Rating Scale (SSRS) [19] was adopted to evaluate social support, which consisted of three dimensions: objective support, subjective support and the degree of social utilization. Coping styles were measured with the Medical Coping Modes Questionnaire (MCMQ) [20, 21], which is a 19-item instrument that addresses three forms of coping: confrontation, avoidance and acceptance-resignation.

MRI studies were performed on a 3.0T clinical scanner (HDx, General Electric, USA) with an 8-channel phase-

array coil. Locations of lesions based on MRI were characterized as anterior cortical when identified in the anterior cerebral artery territory, the frontal and parietal areas of the middle cerebral artery, or predominantly in the temporal lobe of the middle cerebral artery territory. Posterior cortical referred to the lesions at the following occipital area or medial temporal area of the posterior cerebral artery territory; internal capsule/corona radiata/basal ganglia of the lenticulostriate artery territory; thalamus; pons, including pure midbrain lesions; medulla; and cerebellum [19]. White matter intensity (leukoaraiosis) derived from fluid-attenuated inversion recovery (FLAIR) imaging was graded from 0 to 3 on Fazeka's scale of deep white matter changes, with scores of 2 and 3 indicating significant leukoaraiosis [20].

Procedure

Each first interview was completed at approximately 14 days after the onset of stroke to ensure that the patients were stable. We tried our best to avoid conducting the interviews under conditions of acute neurologic progression or any other condition that could affect the emotions of patients or assessments made by our team. To ensure that each evaluation was reliable, patients were asked to take the interview accompanied by the caregiver who lived with him/her. The caregiver was present to verify the information and to assist the patient, but could not take part in the assessment. Neurologic and psychological assessments (including BDI, FSS, SSRS, MCMQ, Fazeka's scale and symptom observation) were completed by doctors assigned specifically for that purpose. MMSE was used for cognition screening, excluding dementia. The second patient interview was performed 3 months after the onset of stroke. Neurologic and psychological assessments were completed by the same doctors who conducted the first interview.

Diagnostic criteria

PSD was diagnosed in patients with a BDI score >13 or those who met the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV) [22, 23]. The presence of PSF is defined as an FSS score of 4.0 or more, because fewer than 5 % of healthy controls rate their fatigue at that level. However, 60–90 % of patients with medical disorders experience fatigue at or above that score [3].

Statistical analyses

Ages, years of education, MCMQ and SSRS scores were compared using Mann–Whitney *U* tests, because they were not normally distributed. Variables of frequency were

compared using Chi-square tests. Fisher's exact probability test was used in tests of dichotomous variables when small numbers were involved. Multiple logistic regression analysis was used to explore relationships among PSF/PSD variables. The level of significance was set at $P < 0.05$. All data were analyzed by SPSS version 19.0 (IBM Corp., Armonk, NY).

Results

Prevalence of PSD and PSF

A total of 368 participants was enrolled within 14 days after the onset of stroke and completed their second interview at 3 months after the stroke. PSD was diagnosed in 19.3 % of the patients at admission and in 23.6 % at 3 months. Of the 71 patients who developed PSD during the acute stage of stroke, 23 recovered but 48 still had PSD after 3 months. In comparison, 39 patients without PSD at admission had developed PSD at 3 months. PSF was present in 23.4 % of the patients at admission and in 29.6 % at 3 months. Of the 86 patients who developed PSF during the acute stage of stroke, 28 recovered and 58 still had PSF at 3 months, whereas 51 patients without PSF at admission developed PSF at 3 months. Further, 75.6 % (65/86) of the patients who had PSF at admission presented with PSD 3 months after stroke. More than half (38/71, 53.5 %) of the patients who had PSD at admission presented with PSF 3 months after stroke. In addition, 10.9 % (40/368) and 12.2 % (45/368) of the patients had PSD accompanied by PSF at admission and 3 months later, respectively.

Factors related to PSD and PSF

Both motor and sensory dysfunction at admission ($P < 0.01$) compared with only motor or only sensory dysfunction was related to PSD at admission. Patients who prefer acceptance-resignation (MCMQ; $P < 0.01$) were prone to PSD at admission. Three months later, none of the above risk factors for PSD changed significantly. Moreover, the patients who manifested a lower degree of social utilization (SSRS; $P < 0.05$) and chose avoidance coping styles (MCMQ; $P < 0.01$) were more prone to PSD. We also found that the patients who had only sensory dysfunction at admission ($P < 0.05$) were not susceptible to PSD (Table 1).

Patients with both motor and sensory dysfunction ($P < 0.01$) were more susceptible to PSF at admission compared to individuals with motor or only sensory dysfunction ($P < 0.01$). Moreover, acceptance-resignation (MCMQ; $P < 0.01$) styles, low objective support (SSRS; $P < 0.05$) and degree of social utilization (SSRS;

Table 1 Factors associated with PSD at admission and 3 months later

Variable	Admission		3 months after stroke	
	Present (<i>n</i> = 71)	Absent (<i>n</i> = 297)	Present (<i>n</i> = 87)	Absent (<i>n</i> = 281)
Age (y), mean ± SD	61.3 ± 9.4	61.6 ± 9.8	62.0 ± 9.7	60.0 ± 9.6
Female, <i>n</i> (%)	21 (29.6)	86 (29.0)	29 (33.3)	78 (27.8)
Education (y), mean ± SD	10.7 ± 3.5	10.8 ± 3.4	11.0 ± 3.7	10.7 ± 3.3
Previous stroke, <i>n</i> (%)	33 (46.5)	147 (49.5)	38 (43.7)	142 (50.5)
Weekly working time, <i>n</i> (%)				
0 h	52 (73.2)	164 (55.2)	56 (64.4)	160 (56.9)
1–20 h	1 (1.4)	10 (3.4)	1 (1.1)	10 (35.6)
21–30 h	2 (2.8)	20 (6.7)	2 (2.3)	20 (7.1)
31–40 h	7 (9.9)	64 (21.5)	15 (17.2)	56 (19.9)
41–50 h	5 (7.0)	23 (7.7)	6 (6.9)	22 (7.8)
51–60 h	3 (4.2)	9 (3.0)	5 (5.7)	7 (2.5)
>60 h	1 (1.4)	7 (2.4)	2 (2.3)	6 (2.1)
Lesion location, <i>n</i> (%)				
Anterior cortex	15 (21.1)	71 (23.9)	24 (27.6)	62 (22.1)
CR + BG + IC	18 (25.3)	92 (31.0)	25 (28.7)	85 (30.2)
Thalamus	5 (7.0)	35 (11.8)	8 (9.2)	32 (11.4)
Pons + midbrain	20 (28.2)	42 (14.1)	18 (20.7)	44 (15.7)
Medulla	1 (1.4)	8 (2.7)	1 (1.1)	8 (2.8)
Cerebellum	4 (5.6)	17 (5.7)	4 (4.6)	17 (6.0)
Posterior cortex	8 (11.3)	32 (10.8)	7 (8.0)	33 (11.7)
Laterality, <i>n</i> (%)				
Left	33 (46.5)	133 (44.8)	40 (46.0)	126 (44.8)
Right	24 (33.8)	128 (43.1)	33 (37.9)	119 (42.3)
Bilateral	14 (19.7)	36 (12.1)	14 (16.1)	36 (12.8)
White matter change, <i>n</i> (%)				
Severe	41 (57.7)	144 (48.5)	44 (50.6)	141 (50.2)
Mild	30 (42.3)	153 (51.5)	43 (23.8)	140 (49.8)
Dysfunction at admission, <i>n</i> (%)				
Motor dysfunction	35 (49.3)	165 (55.6)	45 (51.7)	155 (55.2)
Sensory dysfunction	7 (9.9)	49 (16.5)	6 (6.9)	50 (17.8)*
Both motor and sensory dysfunction	25 (35.2)	46 (15.5)**	33 (37.9)	38 (13.5)**
MCMQ, mean ± SD				
Confrontation	20.4 ± 2.5	20.1 ± 2.5	20.4 ± 2.8	20.1 ± 2.4
Avoidance	14.5 ± 1.6	14.2 ± 1.5	14.9 ± 1.8	14.1 ± 1.4**
Acceptance-resignation	9.5 ± 2.0	7.9 ± 1.8**	10.3 ± 1.7	7.6 ± 1.6**
SSRS, mean ± SD				
Subjective support	23.6 ± 2.2	23.8 ± 2.1	23.6 ± 2.2	23.8 ± 2.1
Objective support	9.9 ± 0.9	9.7 ± 1.0	9.8 ± 1.1	9.7 ± 0.9
Degree of social utilization	6.0 ± 1.0	6.0 ± 1.0	5.8 ± 0.9	6.1 ± 1.1*

BG basal ganglia, CR corona radiata, IC internal capsule, MCMQ Medical Coping Modes Questionnaire, *n* number, PSD post-stroke depression, SSRS Social Support Rating Scale, *y* years

* $P \leq 0.05$

** $P \leq 0.01$

($P < 0.05$) were also predictive factors for PSF. Basal ganglia (BG), corona radiata (CR) or internal capsule (IC) infarction ($P < 0.05$), low degree of social utilization

(SSRS; $P < 0.01$), confrontation (MCMQ; $P < 0.05$), and acceptance-resignation (MCMQ; $P < 0.01$) were all significant risk factors associated with PSF at 3 months

Table 2 Factors associated with PSF at admission and 3 months later

Variable	Admission		3 months after stroke	
	Present (n = 86)	Absent (n = 282)	Present (n = 109)	Absent (n = 259)
Age (y), mean ± SD	61.8 ± 9.5	60.5 ± 10.1	61.7 ± 9.2	61.5 ± 9.8
Female, n (%)	22 (25.6)	85 (30.1)	26 (23.9)	81 (31.3)
Education (y), mean ± SD	10.7 ± 3.7	10.8 ± 3.4	10.7 ± 3.4	10.8 ± 3.4
Previous stroke, n (%)	40 (46.5)	140 (49.6)	54 (49.5)	126 (48.6)
Weekly working hours, n (%)				
0 h	54 (62.8)	162 (57.4)	62 (56.9)	154 (59.5)
1–20 h	0 (0.0)	11 (3.9)	0 (0.0)	11 (4.2)*
21–30 h	4 (4.7)	18 (6.4)	5 (4.6)	17 (6.6)
31–40 h	14 (16.3)	57 (20.2)	21 (19.3)	50 (19.3)
41–50 h	8 (9.3)	20 (7.1)	11 (10.1)	17 (6.6)
51–60 h	4(4.7)	8(2.8)	8(7.3)	4(15.4)
More than 60 h	2 (2.3)	6 (2.1)	2 (1.8)	6 (2.3)
Lesion location, n (%)				
Anterior cortex	19 (22.1)	67 (23.8)	22 (20.2)	64 (24.7)
CR + BG + IC	29 (33.7)	81 (28.7)	42 (38.5)	68 (26.3)*
Thalamus	9 (10.5)	31 (11.0)	12 (11.0)	28 (10.8)
Pons + midbrain	19 (22.1)	43 (15.2)	20 (18.3)	42 (16.2)
Medulla	1 (1.2)	8 (2.8)	4 (3.7)	5 (1.9)
Cerebellum	3 (3.5)	18 (6.4)	2 (1.8)	19 (7.3)*
Posterior cortex	6 (7.0)	34 (12.1)	7 (6.4)	33 (12.7)
Laterality, n (%)				
Left	38 (32.6)	128 (45.4)	47 (43.1)	119 (45.9)
Right	35 (40.7)	117(41.5)	48(44.0)	104(40.2)
Bilateral	13 (15.1)	37 (13.1)	14 (12.8)	36 (13.9)
White matter change, n (%)				
Severe	44 (51.2)	141 (50.0)	56 (51.4)	129 (49.8)
Mild	42 (48.8)	141 (50.0)	53 (48.6)	130 (50.2)
Dysfunction at admission, n (%)				
Motor dysfunction	48 (55.8)	152 (53.9)	55 (50.5)	145 (56.0)
Sensory dysfunction	6 (7.0)	50 (17.7)*	17 (15.6)	39 (15.1)
Both motor and sensory dysfunction	29 (33.7)	42 (14.9)**	29 (26.6)	42 (16.2)
MCMQ score, mean ± SD				
Confrontation	20.4 ± 2.4	20.1 ± 2.5	20.7 ± 2.5	20.0 ± 2.5*
Avoidance	14.3 ± 2.0	14.0 ± 1.6	14.2 ± 2.1	14.0 ± 1.5
Acceptance-resignation	9.8 ± 2.2	7.7 ± 1.7**	8.6 ± 2.4	8.0 ± 1.8*
SSRS score, mean ± SD				
Subjective support	23.6 ± 2.2	23.8 ± 2.1	23.8 ± 2.1	23.7 ± 2.1
Objective support	9.1 ± 1.1	9.7 ± 1.0*	9.8 ± 0.9	9.7 ± 1.0
Degree of social utilization	5.8 ± 0.9	6.1 ± 1.1*	5.7 ± 0.8	6.1 ± 1.1**

BG basal ganglia, CR corona radiata, IC internal capsule, MCMQ Medical Coping Modes Questionnaire, n = number, PSF post-stroke fatigue, SSRS Social Support Rating Scale, y years

* P ≤ 0.05

** P ≤ 0.01

(Table 2). We also found that patients with 1–20 h weekly working time and cerebellum infarction were not susceptible to PSF (P < 0.05; Table 2).

As presented in tabular form (Table 3), multivariate logistic regression analysis indicated that initial presentation with PSD at admission was related to both motor and

sensory dysfunction ($P < 0.05$) and acceptance-resignation (MCMQ; $P < 0.01$), whereas PSD at 3 months was associated with degree of social utilization (SSRS; $P < 0.01$), avoidance (MCMQ; $P < 0.01$) and acceptance-resignation (MCMQ; $P < 0.01$). PSF at admission was also associated with acceptance-resignation (MCMQ; $P < 0.01$) and low degree of social utilization (SSRS; $P < 0.01$), whereas it was related to BG, CR or IC infarction ($P < 0.05$), the confrontation (MCMQ; $P < 0.05$) and acceptance-resignation (MCMQ; $P < 0.01$) styles, and low degree of social utilization (SSRS; $P < 0.01$) (Table 3).

Discussion

Here as we tracked likely predictors of PSD and PSF in stroke patients to promote early identification and prevention, both conditions changed constantly over time. Fatigue and depression often accompanied occurrence after stroke. Compared with previous studies in which PSF ranged from

36 to 77 % of stroke-afflicted subjects [24], the extent of morbidity documented here represented by was lower, that is, of 71 stroke patients we reviewed. PSF was present in 23.4 % at admission and 29.6 % at 3 months. However, throughout this more than 2-year study, we observed that PSD and PSF changed constantly as time passed.

Dopamine and serotonin are considered responsible for fatigue in patients with Parkinson's disease (PD) [25]. With respect to localization of brain damage, basal ganglia, corona radiata or internal capsule infarctions were the independent factors associated with PSF 3 months after stroke occurrence, whereas cerebellum infarction was not susceptible to PSF. Previously, dopamine reward neurons localized in the ventral tegmental area were found projecting into the ventral striatum [26]. Animal experiments have shown that profuse serotonergic fibers from the brainstem raphe nuclei project to the basal ganglia and the cerebellum [27]. In accord, our results suggested that, PSF was related to derangement of dopaminergic system secondary to strokes occurring in strategic areas. Elsewhere,

Table 3 Factors associated with PSD and PSF as analyzed using multiple logistic regression

Variable	<i>B</i>	SE	<i>P</i> value	Exp (<i>B</i>)
PSD at admission				
Both motor and sensory dysfunction at admission: yes	0.709	0.317	0.026	2.031
MCMQ score: acceptance-resignation, mean \pm SD	0.385	0.075	0.001	1.470
Constant	-4.945	0.680	0.000	0.007
PSD 3 months later				
Sensory dysfunction at admission: yes	-0.638	0.602	0.289	0.529
Both motor and sensory dysfunction at admission: yes	0.537	0.387	0.165	1.711
MCMQ: avoidance, mean \pm SD	0.404	0.118	0.001	1.498
MCMQ: acceptance-resignation, mean \pm SD	1.061	0.130	0.000	2.888
SSRS: degree of social utilization, mean \pm SD	-0.558	0.179	0.001	0.555
Constant	-13.001	2.289	0.000	0.000
PSF at admission				
Sensory dysfunction at admission: yes	-0.611	0.511	0.232	0.543
Both motor and sensory dysfunction at admission: yes	0.321	0.340	0.345	1.379
MCMQ: acceptance-resignation, mean \pm SD	0.624	0.087	0.000	1.866
SSRS: objective support, mean \pm SD	0.169	0.144	0.238	1.185
SSRS: degree of social utilization, mean \pm SD	-0.401	0.154	0.009	0.670
Constant	-5.849	1.988	0.003	0.003
PSF 3 month later				
Weekly working hours: 1–20 h	-20.220	11,799.472	0.999	0.000
Lesion location: CR + BG + IC	0.634	0.260	0.015	1.885
Lesion location: cerebellum	-1.060	0.773	0.170	0.346
MCMQ score: confrontation, mean \pm SD	0.118	0.051	0.021	1.125
MCMQ score: acceptance-resignation, mean \pm SD	0.214	0.061	0.000	1.239
SSRS score: degree of social utilization, mean \pm SD	-0.410	0.133	0.002	0.664
Constant	-2.727	1.549	0.078	0.065

MCMQ Medical Coping Modes Questionnaire, *PSD* post-stroke depression, *PSF* post-stroke fatigue, *SSRS* Social Support Rating Scale

pharmacological studies yielded a reduction in fatigue following the administration of a dopamine agonist [28]. High, moderate and low concentrations of extracellular dopamine induced euphoric, seeking and aversive states, respectively. Still other findings identified circuit loops involving the cerebral cortex, basal ganglia, thalamus, epithalamus, and midbrain through which dopaminergic activity affected motivation [29]. In that study, lack of energy or motivation, boredom, adynamia and lassitude were the main symptoms of fatigue after stroke. Overall, in our hands, the relationship between PSF and lesion location was not significant during the acute stage indicating that the redistribution of neurotransmitters may take some time. Moreover, there was no evidence that PSD was related to lesion location, despite the correlation between PSF and PSD. Quite possibly though, PSD is an emotional disturbance associated with social stress, premorbid personality or disability caused by stroke. In support, our study confirmed that types of neurologic dysfunction, social support and coping styles were closely associated with PSD. Since neurologic dysfunction was not an independent factor related to PSF, correct cognitive appraisal may be necessary to determine the pathogenesis of PSF.

Although we recorded age, gender, extent of formal education and stroke history as background factors for the patients included here, no significant differences emerged, either at admission or at 3 months later. Although patients with 1–20 h weekly working time were not susceptible to PSF, this feature was not an independent factor involved in development of PSF as judged by multivariate analyses. Social support is often found to protect against the effects of stress, with most studies finding either a main effect or a buffering effect [30]. In our study, a low degree of social utilization as a cognitive appraisal variable was an independent risk factor associated with PSD at 3 months and PSF at both time points after stroke. So, regardless of how much social support was present, patients were routinely encouraged to seek and use those opportunities during the stroke recovery period. In addition, coping styles were associated with the ability to seek social support. Feifel [20] pointed out that acceptance-resignation indicated that the expression of negative affect, lack of focus, and gloomy expectations about the future always induced longer illness. Records of acceptance-resignation related to PSD and PSF both at admission and at 3 months later revealed that these patients focused on their limited power to influence the course of illness and lost confidence for recovery. High avoidance scores of PSD patients denoted less self-directed life orientation and negative self-perception [20]. Avoidance only associated PSD for individuals with subacute stroke indicated that this style may be effective for managing short-term threats, but for the long-term problem-solving activity was more effective [14]. Higher scores

for confrontation by PSF patients indicated the presence of extroversion, negative self-perception, a view of their illness as serious, religiosity, concern about life direction, the situational variables of fearing death, and positive expectations about recovery and the future. So apart from effective cognitive ability, a confrontational coping style was more important for patients with stroke. The results of this study have potentially important implications for intervention. Possibly, a change in strategies for coping with stroke would reverse HPA-axis dysfunction, thereby avoiding the symptoms of emotional disturbances and the related pathogenic effects. Finally, the presence of a significant relationship between coping and cortisol has been noted in other studies [31]. With strategies for a new direction to predict and to treat PSD and PSF, and by engendering a greater willingness on the part of post-stroke patients to be participants in their own recovery, medical practitioners might improve outcomes.

The main limitations of our study are the size of our sample, the single medical center used as a source of patients and the short time period of only 3 months. Because this is the first study that combines the specific attributes measured here, caution dictated applying these limitations as a foundation before engaging in a larger, longer investigation. Further, we excluded patients with severe neurologic conditions, aphasia, and cognitive impairment and those who lived alone. The patients who dropped out of the study before 3 months were older and had more severe neurologic dysfunction at admission, suggesting that occurrence of emotional disturbances may have been underestimated.

Conclusion

Basal ganglia, corona radiata or internal capsule infarction were the independent factors associated with PSF 3 months after stroke occurrence. The appearance of PSD was not related to lesion location. Both motor and sensory dysfunction was independently associated with PSD at admission. A low degree of social utilization was the independent factor associated with PSD and PSF at 3 months after stroke. Acceptance-resignation independently related to PSD and PSF both at admission and 3 months after stroke. Avoidance was the independent factor related to PSD at 3 months, whereas confrontation was the independent factor related to PSF at 3 months after stroke onset.

Acknowledgments We thank our patients for participating in this study; Drs. C. Chen, M. Xing for facilitating recruitment of the patients; Dr. C. Zhang for advice; Drs. R. Xue for technical support. This work was financially supported by the National Natural Science Foundation of China (81100887 and 81273287 to Junwei Hao) and the National Key Clinical Specialty Construction Project of China.

Compliance with ethical standards

Conflicts of interest The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the paper.

Ethical standard The study was approved by the ethics committees of Tianjin Medical University General Hospital.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- WHO publishes definitive atlas on global heart disease and stroke epidemic (2004). *Indian J Med Sci* 58 (9):405–406
- Brodthmann A, van de Port IG (2013) Fitness, depression, and poststroke fatigue: worn out or weary? *Neurology* 81(18):1566–1567. doi:10.1212/WNL.0b013e3182a9f59b
- Tang WK, Chen YK, Mok V, Chu WC, Ungvari GS, Ahuja AT, Wong KS (2010) Acute basal ganglia infarcts in poststroke fatigue: an MRI study. *J Neurol* 257(2):178–182. doi:10.1007/s00415-009-5284-2
- Li M, Zhang XW, Hou WS, Tang ZY (2015) Impact of depression on incident stroke: a meta-analysis. *Int J Cardiol* 180:103–110. doi:10.1016/j.ijcard.2014.11.198
- De Ryck A, Brouns R, Geurden M, Elseviers M, De Deyn PP, Engelborghs S (2014) Risk factors for poststroke depression: identification of inconsistencies based on a systematic review. *J Geriatr Psychiatry Neurol* 27(3):147–158. doi:10.1177/0891988714527514
- Pompili M, Venturini P, Campi S, Seretti ME, Montebovi F, Lamis DA, Serafini G, Amore M, Girardi P (2012) Do stroke patients have an increased risk of developing suicidal ideation or dying by suicide? An overview of the current literature. *CNS Neurosci Ther* 18(9):711–721. doi:10.1111/j.1755-5949.2012.00364.x
- Choi-Kwon S, Han SW, Kwon SU, Kim JS (2005) Poststroke fatigue: characteristics and related factors. *Cerebrovascular diseases (Basel, Switzerland)* 19(2):84–90. doi:10.1159/000082784
- Carlsson GE, Moller A, Blomstrand C (2003) Consequences of mild stroke in persons < 75 years—a 1-year follow-up. *Cerebrovasc Dis* 16(4):383–388. doi:10.1159/000072561
- Astrom M, Adolfsson R, Asplund K (1993) Major depression in stroke patients. A 3-year longitudinal study. *Stroke* 24(7):976–982
- Mohsenin V, Valor R (1995) Sleep apnea in patients with hemispheric stroke. *Arch Phys Med Rehabil* 76(1):71–76
- Glader EL, Stegmayr B, Asplund K (2002) Poststroke fatigue: a 2-year follow-up study of stroke patients in Sweden. *Stroke* 33(5):1327–1333
- Folkman S, Lazarus RS, Gruen RJ, DeLongis A (1986) Appraisal, coping, health status, and psychological symptoms. *J Pers Soc Psychol* 50(3):571–579
- Moos R, Tsu V (1977) The Crisis of physical illness: an overview. In: Moos R (ed) *Coping with physical illness*. Springer, US, pp 3–21. doi:10.1007/978-1-4615-9089-7_1
- Donnellan C, Hevey D, Hickey A, O’Neill D (2006) Defining and quantifying coping strategies after stroke: a review. *J Neurol Neurosurg Psychiatry* 77(11):1208–1218. doi:10.1136/jnnp.2005.085670
- Shek DT (1990) Reliability and factorial structure of the Chinese version of the Beck depression inventory. *J Clin Psychol* 46(1):35–43
- Schwartz JE, Jandorf L, Krupp LB (1993) The measurement of fatigue: a new instrument. *J Psychosom Res* 37(7):753–762
- Chen SX, Yu FM (2008) Syndrome differentiation and treatment of post-stroke fatigue. *J Pract Tradit Chin Int Med* 22:17–19
- Folstein MF, Folstein SE, McHugh PR (1975) “Mini-mental state”. A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res* 12(3):189–198
- Xiao SY (1994) The theoretical basis and research applications of “Social Support Rating Scale”. *J Clin Psychiatry* 4(2):98–100
- Feifel H, Strack S, Nagy VT (1987) Coping strategies and associated features of medically ill patients. *Psychosom Med* 49(6):616–625
- Wang XD, Wang XL, Ma H (1999) Rating scales for mental health. *Chinese Mental Health J (Chin)*. 250–256. doi:10.1111/j.1365-2230.2010.03831.x
- Choi-Kwon S, Han SW, Kwon SU, Kang DW, Choi JM, Kim JS (2006) Fluoxetine treatment in poststroke depression, emotional incontinence, and anger proneness: a double-blind, placebo-controlled study. *Stroke* 37(1):156–161. doi:10.1161/01.STR.0000190892.93663.e2
- Choi-Kwon S, Han K, Choi S, Suh M, Kim YJ, Song H, Cho KH, Nah HW, Kwon SU, Kang DW, Kim JS (2012) Poststroke depression and emotional incontinence: factors related to acute and subacute stages. *Neurology* 78(15):1130–1137. doi:10.1212/WNL.0b013e31824f8090
- Chen YK, Qu JF, Xiao WM, Li WY, Weng HY, Li W, Liu YL, Luo GP, Fang XW, Ungvari GS, Xiang YT (2015) Poststroke fatigue: risk factors and its effect on functional status and health-related quality of life. *Int J Stroke* 10(4):506–512. doi:10.1111/ij.12409
- Abudi S, Bar-Tal Y, Ziv L, Fish M (1997) Parkinson’s disease symptoms—patients’ perceptions. *J Adv Nurs* 25(1):54–59
- Ikemoto S (2007) Dopamine reward circuitry: two projection systems from the ventral midbrain to the nucleus accumbens—olfactory tubercle complex. *Brain Res Rev* 56(1):27–78. doi:10.1016/j.brainresrev.2007.05.004
- Bishop GA, Ho RH (1985) The distribution and origin of serotonin immunoreactivity in the rat cerebellum. *Brain Res* 331(2):195–207
- Richard PO, Fleshner NE, Bhatt JR, Hersey KM, Chahin R, Alibhai SM (2014) Phase II, randomised, double-blind, placebo-controlled trial of methylphenidate for reduction of fatigue levels in patients with prostate cancer receiving LHRH-agonist therapy. *BJU Int*. doi:10.1111/bju.12755
- Ikemoto S, Yang C, Tan A (2015) Basal ganglia circuit loops, dopamine and motivation: a review and enquiry. *Behav Brain Res* 290:17–31. doi:10.1016/j.bbr.2015.04.018
- Lakey B, Orehek E (2011) Relational regulation theory: a new approach to explain the link between perceived social support and mental health. *Psychol Rev* 118(3):482–495. doi:10.1037/a0023477
- Hori H, Teraishi T, Ota M, Hattori K, Matsuo J, Kinoshita Y, Ishida I, Nagashima A, Koga N, Higuchi T, Kunugi H (2014) Psychological coping in depressed outpatients: association with cortisol response to the combined dexamethasone/CRH test. *J Affect Disord* 152–154:441–447. doi:10.1016/j.jad.2013.10.013