



PUBLISHED FOR SISSA BY SPRINGER

RECEIVED: May 13, 2015

REVISED: July 29, 2015

ACCEPTED: August 5, 2015

PUBLISHED: September 3, 2015

Non-perturbative test of the Witten-Veneziano formula from lattice QCD



The ETM collaboration

Krzysztof Cichy,^{a,b,c} Elena Garcia-Ramos,^{b,d} Karl Jansen,^b Konstantin Ottnad^e and Carsten Urbach^{e,f}

^aGoethe-Universität, Institut für Theoretische Physik,
Max-von-Laue-Straße 1, Frankfurt a.M., D-60438 Germany

^bNIC, DESY,
Platanenallee 6, Zeuthen, D-15738 Germany

^cAdam Mickiewicz University, Faculty of Physics,
Umultowska 85, Poznan, 61-614 Poland

^dHumboldt Universität zu Berlin,
Newtonstr. 15, Berlin, D-12489 Germany

^eInstitut für Strahlen- und Kernphysik (Theorie),
Nussallee 14-16, Bonn 53115 Germany

^fBethe Center for Theoretical Physics,
Nussallee 12, Universität Bonn, Bonn, D-53115 Germany

E-mail: krzysztof.cichy@desy.de, elenagr@ifh.de, Karl.Jansen@desy.de,
ottnad@hiskp.uni-bonn.de, urbach@hiskp.uni-bonn.de

ABSTRACT: We compute both sides of the Witten-Veneziano formula using lattice techniques. For the one side we perform dedicated quenched simulations and use the spectral projector method to determine the topological susceptibility in the pure Yang-Mills theory. The other side we determine in lattice QCD with $N_f = 2 + 1 + 1$ dynamical Wilson twisted mass fermions including for the first time also the flavour singlet decay constant. The Witten-Veneziano formula represents a leading order expression in the framework of chiral perturbation theory and we also employ leading order chiral perturbation theory to relate the flavor singlet decay constant to the relevant decay constant parameters in the quark flavor basis and flavor non-singlet decay constants. After taking the continuum and the SU(2) chiral limits we compare both sides and find good agreement within uncertainties.

KEYWORDS: Lattice QCD, Lattice Gauge Field Theories, 1/N Expansion, QCD

ARXIV EPRINT: [1504.07954](https://arxiv.org/abs/1504.07954)

Contents

1	Introduction	1
2	Lattice actions	3
2.1	$N_f = 2 + 1 + 1$ lattice QCD	3
2.2	Quenched action	4
3	Topological susceptibility in the pure Yang-Mills theory	5
3.1	The spectral projectors method	5
3.2	Computation of Z_P/Z_S	7
3.3	Continuum limit of χ_∞	8
3.4	Autocorrelations	10
4	Fermionic contributions to the Witten-Veneziano formula from dynamical simulations	12
4.1	Computations in the flavor singlet sector	12
4.2	Treatment of f_0	14
4.3	Chiral extrapolations	18
5	Discussion	21
6	Summary	23
A	Computation of κ_c	23

1 Introduction

The Witten-Veneziano formula [1, 2] was first derived by Witten for massless quarks

$$\mathring{M}_{\eta'}^2 = \frac{4N_f}{f_0^2} \chi_\infty. \quad (1.1)$$

It aims at providing an explanation for the unexpectedly large mass of the η' meson, by relating its mass in the chiral limit $\mathring{M}_{\eta'}$ to non-trivial topological fluctuations of the gauge fields, which are encoded in the topological susceptibility computed in pure Yang-Mills (YM) theory χ_∞ . Furthermore, the formula involves the singlet decay constant f_0 and the number of quark flavors N_f .

In order to obtain eq. (1.1) one considers the limit of a large number of colors N_c , for which simplifications of the theory occur that allow one to address a variety of problems which are otherwise impossible to investigate. In particular, in the 't Hooft limit ($N_c \rightarrow \infty$, while $g^2 N_c$ and N_f are kept fixed, with g denoting the gauge coupling) the anomalously

broken axial U(1) symmetry is restored and the η' becomes a Goldstone boson, i.e. $\dot{M}_{\eta'} \rightarrow 0$ [1, 3]. The formula itself is valid up to corrections of $\mathcal{O}(1/N_c^2)$ while $\dot{M}_{\eta'}^2$ and $1/f_0^2$ vanish as $\mathcal{O}(1/N_c)$. However, the topological susceptibility which appears on the right-hand side of eq. (1.1) remains finite in the large N_c limit, i.e. $\chi_\infty = \mathcal{O}(1)$, such that in the $N_c \rightarrow \infty$ limit both sides of eq. (1.1) vanish. The question remains whether $N_c = 3$ in QCD is large enough in practice to sufficiently suppress corrections to this formula to higher order in $1/N_c$.

An alternative way to derive the Witten-Veneziano formula is to expand the anomalous flavor-singlet Ward-Takahashi identities of the theory order by order in $u = N_f/N_c$ around $u = 0$ [2, 4]. Again, the Witten-Veneziano formula corresponds to the lowest order relation in this expansion. Besides, we remark that for lattice QCD it is also possible to obtain an unambiguous, theoretical sound implementation of the Witten-Veneziano formula through the study of anomalous flavor-singlet Ward-Takahashi identities in the limit $u \rightarrow 0$ [5, 6].

However, the very first attempts to compute that topological susceptibility in YM theory date back to the early 80s [7, 8] and a first reasonable number was found in [9].

Moving away from the chiral limit leads to corrections to the formula which are linear in the quadratic meson masses

$$\frac{f_0^2}{4N_f}(M_\eta^2 + M_{\eta'}^2 - 2M_K^2) = \chi_\infty. \quad (1.2)$$

This result was first derived in [2] from the aforementioned expansion in u . In the above expression terms have been reshuffled compared to eq. (1.1) to isolate the topological susceptibility in the YM theory on the r.h.s., whereas the l.h.s. of the formula contains only quantities that need to be computed in full QCD.

From a modern point of view, the Witten-Veneziano formula can also be derived from effective field theory employing a combined power counting scheme in quark masses m_q , momenta p and $1/N_c$, given by $m_q = \mathcal{O}(\delta)$, $p^2\mathcal{O}(\delta)$ and $1/N_c = \mathcal{O}(\delta)$, where δ is a small expansion parameter [10–13]. In principle, this approach allows one to systematically calculate higher order corrections to the standard form of eq. (1.2)

$$M_\eta^2 + M_{\eta'}^2 - 2M_K^2 = \frac{4N_f}{f_0^2}\chi_\infty, \quad (1.3)$$

which represents the leading order expression with respect to the expansion in δ used in chiral perturbation theory (χ PT). Note that in general $f_0 \neq f_\pi$ holds, as the singlet decay constant f_0 in eqs. (1.1), (1.2) becomes equal to the octet decay constant f_8 only when simultaneously taking the chiral limit and dropping all corrections in $1/N_c$ [11]. Since on the lattice we do not work in the octet-singlet basis, we cannot compute f_0 directly but need to apply leading order chiral perturbation theory in order to relate f_0 to the corresponding decay constant parameter in the quark flavor basis and the relevant flavor non-singlet decay constants f_π and f_K .

In this paper, we compute both the topological susceptibility in the pure YM theory (or quenched QCD) and the meson masses and the singlet decay constant f_0 in full QCD. After discussing the lattice actions, we will first discuss the determination of χ_∞ using

the so-called spectral projector method [14] based on dedicated quenched simulations.¹ Thereafter, we discuss the determination of $M_\eta, M_{\eta'}, M_K$ and in particular f_0 using $N_f = 2 + 1 + 1$ lattice QCD. Finally, we compare both results for χ_∞ , finding good agreement between quenched and dynamic computation.

2 Lattice actions

2.1 $N_f = 2 + 1 + 1$ lattice QCD

The calculation of the masses presented in this work was performed using configurations with $N_f = 2 + 1 + 1$ dynamical Wilson twisted mass fermions at maximal twist generated by the European Twisted Mass Collaboration (ETMC) [15–17].

The lattice Wilson twisted mass fermion action² for the light sector [19–21], i.e. u and d quarks, is given, in the twisted mass basis, by:

$$S_l[\psi, \bar{\psi}, U] = a^4 \sum_x \bar{\chi}_l(x) (D_W + m_0 + i\mu_l \gamma_5 \tau_3) \chi_l(x), \quad (2.1)$$

where m_0 and μ_l are the bare untwisted and twisted quark masses, respectively. $\chi_l = (\chi_u \chi_d)^T$ is a flavor doublet and τ_3 acts in flavor space. The massless Wilson Dirac operator D_W is defined as

$$D_W = \frac{1}{2} (\gamma_\mu (\nabla_\mu + \nabla_\mu^*) - a \nabla_\mu^* \nabla_\mu), \quad (2.2)$$

where ∇_μ and ∇_μ^* are the forward and backward covariant derivatives.

The lattice Wilson twisted mass action for the heavy doublet $\chi_h = (\chi_c \chi_s)^T$ [21, 22], i.e. strange and charm quarks, is given by:

$$S_h[\psi, \bar{\psi}, U] = a^4 \sum_x \bar{\chi}_h(x) (D_W + m_0 + i\mu_\sigma \gamma_5 \tau_1 + \mu_\delta \tau_3) \chi_h(x). \quad (2.3)$$

The bare twisted mass parameters μ_δ and μ_σ are related to the bare strange and charm quark masses through the following relation:

$$\mu_{c,s} = \mu_\sigma \pm \frac{Z_P}{Z_S} \mu_\delta \quad (2.4)$$

where Z_P/Z_S defines the ratio of pseudoscalar and scalar flavor non-singlet renormalization factors. Both doublets, χ_l and χ_h , are related to their counterparts in the physical basis via chiral rotations.

For the gauge sector the Iwasaki [23, 24] gauge action was used which is defined as:

$$S_G[U] = \frac{\beta}{3} \sum_x \left(b_0 \sum_{\substack{\mu, \nu=1 \\ 1 \leq \mu < \nu}}^4 \text{Re Tr} (1 - P_{x;\mu\nu}^{1 \times 1}) + b_1 \sum_{\substack{\mu, \nu=1 \\ \mu \neq \nu}}^4 \text{Re Tr} (1 - P_{x;\mu\nu}^{1 \times 2}) \right), \quad (2.5)$$

¹For early attempts to compute the topological susceptibility in YM theory we refer to [7, 8], while a first reasonable number was found in [9].

²For a review on twisted mass fermions we refer to e.g. [18].

ensemble	β	$T/a \times (L/a)^3$	$a\mu_\ell$	$a\mu_\sigma$	$a\mu_\delta$	N	N_s	N_b
A30.32	1.90	64×32^3	0.0030	0.150	0.190	1367	24	5
A40.24	1.90	48×24^3	0.0040	0.150	0.190	2630	32	10
A40.32	1.90	64×32^3	0.0040	0.150	0.190	863	24	4
A60.24	1.90	48×24^3	0.0060	0.150	0.190	1251	32	5
A80.24	1.90	48×24^3	0.0080	0.150	0.190	2449	32	10
A100.24	1.90	48×24^3	0.0100	0.150	0.190	2493	32	10
A80.24 _s	1.90	48×24^3	0.0080	0.150	0.197	2517	32	10
A100.24 _s	1.90	48×24^3	0.0100	0.150	0.197	2312	32	10
B25.32	1.95	64×32^3	0.0025	0.135	0.170	1484	24	5
B35.32	1.95	64×32^3	0.0035	0.135	0.170	1251	24	5
B55.32	1.95	64×32^3	0.0055	0.135	0.170	1545	24	5
B75.32	1.95	64×32^3	0.0075	0.135	0.170	922	24	4
B85.24	1.95	48×24^3	0.0085	0.135	0.170	573	32	2
D15.48	2.10	96×48^3	0.0015	0.120	0.1385	1045	24	10
D30.48	2.10	96×48^3	0.0030	0.120	0.1385	474	24	3
D45.32 _{sc}	2.10	64×32^3	0.0045	0.0937	0.1077	1887	24	10

Table 1. The dynamical $N_f = 2 + 1 + 1$ simulations that are included in our investigations. The notation that is used to label the ensembles is the same as in [15]. In addition to the value of β , the lattice volumes and the bare quark masses μ_ℓ , μ_σ , μ_δ we give the number of configurations N , the number of stochastic samples N_s and the bootstrap block length N_b that were used in the determination of flavor singlet quantities [26–28].

where $b_1 = -0.331$ and $b_0 = 1 - 8b_1$. The dynamical ensembles used in this work are compiled in table 1 with the labeling adopted from ref. [15]. For each β -value, there are several quark masses available, which allows one to study the chiral extrapolation. The input parameters of our simulations are supplemented by table 2, which contains the chirally extrapolated values of the Sommer parameter r_0/a for each value of β , which we use to set the scale in our dynamical simulations. In addition, we quote in this table the chirally extrapolated values of the renormalization constant ratio $Z \equiv Z_P/Z_S$ from the two definitions discussed in ref. [25]. These two definitions differ by lattice artefacts and are, therefore, helpful in understanding the corresponding systematic effects.

2.2 Quenched action

For the determination of χ_∞ in the pure YM theory, we have generated four quenched ensembles at four different values of the lattice spacing. For the gauge action, we used again the Iwasaki action — eq. (2.5).

We emphasize that, in order to test the Witten-Veneziano formula, we tried to achieve a very similar setup for the quenched and dynamical situation — we took the same gauge

β	r_0/a	a [fm]	Z_P/Z_S (M1)	Z_P/Z_S (M2)
1.90	5.31(8)	0.0885(36)	0.699(13)	0.651(6)
1.95	5.77(6)	0.0815(3)	0.697(7)	0.666(4)
2.10	7.60(8)	0.0619(18)	0.740(5)	0.727(3)

Table 2. Chirally extrapolated values of r_0/a and Z_P/Z_S from the two methods M1 and M2 as discussed in [25] for our dynamical simulations.

β	$T/a \times (L/a)^3$	r_0/a	a [fm]	$a\mu$	$r_0\mu$	κ_c^X
2.37	40×20^3	3.59(2)(3)	0.1393(14)	0.0087	0.0312(3)	0.158738
2.48	48×24^3	4.28(1)(5)	0.1182(14)	0.0073	0.0309(4)	0.154928
2.67	64×32^3	5.69(2)(3)	0.0879(6)	0.0055	0.0314(2)	0.150269
2.85	80×40^3	7.29(7)(1)	0.0686(7)	0.0043	0.0313(3)	0.147180

Table 3. All relevant parameters of the pure gauge ensembles for $\beta = 2.37, 2.48, 2.67$ and 2.85 . The errors of r_0/a correspond to statistical and systematic uncertainties, respectively.

action, we matched the physical volume and took a fixed value of $r_0\mu$ in the valence Dirac operator used for spectral projectors, equal to $r_0\mu$ of the dynamical simulations.³ The details of determining κ_c needed for $\mathcal{O}(a)$ improvement can be found in A. All the details of the quenched simulations are summarized in table 3.

The quenched simulations have been performed using the HMC algorithm implemented in the tmLQCD package [29]. The usage of this algorithm might introduce somewhat larger autocorrelation times compared to e.g. the heatbath with overrelaxation algorithm. However, compared to the dynamical simulations, the generation of quenched ensembles was still only a small effort.

3 Topological susceptibility in the pure Yang-Mills theory

3.1 The spectral projectors method

In order to compute the topological susceptibility (χ_{top}) in the pure YM theory (for this case, we will denote it specifically by χ_∞), we used the method of spectral projectors [14]. This method was originally applied to the pure gauge theory in ref. [30] and then to the dynamical case for twisted mass fermions in ref. [31].

We introduce the definition of the topological susceptibility in terms of the spectral projector \mathbb{R}_M and refer to the original papers for further details about the method [14, 30]:

$$\chi_{\text{top}} = \frac{Z_S^2}{Z_P^2} \frac{1}{V} \langle \text{Tr}\{\gamma_5 \mathbb{R}_M^2\} \text{Tr}\{\gamma_5 \mathbb{R}_M^2\} \rangle, \quad (3.1)$$

³We do not have available the value of the renormalization constant Z_P for the quenched simulations, so this implies only approximate matching of valence quark masses, assuming that with the same gauge action Z_P is not very different.

where the ratio of renormalization constants Z_P/Z_S differs from unity due to the use of non-chirally symmetric Wilson-type fermions. The spectral projector \mathbb{R}_M is an orthogonal projector to the subspace of fermion fields spanned by the eigenvectors of the massive Hermitian Dirac operator $D^\dagger D$ (with $D = D_W + m_0 + i\mu_l \gamma_5 \tau^3$ in our case) with eigenvalues not larger than M^2 . To achieve $\mathcal{O}(a^2)$ scaling towards the continuum limit, the renormalized value of the threshold M , denoted by M_R , has to be fixed for all ensembles. The value of M_R can be chosen arbitrarily, but it is advisable to avoid too small values (close to the renormalized quark mass) and choose $aM_R \ll 1$ to avoid enhanced cut-off effects [14].

In practice, we compute the following spectral observables:

$$\mathcal{A} = \frac{1}{N} \sum_{k=1}^N (\mathbb{R}_M^2 \eta_k, \mathbb{R}_M^2 \eta_k), \quad (3.2)$$

$$\mathcal{B} = \frac{1}{N} \sum_{k=1}^N (\mathbb{R}_M \gamma_5 \mathbb{R}_M \eta_k, \mathbb{R}_M \gamma_5 \mathbb{R}_M \eta_k), \quad (3.3)$$

$$\mathcal{C} = \frac{1}{N} \sum_{k=1}^N (\mathbb{R}_M \eta_k, \gamma_5 \mathbb{R}_M \eta_k), \quad (3.4)$$

where N is the number of stochastic sources η_k ($k = 1, \dots, N$) used for the construction of the spectral projector, solving the Dirac equation $(D^\dagger D + M^2)\psi = \eta_k$ an appropriate number of times. The above observables are directly related to the right hand side of eq. (3.1) through the following expression:

$$\langle \text{Tr}\{\gamma_5 \mathbb{R}_M^2\} \text{Tr}\{\gamma_5 \mathbb{R}_M^2\} \rangle = \langle \mathcal{C}^2 \rangle - \frac{\langle \mathcal{B} \rangle}{N} = \langle \mathcal{C}'^2 \rangle. \quad (3.5)$$

Since the use of two independent sets of sources to compute the square of \mathcal{C} would duplicate the cost of the calculation, we need to introduce the correction given by $\langle \mathcal{B} \rangle / N$ to correct for the bias introduced by the usage of the same set of stochastic sources to compute the square of \mathcal{C} .

In order to compute the ratio of renormalization constants Z_P/Z_S , we also use the spectral projector method, since the low cost of the calculation allows us to obtain reliable estimates of Z_P/Z_S not available a priori, since the quenched ensembles were generated specifically for this project. The ratio can be obtained through the following expression:

$$\frac{Z_P^2}{Z_S^2} = \frac{\langle \mathcal{A} \rangle}{\langle \mathcal{B} \rangle}. \quad (3.6)$$

This application was first proposed in ref. [14] and applied for dynamical simulations of twisted mass fermions in ref. [32]. In the latter reference, a comparison of the result given by spectral projectors and more standard methods like RI-MOM [33] and x -space [34] was done and compatible results were found. A study of discretization effects and finite volume effects was also presented.




Figure 1. Results of Z_P/Z_S as a function of $r_0 M$ for quenched ensembles. The straight lines correspond to the final result and the shaded areas to the systematic error. For further details we refer to the text.

3.2 Computation of Z_P/Z_S

As we mentioned in previous section, in order to compute the topological susceptibility using twisted mass fermions, the final value has to be renormalized using the ratio Z_P/Z_S , which is only equal to unity in a chirally symmetric theory.

In figure 1, the value of Z_P/Z_S for different values of $r_0 M$ for the four quenched ensembles listed in table 3 is shown. We follow the strategy presented in ref. [31] to extract the value of Z_P/Z_S for each ensemble, i.e. we take the central value to be the one corresponding to some large value of $r_0 M$, chosen as 1.5, in order to avoid contamination by non-perturbative effects appearing for small $r_0 M$. However, due to cut-off effects, even at large $r_0 M$, we observe some residual dependence on $r_0 M$. To quantify the systematic error coming from this dependence, we consider the spread of Z_P/Z_S in the interval $r_0 M \in [1, 2]$, i.e. this systematic uncertainty is taken as the larger of the deviations of

β	Z_P/Z_S
2.37	0.680(1)(27)
2.48	0.707(1)(19)
2.67	0.752(1)(7)
2.85	0.787(1)(3)

Table 4. Results for Z_P/Z_S values using spectral projectors for the quenched ensembles. The errors quoted correspond to, first, statistical and, second, systematic uncertainty (coming from the dependence on M).

$Z_P/Z_S(r_0M = 1.5)$ with respect to $Z_P/Z_S(r_0M = 1)$ and $Z_P/Z_S(r_0M = 2)$. As expected, this uncertainty drops down significantly with decreasing lattice spacing. Thus, we obtain the results presented in table 4. The first error corresponds to the statistical error given by the spectral projectors method, where autocorrelations were taken into account. The second error corresponds to the systematic error introduced in our calculation through the residual dependence of Z_P/Z_S on M .

3.3 Continuum limit of χ_∞

The main aim of this work is to test the Witten-Veneziano formula. To this end, we need to compute a reliable continuum limit of the topological susceptibility in the quenched case. The physical conditions, such as the volume or the action used, were matched to the situation of the dynamical simulations used to compute the masses.

Following this strategy, we generated the first ensemble using the Iwasaki gauge action at $\beta = 2.67$ with a value of r_0/a matching the one of the dynamical ensemble at $\beta = 1.95$. The other characteristics, such as the volume was matched to the ensemble B55.32 (see table 1) with a $32^3 \times 64$ lattice and the value of the valence quark mass for the spectral projector method was set to $a\mu = 0.0055$.

To obtain the continuum limit we needed to extend the simulations to different lattice spacings. Thus, we kept the physical volume constant by imposing a constant the physical extent of the lattice. In particular, if we assume $r_0 = 0.5$ fm, the physical volume was kept at $L \approx 2.8$ fm. Similarly, for the quark mass, we demanded the product $r_0\mu$ to remain invariant. As already pointed out in the previous section, we do not have any estimate of Z_P and hence we can only match the bare product $r_0\mu$, assuming that the changes in Z_P are small. All the exact values for each ensemble can be found in table 3.

As we mentioned in the introduction, the spectral projector method requires a mass input parameter M . In principle, the continuum limit result should be independent of this value, as long as it is kept fixed for all the ensembles entering the calculation. As the value of the renormalization constant Z_P needed to renormalize the input parameter M is not available, an alternative strategy was followed to match the values of M_R . We used the fact that the mode number remains constant at a fixed value of M_R and in a constant physical volume [14], i.e.

$$\frac{a_1}{a_2} = \left(\frac{\nu_1 n_2}{\nu_2 n_1} \right), \quad (3.7)$$

β	$\langle \mathcal{A} \rangle$	τ_{int}	$\langle \mathcal{B} \rangle$	τ_{int}	$\langle \mathcal{C} \rangle$	τ_{int}
2.37	79.4(2)	0.5(1)	17.51(5)	0.5(1)	0.19(19)	0.5(1)
2.48	78.7(2)	0.5(1)	22.00(8)	0.5(1)	0.29(32)	0.4(1)
2.67	78.5(3)	0.5(1)	29.4(2)	0.5(1)	-0.61(64)	0.7(2)
2.85	78.1(4)	0.5(1)	36.5(2)	0.4(1)	0.93(93)	0.8(3)

Table 5. Results of $\langle \mathcal{A} \rangle$, $\langle \mathcal{B} \rangle$ and $\langle \mathcal{C} \rangle$ their corresponding values of τ_{int} for quenched ensembles at four different lattice spacings. Errors are statistical only. τ_{int} is in the units of the taken step between measured configurations, which was increasing for increasing β (see also table 6).

where a_i is the lattice spacing and n_i the number of lattice points of the ensembles. Hence, the condition In table 5, the values of the mode number given by the spectral observable \mathcal{A} are shown.

Once we know the value of the ratio Z_P/Z_S for each β value, we are prepared to compute the topological susceptibility χ_{top} .

The continuum limit is a crucial aspect of lattice QCD. In fact, the rate at which a quantity approaches the continuum limit plays a fundamental role in lattice calculations, since it is directly related to the accuracy of the final computation of physical observables.

The twisted mass action at maximal twist guarantees the $\mathcal{O}(a)$ improvement for on shell observables [20]. The problem can arise if our observable is affected by short distance singularities, since it can spoil the, otherwise guaranteed, $\mathcal{O}(a^2)$ scaling.

Our observable, as discussed in detail in refs. [35, 36], is affected by short distance singularities. However, all terms linear in the lattice spacing that are consequently arising in the Symanzik expansion vanish at maximal twist [35]. Thus, our observable remains $\mathcal{O}(a)$ improved, guaranteeing an $\mathcal{O}(a^2)$ scaling towards the continuum limit.

Consequently, we have performed a linear extrapolation in $(a/r_0)^2$ to extract the continuum limit which leads to the following continuum result:

$$r_0^4 \chi_\infty = 0.049(6)_{\text{stat+sys}}, \quad (3.8)$$

where the error is dominated by statistical uncertainty, but takes into account also the systematic ones (combined in quadrature with the statistical errors). In figure 2, the results of the topological susceptibility for different values of the lattice spacing together with the continuum limit extrapolation are plotted. All the intermediate and final results are compiled in table 5 and table 6, respectively, as well as other relevant details.

The continuum value for the topological susceptibility can be compared to earlier results using chiral invariant lattice fermions [37], $r_0^4 \chi_\infty = 0.059(3)_{\text{stat}}$ and from spectral projector methods which avoid short distance singularities [30], $r_0^4 \chi_\infty = 0.061(6)$. We note that our result is in agreement with the approach used in [38], $r_0^4 \chi_\infty = 0.0524(7)_{\text{stat}}(6)_{\text{sys}}$, which has been obtained from a combined continuum and infinite volume limit. For the older result in [37] only statistical errors have been quoted and finite size effects could not be resolved within the statistical accuracy. However, the lattices in the present study exhibit a significantly larger volume (the physical value of L is about a factor two larger),




Figure 2. Continuum limit extrapolation of χ_∞ as a function of $(a/r_0)^2$ for our quenched ensembles.

β	N_s	N_{traj}	N_{meas}	$\langle \mathcal{C}^2 \rangle$	$r_0^4 \chi_{\text{top}}$
2.37	6	78000	769	25.9(1.3)	0.0250(14)(22)(8)
2.48	6	98000	412	32.8(2.4)	0.0295(24)(16)(8)
2.67	6	132000	157	47.5(5.1)	0.0376(45)(7)(8)
2.85	6	380000	119	60.9(7.9)	0.0485(70)(4)(18)

Table 6. Results of χ_{top} for the continuum limit in the pure gauge theory. N_s represents the number of stochastic sources whereas N_{meas} is the number of evaluated independent configurations and N_{traj} gives the corresponding number of thermalized MC trajectories. The errors quoted for χ_{top} are statistical, coming from Z_P/Z_S and from r_0/a , respectively.

while the range of lattice spacings is similar which might explain the difference in the final values for $r_0^4 \chi_\infty$. In general, older studies [39–42] tend to favor larger values for the topological susceptibility as has already been remarked in [38].

3.4 Autocorrelations

We close this section by a discussion of autocorrelation effects in the quenched simulations, which might affect significantly the errors.

The topological charge is a quantity highly affected by autocorrelations when the continuum limit is approached due to appearance of topological barriers, i.e. the transitions between topological sectors are suppressed. For this reason, particularly long Monte Carlo




Figure 3. Histograms of the observable \mathcal{C} for four quenched ensembles at $\beta = 2.37$ (a.), $\beta = 2.48$ (b.), $\beta = 2.67$ (c.) and $\beta = 2.85$ (d.).

simulations are needed in order to guarantee that all topological sectors are sampled adequately.

The topological charge is expected to follow a Gaussian distribution centered at zero in the large volume regime. Consequently, due to the fact that the stochastic observable \mathcal{C} is closely related to the topological charge Q , we expect the same behavior for the distribution of \mathcal{C} and, equally, $\langle \mathcal{C} \rangle = 0$.

In figure 3, the histograms of the stochastic observable \mathcal{C} are shown for all the quenched ensembles at four different values of the lattice spacing. In all cases we were able to construct a histogram compatible with a Gaussian distribution within errors.

Figure 4 shows the Monte Carlo history of the observable \mathcal{C} again for all the ensembles. It is clear, even from visual inspection of the plots, that autocorrelations become significant for the finest lattice spacing. Therefore, even though we generated a few times more trajectories for this ensemble than for the other ones, we only obtained an estimate of the topological susceptibility with a much higher statistical error, since we had to take a step of around 3200 trajectories between measurements to obtain τ_{int} compatible with 0.5, i.e. we had fewer than 120 independent gauge field configurations (see also table 6).




Figure 4. Monte Carlo histories of the observable \mathcal{C} for four quenched ensembles at $\beta = 2.37$ (a.), $\beta = 2.48$ (b.), $\beta = 2.67$ (c.) and $\beta = 2.85$ (d.).

4 Fermionic contributions to the Witten-Veneziano formula from dynamical simulations

Having established a reliable way to extract χ_∞ from the pure Yang-Mills theory, we still need to tackle the l.h.s. of eq. (1.2). In this section, we discuss how to compute the relevant masses and the flavor singlet decay constant parameter f_0 . The computations in the flavor singlet sector are very demanding, because of large contributions from quark disconnected diagrams. Moreover, the computation of f_0 on the lattice turns out to be technically involved, due to the fact that the axial vector matrix elements cannot be computed with sufficient signal-to-noise ratio. Therefore, we need to consider pseudoscalar matrix elements and resort to χ PT to relate them to the desired axial vector ones. Further complications arise from the fact that the interpolating operators on the lattice are defined in the so-called quark flavor basis, whereas decay constants and the corresponding matrix elements are commonly defined in the singlet-octet flavor basis. This issue can be circumvented in the framework of χ PT as well.

4.1 Computations in the flavor singlet sector

The masses for η and η' mesons are computed as described in [26, 28]. In particular, we employed the method of subtracting excited states in the quark connected part of the correlation functions [43]. Together with the application of the one-end trick for the

evaluation of quark disconnected diagrams in the light quark sector [44], this yields a substantial improvement of the resulting statistical errors [28, 45]. Here, we will only outline some basic details that are relevant for the discussion of the flavor singlet decay constant parameter f_0 in the next section. We use pseudoscalar operators in the physical basis:

$$\mathcal{P}_l^{0,\text{phys}} = \frac{1}{\sqrt{2}} \bar{\psi}_l i\gamma_5 \psi_l, \quad (4.1)$$

$$\mathcal{P}_h^{\pm,\text{phys}} = \bar{\psi}_h i\gamma_5 \frac{1 \pm \tau^3}{2} \psi_h, \quad (4.2)$$

where $\bar{\psi}_l, \psi_l$ and $\bar{\psi}_h, \psi_h$ refer to degenerate light and non-degenerate heavy quark doublets, respectively. The doublet structure and the flavor projector $(1 \pm \tau^3)/2$ is required due to the twisted mass formulation and the flavor projector in the second lines allows to consider the non-degenerate charm and strange components separately. At maximal twist the corresponding operators read

$$\mathcal{P}_l^{0,\text{phys}} \rightarrow -\frac{1}{\sqrt{2}} \bar{\chi}_l \tau^3 \chi_l \equiv \mathcal{S}_l^{3,tm}, \quad (4.3)$$

$$\mathcal{P}_h^{\pm,\text{phys}} \rightarrow \frac{1}{2} \bar{\chi}_h (-\tau^1 \pm i\gamma_5 \tau^3) \chi_h \equiv \mathcal{P}_h^{\pm,tm}, \quad (4.4)$$

with \mathcal{S} the scalar density. In the following, we will neglect the charm operator, as it does neither contribute to the η nor the η' within errors, i.e. we consider only the strange component $\mathcal{P}_h^{-,tm} \equiv \mathcal{P}_s^{tm}$. The mixing between the flavor non-singlet scalar and pseudoscalar currents in the heavy sector of the twisted basis introduces a relative factor of Z_S/Z_P under renormalization:

$$\mathcal{P}_s^{tm,r} = \frac{1}{2} Z_P \bar{\chi}_h \left(-\frac{Z_S}{Z_P} \tau^1 - i\gamma_5 \tau^3 \right) \chi_h. \quad (4.5)$$

Similarly, we apply a factor Z_S/Z_P instead of Z_S to the light operator $\mathcal{S}_l^{3,tm}$ allowing to pull out a global factor of Z_P^2 from the resulting, renormalized correlation function matrix:

$$\mathcal{C}^r(t) = Z_P^2 \tilde{\mathcal{C}}(t), \quad (4.6)$$

where

$$\tilde{\mathcal{C}}(t) = \begin{pmatrix} \langle \tilde{\mathcal{S}}_l^{3,tm}(t) \tilde{\mathcal{S}}_l^{3,tm}(0) \rangle & \langle \tilde{\mathcal{S}}_l^{3,tm}(t) \tilde{\mathcal{P}}_s^{tm}(0) \rangle \\ \langle \tilde{\mathcal{P}}_s^{tm}(t) \tilde{\mathcal{S}}_l^{3,tm}(0) \rangle & \langle \tilde{\mathcal{P}}_s^{tm}(t) \tilde{\mathcal{P}}_s^{tm}(0) \rangle \end{pmatrix} \quad (4.7)$$

is build from operators $\tilde{\mathcal{S}}_l^{3,tm}, \tilde{\mathcal{P}}_s^{tm}$ that are renormalized up to Z_P and apart from that correspond to the ones in eqs. (4.3), (4.5).

When building this matrix, we subtract the excited states in the connected contributions and apply the required factors of Z_S/Z_P , as given in table 2. We ignore the global factor Z_P^2 as it will cancel analytically in all observables. Then, we proceed by solving the generalized eigenvalue problem [46–48]:

$$\tilde{\mathcal{C}}(t) v^{(n)}(t, t_0) = \lambda^{(n)}(t, t_0) \tilde{\mathcal{C}}(t_0) v^{(n)}(t, t_0), \quad t_0 < t, \quad (4.8)$$

which gives access to masses through

$$\frac{\lambda^{(n)}(t, t_0)}{\lambda^{(n)}(t+1, t_0)} = \frac{\exp(-m^{(n)}t) + \exp(-m^{(n)}(T-t))}{\exp(-m^{(n)}(t+1)) + \exp(-m^{(n)}(T-(t+1)))} \quad (4.9)$$

and amplitudes

$$A_i^{(n)} = \frac{\sum_{j=1}^N \tilde{\mathcal{C}}_{ij}(t) v_j^{(n)}(t, t_0)}{\sqrt{(v^{(n)}(t, t_0), \mathcal{C}(t) v^{(n)}(t, t_0)) (\exp(-m^{(n)}t) \pm \exp(m^{(n)}(t-T)))}} \quad (4.10)$$

respectively. Since masses are renormalization group invariants, they are not affected by the actual choice of Z_S/Z_P , which is only relevant for the computation of amplitudes leading to different lattice artifacts for the methods M1 and M2. Note that the amplitudes computed in this way are only renormalized up to a factor of Z_P as well.

In table 1, we give the number of configurations N used on each gauge ensemble to extract observables in the flavor singlet sector. Errors are computed using bootstrapping with 1000 samples and we have used blocking to deal with autocorrelations. The numbers of configurations per block N_b are given in table 1 and have been chosen to correspond to a block length of at least 20 HMC trajectories. In addition, we give the number of stochastic samples that has been employed in the computation of quark disconnected diagrams. It has been chosen such that the resulting statistical errors are dominated by gauge noise.

The results for all ensembles are given in table 7. Clearly, the statistical error for a test of the Witten-Veneziano formula will be dominated by flavor singlet related quantities as can be inferred from the relative errors on the masses. This is caused by large contributions from quark disconnected diagrams in the flavor singlet sector which are intrinsically more noisy compared to connected contributions and also introduce sizable autocorrelations in case of the η' mass.

For the computation of the kaon masses, we refer to refs. [15, 26]. The values are also listed in table 7, together with the corresponding charged pion masses M_{PS} . Note that some of these values were calculated at smaller statistics than the flavor singlet sector (cf. table 1). Nevertheless, their relative statistical error is still at least one order of magnitude smaller than for quantities in the flavor singlet sector.

4.2 Treatment of f_0

Consider the general definition for the decay constant f_P ,⁴ of any pseudoscalar meson P :

$$\langle 0 | A_\mu^a(0) | P(p) \rangle = i f_P^a p_\mu, \quad (4.11)$$

where $A_\mu^a(0)$ denotes the axial vector current with flavor structure denoted by the index a .

In the charged sector for mass-degenerate, light quarks, the axial vector current in the physical basis transforms into the vector current in the twisted basis at maximal twist.

⁴The normalization of decay constants in this work has been chosen consistently s.t. $f_{PS} \approx 130 \text{ MeV}$ holds for the decay constant of the charged pion.

ensemble	aM_{PS}	aM_K	aM_η	$aM_{\eta'}$
A30.32	0.12358(30)	0.25150(29)	0.2800(55)	0.480(17)
A40.24	0.14484(44)	0.25884(43)	0.2834(35)	0.427(14)
A40.32	0.14140(30)	0.25666(23)	0.2809(28)	0.458(22)
A60.24	0.17277(48)	0.26695(52)	0.2908(46)	0.471(16)
A80.24	0.19870(35)	0.27706(61)	0.3004(17)	0.479(23)
A100.24	0.22127(32)	0.28807(34)	0.3063(24)	0.454(15)
A80.24s	0.19822(33)	0.25503(33)	0.2686(30)	0.463(14)
A100.24s	0.22118(33)	0.26490(74)	0.2763(13)	0.518(30)
B25.32	0.10685(43)	0.21240(50)	0.2348(41)	0.414(19)
B35.32	0.12496(45)	0.21840(28)	0.2363(23)	0.435(24)
B55.32	0.15396(31)	0.22799(34)	0.2469(28)	0.480(26)
B75.32	0.18036(39)	0.23753(32)	0.2537(23)	0.430(23)
B85.24	0.19373(64)	0.24392(59)	0.2640(28)	0.453(28)
D15.48	0.06954(26)	0.16897(85)	0.1891(64)	0.295(21)
D30.48	0.09801(25)	0.17760(23)	0.1969(70)	0.288(21)
D45.32 _{sc}	0.11991(37)	0.17570(84)	0.1898(22)	0.276(15)

Table 7. Results for meson masses on all dynamical ensembles [15, 26, 28]. In addition to the relevant masses for the Witten-Veneziano formula in eq. (1.2), we also give the corresponding values of the charged pion mass M_{PS} .

This feature can be exploited together with the PCVC relation to derive an expression for the charged pion decay constant f_{PS} [49–51]:

$$f_{\text{PS}} = 2\mu_l \frac{\langle 0 | P_l^a | \pi^\pm \rangle}{M_{\text{PS}}^2}, \quad (4.12)$$

where $a = 1, 2$. Since this formula depends on a pseudoscalar matrix element, it can be calculated with very high statistical accuracy. Moreover, this relation allows one to compute f_{PS} in the twisted mass formulation without the need for any renormalization factor at all, because a factor Z_{P}^{-1} for the bare quark mass μ_l cancels the factor for the pseudoscalar matrix element in the twisted basis.

Similar considerations apply for the kaon sector, although a complication arises from the fact that one has to employ interpolating operators made of light and heavy quarks. The relation for the kaon decay constant f_K in the twisted mass formulation is given by:

$$f_K = (\mu_l + \mu_s) \frac{\langle 0 | \tilde{\mathcal{P}}_{\text{neutral}}^{+,tm} | K \rangle}{M_K^2}, \quad (4.13)$$

where $\tilde{\mathcal{P}}_{\text{neutral}}^{+,tm} = \frac{1}{2}(\frac{Z_{\text{S}}}{Z_{\text{P}}}(-\bar{\chi}_d \chi_c + \bar{\chi}_d \chi_s) + \bar{\chi}_d i \gamma_5 \chi_c + \bar{\chi}_d i \gamma_5 \chi_s)$. In this case the factor $Z_{\text{S}}/Z_{\text{P}}$ is required again, because it enters μ_s (cf. eq. (2.4)) as well as relative factors due to mixing between scalar and pseudoscalar currents.

Assuming exact isospin symmetry and neglecting possible contributions from the charm quark and mixing with further states such as glueballs, the most general parametrization for decay constants of the η, η' system reads

$$\begin{pmatrix} f_\eta^8 & f_\eta^0 \\ f_{\eta'}^8 & f_{\eta'}^0 \end{pmatrix} = \begin{pmatrix} f_8 \cos \phi_8 & -f_0 \sin \phi_0 \\ f_8 \sin \phi_8 & f_0 \cos \phi_0 \end{pmatrix} \equiv \Xi(\phi_8, \phi_0) \text{diag}(f_8, f_0) . \quad (4.14)$$

The choice of A_μ^0, A_μ^8 together with $P = \eta, \eta'$ in eq. (4.11) defines the so-called *singlet-octet basis*.

Moreover, employing χ PT, it is possible to relate the decay constant parameters in the η - η' system to the remaining octet decay constants f_π, f_K and a low energy constant $\Lambda_1 = \mathcal{O}(1/N_c)$ occurring at next-to-leading order in the chiral expansion [11, 12, 52] by:

$$f_0^2 = (f_\eta^0)^2 + (f_{\eta'}^0)^2 = \frac{1}{3} (2f_K^2 + f_\pi^2) + \Lambda_1 f_\pi^2, \quad (4.15)$$

$$f_8^2 = (f_\eta^8)^2 + (f_{\eta'}^8)^2 = \frac{1}{3} (4f_K^2 - f_\pi^2), \quad (4.16)$$

$$f_0 f_8 \sin(\phi_8 - \phi_0) = f_\eta^0 f_\eta^8 + f_{\eta'}^0 f_{\eta'}^8 = -\frac{2\sqrt{2}}{3} (f_K^2 - f_\pi^2). \quad (4.17)$$

In the chosen basis and to the given chiral order, the additional, OZI-violating corrections are specific to the singlet sector, i.e. the term $\sim \Lambda_1 = \mathcal{O}(N_c^{-1})$ affects neither f_8 , nor any of the angles ϕ_0, ϕ_8 , but only the parameter f_0 . From the last relation, it can be inferred that in the octet singlet basis, the difference between the two angles ϕ_0, ϕ is given by $SU(3)_F$ -violating effects, leading to the expectation

$$\left| \frac{\phi_0 - \phi_8}{\phi_0 + \phi_8} \right| \ll 1. \quad (4.18)$$

The fact that there are different types of contributions to the mixing, i.e. $SU(3)_F$ -breaking and OZI-violating effects $\sim \Lambda_1$, can be exploited in order to choose a basis in which the two resulting mixing angles do not exhibit a sizable splitting. To this end, one introduces the *quark flavor basis* with the axial vector currents A_μ^0 and A_μ^8 replaced by the combinations

$$A_\mu^l = \frac{2}{\sqrt{3}} A_\mu^0 + \sqrt{\frac{2}{3}} A_\mu^8 = \frac{1}{\sqrt{2}} (\bar{u} \gamma_\mu \gamma_5 u + \bar{d} \gamma_\mu \gamma_5 d), \quad (4.19)$$

$$A_\mu^s = \sqrt{\frac{2}{3}} A_\mu^0 - \frac{2}{\sqrt{3}} A_\mu^8 = \bar{s} \gamma_\mu \gamma_5 s, \quad (4.20)$$

in which the light quarks and the strange quark contributions are disentangled. This is the reason why this basis is more convenient in lattice simulations, as this is also the preferred flavor structure for interpolating operators on the lattice, allowing one to directly access the corresponding matrix elements. In exact analogy to the singlet-octet basis, this basis again allows for a parametrization in terms of two decay constants and two mixing angles

$$\begin{pmatrix} f_\eta^l & f_\eta^s \\ f_{\eta'}^l & f_{\eta'}^s \end{pmatrix} = \Xi(\phi_l, \phi_s) \text{diag}(f_l, f_s), \quad (4.21)$$

where the mixing matrix Ξ has the same form as the one defined in eq. (4.14). In this basis, the relations between mixing parameters in the η - η' system and f_π , f_K , Λ_1 read:

$$f_l^2 = (f_\eta^l)^2 + (f_{\eta'}^l)^2 = f_\pi^2 + \frac{2}{3}\Lambda_1 f_\pi^2, \quad (4.22)$$

$$f_s^2 = (f_\eta^s)^2 + (f_{\eta'}^s)^2 = 2f_K^2 - f_\pi^2 + \frac{1}{3}\Lambda_1 f_\pi^2, \quad (4.23)$$

$$f_l f_s \sin(\phi_l - \phi_s) = f_\eta^l f_\eta^s + f_{\eta'}^l f_{\eta'}^s = \frac{\sqrt{2}}{3}\Lambda_1 f_\pi^2. \quad (4.24)$$

The most important feature of the quark flavor basis becomes manifest in the last expression, which is now entirely given by an OZI-violating contribution $\sim \Lambda_1 = \mathcal{O}(\delta)$, amounting to additional suppression for the difference $|\phi_l - \phi_s|$ compared to $|\phi_0 - \phi_8|$, which is given by $SU(3)_F$ -breaking effects. Besides, in the $SU(3)_F$ symmetric case, the angles $\phi_l \approx \phi_s$ take the value $\phi_{SU(3)_F} = \arctan \sqrt{2}$, and, hence, their numerical value is not expected to be small. Therefore, one expects:

$$\left| \frac{\phi_l - \phi_s}{\phi_l + \phi_s} \right| \ll 1 \quad (4.25)$$

in the quark flavor basis, which has been numerically confirmed in a previous lattice study [28]. This feature allows one to consider a simplified mixing scheme in the quark flavor basis with only one angle ϕ

$$\begin{pmatrix} f_\eta^l & f_\eta^s \\ f_{\eta'}^l & f_{\eta'}^s \end{pmatrix} = \Xi(\phi) \text{diag}(f_l, f_s) + \mathcal{O}(\Lambda_1), \quad (4.26)$$

where $\Xi(\phi) \equiv \Xi(\phi, \phi)$.

As mentioned at the beginning of this section, we are restricted in our simulations to pseudoscalar operators for practical purposes, because the signal-to-noise ratio for axial vector operators turns out too small for a direct computation of the relevant observables. However, one may instead consider pseudoscalar matrix elements in order to retrieve information on the mixing parameters. This is possible due to the relation between axial vector and pseudoscalar matrix elements, which is given non-perturbatively by

$$\partial^\mu A_\mu^a = \bar{\psi}(x) 2MT^a i\gamma_5 \psi(x) + \delta^{0a} \sqrt{2N_f} \omega(x), \quad (4.27)$$

where for $a = 0$ we have $T^0 = \sqrt{1/(2N_f)} \mathbb{1}_{N_f \times N_f}$ and $\omega(x)$ denotes the winding number density.

Nonetheless, the anomaly equation of QCD itself is not sufficient for any practical purposes here, mainly because it requires knowledge of an additional matrix element involving the topological charge density. Therefore, one needs to gain further insight on how the pseudoscalar matrix elements are linked to the mixing parameters. This is again achieved by the use of χ PT. Consider pseudoscalar currents in the quark flavor basis in analogy to eqs. (4.19), (4.20):

$$P^l = \frac{1}{\sqrt{2}} (\bar{u} i\gamma_5 u + \bar{d} i\gamma_5 d), \quad (4.28)$$

$$P^s = \bar{s} i\gamma_5 s, \quad (4.29)$$

and the corresponding matrix elements for pseudoscalar mesons P that are given by

$$h_P^i = 2m_i \langle 0 | P^i | P \rangle . \quad (4.30)$$

To leading order, one can make contact with the quark flavor basis parametrization for axial vector matrix elements in eq. (4.21), i.e. obtain an expression for h_P^i in terms of decay constants f_l , f_s , the mixing angle ϕ and octet meson masses [12]:

$$\begin{pmatrix} h_\eta^l & h_\eta^s \\ h_{\eta'}^l & h_{\eta'}^s \end{pmatrix} = \Xi(\phi) \text{diag} (M_\pi^2 f_l, (2M_K^2 - M_\pi^2) f_s) . \quad (4.31)$$

Again, formally higher order, OZI-violating contributions are neglected in this expression, as demanded by the so-called FKS-scheme, which allows for the determination of process-independent mixing parameters [12, 52, 53].

Finally, we can combine eqs. (4.15), (4.16) and eqs. (4.22), (4.23) to write down leading order relations for the desired singlet decay constant f_0 in terms of the parameters f_{PS} , f_K , f_l and f_s , which we compute on the lattice:

$$f_0^2 = -7/6 f_{PS}^2 + 2/3 f_K^2 + 3/2 f_l^2 , \quad (4.32)$$

$$f_0^2 = +1/3 f_{PS}^2 - 4/3 f_K^2 + f_l^2 + f_s^2 , \quad (4.33)$$

$$f_0^2 = +8/3 f_{PS}^2 - 16/3 f_K^2 + 3 f_s^2 . \quad (4.34)$$

In general, these relations receive corrections of $\mathcal{O}(\delta^2)$. However, since they were derived from continuum χ PT, they also differ by lattice artifacts of $\mathcal{O}(a^2)$, if applied to our lattice data. We will in the following exploit this ambiguity to choose a definition for f_0 that exhibits particularly small systematic effects. Although the use of pseudoscalar matrix elements and changing the flavor basis requires to resort to χ PT, we point out that this is not a serious drawback, as the above expressions are in general of the same order in the chiral expansion as the Witten-Veneziano formula in eq. (1.2). In the following, we will refer to the three definitions eqs. (4.32)–(4.34) of f_0 as D1, D2 and D3, respectively.

Regarding the computation of decay constants in the charged meson sector, we point out that while the values of f_{PS} have been recalculated with the current statistics (they were first published in [15]), we used less configurations for f_K . However, by far the largest contribution to the overall statistical error in our analysis stems again from the flavor singlet sector. Finally, we remark that a dedicated study of η, η' -related decay constant parameters is currently in preparation [54].

4.3 Chiral extrapolations

Since our lattice simulations employ unphysical quark masses, we need to perform chiral extrapolations of our lattice data when computing the l.h.s. of eq. (1.2). In principle, to the given order in χ PT, this simply corresponds to a constant fit in $(r_0 M_{PS})^2$. However, we have to take into account lattice artifacts which might be different for the three definitions eqs. (4.32)–(4.34).

method	M1D1	M1D2	M1D3	M2D1	M2D2	M2D3
$r_0^4\chi_\infty$	0.019(1)	0.046(3)	0.058(3)	0.053(3)	0.060(3)	0.036(4)
χ^2/dof	4.90	1.71	2.60	2.19	2.04	1.95

Table 8. Results from a constant fit for $r_0^4\chi_\infty$ and corresponding χ^2/dof values ($\text{dof} = 15$) for two different sets of values of Z_P/Z_S (M1,M2) and three definitions for f_0 (D1,D2,D3); cf. eqs. (4.32)–(4.34). Errors are statistical only.

Moreover, the dependence on the choice of Z_P/Z_S is non-trivial, as it affects the relevant decay constants (besides f_{PS}) directly through the renormalization of the corresponding matrix elements, as well as through the relative renormalization factor that enters the quark mass μ_s in eq. (2.4), which appears in the definition of f_K in eq. (4.13).

In figure 5 we show results for $r_0^4\chi_\infty$ calculated from using our results for the relevant meson masses as given in table 7. For the plots (a.), (c.) and (e.) in the left column we used Z_P/Z_S values from method M1 and for the right ones (b.), (d.) and (f.) those from M2; cf. table 2. The three rows in figure 5 correspond to the three definitions of f_0 D1, D2 and D3, respectively.

Clearly, the three definitions of f_0 show different lattice artifacts and systematic effects regarding their Z_P/Z_S (and hence m_s) dependence. Besides these systematic effects, their relative statistical errors differ as well. Within errors, the first definition D1 shows the largest lattice artifacts, as well as the most significant dependence on the choice of Z_P/Z_S . Applying a constant fit in $(r_0M_{PS})^2$ does not provide a good description of the data in this case, as can be seen from panel (a.). The fitted value is much lower than for any other definition and most points are incompatible with the fitted line. However, the data points show a trend towards larger values at smaller lattice spacings, i.e. by fitting only to the data at the finest lattice spacing value we obtain $r_0^4\chi_\infty^D = 0.043(12)_{\text{stat}}$, significantly higher than the result of the fit to all data points.

The data points extracted from definitions D2 and D3 lead to a reasonable agreement with a constant extrapolation in $(r_0M_{PS})^2$, with clearly the best fits stemming from definition D3. In particular, with definition D3, the extrapolated value is merely independent on the choice of Z .

We have also tried to add a term of $\mathcal{O}(a^2)$ to our fit. It turns out that apart from the most extreme case shown in panel (a.) of figure 5, the data does not allow to resolve such a dependence within errors, i.e. the results for the corresponding coefficient are compatible with zero. Regarding the dependence on the strange quark mass, we point out that to the given chiral order, the computed χ_∞ is expected to be a constant function of $(r_0M_K)^2$ as well. Although the data shows a residual dependence on the strange quark mass, no clear picture arises with respect to its functional form.

We have collected numerical results from our constant fits for χ_∞ in table 8 for the three definitions of f_0 and the two sets of values for Z_P/Z_S . In addition, we give the respective χ^2/dof values. Clearly, the fit M1D1 is by far worst, which is expected due to the large cutoff effects in this case.




Figure 5. Results for $r_0^4 \chi_\infty$ calculated from meson masses and f_0 as a function of $(r_0 M_{\text{PS}})^2$. The values of f_0 employed in panels (a.), (b.) are obtained from eq. (4.32), for (c.), (d.) from eq. (4.33) and for (e.), (f.) from eq. (4.34). The plots in the left and right column correspond to different Z_P/Z_S values from M1 and M2, respectively, as listed in table 2. The chirally extrapolated value is obtained from a constant fit in $(r_0 M_{\text{PS}})^2$ and has to be compared with the value computed in the pure Yang-Mills theory.

In order to obtain our final result for χ_∞ from the dynamic simulations, we apply a weight to each fit:

$$w = 1 - 2|p - 0.5|, \quad (4.35)$$

where p denotes the p -values corresponding to the values of χ^2/dof given in table 8 and take the average over all six fits, leading to:

$$r_0^4 \chi_\infty = 0.047(3)_{\text{stat}}(11)_{\text{sys}}.$$

The systematic error has been chosen as the mean absolute deviation from the central value and should reflect the uncertainties from residual cutoff and strange quark mass effects. Since we included the fit M1D1, which suffers from particularly large lattice artifacts, the value of the systematic error should be considered a conservative estimate.

Another possibility to deal with the residual effects of quark mass dependence and the lattice spacing is to include additional, higher order terms in the fit function

$$f(r_0^4 \chi_\infty, (r_0 M_\pi)^2, (r_0 M_K)^2, (a/r_0)^2) = r_0^4 \chi_\infty + c_1 (r_0 M_\pi)^2 + c_2 (r_0 M_K)^2 + c_3 (a/r_0)^2, \quad (4.36)$$

where c_i with $i = 1, 2, 3$ denotes the free fit parameters. Note that from the point of view of chiral perturbation theory including only a subset of the linear terms would be inconsistent with power counting. As can be inferred from table 9 this fit model leads to improved χ^2/dof values. However, most of the additional terms are poorly constrained by the data and are close or compatible with zero. In general, this leads to much larger statistical errors and all results for $r_0^4 \chi_\infty$ are compatible within errors. Taking the average of all six fits weighted by their respective p -value and statistical errors yields

$$r_0^4 \chi_\infty = 0.051(24)_{\text{stat}},$$

which agrees with the previously computed result from constant fits. Since for the fit model in E.q. (4.36) all fit results for $r_0^4 \chi_\infty$ are compatible with the averaged result within statistical errors, we refrain from quoting an additional systematic uncertainty, as it has been done for the constant fits.

We remark that the inclusion of even more complicated terms seems not feasible, as the linear terms in the above model are already rather poorly constrained by the data and close or even compatible with zero in many cases. Regarding the data point M1D1 it appears that the introduction of additional terms does still not lead to a good χ^2/dof value for the corresponding fit, although the resulting value for $r_0^4 \chi_\infty$ agrees well with the other results. Therefore, we conclude that this data point is rather to be considered a statistical outlier than revealing any actual effect related to physics.

5 Discussion

In figure 6, we compare the pure YM topological susceptibility to the left hand side of eq. (1.2) computed in $N_f = 2 + 1 + 1$ lattice QCD. All results are given in units of the Sommer parameter r_0 . The weighted mean of the dynamical results and its statistical error




Figure 6. Results from the pure Yang-Mills theory (YM) and dynamical simulations. Open and closed symbols correspond to the two sets of values for Z_P/Z_S (M1, M2). All three definitions D1, D2 and D3 for f_0 are included and represented by circle, triangle and diamond symbols, respectively. The solid black line represent the final, p -value weighted average from dynamical simulations and its statistical error is indicated by the gray band. In addition, the dotted lines correspond to its systematic error; see text.

method	M1D1	M1D2	M1D3	M2D1	M2D2	M2D3
$r_0^4 \chi_\infty$	0.062(12)	0.066(20)	0.039(31)	0.049(21)	0.061(26)	0.030(38)
χ^2/dof	3.45	1.50	1.34	1.91	1.29	1.23

Table 9. Results for $r_0^4 \chi_\infty$ from the linear fit model in eq. (4.36) and the corresponding χ^2/dof values ($\text{dof} = 12$) for two different sets of values of Z_P/Z_S (M1,M2) and three definitions for f_0 (D1,D2,D3); cf. eqs. (4.32)–(4.34). Errors are statistical only.

is indicated by the gray band. The systematic uncertainty of the weighted average is shown by the dotted horizontal lines. Apart from the outlier M1D1, which is affected by sizable cutoff effects, the results from the dynamical simulations are very close to the one from the pure YM theory. Also the agreement of the quenched and the averaged dynamical result appears to be good within statistical and systematic uncertainties.

A complication arises when one attempts to convert the results to physical units. It stems from the value of r_0 in the YM theory: while in $N_f = 2 + 1 + 1$ QCD direct contact to physical quantities is natural, in the YM theory such a relation is not obvious. In particular, the question arises whether or not the values of r_0 in the YM theory and in $N_f = 2 + 1 + 1$ QCD are expected to be equal. From a general point of view, we do not see a reason for this to be true.

Therefore, one may use the standard value of $r_0 = 0.5$ fm to convert the YM result to physical units and obtain:

$$\chi_\infty^{\text{YM}} = (185.3(5.6)_{\text{stat+sys}} \text{ MeV})^4.$$

For the dynamical simulations, on the other hand, we can use the value $r_0 = 0.474(14)$ fm computed in ref. [25] to convert to physical units. Taking the aforementioned weighted average, we obtain:

$$\chi_\infty^{\text{dyn}} = (193.5(6.2)_{\text{stat}}(13.4)_{\text{sys}} \text{ MeV})^4,$$

where we have included the error on the physical value of r_0 in the statistical uncertainty. We find rather close agreement between quenched (χ_∞^{YM}) and dynamical (χ_∞^{dyn}) results with deviations of only up to $\mathcal{O}(10\%)$. This confirms the validity of the Witten-Veneziano formula to the given order and for the assumed value of the quenched r_0 also in physical units.

6 Summary

In this paper, we have presented a non-perturbative test of the famous Witten-Veneziano formula. This formula relates the large mass value obtained for the η' meson to the anomalously broken axial U(1) symmetry in QCD. It, therefore, provides important insights for our understanding of QCD and the generation of masses.

We have computed the topological susceptibility in the pure YM theory with dedicated quenched lattice simulations using the so-called spectral projector method. In particular, by using four values of the lattice spacing, we were able to perform a reliable continuum extrapolation and, hence, control this major systematic uncertainty.

For the first time, we have computed the flavour singlet decay constant f_0 using $N_f = 2 + 1 + 1$ lattice QCD. Together with the η , η' and kaon meson masses determined in ref. [28], this allowed us to compute also the l.h.s. of the Witten-Veneziano formula. Again, lattice artifacts are controlled by using three values of the lattice spacing. By using a wide range of light quark mass values, also the SU(2) chiral extrapolation was performed in a controlled way. The strange quark mass dependence was found to be not important for the Witten-Veneziano formula.

The comparison of the pure YM topological susceptibility and its Witten-Veneziano counterpart from full QCD leads to agreement within errors. This finding provides clear evidence for the hypothesis that the large mass of the η' meson is due the anomalously broken axial U(1) symmetry in QCD.

A Computation of κ_c

In order to guarantee the $\mathcal{O}(a)$ improvement, we need to tune κ to its critical value for the valence quarks. To do so, we followed the strategy introduced in [55]. Thus, we computed the of κ through the evaluation of m_{pcac} for different values of the quark mass $a\mu$. In particular, we imposed $m_{\text{pcac}} < 0.1a\mu$. In figure 7, we show a particular example of the




Figure 7. Chiral extrapolation of κ_c for the quenched ensemble at $\beta = 2.48$. The red point corresponds to the linearly extrapolated value.

chiral behavior of κ_c . In all cases, we perform a chiral fit considering only the lowest masses $a\mu < 0.01$ since the larger masses deviate from the linear behavior. Notice that the data is highly correlated and the fit needed to take this correlation into account.

Acknowledgments

We thank all members of ETMC for the most enjoyable collaboration. We are grateful to R. Frezzotti, U.-G. Meißner and G.C. Rossi for helpful comments and discussion. Furthermore, we would like to thank F. Zimmermann for contributions at an early stage of this project. The computer time for this project was made available to us by the John von Neumann-Institute for Computing (NIC) on the JUDGE and Jugene systems in Jülich and the IDRIS (CNRS) computing center in Orsay. In particular we thank U.-G. Meißner for granting us access on JUDGE. Further computational resources were provided by SuperMUC at LRZ in Garching and the PC cluster in Zeuthen. This project was funded by the DFG as a project in the SFB/TR 16. K.C. has been supported in part by Foundation for Polish Science fellowship “Kolumb” and by the Helmholtz International Center for FAIR within the framework of the LOEWE program launched by the State of Hessen. Two of the authors (K.O. and C.U.) were supported by the Bonn-Cologne Graduate School (BCGS) of Physics and Astronomie. The open source software packages tmLQCD [29], Lemon [56] and R [57] have been used.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] E. Witten, *Current Algebra Theorems for the $U(1)$ Goldstone Boson*, *Nucl. Phys. B* **156** (1979) 269 [[INSPIRE](#)].
- [2] G. Veneziano, *$U(1)$ Without Instantons*, *Nucl. Phys. B* **159** (1979) 213 [[INSPIRE](#)].
- [3] G. 't Hooft, *A Planar Diagram Theory for Strong Interactions*, *Nucl. Phys. B* **72** (1974) 461 [[INSPIRE](#)].
- [4] G. Veneziano, *Goldstone Mechanism From Gluon Dynamics*, *Phys. Lett. B* **95** (1980) 90 [[INSPIRE](#)].
- [5] L. Giusti, G.C. Rossi, M. Testa and G. Veneziano, *The $U_A(1)$ problem on the lattice with Ginsparg-Wilson fermions*, *Nucl. Phys. B* **628** (2002) 234 [[hep-lat/0108009](#)] [[INSPIRE](#)].
- [6] L. Giusti, G.C. Rossi, M. Testa and G. Veneziano, *The $U_A(1)$ problem on the lattice*, *Nucl. Phys. Proc. Suppl.* **106** (2002) 1001 [[hep-lat/0110036](#)] [[INSPIRE](#)].
- [7] P. Di Vecchia, K. Fabricius, G.C. Rossi and G. Veneziano, *Preliminary Evidence for $U_A(1)$ Breaking in QCD from Lattice Calculations*, *Nucl. Phys. B* **192** (1981) 392 [[INSPIRE](#)].
- [8] P. Di Vecchia, K. Fabricius, G.C. Rossi and G. Veneziano, *Numerical Checks of the Lattice Definition Independence of Topological Charge Fluctuations*, *Phys. Lett. B* **108** (1982) 323 [[INSPIRE](#)].
- [9] A. Di Giacomo, H. Panagopoulos and E. Vicari, *The Topological Susceptibility and Lattice Universality*, *Nucl. Phys. B* **338** (1990) 294 [[INSPIRE](#)].
- [10] J. Gasser and H. Leutwyler, *Chiral Perturbation Theory: Expansions in the Mass of the Strange Quark*, *Nucl. Phys. B* **250** (1985) 465 [[INSPIRE](#)].
- [11] R. Kaiser and H. Leutwyler, *Pseudoscalar decay constants at large N_c* , [hep-ph/9806336](#) [[INSPIRE](#)].
- [12] T. Feldmann, *Quark structure of pseudoscalar mesons*, *Int. J. Mod. Phys. A* **15** (2000) 159 [[hep-ph/9907491](#)] [[INSPIRE](#)].
- [13] R. Kaiser and H. Leutwyler, *Large- N_c in chiral perturbation theory*, *Eur. Phys. J. C* **17** (2000) 623 [[hep-ph/0007101](#)] [[INSPIRE](#)].
- [14] L. Giusti and M. Lüscher, *Chiral symmetry breaking and the Banks-Casher relation in lattice QCD with Wilson quarks*, *JHEP* **03** (2009) 013 [[arXiv:0812.3638](#)] [[INSPIRE](#)].
- [15] R. Baron et al., *Light hadrons from lattice QCD with light (u, d), strange and charm dynamical quarks*, *JHEP* **06** (2010) 111 [[arXiv:1004.5284](#)] [[INSPIRE](#)].
- [16] EUROPEAN TWISTED MASS collaboration, R. Baron et al., *Computing K and D meson masses with $N_f = 2 + 1 + 1$ twisted mass lattice QCD*, *Comput. Phys. Commun.* **182** (2011) 299 [[arXiv:1005.2042](#)] [[INSPIRE](#)].
- [17] ETM collaboration, R. Baron et al., *Light Hadrons from $N_f = 2 + 1 + 1$ Dynamical Twisted Mass Fermions*, *PoS(LATTICE 2010)* 123 [[INSPIRE](#)].

- [18] A. Shindler, *Twisted mass lattice QCD*, *Phys. Rept.* **461** (2008) 37 [[arXiv:0707.4093](#)] [[INSPIRE](#)].
- [19] ALPHA collaboration, R. Frezzotti, P.A. Grassi, S. Sint and P. Weisz, *Lattice QCD with a chirally twisted mass term*, *JHEP* **08** (2001) 058 [[hep-lat/0101001](#)] [[INSPIRE](#)].
- [20] R. Frezzotti and G.C. Rossi, *Chirally improving Wilson fermions. 1. $O(a)$ improvement*, *JHEP* **08** (2004) 007 [[hep-lat/0306014](#)] [[INSPIRE](#)].
- [21] R. Frezzotti and G.C. Rossi, *Chirally improving Wilson fermions. II. Four-quark operators*, *JHEP* **10** (2004) 070 [[hep-lat/0407002](#)] [[INSPIRE](#)].
- [22] R. Frezzotti and G.C. Rossi, *Twisted mass lattice QCD with mass nondegenerate quarks*, *Nucl. Phys. Proc. Suppl.* **128** (2004) 193 [[hep-lat/0311008](#)] [[INSPIRE](#)].
- [23] Y. Iwasaki, *Renormalization Group Analysis of Lattice Theories and Improved Lattice Action: Two-Dimensional Nonlinear $O(N)$ σ -model*, *Nucl. Phys. B* **258** (1985) 141 [[INSPIRE](#)].
- [24] Y. Iwasaki, K. Kanaya, T. Kaneko and T. Yoshie, *Scaling in $SU(3)$ pure gauge theory with a renormalization group improved action*, *Phys. Rev. D* **56** (1997) 151 [[hep-lat/9610023](#)] [[INSPIRE](#)].
- [25] EUROPEAN TWISTED MASS collaboration, N. Carrasco et al., *Up, down, strange and charm quark masses with $N_f = 2 + 1 + 1$ twisted mass lattice QCD*, *Nucl. Phys. B* **887** (2014) 19 [[arXiv:1403.4504](#)] [[INSPIRE](#)].
- [26] ETM collaboration, K. Ottnad et al., *η and η' mesons from $N_f = 2 + 1 + 1$ twisted mass lattice QCD*, *JHEP* **11** (2012) 048 [[arXiv:1206.6719](#)] [[INSPIRE](#)].
- [27] EUROPEAN TWISTED MASS collaboration, C. Michael, K. Ottnad and C. Urbach, *η and η' masses and decay constants from lattice QCD with $N_f = 2 + 1 + 1$ quark flavours*, *PoS(LATTICE 2013)253* [[arXiv:1311.5490](#)] [[INSPIRE](#)].
- [28] ETM collaboration, C. Michael, K. Ottnad and C. Urbach, *η and η' mixing from Lattice QCD*, *Phys. Rev. Lett.* **111** (2013) 181602 [[arXiv:1310.1207](#)] [[INSPIRE](#)].
- [29] K. Jansen and C. Urbach, *tmLQCD: A Program suite to simulate Wilson Twisted mass Lattice QCD*, *Comput. Phys. Commun.* **180** (2009) 2717 [[arXiv:0905.3331](#)] [[INSPIRE](#)].
- [30] M. Lüscher and F. Palombi, *Universality of the topological susceptibility in the $SU(3)$ gauge theory*, *JHEP* **09** (2010) 110 [[arXiv:1008.0732](#)] [[INSPIRE](#)].
- [31] ETM collaboration, K. Cichy, E. Garcia-Ramos and K. Jansen, *Topological susceptibility from the twisted mass Dirac operator spectrum*, *JHEP* **02** (2014) 119 [[arXiv:1312.5161](#)] [[INSPIRE](#)].
- [32] K. Cichy, E. Garcia-Ramos and K. Jansen, *Chiral condensate from the twisted mass Dirac operator spectrum*, *JHEP* **10** (2013) 175 [[arXiv:1303.1954](#)] [[INSPIRE](#)].
- [33] C. Alexandrou, M. Constantinou, T. Korzec, H. Panagopoulos and F. Stylianou, *Renormalization constants of local operators for Wilson type improved fermions*, *Phys. Rev. D* **86** (2012) 014505 [[arXiv:1201.5025](#)] [[INSPIRE](#)].
- [34] K. Cichy, K. Jansen and P. Korcyl, *Non-perturbative renormalization in coordinate space for $N_f = 2$ maximally twisted mass fermions with tree-level Symanzik improved gauge action*, *Nucl. Phys. B* **865** (2012) 268 [[arXiv:1207.0628](#)] [[INSPIRE](#)].

- [35] K. Cichy, E. Garcia-Ramos and K. Jansen, *Short distance singularities and automatic $O(a)$ improvement: the cases of the chiral condensate and the topological susceptibility*, *JHEP* **04** (2015) 048 [[arXiv:1412.0456](#)] [[INSPIRE](#)].
- [36] K. Cichy, E. Garcia-Ramos, K. Jansen and A. Shindler, *Topological susceptibility from twisted mass fermions using spectral projectors*, *PoS(LATTICE 2013)* 129 [[arXiv:1312.3535](#)] [[INSPIRE](#)].
- [37] L. Del Debbio, L. Giusti and C. Pica, *Topological susceptibility in the $SU(3)$ gauge theory*, *Phys. Rev. Lett.* **94** (2005) 032003 [[hep-th/0407052](#)] [[INSPIRE](#)].
- [38] S. Dürr, Z. Fodor, C. Hölbling and T. Kurth, *Precision study of the $SU(3)$ topological susceptibility in the continuum*, *JHEP* **04** (2007) 055 [[hep-lat/0612021](#)] [[INSPIRE](#)].
- [39] B. Lucini and M. Teper, *$SU(N)$ gauge theories in four-dimensions: Exploring the approach to $N = \infty$* , *JHEP* **06** (2001) 050 [[hep-lat/0103027](#)] [[INSPIRE](#)].
- [40] L. Del Debbio, H. Panagopoulos and E. Vicari, *θ dependence of $SU(N)$ gauge theories*, *JHEP* **08** (2002) 044 [[hep-th/0204125](#)] [[INSPIRE](#)].
- [41] L. Del Debbio and C. Pica, *Topological susceptibility from the overlap*, *JHEP* **02** (2004) 003 [[hep-lat/0309145](#)] [[INSPIRE](#)].
- [42] L. Giusti, M. Lüscher, P. Weisz and H. Wittig, *Lattice QCD in the ϵ -regime and random matrix theory*, *JHEP* **11** (2003) 023 [[hep-lat/0309189](#)] [[INSPIRE](#)].
- [43] H. Neff, N. Eicker, T. Lippert, J.W. Negele and K. Schilling, *On the low fermionic eigenmode dominance in QCD on the lattice*, *Phys. Rev. D* **64** (2001) 114509 [[hep-lat/0106016](#)] [[INSPIRE](#)].
- [44] ETM collaboration, P. Boucaud et al., *Dynamical Twisted Mass Fermions with Light Quarks: Simulation and Analysis Details*, *Comput. Phys. Commun.* **179** (2008) 695 [[arXiv:0803.0224](#)] [[INSPIRE](#)].
- [45] ETM collaboration, K. Jansen, C. Michael and C. Urbach, *The η' meson from lattice QCD*, *Eur. Phys. J. C* **58** (2008) 261 [[arXiv:0804.3871](#)] [[INSPIRE](#)].
- [46] C. Michael and I. Teasdale, *Extracting Glueball Masses From Lattice QCD*, *Nucl. Phys. B* **215** (1983) 433 [[INSPIRE](#)].
- [47] M. Lüscher and U. Wolff, *How to Calculate the Elastic Scattering Matrix in Two-dimensional Quantum Field Theories by Numerical Simulation*, *Nucl. Phys. B* **339** (1990) 222 [[INSPIRE](#)].
- [48] B. Blossier, M. Della Morte, G. von Hippel, T. Mendes and R. Sommer, *On the generalized eigenvalue method for energies and matrix elements in lattice field theory*, *JHEP* **04** (2009) 094 [[arXiv:0902.1265](#)] [[INSPIRE](#)].
- [49] R. Frezzotti and S. Sint, *Some remarks on $O(a)$ improved twisted mass QCD*, *Nucl. Phys. Proc. Suppl.* **106** (2002) 814 [[hep-lat/0110140](#)] [[INSPIRE](#)].
- [50] M. Della Morte, R. Frezzotti and J. Heitger, *Quenched twisted mass QCD at small quark masses and in large volume*, *Nucl. Phys. Proc. Suppl.* **106** (2002) 260 [[hep-lat/0110166](#)] [[INSPIRE](#)].
- [51] XLF collaboration, K. Jansen, A. Shindler, C. Urbach and I. Wetzorke, *Scaling test for Wilson twisted mass QCD*, *Phys. Lett. B* **586** (2004) 432 [[hep-lat/0312013](#)] [[INSPIRE](#)].

- [52] T. Feldmann, P. Kroll and B. Stech, *Mixing and decay constants of pseudoscalar mesons*, *Phys. Rev. D* **58** (1998) 114006 [[hep-ph/9802409](#)] [[INSPIRE](#)].
- [53] T. Feldmann, P. Kroll and B. Stech, *Mixing and decay constants of pseudoscalar mesons: The Sequel*, *Phys. Lett. B* **449** (1999) 339 [[hep-ph/9812269](#)] [[INSPIRE](#)].
- [54] K. Ottnad and C. Urbach, *Decay constants of η, η' -mesons from lattice qcd*, in preparation.
- [55] XLF collaboration, K. Jansen, M. Papinutto, A. Shindler, C. Urbach and I. Wetzorke, *Light quarks with twisted mass fermions*, *Phys. Lett. B* **619** (2005) 184 [[hep-lat/0503031](#)] [[INSPIRE](#)].
- [56] FOR THE ETM collaboration, A. Deuzeman, S. Reker and C. Urbach, *Lemon: an MPI parallel I/O library for data encapsulation using LIME*, *Comput. Phys. Commun.* **183** (2012) 1321 [[arXiv:1106.4177](#)] [[INSPIRE](#)].
- [57] R Development Core Team, *R: A language and environment for statistical computing*, R Foundation for Statistical Computing, Vienna, Austria (2005) [ISBN: 3-900051-07-0].