Biochimica et Biophysica Acta 1837 (2014) 782



Contents lists available at ScienceDirect

Biochimica et Biophysica Acta

journal homepage: www.elsevier.com/locate/bbabio



Erratum

Erratum to "The *Saccharomyces cerevisiae* gene YPR011c encodes a mitochondrial transporter of adenosine 5′-phosphosulfate and 3′-phospho-adenosine 5′-phosphosulfate" [Biochim. Biophys. Acta Bioenerg. (2014): 326, 334]



Simona Todisco ^{a,b,1}, Maria Antonietta Di Noia ^{c,1}, Alessandra Castegna ^{a,b}, Francesco Massimo Lasorsa ^{a,d}, Eleonora Paradies ^{a,d}, Ferdinando Palmieri ^{a,b,*}

- ^a Department of Biosciences, Biotechnologies and Biopharmaceutics, University of Bari, via Orabona 4, 70125 Bari, Italy
- ^b Center of Excellence in Comparative Genomics, University of Bari, Italy
- ^c Department of Sciences, University of Basilicata, via N. Sauro 85, 85100 Potenza, Italy
- ^d CNR Institute of Biomembranes and Bioenergetics, via Amendola 165/A, 70126 Bari, Italy

For the reader's convenience, the full legend is reprinted below.

Thermotolerance of wild-type, $YPR011c\Delta$, $MET3\Delta$ -mtMet3p, and $YPR011c\Delta$ MET3 Δ -mtMet3p cells shifted from 30 to 45 °C. Wild-type and deletion strains were pre-incubated in glucose-supplemented SC medium until the stationary phase was reached and washed three times with methionine-less SC medium. Then wild-type (\bullet), $YPR011c\Delta$ (\blacksquare), $MET3\Delta$ -mtMet3p (\square) and $YPR011c\Delta$ MET3 Δ -mtMet3p (\square) cells were grown at 30 °C in methionine-less SC medium supplemented with 2% ethanol and 0.2% galactose to 2 × 10⁷ cells ml⁻¹ and shifted to 45 °C. Wild-type cells were also treated as above except that the medium was supplemented with 3 mM methionine (\bullet). At the indicated times after the temperature shift to 45 °C the same portion of each culture was diluted in distilled water and immediately spread onto solid YEPD.

DOI of original article: http://dx.doi.org/10.1016/j.bbabio.2013.11.013.

^{*} Corresponding author at: Department of Biosciences, Biotechnologies and Biopharmaceutics, University of Bari, via Orabona 4, 70125 Bari, Italy. Tel.: +39 080 5443323. E-mail addresses: ferdpalmieri@gmail.com, fpalm@farmbiol.uniba.it (F. Palmieri).

¹ Contributed equally to this work.