ISSN 0735-1097/07/\$32.00 doi:10.1016/j.jacc.2007.01.072

#### **Heart Rhythm Disorders**

# Management and Outcome of Permanent Pacemaker and Implantable Cardioverter-Defibrillator Infections

Muhammad R. Sohail, MD,\* Daniel Z. Uslan, MD,\* Akbar H. Khan, MD,‡ Paul A. Friedman, MD,† David L. Hayes, MD,† Walter R. Wilson, MD,\* James M. Steckelberg, MD,\* Sarah Stoner, MS,§ Larry M. Baddour, MD\*

Rochester, Minnesota

Objectives	We describe the management and outcome of permanent pacemaker (PPM) and implantable cardioverter- defibrillator (ICD) infections in a large cohort of patients seen at a tertiary care facility with expertise in device lead extraction.
Background	Infection is a serious complication of PPM and ICD implantation. Optimal care of patients with these cardiac device infections (CDI) is not well defined.
Methods	A retrospective review of all patients with CDI admitted to Mayo Clinic Rochester between January 1, 1991, and December 31, 2003, was conducted. Demographic and clinical data were collected, and descriptive analysis was performed.
Results	A total of 189 patients met the criteria for CDI (138 PPM, 51 ICD). The median age of the patients was 71.2 years. Generator pocket infection (69%) and device-related endocarditis (23%) were the most common clinical presentations. Coagulase-negative staphylococci and <i>Staphylococcus aureus</i> , in 42% and 29% of cases, respectively, were the leading pathogens for CDI. Most patients (98%) underwent complete device removal. Duration of antibiotic therapy after device removal was based on clinical presentation and causative organism (median duration of 18 days for pocket infection vs. 28 days for endocarditis; 28 days for <i>S. aureus</i> infection vs. 14 days for coagulase-negative staphylococci infection [p < 0.001]). Median follow-up after hospital discharge was 175 days. Ninety-six percent of patients were cured with both complete device removal and antibiotic administration.
Conclusions	Cure of CDI is achievable in the large majority of patients treated with an aggressive approach of combined anti- microbial treatment and complete device removal. Based on findings of our large retrospective institutional sur- vey and previously published data, we submit proposed management guidelines of CDI. (J Am Coll Cardiol 2007;49:1851-9) © 2007 by the American College of Cardiology Foundation

With an increasing number of indications for use of permanent pacemakers (PPM) and implantable cardioverter-defibrillators (ICD), there has been an accelerated rate of device implantation (1,2). According to one estimate in the U.S., there was a 42% increase in the implantation rate of cardiac devices among Medicare beneficiaries from 1990 to 1999 (3). This was associated with a 124% increase in the rate of cardiac device infection (CDI) among the study population.

Management of CDI is a difficult challenge for both cardiology and infectious diseases specialists. Reported incidence rates of CDI range from 0.13% to 19.9% and 0.0% to 0.8% for PPMs and ICDs, respectively (4–9). A CDI can present with a pulse-generator pocket infection or blood-stream infection (BSI) with or without device-related endocarditis. A CDI is complicated by increased morbidity, mortality, and financial cost (10). Reported mortality rates

From the \*Division of Infectious Diseases, †Division of Cardiovascular Diseases, ‡Department of Medicine, and \$Department of Health Sciences Research, Mayo Clinic College of Medicine, Rochester, Minnesota. Dr. Sohail is currently at Tawam Hospital/Johns Hopkins Medicine, Al Ain, United Arab Emirates; Dr. Khan is currently at Emory University School of Medicine, Atlanta, Georgia; and Dr. Uslan is currently at David Geffen School of Medicine, University of California at Los Angeles, California. This work was supported in part by the Enhance Award (Department of Medicine), Small Grants Program (Division of Infectious Diseases), and research funds from the Division of Cardiology, Mayo Clinic College of Medicine. Dr. Hayes received honoraria from Medtronic, Guidant, St. Jude Medical, ELA Medical, and Biotronik; sponsored research from Medtronic, Guidant, and St. Jude

Medical; is on the medical advisory board of Guidant; and is a steering committee member of Medtronic. Dr. Friedman received honoraria from or is a consultant for Medtronic, Guidant, and AstraZeneca; sponsored research from Medtronic, AstraZeneca via Beth Israel, Guidant, St. Jude, and Bard; and holds intellectual property rights with Bard EP, Hewlett Packard, and Medical Positioning, Inc. Dr. Baddour received royalty payments from Elsevier and UpToDate and is an ACP/PIER editorial consultant. Data presented in part at the 43rd Annual Meeting of the Infectious Diseases Society of America, October 6 to 9, 2005, San Francisco, California (abstract no. 387).

Manuscript received October 2, 2006; revised manuscript received December 11, 2006, accepted January 2, 2007.

Abbreviations and Acronyms	
<b>BSI</b> = blood stream infection	1
<b>CDI</b> = cardiac device infection	,
CoNS = coagulase-negative staphylococci	-
ICD = implantable cardioverter-defibrillator	ן נ
PPM = permanent pacemaker	t - -
	1

of cardiac device-related endocarditis range from 31% to 66% if the infected device is not removed, and 18% or less with a combined management approach with complete device removal and antimicrobial therapy (11,12). Estimated average cost of combined medical and surgical treatment of an infected device ranges from \$25,000 to \$50,000 for PPM and ICD, respectively (13,14). Life-threatening complications of device removal are

well-reported (12,15,16).

To characterize the management and outcome of CDI, we retrospectively reviewed 189 cases of CDI that presented to Mayo Clinic Rochester between January 1991 and December 2003.

## **Methods**

We performed a retrospective review of the medical records of patients with CDI treated at Mayo Clinic Rochester between January 1, 1991, and December 31, 2003. The CDI cases were identified using several Mayo Clinic resources that included the Cardiology Device Database, the Surgical Index, and the computerized central diagnostic index. All patients had consented to use of their medical records for research purposes. The study proposal was approved by the Mayo Foundation Institutional Review Board.

**Definitions.** A CDI was defined as previously described by our group and others (17–19). Clinical evidence of CDI included local signs of inflammation at the generator pocket, including erythema, warmth, fluctuance, wound dehiscence, erosion, tenderness, or purulent drainage. Presence of PPM/ICD-related endocarditis was clinically confirmed when valvular or lead vegetations were detected by echocardiography, or if the Duke criteria for infective endocarditis were met (12,20). Vegetation was defined as an oscillating intracardiac mass on the electrode leads, cardiac valve leaflets, or endocardial surface in the setting of valve or lead infection confirmed by imaging in more than 1 echocardiographic plane, and positive blood and/or lead tip cultures (12,21–24).

A CDI was microbiologically confirmed based on positive cultures from the generator pocket, lead(s), or blood (in the presence of local inflammatory signs at generator pocket or absence of another source of bacteremia and resolution of blood stream infection after device explantation). Relapse was defined as the recurrence of the device infection with same organism, based on similar antibiogram.

Follow-up. Patients who presented with CDI and underwent device replacement procedure as part of their CDI treatment at Mayo Clinic Rochester had the option of being followed up at our institution or by their local health care providers. Of the 182 patients who survived index hospitalization for CDI, 142 patients (78%) chose follow-up at our institution, whereas 40 (22%) patients had subsequent care by their referring physician after discharge from Mayo Clinic. Median follow-up duration for cases of CDI was 175 days (mean 657 days).

**Statistical evaluation.** The Fisher exact test was used to test for differences in proportions for categorical data, and the Wilcoxon rank-sum test was used to test for differences in medians for continuous data. All p values were 2-sided with a level of  $\leq 0.05$  considered statistically significant. No adjustments for multiple comparisons were made. All analyses were performed using SAS version 8.2 software (SAS Institute Inc., Cary, North Carolina).

## Results

Baseline characteristics. A total of 189 cases of CDI met the case definition and were included in the study. Demographic and clinical characteristics of study subjects are summarized in Table 1. The median age of patients was 71.2 years (range 17 to 95 years). Seventy-eight percent were male. Devices included 138 PPMs and 51 ICDs. Most patients (75%) had dual-chamber devices. Fifty-eight (31%) patients had their cardiac devices implanted at Mayo Clinic Rochester, and 131 (69%) had device placement at another institution. Transvenous lead placement (93%) was the most commonly used lead insertion technique. The pulse generator was placed in the pectoral area in 172 (91%) patients and in the abdominal wall in 17 (9%) patients. Indications for initial device placement included heart block (38%), sinus node dysfunction (27%), and ventricular arrhythmias (21%). A CDI occurred after initial device implantation in 79 (42%) patients and after a revision (i.e., system upgrade, lead revision, or generator exchange, and so on) in 110 (58%).

**Clinical presentation.** Median time from device implantation to infection was 415 days for PPM and 125 days for ICDs (p = 0.009). Eighty-six (46%) patients initially presented to Mayo Clinic Rochester, and 103 (54%) were referred to Mayo Clinic Rochester because of lack of available expertise for lead extraction at the referring facility. Pocket infection, with (17%) or without (52%) BSI, was the most common clinical presentation, followed by devicerelated endocarditis (23%). Most patients had localized inflammatory signs at the pocket site (Table 2). Systemic signs of sepsis were present in less than one-half of the cases. Ten (5%) cases presented with erosion of lead or device generator, without accompanying inflammatory signs at the generator pocket site.

**Microbiology.** Coagulase-negative staphylococci (CoNS, 42%) and *Staphylococcus aureus* (29%) were the most common causes of CDI, followed by gram-negative bacilli (9%), including *Klebsiella pneumoniae*, *Serratia marcescens*, *Pseudo*-

monas aeruginosa, Stenotrophomonas maltophilia, Acinetobacter xylosoxidans, Acinetobacter baumanii, Citrobacter koseri, Morganella morganii, Hemophilus influenzae, and Moraxella catarrhalis (Fig. 1). Two patients had fungal (Candida albicans and Aspergillus fumigatus) infection. Polymicrobial infection was present in 7% of cases. Fourteen (7%) patients had localized inflammatory signs at generator pocket or erosion of device/lead but pocket cultures were negative. Purulent material was encountered (intraoperative) within the pulse-generator pocket in 151 (80%) of the cases (11% of these patients had no inflammatory signs at the generator pocket on physical examination). Seventy-five percent of the

Table 1

Demographic and Clinical Characteristics of Patients With CDI Who Were Treated at Mayo Clinic Rochester Between 1991 and 2003 (n = 189)

Age (yrs), median (range)71 (17-95)Gender, male148 (78%)Device138 (73%)ICD51 (27%)Chambers44 (23%)Dual142 (75%)Lead placement71 (6 93%)Epicardial5 (3%)Both8 (4%)Indication71 (38%)Kanse dystration51 (27%)Ventricular tachycardia/fibrillation93 (21%)Other28 (14%)De novo implantation79 (42%)System revision/insertion19 (10%)Generator replacement49 (26%)Time from implant/last intervention to infection, median days (mean)114 (60%)PPM415 (920)ICD125 (399)Coronary artery disease114 (60%)Heart failure96 (51%)Anticoagulation67 (35%)Diabetes mellitus46 (24%)Chronic obstructive pulmonary disease34 (18%)Malignancy29 (52%)Pocket infection99 (52%)Pocket infection with bacteremia32 (17%)Device-related endocarditis44 (23%)Bacteremia without localizing signs at pocket21 (11%)Generator re lead erosion10 (5%)	Characteristic	Value
Gender, male148 (78%)Device138 (73%)ICD51 (27%)Chambers44 (23%)Dual124 (75%)Lead placement5 (3%)Epicardial5 (3%)Both8 (4%)Indication71 (38%)Metricular tachycardia/fibrillation39 (21%)Other28 (14%)Event procedure before CDI28 (14%)De novo implantation79 (42%)Generator replacement49 (26%)Indication19 (10%)Coroary artery disease114 (60%)PPM415 (920)ICD125 (399)Dernovo implantation to infection, median days (mean)114 (60%)PPM415 (920)ICD125 (399)Coronary artery disease114 (60%)Heart failure96 (51%)Diabetes mellitus46 (24%)Inticagulation67 (35%)Diabetes mellitus46 (24%)Chronic obstructive pulmonary disease34 (18%)Malignancy29 (52%)Pocket infection99 (52%)Pocket infection99 (52%)Pocket infection with bacteremia32 (17%)Device-related endocarditis44 (23%)Bacteremia without localizing signs at pocket21 (11%)Generator relad erosion10 (5%)	Age (yrs), median (range)	71 (17-95)
DevicePPM138 (73%)ICD51 (27%)Chambers51 (27%)Single44 (23%)Dual142 (75%)Lead placement176 (93%)Epicardial5 (3%)Both8 (4%)Indication171 (38%)Heart block71 (38%)Sinus node dysfunction51 (27%)Other28 (12%)Other39 (21%)Other39 (21%)Other29 (21%)Everodure before CDI20 (10%)De novo implantation79 (42%)System revision/upgrade41 (22%)Lead revision/insertion19 (10%)Generator replacement49 (26%)ID125 (399)Coronary artery disease114 (60%)Heart failure96 (51%)Anticoagulation67 (35%)Diabetes mellitus44 (24%)Diabetes mellitus44 (24%)Malignancy24 (15%)Chronic obstructive pulmonary disease34 (18%)Malignancy32 (17%)Pocket infection with bacteremia32 (11%)Bacteremia without localizing signs at pocket21 (11%)Generator re lead erosion10 (5%)	Gender, male	148 (78%)
PPM138 (73%)ICD51 (27%)Chambers44 (23%)Dual142 (75%)Lead placement176 (93%)Epicardial5 (3%)Both8 (4%)Indication171 (38%)Heart block71 (38%)Ventricular tachycardia/fibrillation39 (21%)Other28 (14%)Indication21 (13%)Least procedure before CDI28 (14%)De novo implantation79 (42%)System revision/upgrade41 (22%)Lead revision/insertion19 (10%)Generator replacement49 (26%)ICD125 (399)Coronary artery disease114 (60%)Anticoagulation67 (35%)Diabetes mellitus46 (24%)Malignancy24 (13%)Chronic obstructive pulmonary disease34 (18%)Malignancy32 (17%)Poket infection with bacteremia32 (17%)Pocket infection with bacteremia32 (17%)Device-related endocarditis44 (23%)Bacteremia without localizing signs at pocket21 (11%)Generator or lead erosion10 (5%)	Device	
ICD51 (27%)ChambersSingle44 (23%)Dual142 (75%)Lead placementTransvenous176 (93%)Epicardial5 (3%)Both8 (4%)Indication11 (38%)Heart block71 (38%)Sinus node dysfunction51 (27%)Ventricular tachycardia/fibrillation39 (21%)Other28 (14%)East procedure before CDI28 (14%)De novo implantation79 (42%)System revision/upgrade41 (22%)Lead revision/insertion19 (10%)Generator replacement49 (26%)ICD125 (399)Coronary artery disease114 (60%)Heart failure96 (51%)IDabetes mellitus46 (24%)Malignancy24 (13%)Pocket infection99 (52%)Pocket infection with bacteremia32 (17%)Pocket infection with bacteremia32 (17%)Device-related endocarditis44 (23%)Bacteremia without localizing signs at pocket21 (11%)Generator re lead erosion10 (5%)	PPM	138 (73%)
ChambersSingle44 (23%)Dual142 (75%)Lead placement176 (93%)Epicardial5 (3%)Both8 (4%)Indication11 (38%)Heart block71 (38%)Sinus node dysfunction51 (27%)Ventricular tachycardia/fibrillation39 (21%)Other28 (14%)Indication99 (21%)De novo implantation79 (42%)System revision/upgrade41 (22%)Lead revision/insertion19 (10%)Generator replacement49 (26%)ICD125 (399)Coronary artery disease114 (60%)Indicagulation67 (35%)Diabetes mellitus66 (24%)Indicagulation67 (35%)Diabetes mellitus44 (23%)Indicagulation67 (35%)Diabetes mellitus44 (23%)Indicagulation67 (35%)Diabetes mellitus44 (23%)Diabetes mellitus44 (23%)Malignancy24 (13%)Pocket infection with bacteremia32 (17%)Pocket infection with bacteremia32 (17%)Pocket infection with bacteremia32 (17%)Device-related endocarditis44 (23%)Bacteremia without localizing signs at pocket21 (11%)Generator relad erosion10 (5%)	ICD	51 (27%)
Single44 (23%)Dual142 (75%)Lead placement176 (93%)Epicardial5 (3%)Both8 (4%)Indication110 (2000)Heart block71 (38%)Sinus node dysfunction51 (27%)Ventricular tachycardia/fibrillation39 (21%)Other28 (14%)Last procedure before CDI28 (14%)De novo implantation79 (42%)System revision/upgrade41 (22%)Lead revision/insertion19 (10%)Generator replacement49 (26%)ICD125 (399)Coronary artery disease114 (60%)Heart failure96 (51%)Anticoagulation67 (35%)Diabetes mellitus46 (24%)Anticoagulation67 (35%)Diabetes mellitus46 (24%)Chronic obstructive pulmonary disease34 (18%)Malignancy24 (13%)Pocket infection with bacteremia32 (17%)Pocket infection with bacteremia32 (17%)Device-related endocarditis44 (23%)Bacteremia without localizing signs at pocket21 (11%)Generator re lead erosion10 (5%)	Chambers	
Dual142 (75%)Lead placementTransvenous176 (93%)Epicardial5 (3%)Both8 (4%)IndicationHeart block71 (38%)Sinus node dysfunction51 (27%)Ventricular tachycardia/fibrillation39 (21%)Other28 (14%)East procedure before CDI28 (14%)De novo implantation79 (42%)System revision/upgrade41 (22%)Lead revision/insertion19 (10%)Generator replacement49 (26%)ICD125 (399)Corronary artery disease114 (60%)Heart failure96 (51%)Anticoagulation67 (35%)Diabetes mellitus46 (24%)Malignancy24 (13%)Clinical presentation99 (52%)Pocket infection with bacteremia32 (17%)Device-related endocarditis44 (23%)Bacteremia without localizing signs at pocket21 (11%)Generator or lead erosion10 (5%)	Single	44 (23%)
Lead placementTransvenous176 (93%)Epicardial5 (3%)Both8 (4%)Indication11 (38%)Heart block71 (38%)Sinus node dysfunction51 (27%)Ventricular tachycardia/fibrillation39 (21%)Other28 (14%)Last procedure before CDI79 (42%)Lead revision/upgrade41 (22%)Lead revision/insertion19 (10%)Generator replacement49 (26%)ICD125 (399)Coronary artery disease114 (60%)Heart failure96 (51%)Anticoagulation67 (35%)Diabetes mellitus46 (24%)Chronic obstructive pulmonary disease34 (18%)Malignancy24 (13%)Coroneriation99 (52%)Pocket infection with bacteremia32 (17%)Device-related endocarditis44 (23%)Bacteremia without localizing signs at pocket21 (11%)Generator or lead erosion10 (5%)	Dual	142 (75%)
Transvenous176 (93%)Epicardial5 (3%)Both8 (4%)Indication71 (38%)Heart block71 (38%)Sinus node dysfunction51 (27%)Ventricular tachycardia/fibrillation39 (21%)Other28 (14%)Last procedure before CDI28 (14%)De novo implantation79 (42%)Lead revision/upgrade41 (22%)Lead revision/insertion19 (10%)Generator replacement49 (26%)Time from implant/last intervention to infection, median days (mean)125 (399)Coronary artery disease114 (60%)Heart failure96 (51%)Othes mellitus46 (24%)Chronic obstructive pulmonary disease34 (18%)Malignancy24 (13%)Clinical presentation99 (52%)Pocket infection with bacteremia Device-related endocarditis44 (23%)Bacteremia without localizing signs at pocket21 (11%)Generator or lead erosion10 (5%)	Lead placement	
Epicardial5 (3%)Both8 (4%)Indication71 (38%)Heart block71 (38%)Sinus node dysfunction51 (27%)Ventricular tachycardia/fibrillation39 (21%)Other28 (14%)Last procedure before CDI28 (14%)De novo implantation79 (42%)System revision/upgrade41 (22%)Lead revision/insertion19 (10%)Generator replacement49 (26%)Time from implant/last intervention to infection, median days (mean)125 (399)Comorbid conditions114 (60%)Heart failure96 (51%)Anticoagulation67 (35%)Diabetes mellitus46 (24%)Malignancy24 (13%)Clinical presentation24 (13%)Pocket infection with bacteremia32 (17%)Pocket infection with bacteremia32 (17%)Bacteremia without localizing signs at pocket21 (11%)Generator or lead erosion10 (5%)	Transvenous	176 (93%)
Both8 (4%)IndicationHeart block71 (38%)Sinus node dysfunction51 (27%)Sinus node dysfunction39 (21%)Other28 (14%)Last procedure before CDI28 (14%)De novo implantation79 (42%)System revision/upgrade41 (22%)Lead revision/insertion19 (10%)Generator replacement49 (26%)Time from implant/last intervention to infection, median days (mean)125 (399)Comorbid conditions114 (60%)Heart failure96 (51%)Anticoagulation67 (35%)Diabetes mellitus46 (24%)Malignancy24 (13%)Clinical presentation99 (52%)Pocket infection with bacteremia32 (17%)Device-related endocarditis44 (23%)Bacteremia without localizing signs at pocket21 (11%)Generator or lead erosion10 (5%)	Epicardial	5 (3%)
IndicationHeart block71 (38%)Sinus node dysfunction51 (27%)Ventricular tachycardia/fibrillation39 (21%)Other28 (14%)Last procedure before CDI28 (14%)De novo implantation79 (42%)System revision/upgrade41 (22%)Lead revision/insertion19 (10%)Generator replacement49 (26%)Time from implant/last intervention to infection, median days (mean)125 (399)Comorbid conditions114 (60%)Heart failure96 (51%)Anticoagulation67 (35%)Diabetes mellitus46 (24%)Malignancy24 (13%)Clinical presentation99 (52%)Pocket infection with bacteremia32 (17%)Device-related endocarditis44 (23%)Bacteremia without localizing signs at pocket21 (11%)Generator or lead erosion10 (5%)	Both	8 (4%)
Heart block71 (38%)Sinus node dysfunction51 (27%)Ventricular tachycardia/fibrillation39 (21%)Other28 (14%)Last procedure before CDI2De novo implantation79 (42%)System revision/upgrade41 (22%)Lead revision/insertion19 (10%)Generator replacement49 (26%)Time from implant/last intervention to infection, median days (mean)125 (399)ICD125 (399)Corronzry artery disease114 (60%)Heart failure96 (51%)Anticoagulation67 (35%)Diabetes mellitus46 (24%)Malignancy24 (13%)Clinical presentation32 (17%)Pocket infection with bacteremia32 (17%)Pocket infection with bacteremia32 (17%)Bacteremia without localizing signs at pocket21 (11%)Generator or lead erosion10 (5%)	Indication	
Sinus node dysfunction51 (27%)Ventricular tachycardia/fibrillation39 (21%)Other28 (14%)Last procedure before CDI79 (42%)De novo implantation79 (42%)System revision/upgrade41 (22%)Lead revision/insertion19 (10%)Generator replacement49 (26%)Time from implant/last intervention to infection, median days (mean)125 (399)ICD125 (399)Coronary artery disease114 (60%)Heart failure96 (51%)Diabetes mellitus46 (24%)Malignancy24 (13%)Clinical presentation32 (17%)Pocket infection with bacteremia32 (17%)Device-related endocarditis44 (23%)Bacteremia without localizing signs at pocket21 (11%)Generator or lead erosion10 (5%)	Heart block	71 (38%)
Ventricular tachycardia/fibrillation39 (21%)Other28 (14%)Last procedure before CDIDe novo implantation79 (42%)System revision/upgrade41 (22%)Lead revision/insertion19 (10%)Generator replacement49 (26%)Time from implant/last intervention to infection, median days (mean)125 (399)ICD125 (399)Coronary artery disease114 (60%)Heart failure96 (51%)Diabetes mellitus46 (24%)Malignancy34 (18%)Chronic obstructive pulmonary disease34 (18%)Malignancy24 (13%)Pocket infection with bacteremia32 (17%)Pocket infection with bacteremia32 (17%)Bacteremia without localizing signs at pocket21 (11%)Generator or lead erosion10 (5%)	Sinus node dysfunction	51 (27%)
Other28 (14%)Last procedure before CDIDe novo implantation79 (42%)System revision/upgrade41 (22%)Lead revision/insertion19 (10%)Generator replacement49 (26%)Time from implant/last intervention to infection, median days (mean)415 (920)ICD125 (399)Coronary artery disease114 (60%)Heart failure96 (51%)Diabetes mellitus46 (24%)Diabetes mellitus46 (24%)Malignancy24 (13%)Clinical presentation99 (52%)Pocket infection with bacteremia32 (17%)Device-related endocarditis44 (23%)Bacteremia without localizing signs at pocket21 (11%)Generator or lead erosion10 (5%)	Ventricular tachycardia/fibrillation	39 (21%)
Last procedure before CDI   De novo implantation 79 (42%)   System revision/upgrade 41 (22%)   Lead revision/insertion 19 (10%)   Generator replacement 49 (26%)   Time from implant/last intervention to infection, median days (mean) 49 (26%)   PPM 415 (920)   ICD 125 (399)   Coronary artery disease 114 (60%)   Heart failure 96 (51%)   Anticoagulation 67 (35%)   Diabetes mellitus 46 (24%)   Chronic obstructive pulmonary disease 34 (18%)   Malignancy 24 (13%)   Clinical presentation 99 (52%)   Pocket infection with bacteremia 32 (17%)   Device-related endocarditis 44 (23%)   Bacteremia without localizing signs at pocket 21 (11%)   Generator or lead erosion 10 (5%)	Other	28 (14%)
De novo implantation79 (42%)System revision/upgrade41 (22%)Lead revision/insertion19 (10%)Generator replacement49 (26%)Time from implant/last intervention to infection, median days (mean)415 (920)ICD125 (399)Cornorbid conditions114 (60%)Heart failure96 (51%)Anticoagulation67 (35%)Diabetes mellitus46 (24%)Chronic obstructive pulmonary disease34 (18%)Malignancy24 (13%)Clinical presentation99 (52%)Pocket infection with bacteremia32 (17%)Device-related endocarditis44 (23%)Bacteremia without localizing signs at pocket21 (11%)Generator or lead erosion10 (5%)	Last procedure before CDI	
System revision/upgrade41 (22%)Lead revision/insertion19 (10%)Generator replacement49 (26%)Time from implant/last intervention to infection, median days (mean)125 (399)PPM415 (920)ICD125 (399)Cornorbid conditions114 (60%)Heart failure96 (51%)Anticoagulation67 (35%)Diabetes mellitus46 (24%)Chronic obstructive pulmonary disease34 (18%)Malignancy24 (13%)Clinical presentation99 (52%)Pocket infection with bacteremia32 (17%)Pocket infection with bacteremia32 (17%)Bacteremia without localizing signs at pocket21 (11%)Generator or lead erosion10 (5%)	De novo implantation	79 (42%)
Lead revision/insertion19 (10%)Generator replacement49 (26%)Time from implant/last intervention to infection, median days (mean)1PPM415 (920)ICD125 (399)Comorbid conditions114 (60%)Heart failure96 (51%)Anticoagulation67 (35%)Diabetes mellitus46 (24%)Malignancy24 (13%)Clinical presentation99 (52%)Pocket infection with bacteremia32 (17%)Pocket infection with bacteremia32 (17%)Bacteremia without localizing signs at pocket21 (11%)Generator or lead erosion10 (5%)	System revision/upgrade	41 (22%)
Generator replacement49 (26%)Time from implant/last intervention to infection, median days (mean)PPM415 (920)ICD125 (399)Comorbid conditionsCoronary artery disease114 (60%)Heart failure96 (51%)Anticoagulation67 (35%)Diabetes mellitus46 (24%)Malignancy24 (13%)Clinical presentation99 (52%)Pocket infection with bacteremia32 (17%)Pocket infection with bacteremia32 (17%)Bacteremia without localizing signs at pocket21 (11%)Generator or lead erosion10 (5%)	Lead revision/insertion	19 (10%)
Time from implant/last intervention to infection, median days (mean)   PPM 415 (920)   ICD 125 (399)   Corronary artery disease 114 (60%)   Heart failure 96 (51%)   Anticoagulation 67 (35%)   Diabetes mellitus 46 (24%)   Chronic obstructive pulmonary disease 34 (18%)   Malignancy 24 (13%)   Clinical presentation 99 (52%)   Pocket infection with bacteremia 32 (17%)   Device-related endocarditis 44 (23%)   Bacteremia without localizing signs at pocket 21 (11%)   Generator or lead erosion 10 (5%)	Generator replacement	49 (26%)
PPM 415 (920)   ICD 125 (399)   Comorbid conditions 114 (60%)   Heart failure 96 (51%)   Anticoagulation 67 (35%)   Diabetes mellitus 46 (24%)   Chronic obstructive pulmonary disease 34 (18%)   Malignancy 24 (13%)   Pocket infection 99 (52%)   Pocket infection with bacteremia 32 (17%)   Device-related endocarditis 44 (23%)   Bacteremia without localizing signs at pocket 21 (11%)   Generator or lead erosion 10 (5%)	Time from implant/last intervention to infection, median days (mean)	
ICD125 (399)Comorbid conditions124 (60%)Coronary artery disease114 (60%)Heart failure96 (51%)Anticoagulation67 (35%)Diabetes mellitus46 (24%)Chronic obstructive pulmonary disease34 (18%)Malignancy24 (13%)Clinical presentation99 (52%)Pocket infection with bacteremia32 (17%)Device-related endocarditis44 (23%)Bacteremia without localizing signs at pocket21 (11%)Generator or lead erosion10 (5%)	PPM	415 (920)
Comorbid conditionsCoronary artery disease114 (60%)Heart failure96 (51%)Anticoagulation67 (35%)Diabetes mellitus46 (24%)Chronic obstructive pulmonary disease34 (18%)Malignancy24 (13%)Clinical presentation99 (52%)Pocket infection with bacteremia32 (17%)Device-related endocarditis44 (23%)Bacteremia without localizing signs at pocket21 (11%)Generator or lead erosion10 (5%)	ICD	125 (399)
Coronary artery disease114 (60%)Heart failure96 (51%)Anticoagulation67 (35%)Diabetes mellitus46 (24%)Chronic obstructive pulmonary disease34 (18%)Malignancy24 (13%)Clinical presentation99 (52%)Pocket infection with bacteremia32 (17%)Device-related endocarditis44 (23%)Bacteremia without localizing signs at pocket21 (11%)Generator or lead erosion10 (5%)	Comorbid conditions	
Heart failure96 (51%)Anticoagulation67 (35%)Diabetes mellitus46 (24%)Chronic obstructive pulmonary disease34 (18%)Malignancy24 (13%)Clinical presentation99 (52%)Pocket infection with bacteremia32 (17%)Device-related endocarditis44 (23%)Bacteremia without localizing signs at pocket21 (11%)Generator or lead erosion10 (5%)	Coronary artery disease	114 (60%)
Anticoagulation67 (35%)Diabetes mellitus46 (24%)Diabetes mellitus46 (24%)Chronic obstructive pulmonary disease34 (18%)Malignancy24 (13%)Clinical presentation24 (13%)Pocket infection99 (52%)Pocket infection with bacteremia32 (17%)Device-related endocarditis44 (23%)Bacteremia without localizing signs at pocket21 (11%)Generator or lead erosion10 (5%)	Heart failure	96 (51%)
Diabetes mellitus46 (24%)Diabetes mellitus46 (24%)Chronic obstructive pulmonary disease34 (18%)Malignancy24 (13%)Clinical presentation99 (52%)Pocket infection99 (52%)Pocket infection with bacteremia32 (17%)Device-related endocarditis44 (23%)Bacteremia without localizing signs at pocket21 (11%)Generator or lead erosion10 (5%)	Anticoagulation	67 (35%)
Chronic obstructive pulmonary disease34 (18%)Malignancy24 (13%)Clinical presentation24 (13%)Pocket infection99 (52%)Pocket infection with bacteremia32 (17%)Device-related endocarditis44 (23%)Bacteremia without localizing signs at pocket21 (11%)Generator or lead erosion10 (5%)	Diabetes mellitus	46 (24%)
Malignancy24 (13%)Clinical presentation99 (52%)Pocket infection99 (52%)Pocket infection with bacteremia32 (17%)Device-related endocarditis44 (23%)Bacteremia without localizing signs at pocket21 (11%)Generator or lead erosion10 (5%)	Chronic obstructive pulmonary disease	34 (18%)
Clinical presentation 99 (52%)   Pocket infection 99 (52%)   Pocket infection with bacteremia 32 (17%)   Device-related endocarditis 44 (23%)   Bacteremia without localizing signs at pocket 21 (11%)   Generator or lead erosion 10 (5%)	Malignancy	24 (13%)
Pocket infection99 (52%)Pocket infection with bacteremia32 (17%)Device-related endocarditis44 (23%)Bacteremia without localizing signs at pocket21 (11%)Generator or lead erosion10 (5%)	Clinical presentation	. ,
Pocket infection with bacteremia32 (17%)Device-related endocarditis44 (23%)Bacteremia without localizing signs at pocket21 (11%)Generator or lead erosion10 (5%)	Pocket infection	99 (52%)
Device-related endocarditis44 (23%)Bacteremia without localizing signs at pocket21 (11%)Generator or lead erosion10 (5%)	Pocket infection with bacteremia	32 (17%)
Bacteremia without localizing signs at pocket21 (11%)Generator or lead erosion10 (5%)	Device-related endocarditis	44 (23%)
Generator or lead erosion 10 (5%)	Bacteremia without localizing signs at pocket	21 (11%)
	Generator or lead erosion	10 (5%)

 $\label{eq:cDI} \mbox{CDI} = \mbox{cardioverter-defibrillator; } \mbox{PPM} = \mbox{permanent} \\ \mbox{pacemaker.} \\$ 

Table 2	Clinical Presentation of Patients With CDI	(n = <b>189</b> )
	Presenting Signs/Symptoms	n (%)
Systemic sy	mptoms	
Fever (>3	38°C)	82 (43)
Chills		73 (39)
Malaise		79 (42)
Anorexia		32 (17)
Nausea		16 (8)
Sweating		34 (18)
Hypotens	on (systolic blood pressure $<$ 90 mm Hg)	18 (10)
Murmur o	n examination	66 (35)
Symptom	atic heart failure	52 (28)
Local findin	gs at generator site	
Erythema		128 (68)
Pain		93 (49)
Swelling		127 (67)
Warmth		71 (38)
Tenderne	SS	86 (46)
Drainage		95 (50)
Purulent	drainage	65 (34)
Skin ulce	ration	59 (31)
Generato	r/lead erosion	48 (25)
Intraopera	ative purulence at generator pocket	151 (80)
Laboratory a	abnormalities	
Leukocyto	osis (WBC $>$ 10 $ imes$ 10 $^9$ /l), n = 188	82 (43)
Anemia (	HCT ${<}38\%$ in men and ${<}35\%$ in women), n = 188	94 (50)
High erytl and $\geq$	rocyte sedimentation rate (ESR $>$ 22 mm/h in men 29 mm/h in women), n = 76	47 (25)
Positive b	lood culture (n = $188$ )	76 (40)

 $\label{eq:cd} \textbf{CDI} = \textbf{cardiac} \ \textbf{device} \ \textbf{infection}; \ \textbf{ESR} = \textbf{erythrocyte} \ \textbf{sedimentation} \ \textbf{rate;} \ \textbf{HCT} = \textbf{hematocrit;} \ \textbf{WBC} = \textbf{white} \ \textbf{blood} \ \textbf{cell} \ \textbf{count.}$ 

above cases had positive cultures from the generator pocket. Blood cultures were positive in only 76 (40%) of all cases. **Complications of device infection.** Complications of CDI included septic arthritis (n = 6), vertebral osteomyelitis (n = 2), sternal osteomyelitis (n = 2), femoral osteomyelitis (n = 1), lung abscesses (n = 5), splenic abscess (n = 1),



brain abscess (n = 1), liver abscess (n = 1), and perinephric abscess (n = 1). Twenty-two (11.6%) patients had thrombosis of a vein where leads were in place (subclavian vein or superior vena cava), and in 6 of these cases, they presented with symptomatic pulmonary embolism. The majority of patients (n = 147, 78%) had an echocardiogram (transthoracic or transesophageal) performed, and 44 (23%) had either lead and/or valvular vegetations. The diameter of vegetations ranged from 0.3 to 7.0 cm in the longest dimension. It is noteworthy that none of these patients developed clinical manifestations of pulmonary embolism as a complication of percutaneous lead extraction.

**Explantation of infected devices.** The cardiac device was explanted in 182 (96%) patients at the time of initial presentation. Three (2%) other patients suffered relapsing CDI and had explantation after failure of conservative (medical) treatment. A cardiac device was not explanted in 4 patients because of high operative risk (advanced age and decompensated heart failure with ejection fraction  $\leq$ 20%). The entire device system was explanted in 178 (94%) patients, whereas lead extraction was unsuccessful in 7 (4%) cases. Most patients (90%) underwent percutaneous lead extraction with use of manual traction (15%), locking stylet (34%), or laser sheath (41%). Nineteen patients (10%) had lead extraction via median sternotomy.

**Complications of lead extraction.** Twenty patients suffered complications of percutaneous lead extraction (3 manual traction, 7 locking stylet, and 10 cases with laser

sheaths), which included damage to the tricuspid valve (n =3) that required valve annuloplasty in 2 patients, subclavian vein laceration (n = 3), hemothorax (n = 4), pocket hematoma (n = 3), fracture of lead tip requiring surgical intervention (n = 6), and massive hemorrhage that required surgical intervention (n = 5) and resulted in the death of 1 patient. Surgical removal of epicardial leads was complicated by massive hemorrhage (n = 2) that was lethal in 1 patient, postoperative cardiac arrest requiring intensive care unit stay (n = 1), subclavian vein laceration (n = 1), and ventriculotomy requiring operative repair (n = 1). Complications during lead extraction were more common in patients with history of multiple device-related procedures (13 of 87, 14.9%) compared with those with initial implants (7 of 77, 9.1%). However, this difference was not statistically significant (p = 0.25).

Antimicrobial treatment. All patients received antimicrobials after removal or debridement of infected device system. Most patients received at least 2 weeks of antibiotics after removal of an infected device (Fig. 2). Median duration of antibiotic treatment was significantly longer in patients with *S. aureus* infection as compared to those with CoNS (28 days vs. 14 days, p < 0.001). Patients with device-related endocarditis (valvular or lead vegetation) and BSI received a median of 28 days of antibiotics after device removal. Cases with ICD infection had a longer duration of antibiotics (median 22 days) as compared to those with PPM infection (median 16 days), but this difference was not statistically



significant (p = 0.27). Most patients (97%) received a combination of intravenous and oral antibiotics. Only 3% were treated with oral antibiotics alone. Adverse effects of antibiotic treatment included diarrhea (8%), nephrotoxicity (5%), neutropenia (2%), Clostridium difficile colitis (1%), and oral candidiasis (1%).

Reimplantation of a new device. All patients were assessed for ongoing need of a cardiac device after extraction of an infected system. Reimplantation of a new device was considered necessary in only 127 (67%) of the cases. Median time from explantation to reimplantation was significantly longer for patients with BSI compared with nonbacteremic cases (13 days vs. 7 days, p < 0.0001). A new system was placed as early as 3 days in cases of negative blood cultures (Fig. 2). The median time from explantation to reimplantation in patients with CoNS was 7 days compared with a delay of 12 days in cases of S. aureus infection (p = 0.09). In cases in which a new device system was implanted, most patients (94%) had transvenous lead insertion with implantation of a generator on the opposite side of the infected pocket. Epicardial electrodes were placed in 7 (5%) patients. Generator or lead erosion. Among our study subjects with CDI, 10 patients presented with generator or lead erosion only. Despite the absence of local inflammatory signs at the pocket site, purulent material in the pulse-generator pocket was noted intraoperatively in 3 (30%) of these cases, and 9 (90%) had positive cultures from an intraoperative swab of the pocket. The causative micro-organism for CDI in this group of patients included common skin flora such as CoNS (n = 4), S. aureus (n = 3), and Propionibacterium spp. (n = 3)1), whereas 1 patient had polymicrobial infection with CoNS, Propionibacterium spp., and Corynebacterium spp. All patients had explanation of the infected device at the initial presentation (complete hardware successfully removed in 9 cases, whereas percutaneous lead extraction was unsuccessful in 1 case). Median time from removal of an infected device to insertion of a new system was 3 days. Among cases (n = 7) with no intraoperative evidence of purulence in the generator pocket, 4 (57%) had a new system placed on the same day when the infected device was removed. All patients were cured and none had infection relapse at their last follow-up visit (mean duration 78 weeks).

Outcome. The average duration of hospitalization for CDI was 17 days (median 13 days). Seven patients died during index hospitalization for CDI. Of the remaining 182 cases, follow-up data were available for 142 cases (40 patients were followed up elsewhere). Most patients (135, 95%) remained free of infection, after the initial treatment, at the last follow-up (median follow-up duration 175 days; mean 657 days). Two patients had persistent pocket infection caused by retained hardware. Five cases presented with relapse of infection. Clinical characteristics, microbiology, management, and outcome of these 5 cases are summarized in Table 3. Two of these patients were subsequently cured with complete removal of the infected device.

Table 3	Characteristics	of Patients With Relapse of	CDI					
Case	Device	Comorbid Conditions	Presentation	Microorganism	Initial Hardware Removal	Reinserted New Device	Timing of Relapse (weeks)	Relapse/Outcome
H	ICD (TVL + EPL)	DM, CHF	Pocket + BSI	CoNS	Partial*	No	9	EPL surgically removed at relapse. However, patient died of stroke.
5	ICD (TVL + EPL)	CHF	Pocket infection	MSSA	Partial†	Yes	4	Relapsed with device-related TV endocarditis. Cured with removal of epicardial patches.
ę	ICD (EPL)	CHF	Pocket + BSI	CONS	Partial*	Yes	16	Persistent pocket infection at last follow-up.
4	PPM (TVL)	Liver TX, Hodgkins lymphoma, HD, CHF	Endocarditis	MRSA	Complete	Yes‡	12	Died of MRSA sepsis and small bowel perforation despite removal of new device.
ъ	PPM (TVL)	CHF	Endocarditis	CoNS	Complete	Yes§	24	Cured with removal of replacement device.
*Patient had a device. §Temp RSI = bloods	handoned EPL from previc orary pacing wires before - stream infection. CHE = or	us devices, which were not removed beca insertion of new device.	use of high operative risk.	†Epicardial patches were dia diabatas mollitus. EDI	difficult to remove at initia = oniverdial leads HD =	l surgery and were left hemodialvsis- ICD =	behind. ‡Patient had dialysi mulantable cardioverter-defi	s catheter-related MRSA bacteremia with seeding of new builletor-MBSA = mothicillin-resistant C arrane: MSSA =

permanent pacemaker; TV = tricuspid valve; TVL = transvenous leads; TX = transplant

methicillin-sensitive S. aureus; PPM =

There were 9 in-hospital deaths, 7 during index hospitalization and 2 in patients who were subsequently hospitalized for relapse of infection. Five of these patients had device-related endocarditis (4 with staphylococci and 1 with *Aspergillus fumigatus*). Staphylococcal BSI with sepsis (n =5) was the most common cause of death. Two patients died secondary to complications of device extraction surgery. One patient died of ventilator-associated pneumonia during an intensive care unit stay after device removal. One severely immunocompromised patient with device-related fungal endocarditis and metastatic abscesses died.

Seven patients had partial device explantation at initial presentation. Two were lost to follow-up after the index hospitalization (both cases had abandoned epicardial leads from previous devices, and the currently infected devices had transvenous leads that were removed along with generators). One patient died of surgical complications during a second attempt at complete hardware removal. Of the remaining 4 patients, 2 had a relapse (Table 3, cases 2 and 3). The other 2 patients remained infection-free despite only partial device removal. Both had negative blood cultures and no evidence of endocarditis on echocardiography. Percutaneous lead extraction was attempted but was unsuccessful in first patient with pocket infection. Transvenous lead removal was not attempted in the second case because of absence of inflammatory changes during generator pocket exploration. In both cases, there was no evidence of infection at their last follow-up visit (5 months in the case of pocket infection and 4 years in the case of generator erosion only).

### Discussion

Despite improvements in the design and implantation techniques, infection of the cardiac devices remains a serious problem. The current study is one of the largest reported series of CDI in the United States and the only one to propose management guidelines.

**Clinical presentation and microbiology.** There was considerable variation in time from device implantation to onset of infection in our survey. Median time from implantation to infection was shorter for ICDs (125 days) than for PPMs (414 days). This time difference between the 2 groups may in part be caused by an increased proportion of *S. aureus* infection in patients with an ICD (35%) as compared to those with PPM (26%). Patients with PPM infection were more likely to have CoNS isolated (46%) from blood or pocket cultures than those with an ICD (31%).

Most patients with CDI present with only localized inflammatory signs at the generator pocket (Table 2), and a lack of systemic signs should not sway clinicians away from a suspicion of CDI (19). Nonspecific laboratory abnormalities such as leukocytosis, anemia, and high sedimentation rate were present in less than one-half of the cases in the current study. Staphylococcal species account for more than two-thirds of CDI cases in most published series (11,12,15,19,25) and were predominant in the present investigation (Fig. 1). Therefore, empiric antibiotics for suspected CDI should include coverage for staphylococci while awaiting microbiology results.

**Conservative treatment versus complete device removal.** Some investigators have advocated conservative treatment with antibiotics and generator pocket debridement without hardware removal (26–28). Most previously published studies, however, have shown unacceptably high failure rates with conservative treatment (11–13,19,22,23,29–35).

The Heart Rhythm Society guidelines state that extraction of an entire lead may not be necessary if the lead can be cut through a sterile incision, totally separate from an infected pocket (36). However, the intracardiac portion of a lead may be colonized in cases in which CDI findings are limited to the pulse-generator pocket only (13,25,37), and there is a high risk of treatment failure with a lead-retention approach as recommended by the Heart Rhythm Society (25,34,38-41). In the current study, 5 patients had CDI relapse (2.6%), and 3 of them had only partial explantation of the device (generator only) at initial presentation. The other 2 patients had a temporary pacing wire as a bridge between explantation of an infected device and reimplantation of a new device. Use of the temporary pacing wires has been identified as a risk factor for subsequent CDI in 2 recent studies (42,43). Based on the above findings, we recommend complete hardware removal to achieve CDI cure; this mirrors the recommendations of the American Heart Association (4).

Percutaneous versus surgical lead extraction. Extraction of cardiac device leads is not without risk (16,31,40). Despite advances in techniques for percutaneous lead extraction (44), there is still considerable morbidity and mortality associated with this procedure. In our study, 20 patients (11%) had complications of percutaneous lead extraction that included damage to cardiac valves, venous lacerations, bleeding complications, and lead tip fracture that required surgical intervention. Some investigators have expressed concern about transvenous lead removal when lead vegetation size is >1 cm because of the risk of pulmonary embolization of lead vegetation fragments (12,45-47). No patient had a clinically significant pulmonary embolism in our series despite the presence of vegetation sizes that ranged from 0.3 to 7.0 cm in the longest dimension. Others have reported similar experiences (22,23,48,49). Lead removal by cardiotomy (50,51) has been advocated to prevent symptomatic pulmonary embolism but, as shown in the current case series, this approach can be complicated by serious adverse events. Five patients suffered massive hemorrhages postoperatively, and 1 died in our series.

**Duration of antimicrobial treatment.** The duration of antimicrobial treatment for CDI depended on the clinical presentation and the causative agent. In a large case series from the Cleveland Clinic Foundation (19), the median duration of antibiotic treatment in CDI cases with pocket infection and those with bacteremia was 26 days and 41 days, respectively. However, in our practice, cases of CDI with only pocket infection were treated with 10 to 14 days of antimicrobials and those with BSI for 4 weeks after device removal. Similarly, patients with cardiac device-related endocarditis limited to the right heart can be treated with 4 weeks of antibiotics instead of the 6-week treatment course that has been advocated by some (12,13,19). Infection by certain microorganisms may require longer antimicrobial treatment for complete eradication of a CDI. In our study, cases of CDI attributable to *S. aureus* had

significantly longer treatment after device explantation as compared to those with CoNS infection (28 days vs. 14 days, p < 0.001). An *S. aureus* CDI is also associated with a higher mortality rate as compared with that caused by other pathogens. In our study, there were 9 in-hospital deaths (crude mortality rate 4.7%), and 5 had *S. aureus* identified in blood cultures.

Timing and need of a replacement device. Timing of reimplantation of a new device system after extraction of



#### Table 4 Mayo Clinic Guidelines for the Diagnosis and Management of Cardiac Device Infections

- 1. All patients should have at least 2 sets of blood cultures drawn at initial evaluation.
- 2. Generator tissue Gram stain and culture and lead tip culture should be obtained.
- 3. Patients who either have positive blood cultures or have negative blood cultures but had recent antibiotics before obtaining blood cultures should have a
- transesophageal echocardiogram (TEE) to assess for device-related endocarditis.
- 4. Sensitivity of TTE is low and is not recommended to evaluate for device-related endocarditis.
- 5. Patients with negative blood cultures and recent prior antibiotics and valve vegetations on TEE should be managed in consultation with an infectious diseases expert
- 6. All patients with device infection should undergo complete device removal, regardless of clinical presentation.
- 7. A large (>1 cm) lead vegetation is not a stand alone indication for surgical lead removal.
- 8. Blood cultures should be repeated in all patients after device explantation. Patients with persistently positive blood cultures should be treated for at least 4 weeks with antimicrobials even if TEE is negative for vegetations or other evidence of infection.
- Duration of antimicrobial therapy should also be extended to ≥4 weeks in patients with complicated infection (endocarditis, septic venous thrombosis, osteomyelitis, metastatic seeding).
- 10. Adequate debridement and control of infection should be achieved at all sites before reimplantation of a new device.
- **11**. Reevaluation for continued need of the device should be performed before new device placement.
- 12. If an infected cardiac device cannot be removed, than long-term suppressive antibiotic therapy should be administered after completing an initial course of treatment and securing a clinical response to therapy. Infectious diseases expert opinion should be sought.

an infected device remains a subject of debate and is influenced by the causative agent and clinical presentation of CDI. Median time from explantation to reimplantation of a new device was significantly longer for patients with BSI as compared to those with negative blood cultures (13 days vs. 7 days, p < 0.0001) in the current investigation. There was no statistically significant difference in time from explant to reimplant in patients with CoNS infection (median 7 days) as compared to those with S. aureus infection (median 12 days, p = 0.09). Although some investigators (13) have suggested delaying reimplantation of new device for 10 to 14 days in cases of pocket infection (without BSI) and up to 6 weeks in bacteremic patients, our data suggest that devices can be safely reimplanted once the pocket has been adequately debrided and blood cultures are negative. These data are consistent with previously published reports (19). Moreover, maintaining a transvenous temporary device increases the risk of subsequent CDI (43,52) and cannot be used indefinitely.

The need for placement of a new device system should be carefully assessed in all patients with CDI because an appreciable number of patients may not require a subsequent cardiac device. Discontinuation of electrophysiologic device use after removal of an infected system has been reported in 13% to 52% of cases (11,19,30,53). In our study, reimplantation of a new device was not required in onethird of patients.

**Proposed Mayo Clinic guidelines for management of CDI.** Based on the findings of our retrospective analysis and review of published literature (1–53), we propose guidelines for CDI (Figs. 3A and 3B, Table 4) to assist clinicians in patient management. These recommendations are not meant to replace individual patient management, however, and consultation with available specialists is advocated.

**Study limitations.** Our investigation has several limitations. First, retrospective studies have inherent limitations, such as potential selection and recall biases. Second, Mayo Clinic Rochester is a tertiary referral center with referral bias, as was shown in the current survey. Patients who are referred to a specialized cardiovascular center may be sicker and have more complex cardiac devices than the general population. Third, the majority of our cohort was white (91%), and this reflected both local population demographics and the referral population. Despite these limitations, data presented herein provide critical information to clinicians who manage CDI.

**Conclusions.** A CDI is an important and serious complication of device use. Because of the expected continued increase in the number of cases of CDI and the current lack of prospective, randomized trial data to assist in defining optimal treatment, we present proposed guidelines to assist clinicians in the management of CDI.

**Reprint requests and correspondence:** Dr. Muhammad R. Sohail, Department of Medicine, Division of Infectious Diseases, Tawam Hospital/Johns Hopkins Medicine, P.O. Box 15258, Al Ain, Abu Dhabi, United Arab Emirates. E-mail: msohail@ tawam-hosp.gov.ae.

#### REFERENCES

- 1. Uslan DZ, Baddour LM. Cardiac device infections: getting to the heart of the matter. Curr Opin Infect Dis 2006;19:345-8.
- Goldberger Z, Lampert R. Implantable cardioverter-defibrillators: expanding indications and technologies. JAMA 2006;295:809–18.
- Cabell CH, Heidenreich PA, Chu VH, et al. Increasing rates of cardiac device infections among Medicare beneficiaries: 1990–1999. Am Heart J 2004;147:582–6.
- Baddour LM, Bettmann MA, Bolger AF, et al. Nonvalvular cardiovascular device-related infections. Circulation 2003;108:2015–31.
- Mela T, McGovern BA, Garan H, et al. Long-term infection rates associated with the pectoral versus abdominal approach to cardioverter-defibrillator implants. Am J Cardiol 2001;88:750–3.
- Frame R, Brodman RF, Furman S, Andrews CA, Gross JN. Surgical removal of infected transvenous pacemaker leads. Pacing Clin Electrophysiol 1993;16:2343–8.
- Lai KK, Fontecchio SA. Infections associated with implantable cardioverter defibrillators placed transvenously and via thoracotomies: epidemiology, infection control, and management. Clin Infect Dis 1998;27:265–9.

- Smith PN, Vidaillet HJ, Hayes JJ, et al. Infections with nonthoracotomy implantable cardioverter defibrillators: can these be prevented? Endotak Lead Clinical Investigators. Pacing Clin Electrophysiol 1998;21:42–55.
- 9. Hill PE. Complications of permanent transvenous cardiac pacing: a 14-year review of all transvenous pacemakers inserted at one community hospital. Pacing Clin Electrophysiol 1987;10:564–70.
- Chu VH, Crosslin DR, Friedman JY, et al. Staphylococcus aureus bacteremia in patients with prosthetic devices: costs and outcomes. Am J Med 2005;118:1416.
- Cacoub P, Leprince P, Nataf P, et al. Pacemaker infective endocarditis. Am J Cardiol 1998;82:480–4.
- 12. Klug D, Lacroix D, Savoye C, et al. Systemic infection related to endocarditis on pacemaker leads: clinical presentation and management. Circulation 1997;95:2098–107.
- Darouiche RO. Treatment of infections associated with surgical implants. N Engl J Med 2004;350:1422–9.
- Ferguson TB Jr., Ferguson CL, Crites K, Crimmins-Reda P. The additional hospital costs generated in the management of complications of pacemaker and defibrillator implantations. J Thorac Cardiovasc Surg 1996;111:742–51, discussion 751–2.
- Arber N, Pras E, Copperman Y, et al. Pacemaker endocarditis. Report of 44 cases and review of the literature. Medicine 1994;73:299–305.
- Lee ME, Chaux A, Matloff JM. Avulsion of a tricuspid valve leaflet during traction on an infected, entrapped endocardial pacemaker electrode. The role of electrode design. J Thorac Cardiovasc Surg 1977;74:433–5.
- 17. Chamis AL, Peterson GE, Cabell CH, et al. Staphylococcus aureus bacteremia in patients with permanent pacemakers or implantable cardioverter-defibrillators. Circulation 2001;104:1029-33.
- Uslan DZ, Sohail MR, Friedman PA, et al. Frequency of permanent pacemaker or implantable cardioverter-defibrillator infection in patients with gram-negative bacteremia. Clin Infect Dis 2006;43:731–6.
- Chua JD, Wilkoff BL, Lee I, Juratli N, Longworth DL, Gordon SM. Diagnosis and management of infections involving implantable electrophysiologic cardiac devices. Ann Intern Med 2000;133:604–8.
- Durack DT, Lukes AS, Bright DK. New criteria for diagnosis of infective endocarditis: utilization of specific echocardiographic findings. Duke Endocarditis Service. Am J Med 1994;96:200–9.
- Sanfilippo AJ, Picard MH, Newell JB, et al. Echocardiographic assessment of patients with infectious endocarditis: prediction of risk for complications. J Am Coll Cardiol 1991;18:1191–9.
- del Rio A, Anguera I, Miro JM, et al. Surgical treatment of pacemaker and defibrillator lead endocarditis: the impact of electrode lead extraction on outcome. Chest 2003;124:1451–9.
- Victor F, De Place C, Camus C, et al. Pacemaker lead infection: echocardiographic features, management, and outcome. Heart 1999; 81:82–7.
- Orsinelli DA, Pearson AC. Detection of prosthetic valve strands by transesophageal echocardiography: clinical significance in patients with suspected cardiac source of embolism. J Am Coll Cardiol 1995;26: 1713–8.
- Klug D, Wallet F, Lacroix D, et al. Local symptoms at the site of pacemaker implantation indicate latent systemic infection. Heart 2004;90:882-6.
- Turkisher V, Priel I, Dan M. Successful management of an infected implantable cardioverter defibrillator with oral antibiotics and without removal of the device. Pacing Clin Electrophysiol 1997;20:2268–70.
- 27. Lee JH, Geha AS, Rattehalli NM, et al. Salvage of infected ICDs: management without removal. Pacing Clin Electrophysiol 1996;19: 437-42.
- Hurst LN, Evans HB, Windle B, Klein GJ. The salvage of infected cardiac pacemaker pockets using a closed irrigation system. Pacing Clin Electrophysiol 1986;9:785–92.
- O'Nunain S, Perez I, Roelke M, et al. The treatment of patients with infected implantable cardioverter-defibrillator systems. J Thorac Cardiovasc Surg 1997;113:121–9.
- Lewis AB, Hayes DL, Holmes DR Jr., Vlietstra RE, Pluth JR, Osborn MJ. Update on infections involving permanent pacemakers. Characterization and management. J Thorac Cardiovasc Surg 1985;89: 758–63.
- Bracke FA, Meijer A, van Gelder LM. Pacemaker lead complications: when is extraction appropriate and what can we learn from published data? Heart 2001;85:254–9.

- Rundstrom H, Kennergren C, Andersson R, Alestig K, Hogevik H. Pacemaker endocarditis during 18 years in Goteborg. Scand J Infect Dis 2004;36:674–9.
- Camus C, Leport C, Raffi F, Michelet C, Cartier F, Vilde JL. Sustained bacteremia in 26 patients with a permanent endocardial pacemaker: assessment of wire removal. Clin Infect Dis 1993;17: 46–55.
- Vogt PR, Sagdic K, Lachat M, Candinas R, von Segesser LK, Turina MI. Surgical management of infected permanent transvenous pacemaker systems: ten year experience. J Card Surg 1996;11:180–6.
- Molina JE. Undertreatment and overtreatment of patients with infected antiarrhythmic implantable devices. Ann Thorac Surg 1997;63: 504–9.
- 36. Love CJ, Wilkoff BL, Byrd CL, et al. Recommendations for extraction of chronically implanted transvenous pacing and defibrillator leads: indications, facilities, training. North American Society of Pacing and Electrophysiology Lead Extraction Conference Faculty. Pacing Clin Electrophysiol 2000;23:544–51.
- Klug D, Wallet F, Kacet S, Courcol RJ. Detailed bacteriologic tests to identify the origin of transvenous pacing system infections indicate a high prevalence of multiple organisms. Am Heart J 2005;149:322–8.
- Harjula A, Jarvinen A, Virtanen KS, Mattila S. Pacemaker infections—treatment with total or partial pacemaker system removal. Thorac Cardiovasc 1985;33:218–20.
- Parry G, Goudevenos J, Jameson S, Adams PC, Gold RG. Complications associated with retained pacemaker leads. Pacing Clin Electrophysiol 1991;14:1251–7.
- Rettig G, Doenecke P, Sen S, Volkmer I, Bette L. Complications with retained transvenous pacemaker electrodes. Am Heart J 1979;98: 587–94.
- Wilhelm MJ, Schmid C, Hammel D, et al. Cardiac pacemaker infection: surgical management with and without extracorporeal circulation. Ann Thorac Surg 1997;64:1707–12.
- Klug D, Balde M, Pavin D, et al. Prospective Evaluation of Pacemaker Lead Endocarditis (PEOPLE study): 12 months incidence and risk factors of pacemaker-related infections (abstr). Eur Heart J 2004; 25 Suppl:27.
- 43. Sohail MR, Khan AH, Freidman PA, et al. Electrophysiologic cardiac device infections: risk factor analysis using a case-control study. Presented at: Infectious Diseases Society of America (IDSA), San Francisco, CA: 2005, abstract no. 343.
- 44. Wilkoff BL, Byrd CL, Love CJ, et al. Pacemaker lead extraction with the laser sheath: results of the pacing lead extraction with the excimer sheath (PLEXES) trial. J Am Coll Cardiol 1999;33:1671–6.
- Voet JG, Vandekerckhove YR, Muyldermans LL, Missault LH, Matthys LJ. Pacemaker lead infection: report of three cases and review of the literature. Heart 1999;81:88–91.
- Dumont E, Camus C, Victor F, et al. Suspected pacemaker or defibrillator transvenous lead infection. Prospective assessment of a TEE-guided therapeutic strategy. Eur Heart J 2003;24:1779–87.
- Meier-Ewert HK, Gray ME, John RM. Endocardial pacemaker or defibrillator leads with infected vegetations: a single-center experience and consequences of transvenous extraction. Am Heart J 2003;146: 339–44.
- Nguyen KT, Neese P, Kessler DJ. Successful laser-assisted percutaneous extraction of four pacemaker leads associated with large vegetations. Pacing Clin Electrophysiol 2000;23:1260–2.
- Ruttmann E, Hangler HB, Kilo J, et al. Transvenous pacemaker lead removal is safe and effective even in large vegetations: an analysis of 53 cases of pacemaker lead endocarditis. Pacing Clin Electrophysiol 2006;29:231–6.
- 50. Kennelly BM, Piller LW. Management of infected transvenous permanent pacemakers. Br Heart J 1974;36:1133–40.
- Mansour KA, Kauten JR, Hatcher CR Jr. Management of the infected pacemaker: explantation, sterilization, and reimplantation. Ann Thorac Surg 1985;40:617–9.
- Klug D, Balde M, Pavin D, et al. Prospective evaluation of pacemaker lead endocarditis: 6 months preliminary results (abstr). Eur Heart J 2003;24:28.
- Bohm A, Banyai F, Preda I, Zamolyi K. The treatment of septicemia in pacemaker patients. Pacing Clin Electrophysiol 1996; 19:1105–11.