



# Invariant tensors and cellular categories

Bruce W. Westbury

*Mathematics Institute, University of Warwick, Coventry CV4 7AL, United Kingdom*

Received 31 January 2008

Available online 8 August 2008

Communicated by Andrew Mathas and Jean Michel

Dedicated to Gus Lehrer on the occasion of his 60th birthday

---

## Abstract

Let  $U$  be the quantised enveloping algebra associated to a Cartan matrix of finite type. Let  $W$  be the tensor product of a finite list of highest weight representations of  $U$ . Then  $\text{End}_U(W)$  has a basis called the dual canonical basis and this gives an integral form for  $\text{End}_U(W)$ . We show that this integral form is cellular by using results due to Lusztig.

© 2008 Elsevier Inc. All rights reserved.

*Keywords:* Cellular algebras; Quantum groups

---

## 1. Introduction

The definition of a cellular algebra was given in [GL96] and the examples of cellular algebras given were the Temperley–Lieb algebras, the Hecke algebras, the Brauer (or Birman–Wenzl) algebras and the Ariki–Koike algebras.

The aim of this paper is to give a general construction of cellular algebras using the dual canonical bases of spaces of invariant tensors defined in [Lus93, Part IV]. This construction includes the Temperley–Lieb algebras, the Hecke algebras, and the Birman–Wenzl algebras. This should not be confused with the observation made in [Gre99] that the results of [Lus93, Chapter 29] imply that the algebra  $\dot{U}$  and its finite-dimensional quotients  $\dot{U}/\dot{U}[P]$  are cellular.

The definition of a cellular algebra has a natural generalisation to cellular categories. An endomorphism algebra in a cellular category is cellular and many of the sequences of cellular algebras arise this way. Examples are the many diagram algebras such as the partition algebras,

---

*E-mail address:* [Bruce.Westbury@warwick.ac.uk](mailto:Bruce.Westbury@warwick.ac.uk).

the blob algebras, the affine Temperley–Lieb algebras, the contour algebras, the two-boundary Temperley–Lieb algebras.

Let  $U$  be the quantised enveloping algebra associated to a Cartan matrix of finite type and let  $V$  be a self-dual finite-dimensional highest weight representation. Then the category of invariant tensors has objects  $n = 0, 1, 2, \dots$  and the morphisms are  $\text{Hom}_U(\otimes^n V, \otimes^m V)$ . The examples which have been studied are

- [FK97], the representations of  $\mathfrak{sl}(2)$ ,
- [BW89], the vector representations of the symplectic and orthogonal Lie algebras,
- [Kup96], the fundamental representations of the rank two simple Lie algebras,
- [BD02], the adjoint representation of  $\mathfrak{sl}(n)$ ,
- [Wes08], the spin representation of  $\mathfrak{so}(7)$ .

In each case there is an integral form of the category of invariant tensors which is constructed using planar graphs and in each case this integral form can be seen to be cellular.

If  $V$  is not self-dual then we need to consider tensor products of copies of both  $V$  and  $V^*$ . The only case that has been studied is for  $V$  the vector representation of  $\mathfrak{sl}(n)$ . Then the walled Brauer algebras are endomorphism algebras in the category of invariant tensors.

In this paper we consider a larger category of invariant tensors. This is the full subcategory of the category of finite-dimensional representations of  $U$  whose objects are tensors products of highest weight representations. In this paper we observe that the results of [Lus93, Part IV] show that the dual canonical basis defines an integral form of this category and that this integral form is cellular. In particular this shows that if  $W$  is any tensor product of highest weight representations then the integral form of  $\text{End}_U(W)$  associated to the canonical basis is cellular.

This work was influenced by the principle that the diagram bases and the dual canonical bases have similar properties. In the examples in which the diagram bases have been defined it is straightforward to see that the category is cellular. It is then natural to look for a cellular structure in the dual canonical basis.

The diagram bases and dual canonical bases for  $\mathfrak{sl}(2)$  are discussed and shown to be the same in [FK97]. The diagram bases and the dual canonical bases for the category of invariant tensors of  $\mathfrak{sl}(3)$  are compared in [KK99] and shown to be different.

The diagram bases and the dual canonical bases have the following properties in common. They each define an integral form of the category of invariant tensors. They each admit a cellular structure with the same poset. In each case the tensor functor  $-\otimes-$  and the duality functor respect the cell structure. This makes it reasonable to conjecture that these categories are equivalent as integral forms and that the equivalences respect the cellular structure, the tensor structure and the duality.

## 2. Cellular categories

First we extend the definition of a cellular algebra given in [GL96] to cellular categories. The motivating examples are the Temperley–Lieb category discussed in [Wes95], and the categories whose endomorphism algebras are the sequences of algebras in the Introduction.

**Definition 2.1.** Let  $R$  be a commutative ring with identity. Let  $\mathcal{A}$  be an  $R$ -linear category with an anti-involution  $*$ . Then cell datum for  $\mathcal{A}$  consists of a partially ordered set  $\Lambda$ , a finite set  $K(n, \lambda)$

for each  $\lambda \in \Lambda$  and each object  $n$  of  $\mathcal{A}$ , and for  $\lambda \in \Lambda$  and  $n, m$  any two objects of  $\mathcal{A}$  we have an inclusion

$$C : K(n, \lambda) \times K(m, \lambda) \rightarrow \text{Hom}_{\mathcal{A}}(n, m),$$

$$C : (S, T) \mapsto C_{S,T}^\lambda.$$

The conditions that this datum is required to satisfy are:

C-1 For all objects  $n$  and  $m$ , the image of the map

$$C : \coprod_{\lambda \in \Lambda} K(n, \lambda) \times K(m, \lambda) \rightarrow \text{Hom}_{\mathcal{A}}(n, m)$$

is a basis for  $\text{Hom}_{\mathcal{A}}(n, m)$  as an  $R$ -module.

C-2 For all objects  $n, m$ , all  $\lambda \in \Lambda$  and all  $S \in K(n, \lambda), T \in K(m, \lambda)$  we have

$$(C_{S,T}^\lambda)^* = C_{T,S}^\lambda.$$

C-3 For all objects  $p, n, m$ , all  $\lambda \in \Lambda$  and all  $a \in \text{Hom}_{\mathcal{A}}(p, n), S \in K(n, \lambda), T \in K(m, \lambda)$  we have

$$aC^\lambda(S, T) = \sum_{S' \in K(p, \lambda)} r_a(S', S)C_{S',T}^\lambda \pmod{\mathcal{A}(< \lambda)}$$

where  $r_a(S', S) \in R$  is independent of  $T$  and  $\mathcal{A}(< \lambda)$  is the  $R$ -linear span of

$$\{C_{S,T}^\mu \mid \mu < \lambda; S \in K(p, \mu), T \in K(m, \mu)\}.$$

Another way of formulating C-3 is that for all objects  $p, n, m$ , all  $\lambda \in \Lambda$  and all  $S \in K(p, \lambda), T \in K(n, \lambda), U \in K(n, \lambda), V \in K(m, \lambda)$  we have

$$C^\lambda(S, T)C^\lambda(U, V) = \langle T, U \rangle C_{S,V}^\lambda \pmod{\mathcal{A}(< \lambda)}$$

where  $\langle T, U \rangle$  is independent of  $S$  and  $V$ .

A consequence of this definition is that for all order ideals  $\pi \subset \Lambda$  we have an ideal  $\mathcal{A}(\pi)$  where for all objects  $n, m$  the subspace  $\mathcal{A}(\pi)(n, m)$  is the  $R$ -linear span of

$$\{C_{S,T}^\mu \mid \mu \in \pi; S \in K(n, \mu), T \in K(m, \mu)\}.$$

This is a generalisation of the definition of a cellular algebra since we can regard any algebra over  $R$  as an  $R$ -linear category with one object.

More significantly, if  $\mathcal{A}$  is a cellular category then  $\text{End}(n)$  is a cellular algebra for any object  $n$  of  $\mathcal{A}$ .

A  $*$ -functor  $\phi : \mathcal{A} \rightarrow \mathcal{A}'$  between cellular categories is cellular if we have a map of partially ordered sets  $\phi : \Lambda \rightarrow \Lambda'$  such that for any order ideal  $\pi \in \Lambda$  we have  $\phi(\mathcal{A}(\pi)) \subset \mathcal{A}'(\phi(\pi))$ .

One reason for considering cellular categories instead of just the cellular endomorphism algebras is that for each object  $n$  and each order ideal  $\pi \subset \Lambda$  we have an  $R$ -linear functor  $\rho(n; \pi)$  from  $\mathcal{A}$  to the category of left  $\text{End}_{\mathcal{A}}(n)$ -modules which on objects is given by

$$\rho(n; \pi) : m \mapsto \text{Hom}_{\mathcal{A}(\pi)}(n, m).$$

Furthermore if  $\pi \subset \pi'$  then we have a natural transformation from  $\rho(n; \pi)$  to  $\rho(n; \pi')$ . This is used in [GL98].

### 3. Canonical bases

The rest of the paper is based on [Lus93, Part IV]. In particular all references are to this book and we have tried to use notation which is the same as or compatible with the notation in this book.

Let  $C$  be a Cartan matrix of finite type and let  $U$  be the Drinfeld–Jimbo quantum group associated to  $C$ . This is a Hopf algebra over the field  $\mathbb{Q}(v)$ . The integral weight lattice is denoted by  $X$  and the dominant integral weights are denoted by  $X^+$ . For each  $\lambda \in X^+$  there is a finite-dimensional representation of  $U$  with highest weight  $\lambda$ ; we denote this representation by  $V_\lambda$ . This representation has a canonical basis which we denote by  $B_\lambda$ . There is an anti-involution,  $\omega$ , of the  $\mathbb{Q}(v)$ -algebra  $U$ . This induces a duality on representations which we denote by  $M \mapsto {}^\omega M$ . Then  ${}^\omega V_\lambda \cong V_{-w_0\lambda}$  where  $w_0$  is the longest element of the Weyl group so we put  ${}^\omega \lambda = -w_0\lambda$ .

A based module is defined in [Lus93, 27.1.2]. Here we consider a category  $\mathcal{B}$ . The objects are the based modules. Let  $(M, B)$  and  $(M', B')$  be based modules. Then a morphism  $(M, B) \rightarrow (M', B')$  is a homomorphism of  $U$ -modules  $\phi : M \rightarrow M'$  which maps  $B$  to the  $\mathbb{Z}[v, v^{-1}]$ -module generated by  $B'$  and which is compatible with the involutions in the sense that  $\phi(\bar{m}) = \overline{\phi(m)}$  for all  $m \in M$ .

Put  $\delta = v + v^{-1}$ . Then observe that  $\mathbb{Z}[\delta]$  is the subring of  $\mathbb{Z}[v, v^{-1}]$  fixed by the bar involution. Then it is clear that  $\mathcal{B}$  is a  $\mathbb{Z}[\delta]$ -linear category. Then [Lus93, 27.3] constructs a tensor product of based modules and shows that this makes  $\mathcal{B}$  a monoidal category. Let  $(M, B)$  and  $(M', B')$  be based modules. Then the tensor product is the based module  $(M \otimes M', B \diamond B')$ . For each  $\lambda \in X^+$  we have a based module  $(V_\lambda, B_\lambda)$  where  $B_\lambda$  is the canonical basis of  $V_\lambda$ . Furthermore [Lus93, Proposition 21.1.2] says that there is also a based module  $({}^\omega V_\lambda, {}^\omega B_\lambda)$ . Let  $\mathcal{T}$  be the full subcategory of  $\mathcal{B}$  whose objects are finite tensor products of based modules isomorphic to these based modules. Then, by construction,  $\mathcal{T}$  is a  $\mathbb{Z}[\delta]$ -linear monoidal subcategory of  $\mathcal{B}$ . This category is also closed under the duality  $\omega$  and is in fact a spherical category, as defined in [BW99].

Now we can state the only result in this paper.

**Theorem 3.1.** *The  $\mathbb{Z}[\delta]$ -linear category  $\mathcal{T}$  with the anti-involution  $\omega$  is a cellular category.*

The remainder of this paper gives the proof of this result. First we give the cell datum. The poset  $(\Lambda, <)$  is the poset  $(X^+, >)$ .

If  $(M, B)$  is a based module [Lus93, 27.2.1] gives a partition of  $B$  into subsets  $B[\lambda]$ . Furthermore [Lus93, 27.2.5] shows that  $B[0]$  can be regarded as a base for the space of coinvariants of  $M$ . There is a canonical identification of  $\text{Hom}_U(M, M')$  with the dual of the space of coinvariants of  $M \otimes {}^\omega M'$ . Hence we may regard  $(B \diamond {}^\omega B')[0]$  as a basis of  $\text{Hom}(M, M')$ .

The subsets  $B[\lambda]^{hi} \subset B[\lambda]$  and  $B[\lambda]^{lo} \subset B[\lambda]$  are defined in [Lus93, 27.2.3] and [Lus93, Proposition 27.2.6] says that for  $\lambda = 0$  both of these inclusions are bijections.

Let  $(M, B)$  be an object of  $\mathcal{T}$ . Then we define the cells by

$$K((M, B), \lambda) = B[\lambda]^{hi}.$$

The map  $C$  is defined by

$$C : (b, b') \mapsto b \diamond {}^\omega b'.$$

This gives cell datum and it remains to check that this satisfies the conditions for a cellular category. The condition C-2 holds by construction.

The partition of  $(B \diamond B')[0]$  as

$$(B \diamond B')[0] = \bigcup_{\mu \in X^+} \{b \diamond b' \mid b \in B[-w_0(\mu)]^{lo}, b' \in B'[\mu]^{hi}\}$$

is given in [Lus93, Proposition 27.3.8]. This shows that condition C-1 holds.

It remains to discuss C-3. This condition holds modulo an ideal  $I$ . This means that, for any  $M$ , that we can replace  $M$  by  $M/MI$  and so we assume that  $MI = 0$ . Equivalently we assume that  $M[> \lambda] = 0$ .

By [Lus93, 27.2.4], each  $b \in B[\lambda]^{hi}$  gives a homomorphism of based modules  $|b\rangle : (M, B) \rightarrow (V_\lambda, B_\lambda)$ . Then  $b \diamond {}^\omega b' : (M, B) \rightarrow (M', B')$  is the composite of the two maps

$$|b\rangle : (M, B) \rightarrow (V_\lambda, B_\lambda) \quad \langle {}^\omega b' | : (V_\lambda, B_\lambda) \rightarrow (M', B')$$

and so we have  $b \diamond {}^\omega b' = |b\rangle \langle {}^\omega b' |$ .

Now  $\text{End}((V_\lambda, B_\lambda))$  is a free  $\mathbb{Z}[\delta]$ -module with basis the identity map  $1_\lambda$ .

## References

- [BD02] Georgia Benkart, Stephen Doty, Derangements and tensor powers of adjoint modules for  $\mathfrak{sl}_n$ , J. Algebraic Combin. 16 (1) (2002) 31–42.
- [BW89] Joan S. Birman, Hans Wenzl, Braids, link polynomials and a new algebra, Trans. Amer. Math. Soc. 313 (1) (1989) 249–273.
- [BW99] John W. Barrett, Bruce W. Westbury, Spherical categories, Adv. Math. 143 (2) (1999) 357–375.
- [FK97] Igor B. Frenkel, Mikhail G. Khovanov, Canonical bases in tensor products and graphical calculus for  $U_q(\mathfrak{sl}_2)$ , Duke Math. J. 87 (3) (1997) 409–480.
- [GL96] J.J. Graham, G.I. Lehrer, Cellular algebras, Invent. Math. 123 (1) (1996) 1–34.
- [GL98] J.J. Graham, G.I. Lehrer, The representation theory of affine Temperley–Lieb algebras, Enseign. Math. (2) 44 (3–4) (1998) 173–218.
- [Gre99] R.M. Green, Completions of cellular algebras, Comm. Algebra 27 (11) (1999) 5349–5366.
- [KK99] Mikhail Khovanov, Greg Kuperberg, Web bases for  $\mathfrak{sl}(3)$  are not dual canonical, Pacific J. Math. 188 (1) (1999) 129–153.
- [Kup96] Greg Kuperberg, Spiders for rank 2 Lie algebras, Comm. Math. Phys. 180 (1) (1996) 109–151.
- [Lus93] George Lusztig, Introduction to Quantum Groups, Progr. Math., vol. 110, Birkhäuser Boston Inc., Boston, MA, 1993.
- [Wes95] B.W. Westbury, The representation theory of the Temperley–Lieb algebras, Math. Z. 219 (4) (1995) 539–565.
- [Wes08] Bruce W. Westbury, Invariant tensors for the spin representation of  $\mathfrak{so}(7)$ , Math. Proc. Cambridge Philos. Soc. 144 (1) (2008) 217–240.