

JOURNAL OF APPROXIMATION THEORY 39, 361–373 (1983)

The Derivatives and Integrals of
Fractional Order in Walsh–Fourier Analysis,
with Applications to Approximation Theory

HE ZELIN

*Department of Mathematics, Nanjing University,
Nanjing, People's Republic of China*

Communicated by P. L. Butzer

Received January 26, 1982

In recent years the derivatives and integrals introduced by P. L. Butzer and H. J. Wagner [1] have been widely discussed in the literature. In Walsh–Fourier analysis and approximation theory they play a very important role. In 1977, C. W. Onneweer [4] introduced a kind of derivative of fractional order and obtained some interesting results, but they are not a generalization of the preceding derivatives. The aim of the present paper is to generalize the Butzer and Wagner derivatives and integrals to the case of fractional order. In this paper we give their definition and, under this new definition, establish some of the basic properties given in the case of integral order by Butzer and Wagner [1], Pal and Simon [6], etc. As applications to approximation theory, we generalize a Jackson–type theorem and Bernstein-type theorem given by Watarai [10], Butzer and Wagner [2, 8] to the case of fractional order, and estimate the degree of approximation by the typical means of the Walsh–Fourier series.

The generalized “Walsh functions” studied in this paper were also considered by Виленкин [13].

1. SYMBOLS

(a) Let $\mathbb{N} := \{1, 2, \dots\}$; $\mathbb{P} := \{0, 1, 2, \dots\}$; $m_j \in \mathbb{N} - \{1\}$, $j \in \mathbb{P}$, $\lim_{j \rightarrow \infty} m_j < \infty$; $M_0 := 1$, $M_r := m_0 m_1 \cdots m_{r-1}$, $r \in \mathbb{N}$; $\mathbb{Z}_j := \{0, 1, \dots, m_j - 1\}$; then each $x \in [0, 1)$ has a unique expansion $x = \sum_{j=1}^{\infty} x_j M_j^{-1}$ ($x_j \in \mathbb{Z}_j$), and each $k \in \mathbb{P}$ has a unique expansion $k = \sum_{j=0}^{\infty} k_j M_j$ ($k_j \in \mathbb{Z}_j$).

(b) Let $X = \sum_{j=1}^{\infty} x_j M_j^{-1}$, $y = \sum_{j=1}^{\infty} y_j M_j^{-1}$ ($x_j, y_j \in \mathbb{Z}_j$).

$$x \oplus y := \sum_{j=1}^{\infty} (x_j + y_j - \alpha_j) M_j^{-1},$$

where

$$\begin{aligned} a_j &= 0 && \text{if } x_j + y_i < m_j, \\ &= m_j && \text{if } x_j + y_i \geq m_j. \end{aligned}$$

If $m_j = m$, $j \in \mathbb{P}$, then the symbol \oplus is called the addition modulo m .

- (c) $\varphi_k(x) := \exp(2\pi i)/(m_k) \chi_{k+1}$ ($x \in [0, 1]$), $i = \sqrt{-1}$, $k \in \mathbb{P}$);
 $\psi_k(x) := \prod_{j=0}^{\lfloor x \rfloor} (\varphi_j(x))^k$; where $k = \sum_{j=0}^{\lfloor x \rfloor} k_j M_j$, $k_j \in \mathbb{Z}_j$. $D_n(x) := \sum_{j=0}^{n-1} \bar{\psi}_j(x)$,
 $F_n(x) := 1/n \sum_{j=1}^n D_j(x)$.
- (d) $X := X[0, 1] := WC[0, 1]$ or $L^p[0, 1]$ ($1 \leq p < \infty$).

$$\begin{aligned} \|f\|_X &:= \sup_{0 \leq x \leq 1} |f(x)|, && \text{if } X = WC, \\ &:= \left(\int_0^1 |f(x)|^p dx \right)^{1/p}, && \text{if } X = L^p, \end{aligned}$$

where $WC[0, 1] := \{f \mid \sup_{0 \leq x \leq 1} |f(x \oplus h) - f(x)| \rightarrow 0 \text{ as } h \rightarrow 0\}$.

- (e) If $f \in X[0, 1]$, $g \in L^1[0, 1]$, $(f * g)(x) := \int_0^1 f(x \oplus t) g(t) dt$.
- (f) For $f \in X[0, 1]$, let $f(k) := \int_0^1 f(t) \overline{\psi_k(t)} dt$.
- (g) $\omega(\delta) := \omega(f, \delta) := \omega(X, f, \delta) := \sup_{0 \leq h \leq \delta} \|f(\cdot \oplus h) - f(\cdot)\|_X$.
 $\text{Lip } \beta := \text{Lip}(X, \beta) := \{f \in X \mid \omega(X, f, \delta) = O(\delta^\beta), \delta \rightarrow 0\}$. $\text{lip } \beta := \text{lip}(X, \beta) := \{f \in X \mid \omega(X, f, \delta) = o(\delta^\beta), \delta \rightarrow 0\}$. $E_n(f) := E_n(X, f) := \inf_{P_n \in W_n} \|f - P_n\|_X$, where W_n denotes the set of all Walsh polynomials of order $\leq n$, i.e., $W_n := \{f \in L'[0, 1] \mid \hat{f}(k) = 0, k \geq n\}$.

2. DEFINITION AND PROPERTIES OF DERIVATIVE (INTEGRAL)

In [1], Butzer and Wagner gave the following definition of the dyadic derivative:

DEFINITION A. Let $f \in X[0, 1]$,

$$d_n f(x) = \frac{1}{2} \sum_{j=0}^n 2^j \left[f(x) - f\left(x \oplus \frac{1}{2^{j+1}}\right) \right], \quad (1)$$

where \oplus is the addition modulo 2; if there exists $g \in X[0, 1]$ such that $\lim_{n \rightarrow \infty} \|d_n f(\cdot) - g(\cdot)\|_X = 0$, then g is called the (strong) derivative of f .

In [7, 11] Zhen Weixing *et al.* gave the following definition of the m -adic derivative:

DEFINITION B. Let $f \in X[0, 1]$,

$$d_n f(x) = \sum_{k=0}^n m^k \sum_{j=0}^{m_j-1} a_j f(x \oplus jm^{-k-1}), \quad (2)$$

where \oplus is the addition modulo m , and

$$\begin{aligned} a_j &= \frac{m-1}{2}, & j &= 0, \\ &= \frac{1}{\exp \frac{-2\pi i}{m} j - 1}, & j &= 1, 2, \dots, m-1. \end{aligned}$$

If there exists $g \in X[0, 1]$ such that $\lim_{n \rightarrow \infty} \|d_n f(\cdot) - g(\cdot)\| = 0$, then g is called the (strong) derivative of f .

In [5] Onneweer gave the following definition of the $\{m_j\}$ -adic derivative:

DEFINITION C. Let $f \in X[0, 1]$,

$$d_n f(x) = \sum_{j=0}^n M_j \sum_{l=0}^{m_j-1} lm_j^{-1} \sum_{k=0}^{m_j-1} \overline{\varphi_j \left(\frac{k}{M_{j+1}} \right)}^l f \left(x \oplus \frac{k}{M_{j+1}} \right); \quad (3)$$

if there exists a g such that $\lim_{n \rightarrow \infty} \|d_n f(\cdot) - g(\cdot)\| = 0$, then g is called the (strong) derivative of f , and denoted by $D^{[1]} f$.

Formula (3) may be reduced to

$$d_n f(x) = \sum_{j=0}^{n-1} M_j \sum_{k=0}^{m_j-1} a_{jk} f \left(x \oplus \frac{k}{M_{j+1}} \right), \quad (4)$$

where

$$\begin{aligned} a_{jk} &= \frac{m_j - 1}{2}, & k &= 0, j = 0, 1, 2, \dots, n-1, \\ &= \frac{1}{\exp \frac{-2\pi i}{m_j} k - 1}, & k &= 1, 2, \dots, m_j - 1, j = 0, 1, \dots, n-1. \end{aligned}$$

In fact,

$$\begin{aligned} \sum_{l=0}^{m_j-1} l/m_j \overline{\varphi_j \left(\frac{k}{M_{j+1}} \right)}^l &= 1/m_j \sum_{l=0}^{m_j-1} le^{-(2\pi i/m_j)kl} = a_{jk}, \\ j &= 0, 1, \dots, n-1, \quad k = 0, 1, \dots, m_j - 1. \end{aligned}$$

Obviously, Definition B is more general than Definition A, and Definition C than Definition B.

In [1] Butzer and Wagner also gave a definition of the integral for the case $m_j = 2, j \in \mathbb{P}$; it is also suited to that of m_j in the general case. The integral of f can be defined as $I^{[1]}f := W_1 * f$, where $W_1 \in L'(0, 1)$ and

$$\begin{aligned} \hat{W}_1(k) &= 1 && \text{if } k = 0, \\ &= k^{-1} && \text{if } k \in \mathbb{N}. \end{aligned}$$

Now we give a definition of derivative and integral of fractional order, which is a generalization of the preceding definitions of derivative and integral of integral order.

To simplify our statements hereafter, in this paper we define $0^\alpha = 1$ if $\alpha \leq 0$.

DEFINITION 1. Let $f \in X(0, 1)$, $\alpha \in \mathbb{R}$, $T_r^{(\alpha)}(t) := \sum_{k=0}^{M_r-1} k^\alpha \overline{\psi_k(t)}$; if there exists $g \in X[0, 1]$ such that $\lim_{r \rightarrow \infty} \| (T_r^{(\alpha)} * f)(\cdot) - g(\cdot) \|_X = 0$, then if $\alpha > 0$, g is called the (strong) derivative of order α of f in $X[0, 1]$; if $\alpha < 0$, g is called the (strong) integral of order $(-\alpha)$ of f in $X[0, 1]$. In both cases, g will be denoted by $T^{(\alpha)}f$.

THEOREM 1.

- (1) $T^{(\alpha)}$ are linear operators for $\alpha \in \mathbb{R}$.
- (2) $T^{(\alpha)}\psi_k = k^\alpha \psi_k, \quad k \in \mathbb{N}$.
- (3) If $T^{(\alpha)}f \in X[0, 1]$, then $(T^{(\alpha)}f)^\wedge(k) = k^\alpha f^\wedge(k), k \in \mathbb{N}$.
- (4) If $\alpha > 0$, $T^{(\alpha)}f = 0 \Leftrightarrow f = \text{const.};$ if $\alpha = 0$, $T^{(\alpha)}f = 0 \Leftrightarrow f = 0$.

Proof. (1) and (2) are trivial in view of the definition.

(3) If $T^{(\alpha)}f \in X[0, 1]$, i.e., $\| (f * T_r^{(\alpha)})(\cdot) - T^{(\alpha)}f(\cdot) \|_X \rightarrow 0$ ($r \rightarrow \infty$), then $(T^{(\alpha)}f)^\wedge(k) = k^\alpha f^\wedge(k), k \in \mathbb{N}$.

(4) Assume $\alpha > 0$; if $T^{(\alpha)}f = 0$, then $k^\alpha f^\wedge(k) = 0, k \in \mathbb{N}$, i.e., $f^\wedge(k) = 0, k \in \mathbb{N}$, thus $f = \text{const.}$ Conversely, if $f = \text{const.}$, then by definition, $T^{(\alpha)}f = 0$. For $\alpha < 0$, the proof is analogous.

LEMMA 1. If $\alpha < 0$, then $T_r^{(\alpha)}(t)$ converges to a function $T_\infty^{(\alpha)}(t)$ in $X[0, 1]$, and

$$(T_\infty^{(\alpha)})^\wedge(k) = k^\alpha, \quad k \in \mathbb{N}; \quad \| T_r^{(\alpha)}(\cdot) - T_\infty^{(\alpha)}(\cdot) \|_{L^1} = O(M_r^\alpha) \quad (r \rightarrow \infty).$$

Proof. Applying Abel's transform twice, we get, assuming $s > r \geq 0$,

$$\begin{aligned} T_s^{(\alpha)}(t) - T_r^{(\alpha)}(t) &= \sum_{k=M_r}^{M_s-1} k^\alpha \psi_k(t) = \sum_{k=M_r}^{M_s-3} [k^\alpha - 2(k+1)^\alpha + (k+2)^\alpha](k+1) F_{k+1}(t) \\ &\quad - [M_r^\alpha - (M_r+1)^\alpha] M_r F_{M_r}(t) \\ &\quad + [(M_s-2)^\alpha - (M_s-1)^\alpha] (M_s-1) F_{M_s-1}(t) \\ &\quad - M_r^\alpha D_{M_r}(t) + (M_s-1)^\alpha D_{M_s}(t). \end{aligned}$$

On the other hand (see [6]), since

$$\begin{aligned} \|D_{M_r}(\cdot)\|_{L^1} &= 1; \quad \|F_k(\cdot)\|_{L^1} = O(1); \\ M_r^\alpha - (M_r+1)^\alpha &= O(M_r^{\alpha-1}); \quad (M_s-2)^\alpha - (M_s-1)^\alpha = O(M_s^{\alpha-1}); \\ \sum_{k=M_r}^{M_s-3} [k^\alpha - 2(k+1)^\alpha + (k+2)^\alpha](k+1) &= O(1) \sum_{k=M_r}^{M_s-3} k^{\alpha-1} = O(M_r^\alpha). \end{aligned}$$

Thus,

$$\|T_s^{(\alpha)}(\cdot) - T_r^{(\alpha)}(\cdot)\|_{L^1} = O(M_r^\infty) \rightarrow 0 \quad \text{as } r, s \rightarrow \infty. \quad (5)$$

Therefore by the completeness of $L'[0, 1]$, there exists $T_\infty^{(\alpha)}(t) \in L'[0, 1]$ such that

$$\|T_r^{(\alpha)}(\cdot) - T_\infty^{(\alpha)}(\cdot)\|_{L^1} \rightarrow 0 \quad \text{as } r \rightarrow \infty. \quad (6)$$

It is obvious by (5) and (6) that $(T_\infty^{(\alpha)})^\wedge(k) = k^\alpha$ and $\|T_r^{(\alpha)}(\cdot) - T_\infty^{(\alpha)}(\cdot)\|_{L^1} = O(M_r^\alpha)$.

The following simple lemma is a counterpart of Lemma 2 in [2]; it plays an important role in this paper.

LEMMA 2. *Assume $f \in X(0, 1)$ and let $W(t)$ be a Walsh polynomial of order $\leq M_r$; then*

$$\left\| \int_0^1 f(\cdot \oplus t) \overline{\phi_r(t)}^k W(t) dt \right\|_X = \omega \left(X, f, \frac{1}{M_r} \right) \|W(t)\|_{L^1}, \quad k \in \{1, 2, \dots, m_r - 1\}.$$

Proof. Let $x \in [0, 1), j \in \mathbb{Z}_r$, then

$$\begin{aligned} I &= \int_0^1 f(x \oplus t) \overline{\phi_r(t)}^k W(t) dt \\ &= \int_0^1 f \left(x \oplus t \oplus \frac{j}{M_{r+1}} \right) \overline{\phi_r \left(t \oplus \frac{j}{M_{r+1}} \right)}^k W \left(t \oplus \frac{j}{M_{r+1}} \right) dt \\ &= \int_0^1 f \left(x \oplus t \oplus \frac{j}{M_{r+1}} \right) \overline{\phi_r(t)}^k e^{-j2\pi i/m_r} W(t) dt, \end{aligned}$$

i.e.,

$$I = \int_0^1 \frac{1}{m_r} \sum_{j=0}^{m_r-1} e^{-(2\pi i/m_r)jk} \overline{\varphi_r(t)}^k f\left(x \oplus t \oplus \frac{j}{M_{n+1}}\right) W(t) dt.$$

Thus,

$$\begin{aligned} \|I\|_X &\leq \frac{1}{m_r} \left\| \sum_{j=0}^{m_r-1} e^{-(2\pi i/m_r)jk} f\left(\cdot \oplus \frac{j}{M_{r+1}}\right) \right\|_X \|W(t)\|_{L^1} \\ &= \frac{1}{m_r} \left\| \sum_{j=0}^{m_r-1} e^{-(2\pi i/m_r)jk} \left[f\left(\cdot \oplus \frac{j}{M_{r+1}}\right) - f(\cdot) \right] \right\|_X \|W(t)\|_{L^1} \\ &\leq \omega\left(X, f, \frac{1}{M_r}\right) \|W\|_{L^1}. \end{aligned}$$

LEMMA 3. Assume $f \in X[0, 1]$, $\alpha \geq 0$, $s > r \geq 0$; then $\|T_s^{(\alpha)} * f - T_r^{(\alpha)} * f\|_X = O(1) \sum_{l=r}^{s-1} \omega(f, 1/M_l) M_l^\alpha$. In particular, $\|T_s^{(\alpha)} * f\|_X = O(1) \sum_{l=0}^{s-1} \omega(f, 1/M_l) M_l^\alpha$.

Proof.

$$\begin{aligned} &\|T_s^{(\alpha)} * f - T_r^{(\alpha)} * f\|_X \\ &= \left\| \int_0^1 f(\cdot \oplus t) \sum_{l=r}^{s-1} \sum_{j=1}^{m_l-1} \sum_{k=jM_l}^{(j+1)M_l-1} k^\alpha \overline{\psi_k(t)} dt \right\|_X \\ &= \left\| \int_0^1 f(\cdot \oplus t) \sum_{l=r}^{s-1} \sum_{j=1}^{m_l-1} \sum_{k=0}^{M_l-1} (jM_l + k)^\alpha \overline{\psi_{jM_l+k}(t)} dt \right\|_X \\ &= \left\| \sum_{l=r}^{s-1} \sum_{j=1}^{m_l-1} \int_0^1 f(\cdot \oplus t) \overline{\varphi_l(t)}^j \sum_{k=0}^{M_l-1} (jM_l + k)^\alpha \overline{\psi_k(t)} dt \right\|_X \\ &\leq \sum_{l=r}^{s-1} \sum_{j=1}^{m_l-1} \omega\left(f, \frac{1}{M_l}\right) \left\| \sum_{k=0}^{M_l-1} (jM_l + k)^\alpha \overline{\psi_k(t)} \right\|_{L^1}. \end{aligned}$$

By applying Abel's transform twice, we have

$$\begin{aligned} &\left\| \sum_{k=0}^{M_l-1} (jM_l + k)^\alpha \overline{\psi_k(t)} \right\|_{L^1} \\ &= \left\| \sum_{k=0}^{M_l-3} [(jM_l + k)^\alpha - 2(jM_l + k + 1)^\alpha + (jM_l + k + 2)^\alpha] (k+1) F_{k+1}(t) \right. \\ &\quad \left. + [(j+1)M_l - 2]^\alpha - [(j+1)M_l - 1]^\alpha (M_l - 1) F_{M_l-1}(t) \right. \\ &\quad \left. - [(j+1)M_l - 1]^\alpha D_{M_l}(t) \right\|_{L^1} = O(M_l^\alpha). \end{aligned} \tag{7}$$

Therefore, $\|T_s^{(\alpha)} * f - T_r^{(\alpha)} * f\|_X = O(1) \sum_{l=r}^{s-1} \omega(f, 1/M_l) M_l^\alpha$.

LEMMA 4. If $f \in X[0, 1]$, $\alpha > 0$, $M_s \leq n < M_{s+1}$, $g(t) = \sum_{k=a_s M_s}^{n-1} (n^\alpha - k^\alpha) \psi_k(t)$, then $\|(g * f)(\cdot)\|_X = O(1) \omega(X, f, 1/M_s) M_s^\alpha$.

Proof. Let $n = \sum_{j=0}^s a_j M_j$ ($a_j \in \mathbb{Z}_j, j = 0, 1, \dots, s, a_s \neq 0$). Then

$$\begin{aligned} g(t) &= \sum_{k=a_s M_s}^{n-1} (n^\alpha - k^\alpha) \overline{\psi_k(t)} \\ &= \sum_{r=1}^s \sum_{j=0}^{a_{s-r}-1} a_s M_s + a_{s-1} M_{s-1} + \dots + a_{s-r+1} M_{s-r+1} + (j+1) M_{s-r-1} (n^\alpha - k^\alpha) \overline{\psi_k(t)} \\ &= \sum_{r=1}^s \overline{\varphi_s(t)}^{a_s} \cdots \overline{\varphi_{s-r+1}(t)}^{a_{s-r+1}} \sum_{j=0}^{a_{s-r}-1} \overline{\varphi_{s-r}(t)}^j \\ &\quad \times \sum_{k=0}^{M_{s-r}-1} |n^\alpha - (a_s M_s + \dots + a_{s-r+1} M_{s-r+1} + j M_{s-r} + k)^\alpha| \overline{\psi_k(t)}. \end{aligned}$$

By Lemma 2, we have

$$\begin{aligned} &\|(g * f)(t)\|_X \\ &\leq \omega \left(X, f, \frac{1}{M_s} \right) \sum_{r=1}^s \sum_{j=0}^{a_{s-r}-1} \\ &\quad \times \left\| \sum_{k=0}^{M_{s-r}-1} |n^\alpha - (a_s M_s + \dots + a_{s-r+1} M_{s-r+1} + j M_{s-r} + k)^\alpha| \overline{\psi_k(t)} \right\|_{L^1} \\ &= \omega \left(X, f, \frac{1}{M_s} \right) \sum_{r=1}^s \sum_{j=0}^{a_{s-r}-1} \\ &\quad \times \left\| \sum_{k=0}^{M_{s-r}-1} |2(a_s M_s + \dots + a_{s-r+1} M_{s-r+1} + j M_{s-r} + k - 1)^\alpha \right. \\ &\quad \left. - (a_s M_s + \dots + a_{s-r+1} M_{s-r+1} + j M_{s-r} + k)^\alpha \right. \\ &\quad \left. - (a_s M_s + \dots + a_{s-r+1} M_{s-r+1} + k + 2)^\alpha |(k+1) F_{k+1}(t) \right. \\ &\quad \left. + |(a_s M_s + \dots + a_{s-r+1} M_{s-r+1} + (j+1) M_{s-r} - 1)^\alpha \right. \\ &\quad \left. - (a_s M_s + \dots + a_{s-r+1} M_{s-r+1} + (j+1) M_{s-r} - 2)^\alpha |(M_{s-r} - 1) F_{M_{s-r}}(t) \right. \\ &\quad \left. + |n^\alpha - (a_s M_s + \dots + a_{s-r+1} M_{s-r+1} + (j+1) M_{s-r} - 1)^\alpha| D_{M_{s-r}}(t) \right\|_{L^1} \\ &= O(1) \omega \left(X, f, \frac{1}{M_s} \right) \sum_{r=1}^s (M_s^{\alpha-2} M_{s-r}^2 + M_s^{\alpha-1} M_{s-r} + M_s^{\alpha+1} M_{s-r}) \\ &= O(1) \omega \left(X, f, \frac{1}{M_s} \right) M_s^\alpha. \end{aligned}$$

LEMMA 5. If $T^{(\alpha)}f \in X[0, 1]$, $\alpha > 0$, then

$$(1) \quad \omega(f, 1/M_r) = O(1) 1/M_r^\alpha \omega(T^{(\alpha)}f, 1/M_r).$$

$$(2) \quad \omega(T^{(\alpha)}f, 1/M_r) = O(1) \sum_{l=r}^{\infty} \omega(f, 1/M_l) M_l^\alpha.$$

Proof. (1) Let $0 \leq h < 1/M_r$. By Theorem 1 (3), we have

$$|f(x \oplus h) - f(x)|^{\hat{}}(k) = |(T_{\infty}^{(-\alpha)} - T_r^{(-\alpha)}) * (T^{(\alpha)}f(x \oplus h) - T^{(\alpha)}f(x))|^{\hat{}}(k), \quad k \in \mathbb{P},$$

i.e.,

$$f(x \oplus h) f(x) = (T_{\infty}^{(-\alpha)} - T_r^{(-\alpha)}) * (T^{(\alpha)}f(x \oplus h) - T^{(\alpha)}f(x)).$$

Therefore, by Lemma 1, we have $\omega(f, 1/M_r) = O(1)(1/M_r^\alpha) \omega(T^{(\alpha)}f, 1/M_r)$.

(2) Since $\omega(T^{(\alpha)}f, 1/M_r) \leq 2E_{M_r}(T^{(\alpha)}f) \leq 2 \|T^{(\alpha)}f - T_r^{(\alpha)} * f\|_X \leq 2 \|T^{(\alpha)}f - T_s^{(\alpha)} * f\|_X + 2 \|T_s^{(\alpha)} * f - T_s^{(\alpha)} * f - T_r^{(\alpha)} * f\|_X$, by Lemma 3, we get for $s \rightarrow \infty$. $\omega(T^{(\alpha)}f, 1/M_r) = O(1) \sum_{l=r}^{\infty} \omega(f, 1/M_l) M_l^\alpha$.

LEMMA 6. If for $f \in X[0, 1]$ there exists $g \in X[0, 1]$ such that $k^\alpha f^{\hat{}}(k) = g^{\hat{}}(k)$, $\alpha > 0$, $k \in \mathbb{P}$, then $f = T_{\infty}^{(-\alpha)} * g + f^{\hat{}}(0)$.

Proof. Since $(T_{\infty}^{(-\alpha)} * g)^{\hat{}}(k) = k^{-\alpha} k^\alpha f^{\hat{}}(k)$, we have $f = T_{\infty}^{(-\alpha)} * g + f^{\hat{}}(0)$.

THEOREM 2. If $a < 0$, $f \in X[0, 1]$, then $T^{(a)}f \in X[0, 1]$ and $T^{(a)}f = T^{(a)}f = T_{\infty}^{(a)} * f$.

Proof. By Lemma 1, $T_{\infty}^{(a)} \in L^1[0, 1]$, thus $T_{\infty}^{(a)} * f \in X[0, 1]$. Moreover,

$$\|T_r^{(a)} * f - T_{\infty}^{(a)} * f\|_X \leq \|T_r^{(a)} - T_{\infty}^{(a)}\|_{L^1} \|f\|_X = O(M_r^\alpha) \rightarrow 0 \quad (r \rightarrow \infty).$$

This completes the proof.

By this theorem and Lemma 1 we get immediately

COROLLARY 2.1. $T^{(-1)}f = I^{(1)}f$, where $I^{(1)}$ is the integral operator introduced by Butzer and Wagner.

THEOREM 3. If $\alpha > 0$, $U^{(\alpha)} := \{f \in X[0, 1] \mid T^{(\alpha)}f \in X[0, 1]\}$;

$$U_1^{(\alpha)} := \left\{ f \in X[0, 1] \mid \omega \left(f, \frac{1}{M_l} \right) = O \left(\frac{a_l}{M_l^\alpha} \right) \right\}, \quad \text{where } a_l > 0, \sum_{l=1}^{\infty} a_l < \infty;$$

$$U_2^{(\alpha)} := \left\{ f \in X[0, 1] \mid \omega \left(f, \frac{1}{M_l} \right) = O(M_l^{-\alpha}) \right\},$$

then $U_1^{(\alpha)} \subset U^{(\alpha)} \subset U_2^{(\alpha)}$.

Proof. Suppose $f \in U_1^{(\alpha)}$. By Lemma 3, we have $\|T_s^{(\alpha)} * f - T_r^{(\alpha)} * f\|_X = O(1) \sum_{l=r}^{s-1} \omega(f, 1/M_l) M_l^\alpha = O(1) \sum_{l=r}^{s-1} a_l \rightarrow 0$ ($r, s \rightarrow \infty$). By the completeness of $X[0, 1]$, there exists $g \in X[0, 1]$ such that $\|T_r^{(\alpha)} * f - g\|_X \rightarrow 0$ ($r \rightarrow \infty$), i.e., $f \in U^{(\alpha)}$.

Suppose $f \in U^{(\alpha)}$. By Lemma 5, we have $\omega(f, 1/M_r) = O(1)(1/M_r^\alpha)$ $\omega(T^{(\alpha)}f, 1/M_r) = O(1/M_r^\alpha)$, i.e., $f \in U_2^{(\alpha)}$.

COROLLARY 3.1. If $0 < \alpha < \beta$, then $U^{(\beta)} \subset \text{lip } \beta \subset U^{(\alpha)} \subset \text{lip } \alpha$.

THEOREM 4. Let $f \in X[0, 1]$, $\int_0^1 f(x) dx = 0$. If one of the following two conditions holds, then $T^{(\alpha)} T^{(\beta)} f = T^{(\alpha+\beta)} f$.

- (1) $\alpha \leq 0$ and $T^{(\beta)} f \in X[0, 1]$.
- (2) $\alpha > 0$ and $T^{(\alpha+\beta)} f \in X[0, 1]$ or $T^{(\beta)} f, T^{(\alpha)} T^{(\beta)} f \in X[0, 1]$

(cf. Theorem 3 and Corollary 4 in [4].)

Proof. (1) If $\alpha \leq 0$, $T^{(\beta)} f \in X[0, 1]$, then by Theorem 2 and Corollary 3.1, we have $T^{(\alpha)} T^{(\beta)} f, T^{(\alpha+\beta)} f \in X[0, 1]$. Therefore, since $(T^{(\alpha)} T^{(\beta)} f) \hat{\cdot}(k) = k^{\alpha+\beta} f \hat{\cdot}(k) = (T^{(\alpha+\beta)} f) \hat{\cdot}(k)$, $k \in \mathbb{P}$, we have $T^{(\alpha)} T^{(\beta)} f = T^{(\alpha+\beta)} f$.

(2) If $\alpha > 0$, $T^{(\alpha+\beta)} f \in X[0, 1]$, then $T^{(\beta)} f \in X[0, 1]$, thus by $T_r^{(\alpha)} * T^{(\beta)} f = T_r^{(\alpha+\beta)} f$ and $T^{(\alpha+\beta)} f \in X[0, 1]$, we get $T^{(\alpha)} T^{(\beta)} f \in X[0, 1]$. If $T^{(\beta)} f, T^{(\alpha)} T^{(\beta)} f \in X[0, 1]$, then by $T_r^{(\alpha)} * T^{(\beta)} f = T_r^{(\alpha+\beta)} f$, we know $T^{(\alpha+\beta)} f \in X[0, 1]$. Therefore, since $(T^{(\alpha)} T^{(\beta)} f) \hat{\cdot}(k) = k^{\alpha+\beta} f \hat{\cdot}(k) = (T^{(\alpha+\beta)} f) \hat{\cdot}(k)$, we get $T^{(\alpha)} T^{(\beta)} f = T^{(\alpha+\beta)} f$.

THEOREM 5. If $\alpha > 0$ and $f \in X[0, 1]$, the following statements are equivalent:

- (1) $T^{(\alpha)} f = g \in X[0, 1]$.
- (2) There exists $g \in X[0, 1]$ such that $\hat{g}(k) = k^\alpha \hat{f}(k)$, $k \in \mathbb{P}$.
- (3) There exists $g \in X[0, 1]$ such that $f = T^{(-\alpha)} g + \hat{f}(0)$

(cf. Corollary 1 and 3 in [4]).

Proof. Assertion (2) follows from (1) by Theorem 1, (3) follows from (2) by Lemma 6, and (1) follows from (3) by Theorem 4.

COROLLARY 5.1. $T^{(1)} f \in X[0, 1] \Leftrightarrow D^{(1)} f \in X[0, 1]$; in this event $T^{(1)} f = D^{(1)} f$.

Proof. By Theorem 5 $T^{(1)} f = g \in X[0, 1]$ is equivalent to $\hat{g}(k) = k \cdot \hat{f}(k)$ ($k \in \mathbb{P}$). On the other hand it is proved in [6] that the last equality is equivalent to $D^{(1)} f = g \in X[0, 1]$.

3. APPLICATIONS

First we give a generalization of the Jackson-type and Bernstein-type theorems given in the case $\alpha = 0$ by Watari [10], and $\alpha \in \mathbb{N}$ by Butzer and Wagner [2, 8].

THEOREM 6. *If $T^{(\alpha)}f \in \text{Lip}(X, \beta)$, $\alpha \geq 0$, $\beta > 0$, then $E_n(X, f) = O(1/n^{\alpha+\beta})$.*

Proof. Let $M_r \leq n < M_{r+1}$; then by [14] and Lemma 5 we have

$$\begin{aligned} E_n(f) &\leq E_{M_r}(f) \leq \omega\left(f, \frac{1}{M_r}\right) = O(1) \frac{1}{M_r^\alpha} \omega\left(T^{(\alpha)}f, \frac{1}{M_r}\right) \\ &= O\left(\frac{1}{M_r^{\alpha+\beta}}\right) = O\left(\frac{1}{n^{\alpha+\beta}}\right). \end{aligned}$$

THEOREM 7. *If $f \in X[0, 1]$, $E_n(X, f) = O(1/n^\beta)$, $\beta > \alpha \geq 0$, then $T^{(\alpha)}f \in \text{Lip}(X, \beta - \alpha)$.*

Proof. Let $M_r \leq n < M_{r+1}$; then $\omega(f, 1/M_r) \leq 2E_{M_r}(f) = O(1/M_r^\beta)$. By Corollary 3.1, we have $T^{(\alpha)}f \in X[0, 1]$. Moreover, by Lemma 5,

$$\begin{aligned} \omega\left(T^{(\alpha)}, \frac{1}{M_r}\right) &= O(1) \sum_{l=r}^{\alpha} \omega\left(f, \frac{1}{M_l}\right) M_l^\alpha = O(1) \sum_{l=r}^{\alpha} E_{M_l}(f) M_l^\alpha \\ &= O(1) \sum_{l=r}^{\alpha} M_l^{\alpha+\beta} = O\left(\frac{1}{M_r^{\beta-\alpha}}\right). \end{aligned}$$

For any $\delta > 0$, let $1/M_{r+1} \leq \delta < 1/M_r$; then $\omega(T^{(\alpha)}f, \delta) \leq \omega(T^{(\alpha)}f, 1/M_r) = O(1/M_r^{\beta-\alpha}) = O(\delta^{\beta-\alpha})$, i.e., $T^{(\alpha)}f \in \text{Lip}(X, \beta - \alpha)$.

Below we discuss the degree of approximation by the typical means of the Walsh–Fourier series.

DEFINITION 2. Let $f \in X[0, 1]$, $K_{n,\lambda}(t) = \sum_{k=0}^{n-1} |1 - (k/n)^\lambda| \overline{\psi_k(t)}$, $\lambda > 0$, then

$$R_{n,\lambda}(f, x) = f * K_{n,\lambda}(x) = \int_0^1 f(x \oplus t) K_{n,\lambda}(t) dt$$

are called the typical means of the Walsh–Fourier series of f .

THEOREM 8. *If $T^{(\alpha)}f \in X[0, 1]$, $\alpha \geq 0$, $\lambda > 0$, then, for $M_s \leq n < M_{s+1}$, $\|R_{n,\lambda}(f, \cdot) - f(\cdot)\|_X = O(1)(1/n^\lambda) \sum_{l=0}^s \omega(T^{(\alpha)}f, 1/M_l) M_l^{\lambda-\alpha}$.*

Proof. Since

$$\begin{aligned}
& \|R_{n,\lambda}(f, \cdot) - f(\cdot)\|_X \\
& \leq \left\| f * \sum_{k=0}^{M_s-1} \bar{\psi}_k - f \right\|_X + \left\| f * \sum_{k=M_s}^{a_s M_s - 1} \bar{\psi}_k \right\|_X + \frac{1}{n^\lambda} \left\| f * \sum_{k=0}^{M_s-1} k^\lambda \bar{\psi}_k \right\|_X \\
& \quad + \frac{1}{n^\lambda} \left\| f * \sum_{k=M_s}^{a_s M_s - 1} k^\lambda \bar{\psi}_k \right\|_X + \frac{1}{n^\lambda} \left\| f * \sum_{k=M_s}^{a_s M_s - 1} (n^\lambda - k^\lambda) \bar{\psi}_k \right\|_X \\
& \leq \omega \left(\frac{1}{M_s} \right) + \left\| f * \sum_{k=M_s}^{a_s M_s - 1} \bar{\psi}_k \right\|_X + \frac{O(1)}{n^\lambda} \sum_{l=0}^{s-1} \omega \left(f, \frac{1}{M_l} \right) M_l^\lambda \\
& \quad + \frac{1}{n^\lambda} \left\| f * \sum_{k=M_s}^{a_s M_s - 1} k^\lambda \bar{\psi}_k \right\|_X + O(1) \omega \left(\frac{1}{M_s} \right), \\
& \left\| f * \sum_{k=M_s}^{a_s M_s - 1} \bar{\psi}_k \right\|_X \\
& = \left\| f * \left(\sum_{j=1}^{a_s - 1} \bar{\phi}_s^j \right) D_{M_s} \right\|_X \leq (a_s - 1) \omega \left(\frac{1}{M_s} \right),
\end{aligned}$$

by (7), as well as

$$\begin{aligned}
& \left\| \sum_{k=M_s}^{a_s M_s - 1} k^\lambda \bar{\psi}_k \right\|_X = \left\| f * \sum_{j=1}^{a_s - 1} \sum_{k=j M_s}^{(j+1) M_s - 1} k^\lambda \bar{\psi}_k \right\|_X \\
& = \left\| \sum_{j=1}^{a_s - 1} f * \bar{\phi}_s^j \sum_{k=0}^{M_s-1} (j M_s + k)^\lambda \psi_k \right\|_X \\
& = O(1) \omega(1/M_s) M_s^\lambda,
\end{aligned}$$

it follows that $\|R_{n,\lambda}(f, \cdot) - f(\cdot)\|_X = (O(1)/n^\lambda) \sum_{l=0}^s \omega(f, 1/M_l) M_l^\lambda = (O(1)/n^\lambda) \sum_{l=0}^s \omega(T^{(\alpha)} f, 1/M_l) M_l^{\lambda-\alpha}$.

THEOREM 9. If $f \in X[0, 1]$, $\lambda > 0$, then $\lambda_n = \|R_{n,\lambda}(f, \cdot) - f(\cdot)\|_X \rightarrow 0$ ($n \rightarrow \infty$).

Proof. Let $M_s \leq n < M_{s+1}$. Since $(1/n^\lambda) \sum_{l=0}^s M_l^\lambda \leq \sum_{l=0}^s (M_l/M_s)^\lambda \leq 1 + (1/2^\lambda) + (1/4^\lambda) + \dots + (1/2^{s\lambda}) \leq c < \infty$, we have $\Delta_n = O(1)(1/n^\lambda) \sum_{l=0}^s \omega(f, 1/M_l) M_l^\lambda = O(1)(1/\sum_{l=0}^s M_l^\lambda) \sum_{l=0}^s \omega(f, 1/M_l) M_l^\lambda$. It is easy to see that the transformation of the sequence $\{\omega(1/M_l)\}$ into the sequence $\{(1/\sum_{l=0}^s M_l^\lambda) \sum_{l=0}^s \omega(f, 1/M_l) M_l^\lambda\}$ is regular. Therefore, since $\omega(1/M_l) \rightarrow 0$ ($l \rightarrow \infty$), we get $\Delta_n \rightarrow 0$ ($n \rightarrow \infty$).

THEOREM 10. *If $T^{(\alpha)}f \in \text{Lip}(X, \beta)$, $\alpha \geq 0$, $\beta > 0$, $\lambda > 0$, then*

$$\begin{aligned} \|R_{n,\lambda}(f, \cdot) - f(\cdot)\|_X &= O\left(\frac{1}{n^{\alpha+\beta}}\right) \quad \text{if } \alpha + \beta < \lambda, \\ &= O\left(\frac{\ln n}{n^\lambda}\right) \quad \text{if } \alpha + \beta = \lambda, \\ &= O\left(\frac{1}{n^\lambda}\right) \quad \text{if } \alpha + \beta > \lambda. \end{aligned}$$

Proof. Let $M_s \leq n < M_{s+1}$; then by Theorem 8, $A_n = \|R_{n,\lambda}(f, \cdot) - f(\cdot)\|_X = (O(1)/n^\lambda) \sum_{l=0}^s \omega(T^{(\alpha)}f, 1/M_l) M_l^{\lambda-\alpha}$. If $\alpha + \beta < \lambda$, then

$$\begin{aligned} A_n &= \frac{O(1)}{n^\lambda} \sum_{l=0}^s \frac{1}{M_l^\beta} M_l^{\lambda-\alpha} = \frac{O(1)}{n^\lambda} \sum_{l=0}^s M_l^{\lambda-\alpha-\beta} \\ &= \frac{O(1)}{M_s^{\alpha+\beta}} \sum_{l=0}^s \left(\frac{M_l}{M_s}\right)^{\lambda-\alpha-\beta} = O\left(\frac{1}{n^{\alpha+\beta}}\right). \end{aligned}$$

If $\alpha + \beta = \lambda$, then

$$A_n = \frac{O(1)}{n^\lambda} \sum_{l=0}^s M_l^\alpha = O\left(\frac{s}{n^\lambda}\right) = O\left(\frac{\ln n}{n^\lambda}\right).$$

If $\alpha + \beta > \lambda$, then

$$A_n = \frac{O(1)}{n^\lambda} \sum_{l=0}^s \frac{1}{M_l^\beta} M_l^{\lambda-\alpha} = \frac{O(1)}{n^\lambda} \sum_{l=0}^s M_l^{\lambda-\alpha-\beta} = O\left(\frac{1}{n^\lambda}\right).$$

The above results have been obtained in the case $\alpha = 0$, $0 < \beta < 1$, $\lambda = 1$ by Yano [9] and Ефимов [14], and for $\alpha = 0$, $\beta = 1$, $\lambda = 1$ by Блюмин [12].

ACKNOWLEDGMENTS

I would like to thank Professor P. L. Butzer and the referees for their valuable suggestions and kind help.

REFERENCES

1. P. L. BUTZER AND H. J. WAGNER, Walsh-Fourier series and the concept of a derivative. *Applicable Anal.* **3** (1973), 29–46.
2. P. L. BUTZER AND H. J. WAGNER, Application by Walsh polynomials and the concept of a derivative, in “Proceedings, Sumpos., Applications of Walsh Function,” Vol. 3, pp. 388–392, 1972.

3. HE ZELIN, An approximation theorem on the p -adic Walsh–Fejer operator with corollaries, *J. Nanjing Univ.* **3** (1982).
4. C. W. OONNEWEER, Fractional differentiation on the group of integers of a p -adic or p -series field, *Anal. Math.* **3** (1977), 119–130.
5. C. W. OONNEWEER, Differentiability for Rademacher series on groups, *Acta Sci. Math. Szeged.* **39** (1977), 121–128.
6. J. PAL AND P. SIMON, On a generalization of the concept of derivative, *Acta Math. Sci. Hungar.* **29** (1977), 155–164.
7. REN FUXIAN, SU WEIYI, AND ZHENG WEIXING, The generalized logical derivative and its application, *J. Nanjing Univ.* **3** (1978), 1–9.
8. H. J. WAGNER, "Ein Differential und Integralkalkül in der Walsh–Fourier Analysis mit Anwendungen; Köln–Opladen, 1973.
9. S. YANO, On approximation by Walsh functions, *Proc. Amer. Math. Soc.* **2** (1951), 962–967.
10. C. WATARI, Best approximation by Walsh polynomials, *Tôhoku Math. J.* (2) **15** (1963), 1–15.
11. ZHENG WEIXING AND SU WEIYI, The logical derivatives and Integrals, *J. Math. Res. Exposition* **1** (1981), 79–90.
12. С. Л. Блюмин, О линейных методах суммирования рядов Фурье по мультиплитивным системам, *Сиб. Мат. Жур.* **IX**, **2** (1968), 449–455.
13. Н. Я. Виленкин, Об одном классе малых ортогоабальных систем, Изв. АН СССР. Серия Матем. **13** (1949), 245–252.
14. А. В. Ефимов, Об некоторых аппроксимативных свойствах периодических мультиплитивных ортформированных систем, *Матем. Сб.* **69** (3) (1966), 354–370.