


Contents lists available at [SciVerse ScienceDirect](http://SciVerse.Sciencedirect.com)

Annals of Pure and Applied Logic

journal homepage: www.elsevier.com/locate/apal

Randomness and lowness notions via open covers

Laurent Bienvenu^{a,*}, Joseph S. Miller^b^a LIAFA, CNRS & Université de Paris 7, France^b University of Wisconsin-Madison, USA

ARTICLE INFO

Article history:

Available online 10 August 2011

AMS subject classifications:

03D28

03D32

Keywords:

Algorithmic randomness

Lowness for randomness

Kolmogorov complexity

ABSTRACT

One of the main lines of research in algorithmic randomness is that of lowness notions. Given a randomness notion \mathcal{R} , we ask for which sequences A does relativization to A leave \mathcal{R} unchanged (i.e., $\mathcal{R}^A = \mathcal{R}$)? Such sequences are called *low for \mathcal{R}* . This question extends to a pair of randomness notions \mathcal{R} and \mathcal{S} , where \mathcal{S} is weaker: for which A is \mathcal{S}^A still weaker than \mathcal{R} ? In the last few years, many results have characterized the sequences that are low for randomness by their low computational strength. A few results have also given measure-theoretic characterizations of low sequences. For example, Kjos-Hanssen (following Kučera) proved that A is low for Martin-Löf randomness if and only if every A -c.e. open set of measure less than 1 can be covered by a c.e. open set of measure less than 1.

In this paper, we give a series of results showing that a wide variety of lowness notions can be expressed in a similar way, i.e., via the ability to cover open sets of a certain type by open sets of some other type. This provides a unified framework that clarifies the study of lowness for randomness notions, and allows us to give simple proofs of a number of known results. We also use this framework to prove new results, including showing that the classes $\text{Low}(\text{MLR}, \text{SR})$ and $\text{Low}(\text{W2R}, \text{SR})$ coincide, answering a question of Nies. Other applications include characterizations of highness notions, a broadly applicable explanation for why low for randomness is the same as low for tests, and a simple proof that $\text{Low}(\text{W2R}, \mathcal{S}) = \text{Low}(\text{MLR}, \mathcal{S})$, where \mathcal{S} is the class of Martin-Löf, computable, or Schnorr random sequences.

The final section gives characterizations of lowness notions using summable functions and convergent measure machines instead of open covers. We finish with a simple proof of a result of Nies, that $\text{Low}(\text{MLR}) = \text{Low}(\text{MLR}, \text{CR})$.

© 2012 Published by Elsevier B.V.

1. Introduction

This paper is organized as follows. In the remainder of this section we review notation, introduce the basic notions, including the relevant randomness classes, and survey what is known about lowness for randomness notions. In Section 2 we consider Kučera's result that X is not Martin-Löf random iff there is a c.e. open set \mathcal{U} of measure less than 1 such that \mathcal{U} covers all tails of X . We prove analogous theorems for computable and Schnorr randomness by placing further restrictions on the c.e. open covers.

In Section 3 we prove our main technical lemma and show that it applies to Martin-Löf randomness, computable randomness and Schnorr randomness. Together with the previous section, the main lemma provides a unified framework to study lowness classes in terms of c.e. open covers. Section 4 gives a number of applications. Kjos-Hanssen [10] showed that A is low for Martin-Löf randomness if and only if every A -c.e. open set of measure less than 1 can be covered by a c.e.

* Corresponding author.

E-mail addresses: laurent.bienvenu@liafa.jussieu.fr, laurent.bienvenu@lif.univ-mrs.fr (L. Bienvenu), jmiller@math.wisc.edu (J.S. Miller).

open set of measure less than 1.¹ In Section 4.1, we show that a wide variety of lowness notions can be expressed in a similar way, i.e., via the ability to cover open sets of a certain type by open sets of another type. Kjos-Hanssen’s result actually gives a characterization of LR-reducibility, and in Section 4.2, we note that similar characterizations could be given for the weak reducibilities associated with computable and Schnorr randomness. In Section 4.3 we give a broadly applicable explanation for why lowness for randomness has, in the cases that have been studied, turned out to be the same as lowness for tests. In Section 4.4 we show that $\text{Low}(\text{WZ2R}, \mathcal{S}) = \text{Low}(\text{MLR}, \mathcal{S})$ for $\mathcal{S} \in \{\text{MLR}, \text{CR}, \text{SR}\}$. Two of these facts were known, but the Schnorr randomness case answers an open question of Nies [18, Problem 8.3.16]. Finally, Section 4.5 applies our framework to highness notions, focusing on the poorly understood class $\text{High}(\text{CR}, \text{MLR})$.

Section 5 departs from the rest of the paper; in it, we reformulate lowness notions using summable functions and convergent measure machines instead of open covers. A final application is given in Section 5.2, where we give a straightforward proof that $\text{Low}(\text{MLR}) = \text{Low}(\text{MLR}, \text{CR})$ (Nies [17,18]).

1.1. Basic notation

We work in Cantor space, in other words, the set 2^ω of infinite binary sequences. We write $2^{<\omega}$ for the set of finite binary strings and $\epsilon \in 2^{<\omega}$ for the empty string. If S is a subset of $2^{<\omega}$, we define

$$S^n = \{\sigma \in 2^{<\omega} : \sigma = \sigma_0\sigma_1\sigma_2 \dots \sigma_{n-1} \text{ s.t. } (\forall i < n) \sigma_i \in S\}, \quad \text{and}$$

$$S^\omega = \{A \in 2^\omega : A = \sigma_0\sigma_1\sigma_2 \dots \text{ s.t. } (\forall i) \sigma_i \in S\}.$$

For a string σ , $[\sigma]$ denotes the cylinder generated by σ , in other words, the set of infinite sequences with prefix σ . For $U \subseteq 2^{<\omega}$, the open set generated by U is $[U] = \bigcup_{\sigma \in U} [\sigma]$. We denote the Lebesgue measure on 2^ω by μ (a.k.a. the uniform measure on 2^ω , which can be defined as the unique probability measure on 2^ω that satisfies $\mu([\sigma]) = 2^{-|\sigma|}$). If U is a prefix-free subset of $2^{<\omega}$, the *measure of U* is the quantity $\mu(U) = \mu([U]) = \sum_{\sigma \in U} 2^{-|\sigma|}$. Note that $\mu(U^n) = \mu(U)^n$, again assuming that $U \subseteq 2^{<\omega}$ is prefix-free. We say that an open set (resp. prefix-free set of strings) is *bounded* if its measure is smaller than 1. A *c.e. open set* (or Σ_1^0 class) is an open set generated by a c.e. prefix-free set of strings. We say that a c.e. open set (resp. c.e. prefix-free set of strings) is a *Schnorr set* if its measure is computable.

If $A \in 2^\omega$, we denote by $A \upharpoonright n$ the prefix A of size n , i.e., $A \upharpoonright n = A(0)A(1) \dots A(n-1)$. Also, we call a *tail of A* any infinite sequence of type $A(k)A(k+1)A(k+2) \dots$ for $k \in \mathbb{N}$ (in other words, any sequence obtained by removing a finite prefix from A). If \mathcal{X} is a subset of 2^ω and σ a finite string, we set

$$(\mathcal{X} \mid \sigma) = \{Z \in 2^\omega : \sigma Z \in \mathcal{X}\}.$$

Similarly, if W is a subset of $2^{<\omega}$ we set

$$(W \mid \sigma) = \{\tau \in 2^{<\omega} : \sigma\tau \in W\}.$$

Note that this is consistent with the conditional probability notation: $\mu(\mathcal{X} \mid \sigma)$ is just the measure of \mathcal{X} conditioned by $[\sigma]$, i.e., $\mu(\mathcal{X} \cap [\sigma])/\mu(\sigma)$. Note also that if \mathcal{U} is a c.e. open set, then so is $(\mathcal{U} \mid \sigma)$ for all σ . If moreover the measure of \mathcal{U} is computable, then so is the measure of $(\mathcal{U} \mid \sigma)$ (uniformly in $\mu(\mathcal{U})$ and σ).

1.2. Randomness notions

In general, a *test* is a non-increasing sequence $(\mathcal{U}_n)_{n \in \mathbb{N}}$ of open sets such that $\bigcap_n \mathcal{U}_n$ has measure 0. We say that a sequence $X \in 2^\omega$ *fails* the test $(\mathcal{U}_n)_{n \in \mathbb{N}}$ if $X \in \bigcap_n \mathcal{U}_n$, and that X *passes the test* otherwise. If \mathcal{X} is a subset of 2^ω , we say that a test $(\mathcal{U}_n)_{n \in \mathbb{N}}$ *covers \mathcal{X}* if $\mathcal{X} \subseteq \bigcap_n \mathcal{U}_n$. We say that a test $(\mathcal{U}'_n)_{n \in \mathbb{N}}$ *covers another test* $(\mathcal{U}_n)_{n \in \mathbb{N}}$ if $\bigcap_n \mathcal{U}'_n \subseteq \bigcap_n \mathcal{U}_n$.

Definition 1. A test $(\mathcal{U}_n)_{n \in \mathbb{N}}$ is a *Martin-Löf test* if $\mu(\mathcal{U}_n) \leq 2^{-n}$ for all n . It is a *Schnorr test* if one further has $\mu(\mathcal{U}_n) = 2^{-n}$. We say that X is *Martin-Löf random* if it passes all Martin-Löf tests, and that X is *Schnorr random* if it passes all Schnorr tests. We denote by MLR the set of Martin-Löf random sequences and by SR the set of Schnorr random sequences.

Remark 2. It should be noticed that the quantity 2^{-n} in the above definition is arbitrary: we would get the same classes MLR and SR if we replaced it by any $f(n)$, with f a computable function that tends to 0. Another important fact is that there exists a *universal* Martin-Löf test, i.e., a Martin-Löf test such that for any sequence X , X passes that test if and only if X is Martin-Löf random. There is no such universal test for Schnorr randomness.

A third important notion of randomness is *computable randomness*, whose definition involves the concept of a *martingale*.

A *martingale* is a function $d : 2^{<\omega} \rightarrow \mathbb{R}^{\geq 0}$ such that for all $\sigma \in 2^{<\omega}$

$$d(\sigma) = \frac{d(\sigma 0) + d(\sigma 1)}{2}.$$

It is said to be *normed* if $d(\epsilon) = 1$. We say that a martingale *succeeds* on a sequence $X \in 2^\omega$ if $\limsup d(X \upharpoonright n) = +\infty$.

¹ The fact that the covering condition implies that A is low for Martin-Löf randomness was proved by Kučera and Terwijn [14]. A straightforward proof of Kjos-Hanssen’s result using open covers is given by Barmpaliadis et al. [1].

For any martingale, the set of sequences on which it succeeds has measure 0. This is a direct consequence of the so-called Ville–Kolmogorov inequality.

Proposition 3. Let d be a martingale, $\sigma \in 2^{<\omega}$ and $q > 1$ a real number. Let

$$\mathcal{U}_{d,\sigma,q} = \{X \in 2^\omega : (\exists n > |\sigma|) d(X \upharpoonright n) \geq q \cdot d(\sigma)\}.$$

Then $\mu(\mathcal{U}_{d,\sigma,q} \mid \sigma) \leq 1/q$.

We can now define the notion of computable randomness.

Definition 4. We say that X is *computably random* if no computable martingale succeeds on X . We denote by CR the set of computably random sequences.

In the above definition, by “computable” we mean computable as a real-valued function. However, it will be more convenient in this paper to work with *exactly computable martingales*, i.e., martingales that are rational-valued and computable as functions from \mathbb{N} to $\mathbb{Q}^{\geq 0}$. This is made possible by a note of Lutz [15], where it is proven that for every computable martingale d , there exists an exactly computable martingale d' and positive real constants α, β such that $\alpha d < d' < \beta d$ (in particular, d and d' succeed on the same set of sequences). Furthermore, an index of d' can be uniformly computed from an index of d . Therefore, we can equivalently define the set CR as being the set of sequences X such that no (normed) exactly computable martingale succeeds on X . We can also rephrase the definition in terms of test.

Definition 5. Let d be a rational-valued normed martingale and q a rational such that $q > 1$. We say that $U \subseteq 2^{<\omega}$ is a (d, q) -winning set if for some rational $q > 1$ we have $U = \{\sigma : \sigma \text{ minimal s.t. } d(\sigma) \geq q\}$. We say that $U \subseteq 2^{<\omega}$ is a winning set if it is a (d, q) -winning set for some exactly computable normed martingale d and rational $q > 1$. We also say that a c.e. open set \mathcal{U} is a winning set if $\mathcal{U} = [U]$ where U is a winning set of strings.

Given a normed exactly computable martingale d , the test induced by d is the sequence $(\mathcal{U}_n)_{n \in \mathbb{N}}$ where \mathcal{U}_n is the $(d, 2^n)$ -winning set.

(in the above definition, and in the rest of the paper, we say that a string σ is *minimal* for a given property P if σ satisfies P and no prefix of σ does).

Now, we immediately see that X is computably random if and only if X passes all tests induced by normed exactly computable martingales.

Remark 6. If d is a normed, exactly computable martingale, any (d, q) -winning set is c.e. open, and (by the Ville–Kolmogorov inequality) has measure at most $1/q$. Thus the test induced by a normed, exactly computable martingale is a Martin–Löf test.

The last randomness notion we will discuss in this paper is a very natural generalization of Martin–Löf randomness. Weak 2-randomness (sometimes called Kurtz 2-randomness) allows tests $(\mathcal{U}_n)_{n \in \mathbb{N}}$ with the looser condition that $\mu(\mathcal{U}_n)$ tends to 0, possibly at a non-computable rate.

Definition 7. A generalized Martin–Löf test is a sequence $(\mathcal{U}_n)_{n \in \mathbb{N}}$ of uniformly c.e. open sets such that $\lim_n \mu(\mathcal{U}_n) = 0$. We say that X is weak 2-random if X passes all generalized Martin–Löf tests and denote by W2R the set of weak 2-random sequences.

Note that a generalized Martin–Löf test is nothing more than a measure zero Π_2^0 class and X is weak 2-random iff it avoids every such class.

1.3. Lowness notions: the state of the art

In computability theory, an oracle A is said to be *low* in a certain context if a relativization to A “does not help”. For example, A is low for the Turing jump (usually referred to simply as “low”) if $A' \equiv_T \mathbf{0}'$. Lowness notions have been very important in the recent development of algorithmic randomness. Take a randomness notion \mathcal{R} that, like all the above, can be defined via tests. One can relativize the notion of test to an oracle A , getting the class \mathcal{R}^A of sequences that pass all A -tests. Since taking A as an oracle gives additional computational power, we have $\mathcal{R}^A \subseteq \mathcal{R}$. We say that A is *low for the randomness notion* \mathcal{R} if, as an oracle, A has so little computational power that $\mathcal{R}^A = \mathcal{R}$. We denote by $\text{Low}(\mathcal{R})$ the set of sequences that are low for \mathcal{R} .

Zambella [20] introduced lowness for Martin–Löf randomness. A beautiful series of results by Nies and others (see [18] for a complete exposition) showed that these oracles have remarkable properties. They proved that $A \in \text{Low}(\text{MLR})$ if and only if A is low for prefix-free Kolmogorov complexity (i.e., $K^A = K + O(1)$), and if and only if A is K -trivial. This latter property states that the initial segments of A have minimal prefix-free Kolmogorov complexity (i.e., $K(A \upharpoonright n) \leq K(n) + O(1)$). Lowness has been studied for other randomness notions. The work of Terwijn and Zambella [19] and of Kjos-Hanssen et al. [12] characterized low for Schnorr randomness as computably traceable (a strengthening of hyperimmune-free; see below). Nies [17] showed that only computable oracles can be low for computable randomness.

One can also study lowness for a pair of randomness notions. If \mathcal{R} and \mathcal{S} are two randomness notions with $\mathcal{R} \subseteq \mathcal{S}$, $\text{Low}(\mathcal{R}, \mathcal{S})$ is the set of oracles A such that $\mathcal{R} \subseteq \mathcal{S}^A$. The task of characterizing the sequences that are low for randomness has attracted a lot of effort in the last few years, and is now nearly completed, as shown in the following diagram.

		\mathcal{S}		
	W2R	MLR	CR	SR
W2R	K-trivial [6,11,18]	K-trivial [6]	K-trivial [18]	c.e. traceable
		MLR	K-trivial [17]	c.e. traceable [12]
		\mathcal{R}	CR	computably traceable [12]
				SR
				computably traceable [19,12]

Low(\mathcal{R}, \mathcal{S}) for various randomness classes.

Note that each class in the diagram is contained in the classes above it and to its right. The gray entry is settled in this paper. It should be also noted that this diagram omits the results obtained by Greenberg and Miller [9] that characterize almost all lowness notions related to *weak 1-randomness*, as we do not discuss weak 1-randomness in the present paper.

Although we will not directly use these notions in this paper, we recall the definitions of the classes that are referred to in this diagram. We defined *K-triviality* above. A sequence A is *computably traceable* if there exists a single computable function h such that for any total function $f : \mathbb{N} \rightarrow \mathbb{N}$ computable in A , there exists a uniformly computable sequence of finite sets $(T_n)_{n \in \mathbb{N}}$, given by their strong index, such that for all $n, f(n) \in T_n$ and $|T_n| < h(n)$. The definition of *c.e. traceability* is the same, except that the sets T_n are given by their index as c.e. sets. Kjos-Hanssen et al. [12] introduced c.e. traceability specifically to characterize Low(MLR, SR), making it one of several examples of interesting computability theoretic properties that have arisen from the study of randomness and lowness notions.

2. Testing randomness via open covers

In this section, we present an alternative way to look at the above randomness notions. Instead of using tests, i.e., sequences of open sets, it is possible to provide equivalent definitions involving a single open set (or c.e. set of strings). The first theorem below is due to Kučera [13] and characterizes Martin-Löf randomness. We prove analogous theorems for computable and Schnorr randomness.

Theorem 8. *Let $X \in 2^\omega$. The following are equivalent:*

- (i) X is not Martin-Löf random.
- (ii) There is a bounded c.e. open set \mathcal{U} such that all tails of X belong to \mathcal{U} .
- (iii) $X \in U^\omega$ for some bounded c.e. prefix-free subset U .

Proof. (i) \Rightarrow (ii) This follows easily from the existence of a universal Martin-Löf test $(\mathcal{U}_n)_{n \in \mathbb{N}}$. Let $\mathcal{U} = \mathcal{U}_1$. So \mathcal{U} is a bounded c.e. open set covering all non-Martin-Löf random sequences. If X is not Martin-Löf random, then none of its tails are Martin-Löf random. Hence they all belong to \mathcal{U} .

(ii) \Rightarrow (iii) Suppose that \mathcal{U} is a bounded c.e. open set and that all of the tails of X belong to \mathcal{U} . Let $U \subset 2^{<\omega}$ be a c.e. prefix-free set such that $\mathcal{U} = [U]$. We show by induction on n that $X \in [U^n]$. This is true for $n = 1$ by assumption. Now assume that $X \in [U^n]$. So for some $\sigma \in U^n$, there is a Z such that $X = \sigma Z$. Since Z is a tail of X , we know that $Z \in \mathcal{U} = [U]$. But this implies that $X \in [U^{n+1}]$. Thus $X \in [U^n]$ for all n . Therefore, $X \in U^\omega = \bigcap_n [U^n]$.

(iii) \Rightarrow (i) Assume that $X \in U^\omega$ for some bounded c.e. prefix-free subset U . For each n , we have $X \in \mathcal{U}_n = [U^n]$. Also, $\mu(\mathcal{U}_n) = \mu(U^n) = \mu(U)^n$, so $(\mathcal{U}_n)_{n \in \mathbb{N}}$ is a Martin-Löf test (in the more general sense of Remark 2). Since $(\mathcal{U}_n)_{n \in \mathbb{N}}$ covers X , it is not Martin-Löf random. \square

Theorem 9. *Let $X \in 2^\omega$. The following are equivalent:*

- (i) X is not Schnorr random.
- (ii) There is a bounded Schnorr open set \mathcal{U} such that all tails of X belong to \mathcal{U} .
- (iii) $X \in U^\omega$ for some bounded Schnorr prefix-free subset U of $2^{<\omega}$.

Proof. (i) \Rightarrow (ii) Suppose that X is not Schnorr random. Then $X \in \bigcap_n \mathcal{V}_n$ where $(\mathcal{V}_n)_{n \in \mathbb{N}}$ is a Schnorr test (say with $\mu(\mathcal{V}_n) = 2^{-n}$). We build the desired set \mathcal{U} from this Schnorr test. For every k , the tail $Y_k = X(k)X(k+1) \dots$ of X belongs to $\bigcap_n (\mathcal{V}_n \mid \tau)$, where $\tau = X(0) \dots X(k-1)$. Thus, for all n , Y_k belongs to

$$\bigcup_{|\sigma|=k} (\mathcal{V}_n \mid \sigma),$$

which, for n large enough, has small measure. For example, for $n = 3k + 2$, the above set has measure at most 2^{-k-2} (indeed each of the 2^k sets of type $(\mathcal{V}_n \mid \sigma)$ has measure at most $\mu(\mathcal{V}_n)/\mu([\sigma]) \leq 2^{-n}2^k$, hence the total measure is at most 2^{-n+2k}). Thus, define:

$$\mathcal{U} = \bigcup_{k \in \mathbb{N}} \bigcup_{|\sigma|=k} (\mathcal{V}_{3k+2} \mid \sigma).$$

We claim that \mathcal{U} is as wanted. Indeed, \mathcal{U} is clearly Σ_1^0 . By the above discussion, \mathcal{U} contains all tails of X , the measure of \mathcal{U} is at most

$$\sum_{k \in \mathbb{N}} 2^{-k-2} \leq 1/2.$$

To see that the measure of \mathcal{U} is computable, note that the measure $\mu(\mathcal{V}_{3k+2} \mid \sigma)$ is computable uniformly in k and σ . So the measure of

$$\bigcup_{k \leq N} \bigcup_{|\sigma| = k} (\mathcal{V}_{3k+2} \mid \sigma),$$

is computable, uniformly in N , and approximates $\mu(\mathcal{U})$ up to $\sum_{k > N} 2^{-k-2} < 2^{-N}$.

The proofs of (ii) \Rightarrow (iii) and (iii) \Rightarrow (i) go exactly as in [Theorem 8](#). \square

Theorem 10. *Let $X \in 2^\omega$. The following are equivalent:*

- (i) X is not computably random.
- (ii) There exists a winning open set \mathcal{U} such that all tails of X belong to \mathcal{U} .
- (iii) $X \in U^\omega$ for some winning subset U of $2^{<\omega}$.

Proof. (i) \Rightarrow (ii). Suppose X is not computably random. Then there exists a computable martingale d that succeeds against X . Up to adding a positive constant to d (preserving the fairness condition) we can assume that d is positive. Now, for all strings σ , one can consider the “translated” version d_σ of d defined by

$$d_\sigma(\tau) = d(\sigma\tau).$$

It is easy to see that if Y is a tail of X , with $X = \sigma Y$, then d_σ succeeds against Y . Therefore, the martingale D defined by:

$$D(\tau) = \sum_{\sigma \in 2^{<\omega}} 2^{-2|\sigma|-1} \cdot \frac{d_\sigma(\tau)}{d_\sigma(\epsilon)}$$

succeeds against all tails of X . Moreover, D is normed and computable as a sum of exponentially decreasing uniformly computable terms. By the result of Lutz [15] mentioned earlier, we can also assume that D is exactly computable. Since D succeeds against all tails of X , this in particular implies that all tails of X belong to the winning open set $[U]$ with $U = \{\sigma : \sigma \text{ minimal s.t. } D(\sigma) \geq 2\}$.

The proof of (ii) \Rightarrow (iii) is as in [Theorem 8](#).

(iii) \Rightarrow (i). If $X \in U^\omega$ where U is a winning set of strings, let d be the normed exactly computable martingale and $q > 1$ a rational such that U is a (d, q) -winning set. We can assume that d is positive, otherwise we set $d' = \frac{1}{2}d + \frac{1}{2}$, which makes U a $(d', \frac{1+q}{2})$ -winning set, and d' is positive. Now, we design a computable martingale D that succeeds on all sequences in U^ω . Basically, D simulates the martingale d and “resets” after reading a block $\sigma \in U$. Formally, D is defined by induction. We set $D(\epsilon) = 1$ and if $D(\sigma)$ is already defined, we write $\sigma = \rho\tau$ where ρ is a concatenation of strings in U and τ has no prefix in U (this decomposition is unique as U is prefix-free) and then set

$$D(\sigma\iota) = D(\sigma) \cdot \frac{d(\tau\iota)}{d(\tau)},$$

for $\iota \in \{0, 1\}$. It is easy to see that D is an exactly computable martingale, and if σ is a concatenation of k strings in U , $D(\sigma) \geq q^k$. Hence D succeeds against all sequences in U^ω . \square

Remark 11. In fact, the proofs of [Theorems 8–10](#) show a little more. What we actually proved is the following equivalence for any subset \mathcal{X} of 2^ω

- (i) \mathcal{X} is covered by a Martin-Löf test (resp. a test induced by a martingale, a Schnorr test).
- (ii) There exists a single bounded c.e. open set (resp. winning open set, bounded Schnorr open set) \mathcal{U} such that for any $X \in \mathcal{X}$, all tails of X are in \mathcal{U} .
- (iii) There exists a single bounded c.e. set of strings (resp. winning set of strings, bounded Schnorr set of strings) U such that $\mathcal{X} \subseteq U^\omega$.

3. The main lemma

The following technical lemma is the cornerstone of this paper. It lets us use the characterizations of Martin-Löf randomness, computable randomness and Schnorr randomness proven in the previous section to study the associated lowness notions. Roughly speaking, it states that if a prefix-free set of strings U is not covered by any of the members of a (reasonably well-behaved) collection \mathbf{C} of open sets, then there exists an $X \in U^\omega$ that passes all tests that can be built from the elements of \mathbf{C} .

Lemma 12. Let \mathbf{C} be a class of bounded open subsets of 2^ω . Let also $(\mathcal{T}_n^{(e)})_{e,n \in \mathbb{N}}$ be a countable family of tests (i.e., for all e , $(\mathcal{T}_n^{(e)})_{n \in \mathbb{N}}$ is a test) such that $\mathcal{T}_n^{(e)}$ belongs to \mathbf{C} for all e, n . Suppose we have the following closure properties.

- (P1) For all $\mathcal{U} \in \mathbf{C}$ and $\sigma \in 2^{<\omega}$, if $\mu(\mathcal{U} \mid \sigma) < 1$, then there exists a $\mathcal{V} \in \mathbf{C}$ such that $(\mathcal{U} \mid \sigma) \subseteq \mathcal{V}$.
- (P2) For all $\mathcal{U} \in \mathbf{C}$, there exists a $\mathcal{V} \in \mathbf{C}$ such that $\mathcal{U} \subseteq \mathcal{V}$, and for all $\sigma \in 2^{<\omega}$, if $\mu(\mathcal{U} \mid \sigma) = 1$, then $[\sigma] \subseteq \mathcal{V}$.
- (P3) For all $\mathcal{U} \in \mathbf{C}$, and $\sigma \in 2^{<\omega}$, if $\mu(\mathcal{U} \mid \sigma) < 1$, then for all $e \in \mathbb{N}$, there exists $n_e \in \mathbb{N}$ and $\mathcal{V} \in \mathbf{C}$ such that $(\mathcal{U} \cup \mathcal{T}_{n_e}^{(e)}) \subseteq \mathcal{V}$ and $\mu(\mathcal{V} \mid \sigma) < 1$.

Finally, let W be a prefix-free subset of $2^{<\omega}$ such that $[W]$ cannot be covered by any open set $\mathcal{U} \in \mathbf{C}$. Then, there exists $X \in W^\omega$ that passes all tests $\mathcal{T}^{(e)}$.

Proof. We build X by a finite extension technique: having built a prefix σ_e of X , during stage e , we build an extension σ_{e+1} of σ_e . This is done as follows. We begin with σ_0 equal to the empty string, and \mathcal{U}_0 equal to the empty subset of 2^ω . At the beginning of stage e , suppose we have already built σ_e and \mathcal{U}_e , satisfying $\mu(\mathcal{U}_e \mid \sigma_e) < 1$. We then use property (P3) to find n_e and $\mathcal{V} \in \mathbf{C}$ such that $(\mathcal{U}_e \cup \mathcal{T}_{n_e}^{(e)}) \subseteq \mathcal{V}$ and $\mu(\mathcal{V} \mid \sigma_e) < 1$. Next, we pick a non-empty string $\tau \in W$ such that $\mu(\mathcal{V} \mid \sigma_e \tau) < 1$. We then set $\mathcal{U}_{e+1} = \mathcal{V}$ and $\sigma_{e+1} = \sigma_e \tau$, finishing the e -th stage.

It remains to verify that this construction works. First, we have to make sure that at stage e of our construction, there is indeed a string $\tau \in W$ such that $\mu(\mathcal{V} \mid \sigma_e \tau) < 1$. Suppose that this is not the case. This means that $\mu(\mathcal{V} \mid \sigma_e \tau) = 1$ for all $\tau \in W$, or equivalently that $\mu((\mathcal{V} \mid \sigma_e) \mid \tau) = 1$ for all $\tau \in W$. By definition of \mathcal{V} , we have $\mu(\mathcal{V} \mid \sigma_e) < 1$, so we can apply property (P1) to get some $\mathcal{V}' \in \mathbf{C}$ such that $(\mathcal{V} \mid \sigma_e) \subseteq \mathcal{V}'$. In particular, \mathcal{V}' is such that $\mu(\mathcal{V}' \mid \tau) = 1$ for all $\tau \in W$. Now by property (P2) there exists $\mathcal{V}'' \in \mathbf{C}$ covering \mathcal{V}' and such that $[\tau] \subseteq \mathcal{V}''$ for all $\tau \in W$, which means that $[W]$ is covered by \mathcal{V}'' . This contradicts the assumption that $[W]$ is not covered by a set that belongs to \mathbf{C} .

Now, let X be the unique element of 2^ω such that all σ_e 's are prefixes of X . It is easy to see from the construction that $X \in W^\omega$. Moreover, suppose X fails a test $\mathcal{T}^{(e)}$. This would imply that $X \in \mathcal{T}_{n_e}^{(e)}$ (the n_e being defined in the above construction). Thus, there would exist $e' > e$ large enough, such that $[\sigma_{e'}] \subseteq \mathcal{T}_{n_e}^{(e)}$. This would be a contradiction since, by construction, on the one hand $[\sigma_{e'}] \not\subseteq \mathcal{U}_{e'+1}$ and on the other hand $\mathcal{T}_{n_e}^{(e)} \subseteq \mathcal{U}_{e+1} \subseteq \mathcal{U}_{e'+1}$. \square

Proposition 13. The hypotheses (P1, P2, P3) of Lemma 12 are satisfied in the following three cases.

(MLR) \mathbf{C} is the class of bounded c.e. open sets and (\mathcal{T}_e) is the family of Martin-Löf tests.

(CR) \mathbf{C} is the class of open sets that are winning sets of exactly computable martingales and (\mathcal{T}_e) is the family of tests induced by those martingales.

(SR) \mathbf{C} is the class of bounded Schnorr open sets and (\mathcal{T}_e) is the family of Schnorr tests.

Proof. (MLR) As we previously observed, for every c.e. open set \mathcal{U} and $\sigma \in 2^{<\omega}$, $(\mathcal{U} \mid \sigma)$ is a c.e. open set and its index can be computed from an index of \mathcal{U} and σ . The property (P1) thus follows immediately. For the property (P3), given a c.e. open set \mathcal{U} and $\sigma \in 2^{<\omega}$, such that $\mu(\mathcal{U} \mid \sigma) < 1 - 2^{-k}$ for some $k > 0$, together with a Martin-Löf test $\mathcal{T}^{(e)}$, take $n_e = |\sigma| + k$. By the definition of a Martin-Löf test, $\mu(\mathcal{T}_{n_e}^{(e)}) < 2^{-n_e} = 2^{-|\sigma|-k}$. So $\mu(\mathcal{T}_{n_e}^{(e)} \mid \sigma) < 2^{-k}$, and the set $\mathcal{V} = \mathcal{U} \cup \mathcal{T}_{n_e}^{(e)}$ is c.e. open and satisfies $\mu(\mathcal{V} \mid \sigma) < (1 - 2^{-k}) + 2^{-k} < 1$. We now check that (P2) holds. Given a bounded c.e. open set \mathcal{U} , let $q < 1$ be a rational such that $\mu(\mathcal{U}) < q$ and set

$$\mathcal{V} = \bigcup \{[\sigma] : \mu(\mathcal{U} \mid \sigma) > q\}.$$

It is clear that \mathcal{V} is c.e. open and if $\mu(\mathcal{U} \mid \sigma) = 1$ then $[\sigma] \subseteq \mathcal{V}$. It remains to check that \mathcal{V} is bounded. Let F be the set of strings σ that are minimal among those satisfying $\mu(\mathcal{U} \mid \sigma) > q$. We have

$$\mu(\mathcal{V}) = \sum_{\sigma \in F} \mu([\sigma]) \leq \sum_{\sigma \in F} \frac{\mu(\mathcal{U} \cap [\sigma])}{q} \leq \frac{\mu(\mathcal{U} \cap [F])}{q} \leq \frac{\mu(\mathcal{U})}{q} < 1.$$

(CR) Let U be a c.e. set of strings such that $U = \{\sigma : \sigma \text{ minimal s.t. } d(\sigma) \geq q\}$ for some exactly computable normed martingale d and rational $q > 1$.

For property (P1), suppose that $\mu(U \mid \sigma) < 1$. We thus have $d(\sigma) < q$ (otherwise $[\sigma] \subseteq [U]$), and we can also assume that $d(\sigma) > 0$ (otherwise $(U \mid \sigma) = \emptyset$ and there is nothing to prove). We have

$$\begin{aligned} (U \mid \sigma) &= \{\tau : \tau \text{ minimal s.t. } d(\sigma \tau) \geq q\} \\ &= \left\{ \tau : \tau \text{ minimal s.t. } \frac{d(\sigma \tau)}{d(\sigma)} \geq \frac{q}{d(\sigma)} \right\}. \end{aligned}$$

It is easy to check that $\tau \mapsto \frac{d(\sigma \tau)}{d(\sigma)}$ is an exactly computable normed martingale, and since $\frac{q}{d(\sigma)} > 1$ we see that $(U \mid \sigma)$ is a winning set of strings.

For property (P2), we will see that the property $(\mu(U \mid \sigma) = 1 \Rightarrow [\sigma] \subseteq [U])$ holds. Indeed, if $\mu(U \mid \sigma) = 1$, this means that for almost all $X \in [\sigma]$, there exists an n such that $d(X \upharpoonright n) \geq q$. By the Ville–Kolmogorov inequality, this implies that $d(\sigma) \geq q$.

For property (P3), let σ be such that $\mu(U \mid \sigma) < 1$. As we have seen, this implies $d(\sigma) < q$. Take the e -th test $\mathcal{T}^{(e)}$ associated to an exactly computable normed martingale d_e (i.e., $\mathcal{T}_n^{(e)}$ is the open set generated by the strings σ such that $d(\sigma) \geq 2^n$). We need to find n_e such that $([U] \cup \mathcal{T}_{n_e}^{(e)})$ is covered by $[V]$, where V is a winning set such that $\mu(V \mid \sigma) < 1$. Let n_e be large, to be specified later. Let D be the exactly computable normed martingale defined by

$$D = (1 - 2^{-n_e+1})d + 2^{-n_e+1}d_e.$$

We have $D(\sigma) = (1 - 2^{-n_e+1})d(\sigma) + 2^{-n_e+1}d_e(\sigma)$. Now, suppose $X \in ([U] \cup \mathcal{T}_{n_e}^{(e)})$. If $X \in [U]$, then $d(X \upharpoonright n) \geq q$ for some n , and then $D(X \upharpoonright n) \geq (1 - 2^{-n_e+1})q$. If $X \in \mathcal{T}_{n_e}^{(e)}$, then $d(X \upharpoonright n) \geq 2^{-n_e+1}2^n = 2$ for some n . We thus consider the set

$$V = \{\tau : \tau \text{ minimal s.t. } D(\tau) \geq \min((1 - 2^{-n_e+1})q, 2)\}.$$

Now, for n_e large enough, we can ensure from the previous calculations that $D(\sigma)$ is as close as we want to $d(\sigma) < q$, and thus that $\min((1 - 2^{-n_e+1})q, 2) > D(\sigma)$. The above set V is then as desired: it is clearly a winning set, it covers $([U] \cup \mathcal{T}_{n_e}^{(e)})$, and $\mu(V \mid \sigma) < 1$ as $D(\sigma) < \min((1 - 2^{-n_e+1})q, 2)$.

(SR) Property (P1) is clearly satisfied. Given \mathcal{U} a c.e. open set of computable measure, and $\sigma \in 2^{<\omega}$, $(\mathcal{U} \mid \sigma)$ is a c.e. open set whose measure is computable (uniformly in σ and an index for \mathcal{U}). Indeed, if $\widehat{\mathcal{U}}$ is a clopen approximation of \mathcal{U} such that $\mu(\mathcal{U} \setminus \widehat{\mathcal{U}}) < \varepsilon$, then $\mu((\mathcal{U} \mid \sigma) \setminus (\widehat{\mathcal{U}} \mid \sigma)) < \varepsilon \cdot 2^{|\sigma|}$. Property (P3) is also clearly satisfied, as it is easy to see that given two c.e. open sets \mathcal{U} and \mathcal{V} of computable measure, $\mathcal{U} \cup \mathcal{V}$ is also a c.e. open set of computable measure.

For property (P2), let \mathcal{U} be a bounded Schnorr open set and let k be large enough that $\mu(\mathcal{U}) < 1 - 2^{-k}$. We define a computable set of strings V such that $[V] \supseteq \mathcal{U}$ is a bounded Schnorr open set. For every $\sigma \in 2^{<\omega}$, look for a stage s such that $\mu(\mathcal{U} \setminus \mathcal{U}_s) < 2^{-2|\sigma|-k-1}$. If $\mu(\mathcal{U}_s \mid \sigma) > 1 - 2^{-|\sigma|-k-1}$, then put σ into V . It is clear that V is computable. If $\mu(\mathcal{U} \mid \sigma) = 1$, then it must be the case that $\mu(\mathcal{U}_s \mid \sigma) > 1 - 2^{-|\sigma|-k-1}$; otherwise

$$\mu(\mathcal{U} \setminus \mathcal{U}_s) \geq \mu([\sigma] \setminus \mathcal{U}_s) \geq 2^{-|\sigma|} - 2^{-|\sigma|}(1 - 2^{-|\sigma|-k-1}) = 2^{-2|\sigma|-k-1},$$

which contradicts the choice of s . Therefore, $\mu(\mathcal{U} \mid \sigma) = 1$ implies that $\sigma \in V$, so $[\sigma] \subseteq [V]$ as required. This also implies that $\mathcal{U} \subseteq [V]$. It remains to show that $[V]$ is bounded and has computable measure. By adding σ to V , we are increasing the measure of $[V] \setminus \mathcal{U}$ by less than $2^{-|\sigma|}2^{-|\sigma|-k-1} = 2^{-2|\sigma|-k-1}$. Therefore,

$$\mu([V] \setminus \mathcal{U}) < \sum_{\sigma \in 2^{<\omega}} 2^{-2|\sigma|-k-1} = \sum_{n \in \omega} 2^n 2^{-2n-k-1} = 2^{-k}.$$

This implies that $\mu([V]) < \mu(\mathcal{U}) + 2^{-k} < 1$, so $[V]$ is a bounded c.e. open set. Similarly, for every m ,

$$\mu([V] \setminus ([V \cap 2^{<m}] \cup \mathcal{U})) < \sum_{\sigma \in 2^{<m}} 2^{-2|\sigma|-k-1} = \sum_{n \geq m} 2^n 2^{-2n-k-1} = 2^{-m-k}.$$

But $\mu([V \cap 2^{<m}] \cup \mathcal{U})$ is computable uniformly in m , so $\mu([V])$ is also computable. \square

4. Applications

4.1. Lowness notions

Together with the results of Section 2, Lemma 12 has interesting consequences. First, it follows from it that all lowness notions involving Martin-Löf randomness, computable randomness, or Schnorr randomness can essentially be reduced to a property of open sets. For example $A \in \text{Low}(\text{MLR})$ if and only if every bounded A -c.e. open set can be covered by a bounded c.e. open set (under this form, this was first stated by Kjos-Hanssen in [10], but most of the ideas are already present in Kučera [13]); $A \in \text{Low}(\text{CR})$ if and only if every A -winning open set can be covered by a winning open set; $A \in \text{Low}(\text{SR})$ if and only if every bounded A -computable open set can be covered by a bounded computable open set. This also works for pairs of randomness notions: for example, $A \in \text{Low}(\text{MLR}, \text{CR})$ if and only if every A -winning open set can be covered by a bounded c.e. open set. Let us briefly present the proof of one such result (the proofs of all the other claims are almost identical).

Corollary 14. *Let $A \in 2^\omega$. The following are equivalent:*

- (i) $A \in \text{Low}(\text{CR}, \text{SR})$
- (ii) Every bounded A -Schnorr open set can be covered by a winning open set.

Proof. (i) \Rightarrow (ii). Suppose that (ii) does not hold, i.e., there exists a bounded A -Schnorr open set \mathcal{U} that cannot be covered by any winning open set. Let U be an A -c.e. prefix-free set of strings generating \mathcal{U} . We can apply Lemma 12 with \mathbf{C} the class of winning open sets and \mathcal{T} the family tests induced by exactly computable martingales (this is allowed by Proposition 13), from which we get the existence of $X \in U^\omega$ that passes all tests induced by exactly computable martingales (hence X is computably random). By Theorem 9 (relativized to A), X is not A -Schnorr random. Thus $A \notin \text{Low}(\text{CR}, \text{SR})$.

(ii) \Rightarrow (i). Suppose (ii) holds and take $X \in 2^\omega$ such that X is not A -Schnorr random. By Theorem 9, there exists a bounded A -Schnorr open set \mathcal{U} such that all tails of X belong to \mathcal{U} . By assumption (ii), \mathcal{U} can be covered by a winning open set \mathcal{V} . Thus, all tails of X belong to \mathcal{V} , which by Theorem 10 implies that X is not computably random. \square

4.2. Weak reducibilities

Nies [17] introduced a weak reducibility generalizing low for Martin-Löf randomness. He defined $A \leq_{LR} B$ to mean that $\text{MLR}^B \subseteq \text{MLR}^A$. So, $A \leq_{LR} \emptyset$ iff $A \in \text{Low}(\text{MLR})$. What Kjos-Hanssen [10] actually proved was that $A \leq_{LR} B$ if and only if every bounded A -c.e. open set can be covered by a bounded B -c.e. open set. This result follows easily from our framework, as do the analogous results for the weak reducibilities associated with computable and Schnorr randomness, although these relations have not received attention. For example, if we write $A \leq_{CR} B$ to mean that $\text{CR}^B \subseteq \text{CR}^A$, then the following characterization follows by a proof identical to that of Corollary 14.

Corollary 15. *The following are equivalent for $A, B \in 2^\omega$:*

- (i) $A \leq_{CR} B$.
- (ii) Every A -winning open set can be covered by a B -winning open set.

4.3. Lowness for randomness vs lowness for tests

When defining lowness for randomness notions, two approaches are possible. The obvious one is the one we have studied so far in this paper: $A \in 2^\omega$ is low for a randomness notion \mathcal{R} if relativizing the notion \mathcal{R} to A leaves the set of random sequences unchanged. Now suppose that the notion \mathcal{R} is described via a family of tests (i.e., a sequence is random for the notion \mathcal{R} if it passes all tests), like all of the notions we have presented above. A second possible lowness condition on A is to require that A -tests are not stronger than unrelativized tests, i.e., that for every A -test \mathcal{T} , there exists a test \mathcal{T}' such that every sequence failing \mathcal{T} also fails \mathcal{T}' . While it is clear that (given a randomness notion defined by tests) lowness for tests implies lowness for randomness, the converse may not hold; a priori, it could be the case that many unrelativized tests are needed to cover a particular A -test. Nonetheless, there is currently no known example of a randomness notion (or a pair of randomness notions) for which lowness for tests is different from lowness for randomness. The results proven above provide a uniform explanation to why this is the case for lowness notions relating to Martin-Löf randomness, computable randomness and Schnorr randomness. Let us prove for example that lowness for Schnorr randomness implies lowness for Schnorr tests (a result originally proven by Kjos-Hanssen et al. [12]). Let A be low for Schnorr randomness, and consider an A -Schnorr test $(\mathcal{V}_n)_{n \in \mathbb{N}}$. Let us set $\mathcal{X} = \bigcap_n \mathcal{V}_n$. By Remark 11 (relativized to A), there exists a bounded A -Schnorr open set \mathcal{U} such that for any $X \in \mathcal{X}$, all tails of X are in \mathcal{U} . But since A is low for Schnorr randomness, we now know that \mathcal{U} must be covered by a bounded Schnorr open set \mathcal{U}' . This implies in particular that for any $X \in \mathcal{X}$, all tails of X are in \mathcal{U}' . Applying Remark 11 again, there must exist a Schnorr test $(\mathcal{V}'_n)_{n \in \mathbb{N}}$ that covers \mathcal{X} . In other words, $(\mathcal{V}'_n)_{n \in \mathbb{N}}$ covers $(\mathcal{V}_n)_{n \in \mathbb{N}}$. Therefore, A is low for Schnorr tests. The same proof works for computable randomness and Martin-Löf randomness, including lowness for pairs. For example, $A \in \text{Low}(\text{CR}, \text{SR})$ if and only if for every A -Schnorr test $(\mathcal{V}_n)_{n \in \mathbb{N}}$, there exists a single computable martingale d that succeeds against all $X \in \bigcap_n \mathcal{V}_n$.

4.4. Partial relativization and lowness for weak 2-randomness

The results we presented above can be extended to the case of weak 2-randomness by a *partial relativization*. A “partial relativization” of a computability concept \mathcal{C} to an oracle A consists in relativizing only some parts of the definition of \mathcal{C} to A . This device has already appeared in Section 4.2: $A \leq_{LR} B$ can be seen as a partial relativization of “ A is low for Martin-Löf randomness” to B . A full relativization would demand that every bounded $A \oplus B$ -c.e. open set can be covered by a bounded B -c.e. open set. Another interesting example was given by Cole and Simpson [3], who define a notion of *boundedly limit recursive in X* by partially relativizing the notion of ω -c.e.; they relativize the approximation function but not the computable bound on the number of mind changes. More examples are given in Barmpalias et al. [2].

The following partial relativization will be central to this section.

Definition 16. Let Z be a given oracle and let $(\mathcal{U}_e)_{e \in \mathbb{N}}$ be an effective enumeration of all c.e. open subsets of 2^ω . A $\langle Z \rangle$ -Martin-Löf test is a sequence $(\mathcal{U}_{f(n)})_{n \in \mathbb{N}}$ such that f is computable in Z and $\mu(\mathcal{U}_{f(n)}) \leq 2^{-n}$ for all n . A sequence $X \in 2^\omega$ is $\langle Z \rangle$ -Martin-Löf random if it passes all $\langle Z \rangle$ -Martin-Löf tests. We denote by $\text{MLR}(Z)$ the set of $\langle Z \rangle$ -Martin-Löf random sequences.

The concept of $\langle Z \rangle$ -Martin-Löf randomness is only a partial relativization of Martin-Löf randomness: a full relativization to Z would also allow, in the above definitions, the sets \mathcal{U}_e to be Z -c.e. open, as opposed to just c.e. open.

It turns out that Proposition 13 extends to partially relativized Martin-Löf tests.

Proposition 17. *For any $Z \in 2^\omega$, the hypotheses (P1, P2, P3) of Lemma 12 are satisfied when \mathbf{C} is the class of bounded c.e. open sets and $(\mathcal{T}^e)_{e \in \mathbb{N}}$ the family of $\langle Z \rangle$ -Martin-Löf tests.*

Proof. The proof is identical to the proof of Proposition 13, as the latter does not use the uniform enumerability of Martin-Löf tests, but solely the fact that $\mu(\mathcal{T}_n^{(e)}) \leq 2^{-n}$ for all e, n . \square

It follows from this proposition that, when it comes to lowness properties, partially relativized Martin-Löf randomness behaves exactly like Martin-Löf randomness (at least for the lowness properties discussed in this paper; it is not the case for some other notions. For example, the class $\text{Low}(\text{MLR}, \text{KR})$ is different from $\text{Low}(\text{W2R}, \text{KR})$, where KR is the class of Kurtz random sequences).

Proposition 18. For any Z in 2^ω , the following equalities hold.

- (i) $\text{Low}(\text{MLR}\langle Z \rangle, \text{MLR}) = \text{Low}(\text{MLR})$
- (ii) $\text{Low}(\text{MLR}\langle Z \rangle, \text{CR}) = \text{Low}(\text{MLR}, \text{CR})$
- (iii) $\text{Low}(\text{MLR}\langle Z \rangle, \text{SR}) = \text{Low}(\text{MLR}, \text{SR})$.

Proof. The proofs of the three items are almost identical. Let us prove for example item (ii). Since $\text{MLR}\langle Z \rangle \subseteq \text{MLR}$, it is clear by definition that $\text{Low}(\text{MLR}, \text{CR}) \subseteq \text{Low}(\text{MLR}\langle Z \rangle, \text{CR})$. Now take $A \notin \text{Low}(\text{MLR}, \text{CR})$. By the discussion of Section 4.1, this means that there exists an A -winning set U such that $[U]$ is covered by no bounded c.e. open set. Therefore, one can apply Lemma 12 with \mathbf{C} the class of bounded c.e. open sets and $(\mathcal{T}^e)_{e \in \mathbb{N}}$ the family of $\langle Z \rangle$ -Martin-Löf tests (which is possible, by Proposition 17), to get an $X \in U^\omega$ that passes all $\langle Z \rangle$ -Martin-Löf tests. So X is $\langle Z \rangle$ -Martin-Löf random, and, by Theorem 10 (relativized to A), X is not A -computably random. Thus $A \notin \text{Low}(\text{MLR}\langle Z \rangle, \text{CR})$. \square

Partial relativization of Martin-Löf randomness is useful to study higher notions of randomness. For example, given an effective enumeration $(\mathcal{U}_e)_{e \in \mathbb{N}}$, the set (e, k) of pairs such that $\mu(\mathcal{U}_e) < 2^{-k}$ is $\mathbf{0}'$ -enumerable (because $\mu(\mathcal{U}_e)$ is $\mathbf{0}'$ -computable, uniformly in e). Thus, given a generalized Martin-Löf test $(\mathcal{V}_n)_{n \in \mathbb{N}}$, one can $\mathbf{0}'$ -compute a sequence $k_1 < k_2 < k_3 < \dots$ such that $\mu(\mathcal{V}_{k_i}) < 2^{-i}$ for all i . Therefore the generalized Martin-Löf test $(\mathcal{V}_n)_{n \in \mathbb{N}}$ is covered by the $\langle \mathbf{0}' \rangle$ -Martin-Löf test $\mu(\mathcal{V}_{k_i})_{i \in \mathbb{N}}$. This shows that $\langle \mathbf{0}' \rangle$ -Martin-Löf randomness implies weak 2-randomness, and yields the following corollary.

Corollary 19. The following equalities hold.

- (i) $\text{Low}(\text{W2R}, \text{MLR}) = \text{Low}(\text{MLR})$
- (ii) $\text{Low}(\text{W2R}, \text{CR}) = \text{Low}(\text{MLR}, \text{CR})$
- (iii) $\text{Low}(\text{W2R}, \text{SR}) = \text{Low}(\text{MLR}, \text{SR})$.

Proof. Immediate from Proposition 18 (with $Z = \mathbf{0}'$) and the fact that $\text{MLR}\langle \mathbf{0}' \rangle \subseteq \text{W2R}$. \square

Item (iii) answers a question of Nies [18] (see also Franklin [7], Greenberg and Miller [9]). Although items (i) and (ii) were already known (proven respectively by Downey et al. [6] and Nies [18]), the proofs presented in this paper are simpler than the original ones. Note that Nies proved (ii) by showing that $\text{Low}(\text{W2R}, \text{CR}) = \text{Low}(\text{MLR})$, which is more than we show above. However, in Theorem 28 we give a short proof that $\text{Low}(\text{MLR}, \text{CR}) = \text{Low}(\text{MLR})$, so together with Corollary 19(ii), we have reproved the stronger result.

4.5. Highness notions

For a given pair \mathcal{R}, \mathcal{S} of randomness notions with $\mathcal{R} \subseteq \mathcal{S}$, the class $\text{Low}(\mathcal{R}, \mathcal{S})$ denotes the set of oracles $A \in 2^\omega$ that are weak enough to have $\mathcal{R} \subseteq \mathcal{S}^A$. It is natural to look at the dual concept of *highness*, i.e. the set of oracles $A \in 2^\omega$ that are powerful enough to have $\mathcal{S}^A \subseteq \mathcal{R}$, which we denote by $\text{High}(\mathcal{S}, \mathcal{R})$. These classes have primarily been studied by Franklin et al. [8] and Barmpalias et al. [2]. Most highness notions involving the classes W2R, MLR, CR, and SR have been characterized: Franklin et al. showed that the classes $\text{High}(\text{SR}, \mathcal{R})$ for $\mathcal{R} = \text{W2R}, \text{MLR}, \text{CR}$ are all equal to the set of A such that $A \geq_T \mathbf{0}'$; Barmpalias et al. proved that the class $\text{High}(\text{MLR}, \text{W2R})$ is equal to the set of sequences A such that no $\mathbf{0}'$ -computable function is diagonally non-computable relative to A . Miller et al. [16] have recently given another characterization of $\text{High}(\text{MLR}, \text{W2R})$; they proved that $A \notin \text{High}(\text{MLR}, \text{W2R})$ iff every partial A -computable function is dominated by a $\mathbf{0}'$ -computable function.

The classes $\text{High}(\text{CR}, \text{MLR})$ and $\text{High}(\text{CR}, \text{W2R})$ on the other hand are not fully understood yet. Franklin et al. proved that every $A \in \text{High}(\text{CR}, \text{MLR})$ computes a Martin-Löf random sequence. Kastermans, Lempp and Miller (unpublished) gave an alternative proof of this fact by showing that if $A \in \text{High}(\text{CR}, \text{MLR})$, then there is an A -computable martingale that succeeds against all non-Martin-Löf random sequences. This result is an easy consequence of Lemma 12; we present the proof here.

Proposition 20. Let $A \in 2^\omega$. The following are equivalent.

- (i) $A \in \text{High}(\text{CR}, \text{MLR})$
- (ii) Every bounded c.e. open set is covered by an A -winning set.
- (iii) The first level \mathcal{U}_1 of a universal Martin-Löf test is covered by an A -winning set.
- (iv) There exists an A -computable martingale that succeeds against all X that are not Martin-Löf random.

Proof. (i) \Rightarrow (ii). If (ii) does not hold, there is a bounded c.e. open set \mathcal{U} that cannot be covered by any A -winning open set. Let U be a c.e. prefix-free set of strings generating \mathcal{U} . We can apply Lemma 12 with \mathbf{C} the class of A -winning open sets and \mathcal{T} the family of tests induced by exactly A -computable martingales. To see that this is allowed, relativize Proposition 13 to A . Lemma 12 gives us a sequence $X \in U^\omega$ that passes all tests induced by exactly A -computable martingales (hence is A -computably random). By Theorem 8, X is not Martin-Löf random, so $A \notin \text{High}(\text{CR}, \text{MLR})$.

(ii) \Rightarrow (iii) is immediate, as is (iv) \Rightarrow (i). This leaves (iii) \Rightarrow (iv). Let U be an A -winning set such that $[U]$ covers \mathcal{U}_1 . Note that U is prefix-free, by definition. The proof of (iii) \Rightarrow (i) in Theorem 10 (relativized to A) produces an A -computable martingale d that succeeds against all $X \in U^\omega$. Now assume that X is not Martin-Löf random. Then all tails of X are contained in \mathcal{U}_1 , hence in $[U]$. This implies that $X \in U^\omega$, so d succeeds on X . \square

To see that **Proposition 20** implies the Franklin, Stephan and Yu result, note that every martingale computes a sequence on which it does not succeed. Hence, if $A \in \text{High}(\text{CR}, \text{MLR})$, then A computes a Martin-Löf random sequence. In fact:

Corollary 21. *If $A \in \text{High}(\text{CR}, \text{MLR})$, then there is an A -Turing functional $\Phi : 2^\omega \rightarrow 2^\omega$ that is total, one-to-one, and such that for all $X \in 2^\omega$, $\Phi(X) \in \text{MLR}$.*

Proof. If $A \in \text{High}(\text{CR}, \text{MLR})$, let (by **Proposition 20**) d be a normed A -computable martingale that succeeds on all X that are not Martin-Löf random. We shall build an A -computable tree, i.e., an A -computable total function $T : 2^{<\omega} \rightarrow 2^{<\omega}$ such that if σ' is a strict extension of σ , then $T(\sigma')$ is a strict extension of $T(\sigma)$ and if σ and σ' are incomparable then so are $T(\sigma)$ and $T(\sigma')$. We ensure that d succeeds on no infinite path in this tree by imposing the condition that for any $\sigma \in 2^{<\omega}$, setting $\tau = T(\sigma)$, we have $d(\tau') \leq 2 - 2^{-|\sigma|}$ for all prefixes τ' of τ . Then, it follows immediately that for any infinite path $Y \in 2^\omega$ in T , we have $d(Y \upharpoonright n) \leq 2$ for all n .

The construction of T is done by induction on the length k of σ . We first set $T(\epsilon) = \epsilon$. As $d(\epsilon) = 1$ (d is normed) this satisfies the requirement. Now, suppose we have defined $T(\sigma)$ for all σ of length k respecting the above requirement. Let σ be of length k and set $\tau = T(\sigma)$. We have by assumption $d(\tau) \leq 2 - 2^{-k}$. Hence, by the Ville-Kolmogorov inequality, the set of sequences $X \in 2^\omega$ that extend τ and, for all n , satisfy $d(X \upharpoonright n) < 2 - 2^{-(k+1)}$ has positive measure. Since d is A -computable, using oracle A we can find two incomparable extensions τ_0 and τ_1 of τ such that $d(\tau') \leq 2 - 2^{-(k+1)}$ for any prefix τ' of τ_0 or τ_1 . We then set $T(\sigma \iota) = \tau_\iota$ for $\iota \in \{0, 1\}$. This concludes the induction.

Finally, the A -functional Φ is defined in a straightforward manner by setting $\Phi^\sigma = T(\sigma)$ for all $\sigma \in 2^{<\omega}$. \square

5. Reformulation in other contexts: converging series and machines

5.1. Randomness via machines and Kolmogorov complexity

Although we had no need for it so far in this paper, it is well known that Kolmogorov complexity provides an alternative and elegant way to characterize randomness. In particular, Levin and Schnorr independently showed that a sequence $X \in 2^\omega$ is Martin-Löf random if and only if $K(X \upharpoonright n) \geq n - O(1)$, where K denotes prefix-free Kolmogorov complexity.

In this last section, we discuss how our results relate to Kolmogorov complexity, and more precisely to prefix-free complexity. A *prefix-free machine* is a partial computable function $M : 2^{<\omega} \rightarrow 2^{<\omega}$ whose domain is prefix-free and K_M is the Kolmogorov complexity associated to M . As usual, we fix an optimal prefix-free machine M_{opt} ; we abbreviate $K_{M_{opt}}$ by K and call it *prefix-free Kolmogorov complexity*.

It is also well known that prefix-free Kolmogorov complexity is tightly related to convergent series. By the Kraft-Chaitin theorem, if $f : \mathbb{N} \rightarrow \mathbb{R}^{\geq 0}$ is summable (i.e., $\sum_n f(n) < +\infty$) and left-c.e., then $K \leq -\log f + O(1)$. Levin's coding theorem proves that there exists a maximal left-c.e. summable function $F : \mathbb{N} \rightarrow \mathbb{R}^{\geq 0}$ (i.e., for any other such function f , $f = O(F)$), and that we precisely have $K = -\log F + O(1)$.

Nies [17] proved that a sequence A is low for K (i.e., K^A , the prefix-free Kolmogorov complexity relativized to the oracle A , is equal to the unrelativized version K , up to an additive constant) if and only if A is low for Martin-Löf randomness. By Levin's coding theorem, this can be rephrased as follows.

Theorem 22. *The following are equivalent:*

- (i) $A \in \text{Low}(\text{MLR})$.
- (ii) For every A -left-c.e. summable function $f : \mathbb{N} \rightarrow \mathbb{R}^{\geq 0}$, there exists a left-c.e. summable function g such that $f \leq g$.

Remark 23. Condition (ii) can be replaced by the seemingly weaker condition on A : "For every A -computable summable function $f : \mathbb{N} \rightarrow \mathbb{R}^{\geq 0}$, there exists a left-c.e. summable function g such that $f \leq g$ ". Indeed, if f is an A -left-c.e. summable function, then defining $\tilde{f}(\langle i, t \rangle)$ to be the increase of $f(i)$ at stage t , the function \tilde{f} is computable and summable (its sum is the same as that of f), and if h is a left-c.e. summable function that dominates \tilde{f} , then for all i , $f(i)$ is dominated by $g(i) = \sum_t h(\langle i, t \rangle)$ and it is clear that g is left-c.e. and summable (its sum is the same as that of h).

Using techniques similar to Nies', we can prove the analogous results for $\text{Low}(\text{SR})$ and $\text{Low}(\text{MLR}, \text{SR})$.

Proposition 24. *The following are equivalent:*

- (i) $A \in \text{Low}(\text{SR})$.
- (ii) For every A -computable function $f : \mathbb{N} \rightarrow \mathbb{R}^{\geq 0}$ whose sum is finite and computable, there exists a computable function g whose sum is finite and computable and such that $f \leq g$.

Proposition 25. *The following are equivalent:*

- (i) $A \in \text{Low}(\text{MLR}, \text{SR})$.
- (ii) For every A -computable function $f : \mathbb{N} \rightarrow \mathbb{R}^{\geq 0}$ whose sum is finite and computable, there exists a left-c.e. summable function g such that $f \leq g$.

Proof. We prove Proposition 25 only, the proof of Proposition 24 being similar.

(ii) \Rightarrow (i). Let $A \in 2^\omega$ satisfying the hypotheses of (ii). Let $(\mathcal{U}_n)_{n \in \mathbb{N}}$ be an A -Schnorr test. We can assume that every \mathcal{U}_n is generated by an A -computable subset U_n of $2^{<\omega}$ (here we use the well-known fact that every c.e. open set \mathcal{U} is generated by a computable subset U of $2^{<\omega}$, and an index for U can be computed from an index of \mathcal{U}). Then, define the function $f : 2^{<\omega} \rightarrow \mathbb{R}^{\geq 0}$ by $f(\sigma) = 2^{-|\sigma|}n$ where n is the largest integer such that $\sigma \in U_n$ and $f(\sigma) = 0$ if σ belongs to no set U_n . Note that f is A -computable since $\sigma \in U_n$ implies $n \leq \sigma$. Moreover,

$$\sum_{\sigma} f(\sigma) \leq \sum_n \sum_{\sigma \in U_n} 2^{-|\sigma|}n \leq \sum_n n \mu(\mathcal{U}_n) \leq \sum_n n2^{-n} < +\infty.$$

We can thus apply the hypothesis of (ii) (where \mathbb{N} and $2^{<\omega}$ are identified) to get a summable left-c.e. function $g : 2^{<\omega} \rightarrow \mathbb{R}^{\geq 0}$ such that $f \leq g$. Then define, for all n ,

$$V_n = \{\sigma : g(\sigma) > 2^{-|\sigma|}(n/2)\}$$

and set $\mathcal{V}_n = [V_n]$. It is clear that \mathcal{V}_n is a c.e. open set, uniformly in n , that covers \mathcal{U}_n . Let S be the sum of g . We have:

$$\mu(\mathcal{V}_n) \leq \sum_{\sigma \in V_n} 2^{-|\sigma|} < \sum_{\sigma \in V_n} \frac{2g(\sigma)}{n} \leq \frac{2S}{n}.$$

Thus $(\mathcal{V}_n)_{n \in \mathbb{N}}$ is a Martin-Löf test (in the general sense of Remark 2) that covers $(\mathcal{V}_n)_{n \in \mathbb{N}}$. We have shown that every A -Schnorr test is covered by a Martin-Löf test, so $A \in \text{Low}(\text{MLR}, \text{SR})$.

(ii) \Rightarrow (i). For this proof it is convenient to identify 2^ω with the space $[0, 1]^\mathbb{N}$ (using the usual identification of 2^ω to $(2^\omega)^\mathbb{N}$ and then of 2^ω to $[0, 1]$; the non-uniqueness of binary expansion for dyadic rationals does not cause any problems here).

For all $n \in \mathbb{N}$ and $\alpha \in [0, 1]$, set

$$\mathcal{B}_{n,\alpha} = \{X \in [0, 1]^\mathbb{N} : X_n \in [0, \alpha]\}.$$

Now, let $f : \mathbb{N} \rightarrow \mathbb{R}^{\geq 0}$ be an A -computable function whose sum is finite and A -computable. Without loss of generality, we can assume that $f(n) \leq 1$ for all n (otherwise we divide f by a constant C to make this true, then find the desired function g that dominates f/C , and observe that $C \cdot g$ dominates f and is as wanted). Consider the set

$$\mathcal{U} = \bigcup_n \mathcal{B}_{n,f(n)}.$$

\mathcal{U} is a c.e. open set. Also, observe that if $n \neq m$, the sets $\mathcal{B}_{n,\alpha}$ and $\mathcal{B}_{m,\beta}$ correspond to independent events. Thus,

$$\mu(\mathcal{U}) = 1 - \prod_n (1 - \mu(\mathcal{B}_{n,f(n)})) = 1 - \prod_n (1 - f(n)).$$

This can be reformulated as

$$\log(1 - \mu(\mathcal{U})) = \sum_n \log(1 - f(n)).$$

Since $f(n)$ tends to 0, we have $\log(1 - f(n)) \sim -f(n)$. This implies that $\sum_n \log(1 - f(n))$ is finite and A -computable (as $\sum_n f(n)$ is). Thus, \mathcal{U} is a bounded A -Schnorr open set. Since $A \in \text{Low}(\text{MLR}, \text{SR})$, we know that \mathcal{U} must be covered by some bounded c.e. open set \mathcal{V} . Having such a set, we define

$$g(n) = \sup\{\alpha \in [0, 1] : \mathcal{B}_{n,\alpha} \subseteq \mathcal{V}\}.$$

It is clear that $f \leq g$, and that g is left-c.e. as \mathcal{V} is a c.e. open set. The sum $\sum_n g(n)$ is bounded because $\prod_n (1 - g(n)) > 0$, the latter being equal to $1 - \mu(\bigcup_n \mathcal{B}_{n,g(n)}) \geq 1 - \mu(\mathcal{V}) > 0$. \square

Downey and Griffiths [5] gave a Levin–Schnorr-like characterization of Schnorr randomness by restricting Kolmogorov complexity to a specific class of prefix-free machines. They proved that a sequence $X \in 2^\omega$ is Schnorr random if and only if for every computable measure machine M , one has $K_M(X \upharpoonright n) \geq n - O(1)$, where a *computable measure machine* is a prefix-free machine whose domain has computable measure (i.e., is a Schnorr set). Furthermore, Downey et al. [4] showed that an analogue of Nies’ result “low for random equals low for K ” holds for Schnorr randomness, as explained in the next proposition.

Proposition 26. *The following are equivalent:*

- (i) $A \in \text{Low}(\text{SR})$
- (ii) For any A -computable measure machine M , there exists a computable measure machine M' such that $K_{M'} \leq K_M + O(1)$.

It turns out that the results we proved earlier in this section allow us to give a short proof of the above.

Proof. The part (ii) \Rightarrow (i) is clear from the Downey–Griffith characterization of Schnorr randomness. For the reverse direction, let M be an A -computable measure machine. Let $f : 2^{<\omega} \rightarrow \mathbb{R}^{\geq 0}$ be the function defined by

$$f(\sigma) = 2^{-K_M(\sigma)}$$

It is easy to see that f is an A -left-c.e. function. Moreover, if we enumerate the domain of M , whenever a new p is found (i.e., at that stage the measure of $\text{dom}(M)$ is increased by $2^{-|p|}$, this increases the sum $\sum_{\sigma} f(\sigma)$ by either 0 or $2^{-|p|}$ (depending on whether we had already enumerated some p' with $|p'| \leq |p|$ and $M(p) = M(p')$ or not before that stage). This shows that the sum $\sum_{\sigma} f(\sigma)$ can be computed from the measure of $\text{dom}(M)$, which is A -computable. Hence, $\sum_{\sigma} f(\sigma)$ is A -computable. We can therefore apply [Proposition 24](#) to get a left-c.e. summable function $g : 2^{<\omega} \rightarrow \mathbb{R}^{\geq 0}$ whose sum is computable and such that $f \leq g$. Let $c \in \mathbb{N}$ be a constant such that $\sum_{\sigma} f(\sigma) \leq 2^c$. We enumerate the Kraft–Chaitin set

$$L = \{(k, \sigma) : g(\sigma) \geq 2^{-k+c+1}\}.$$

We have

$$\sum_{(k,\sigma) \in L} 2^{-k} = \sum_{\sigma} 2^{-\lceil \log g(\sigma) - c - 1 \rceil + 1} \leq 2^{-c} \sum_{\sigma} g(\sigma) \leq 1,$$

so L is indeed a Kraft–Chaitin set. Now, apply the Kraft–Chaitin theorem to construct a machine M' whose domain is a prefix-free set $\{p_{k,\sigma} : (k, \sigma) \in L\}$ with $|p_{k,\sigma}| = k$ and $M'(p_{k,\sigma}) = \sigma$. It follows by construction that M' is a computable measure machine (the measure of its domain is $\sum_{\sigma} 2^{-\lceil \log g(\sigma) - c - 1 \rceil + 1}$, which is computable because $\sum_{\sigma} g(\sigma)$ is computable) and

$$K_{M'} \leq -\log g + c + 1 \leq -\log f + c + 1 \leq K_M + c + 1.$$

This completes the proof. \square

In the same way, we can get the analogous result for the pair (MLR, SR).

Proposition 27. *The following are equivalent:*

- (i) $A \in \text{Low}(\text{MLR}, \text{SR})$
- (ii) For any A -computable measure machine M , we have $K \leq K_M + O(1)$.

The proof is almost identical to the proof of the previous result (using [Proposition 25](#) instead of [Proposition 24](#)) and is left to the reader.

5.2. A final application: $\text{Low}(\text{MLR}) = \text{Low}(\text{MLR}, \text{CR})$

We finish with an alternative proof that $\text{Low}(\text{MLR}) = \text{Low}(\text{MLR}, \text{CR})$. This was shown by Nies [[17,18](#)], but the only known proof is long and technical. We believe that our proof is more comprehensible. It uses the covering characterization of $\text{Low}(\text{MLR}, \text{CR})$ and the characterization of $\text{Low}(\text{MLR})$ via summable series (which as we pointed out, is an easy consequence of the coding theorem). Together with [Corollary 19\(ii\)](#), we in fact have an alternate proof that $\text{Low}(\text{MLR}) = \text{Low}(\text{W2R}, \text{CR})$ (Nies [[18](#)]).

Theorem 28. *The classes $\text{Low}(\text{MLR})$ and $\text{Low}(\text{MLR}, \text{CR})$ coincide.*

Proof. It is clear that $\text{Low}(\text{MLR})$ is contained in $\text{Low}(\text{MLR}, \text{CR})$. We need to show the reverse implication. Let $A \in \text{Low}(\text{MLR}, \text{CR})$. To show that $A \in \text{Low}(\text{MLR})$, we use [Theorem 22](#) and [Remark 23](#): we consider an A -computable summable function $f : \mathbb{N} \rightarrow \mathbb{R}^{\geq 0}$ and we will show that there exists a left-c.e. summable function g such that $f \leq g$. Without loss of generality, we assume that $\sum_i f(i) \leq 1$ and that the $f(i)$ are dyadic rational numbers in virtue of which we set $f(i) = 2^{-a_i}$, the sequence of a_i being A -computable. The proof's strategy is the following: we shall "encode" the function f in an A -winning open set V . Then, since $A \in \text{Low}(\text{MLR}, \text{CR})$, there exists a bounded c.e. open set \mathcal{W} that covers V , and from \mathcal{V} we will build a left-c.e. summable function g that dominates f .

To construct the set \mathcal{U} , we first consider a computable partition of \mathbb{N} into intervals $I_{i,l}$ where for all i, l , the length of $I_{i,l}$ is l (the order in which the intervals are placed does not matter). For all i, l , we consider the set $\mathcal{Z}_{i,l}$ of sequences X such that $X(n) = 0$ for all $n \in I_{i,l}$. Note that $\mu(\mathcal{Z}_{i,l}) = 2^{-l}$ and the $\mathcal{Z}_{i,l}$ are pairwise independent. Then, we set

$$\mathcal{U} = \bigcup_i \mathcal{Z}_{i,a_i}$$

Since the \mathcal{Z}_{i,a_i} are all independent, the measure of \mathcal{U} is $1 - \prod_i (1 - 2^{-a_i})$, which is smaller than 1 as the sum $\sum_i 2^{-a_i}$ is finite. We now show that some A -computable martingale wins money against every member of \mathcal{U} . Let $q > 1$ be a rational number such that $\sum_i 2^{-a_i} < 1/q$. We define an A -computable martingale d as follows. For each i , we reserve an amount $q2^{-a_i}$. When betting on a sequence X , at the start of an interval $I_{i,l}$, the martingale d check whether $l = a_i$. If not, d does not bet on any position of the interval. If so, then d uses the reserved capital $q2^{-a_i}$ to bet that X contains only zeroes on the interval I_{i,a_i} . This is done by first betting an amount $x = q2^{-a_i}$ on 0, then if correct an amount $2x$, then $4x$, etc., stopping if some guess was incorrect. If all of its guesses are correct, the capital of d has increased by $x2^{a_i} = q$ at the end of the interval I_{i,a_i} . Thus, the martingale d is A -computable and reaches a capital of at least q on every element X of \mathcal{U} . Hence \mathcal{U} is contained in the winning set $\mathcal{V} = [V]$ with

$$V = \{\sigma \text{ minimal s.t. } d(\sigma) \geq q\}$$

Now, since $A \in \text{Low}(\text{MLR}, \text{CR})$, there exists an unrelativized bounded c.e. open set \mathcal{W} that covers \mathcal{V} . Then, set for all i , $g(i) = 2^{-b_i}$ with

$$b_i = \min\{l \mid \mathcal{Z}_{i,l} \subseteq \mathcal{W}\}$$

The sequence b_i is right-c.e. and by construction $b_i \leq a_i$ as $\mathcal{Z}_{i,a_i} \subseteq \mathcal{U} \subseteq \mathcal{W}$, so $g(i) \geq f(i)$. Finally, notice that the sum $\sum_i 2^{-b_i}$ is finite as

$$1 > \mu(\mathcal{W}) \geq 1 - \prod_i (1 - 2^{-b_i})$$

hence $\prod_i (1 - 2^{-b_i}) > 0$ so $\sum_i 2^{-b_i}$ converges. Therefore the function g is as wanted. \square

Acknowledgement

Most of this work was done while the first author was a von Humboldt postdoctoral fellow at the Institut für Informatik, Ruprecht-Karls Universität Heidelberg, Germany. The second author was supported by the National Science Foundation under grants DMS-0945187 and DMS-0946325, the latter being part of a Focused Research Group in Algorithmic Randomness.

References

- [1] George Barmpalias, Andrew E.M. Lewis, Mariya Soskova, Randomness, lowness and degrees, *Journal of Symbolic Logic* 73 (2) (2008) 559–577.
- [2] George Barmpalias, Joseph S. Miller, André Nies, Randomness notions and partial relativizations (in press).
- [3] Joshua A. Cole, Stephen G. Simpson, Mass problems and hyperarithmeticality, *J. Math. Log.* 7 (2) (2007) 125–143.
- [4] Rodney Downey, Noam Greenberg, Nenad Mihailovic, André Nies, Lowness for computable machines, in: *Computational Prospects of Infinity: Part II. Presented Talks*, National University of Singapore, Institute for Mathematical Sciences, in: *Lecture Notes Series. Institute for Mathematical Sciences. National University of Singapore*, vol. 15, World Scientific Publishing Company, Singapore, 2008, pp. 79–86.
- [5] Rodney Downey, Evan Griffiths, Schnorr randomness, *Journal of Symbolic Logic* 69 (2) (2004) 553–554.
- [6] Rodney Downey, André Nies, Rebacca Weber, Liang Yu, Lowness and Π_2^0 -nullsets, *Journal of Symbolic Logic* 71 (3) (2006) 1044–1052.
- [7] Johanna N.Y. Franklin, Lowness and highness properties for randomness notions, in: T. Arai, et al. (Eds.), *Proceedings of the 10th Asian Logic Conference*, World Scientific, 2010, pp. 124–151.
- [8] Johanna N.Y. Franklin, Frank Stephan, Liang Yu, Relativizations of randomness and genericity notions, *Bull. London Math. Soc.* 43 (4) (2011) 721–733.
- [9] Noam Greenberg, Joseph S. Miller, Lowness for Kurtz randomness, *Journal of Symbolic Logic* 74 (2) (2009) 665–678.
- [10] Bjørn Kjos-Hanssen, Low for random reals and positive-measure domination, *Proceedings of the American Mathematical Society* 135 (11) (2007) 3703–3709.
- [11] Bjørn Kjos-Hanssen, Joseph S. Miller, Reed Solomon, Lowness notions, measure, and domination (in press).
- [12] Bjørn Kjos-Hanssen, André Nies, Frank Stephan, Lowness for the class of Schnorr random reals, *SIAM Journal of Computing* 35 (3) (2005) 647–657.
- [13] Antonín Kučera, Measure, Π_1^0 classes, and complete extensions of PA, *Lecture Notes in Mathematics* 1141 (1985) 245–259.
- [14] Antonín Kučera, Sebastiaan A. Terwijn, Lowness for the class of random sets, *Journal of Symbolic Logic* 64 (4) (1999) 1396–1402.
- [15] Jack Lutz, Computability versus exact computability of martingales, *Information Processing Letters* 92 (2004) 235–237.
- [16] Joseph S. Miller, Keng Meng Ng, Nicholas Rupprecht, Notions of effectively null and their covering properties (in preparation).
- [17] André Nies, Lowness properties and randomness, *Advances in Mathematics* 197 (1) (2005) 274–305.
- [18] André Nies, *Computability and Randomness*, in: *Oxford Logic Guides*, vol. 51, Oxford University Press, Oxford, 2009.
- [19] Sebastiaan Terwijn, Domenico Zambella, Computational randomness and lowness, *Journal of Symbolic Logic* 66 (3) (2001) 1199–1205.
- [20] Domenico Zambella, *Sequences with simple initial segments*. Technical report ML-1990-05, ILLC Publications, Mathematical Logic and Foundations (ML) Series, 1990.