

Alterations in the Gut Microbiota Associated with HIV-1 Infection

Catherine A. Lozupone,¹ Marcella Li,¹ Thomas B. Campbell,¹ Sonia C. Flores,¹ Derek Linderman,¹ Matthew J. Gebert,³ Rob Knight,^{2,3} Andrew P. Fontenot,¹ and Brent E. Palmer^{1,*}

¹Department of Medicine, University of Colorado Denver, Anschutz Medical Campus, Aurora, CO 80045, USA

²Howard Hughes Medical Institute

³Department of Chemistry and Biochemistry and Biofrontiers Institute, University of Colorado, Boulder, CO 80309, USA

*Correspondence: brent.palmer@ucdenver.edu

<http://dx.doi.org/10.1016/j.chom.2013.08.006>

SUMMARY

Understanding gut microbiota alterations associated with HIV infection and factors that drive these alterations may help explain gut-linked diseases prevalent with HIV. 16S rRNA sequencing of feces from HIV-infected individuals revealed that HIV infection is associated with highly characteristic gut community changes, and antiretroviral therapy does not consistently restore the microbiota to an HIV-negative state. Despite the chronic gut inflammation characteristic of HIV infection, the associated microbiota showed limited similarity with other inflammatory states and instead showed increased, rather than decreased, diversity. Meta-analysis revealed that the microbiota of HIV-infected individuals in the U.S. was most similar to a *Prevotella*-rich community composition typically observed in healthy individuals in agrarian cultures of Malawi and Venezuela and related to that of U.S. individuals with carbohydrate-rich, protein- and fat-poor diets. By evaluating innate and adaptive immune responses to lysates from bacteria that differ with HIV, we explore the functional drivers of these compositional differences.

INTRODUCTION

HIV-1 infection induces rapid and substantial damage to gut-associated lymphoid tissues (GALTs), with massive depletion of Th17 cells, a subset of CD4⁺ T cells that control intestinal bacteria (Brenchley et al., 2004). Loss of this CD4⁺ T cell subset may facilitate disease progression by allowing translocation of bacterial products such as lipopolysaccharide to blood, causing systemic T cell activation (Brenchley et al., 2006).

In healthy individuals, intestinal bacteria help control harmful pathogens, educate the immune system, and aid in digestion (Lozupone et al., 2012), but distinctive compositions of this complex community have been associated with pathologies that are of increased prevalence with HIV infection, including metabolic disorders (Madge et al., 1999; Vijay-Kumar et al., 2010), chronic inflammation (Bjarnason et al., 1996; Willing et al., 2010), wasting/malnutrition (Smith et al., 2013; Wanke et al., 2000),

atherosclerosis (Hsue et al., 2009; Koeth et al., 2013), and susceptibility to opportunistic infections (Chang et al., 2008). Considering the important role that both innate and adaptive immunity play in shaping gut microbiota composition (Carvalho et al., 2012; Hepworth et al., 2013), understanding how composition differs with HIV infection, immunologic driving factors of these differences, and their implication for gut-linked diseases that increase in prevalence with HIV is of paramount importance.

Using 16S rRNA sequencing of feces, we show that HIV infection is associated with highly characteristic changes in gut community structure and that antiretroviral therapy (ART) does not consistently restore the microbiota to an HIV-negative like state. Using meta-analysis, we demonstrate that although the HIV-associated microbiota bears some resemblance to that of other chronic inflammatory states, the differences between HIV-associated and healthy gut microbiota are also related to compositional changes previously associated with diet (Wu et al., 2011) and Western compared to agrarian cultures (De Filippo et al., 2010; Yatsunenken et al., 2012), suggesting an interplay between diet and the immune system in shaping gut microbiota composition. By comparing T cell proliferation and cytokine responses to lysates from bacteria that differ with HIV infection in blood from HIV-positive and -negative individuals, we begin to explore the relationship between immune dysfunction and HIV-associated microbiota change.

RESULTS

Stool Microbiota in Chronic Untreated HIV Infection Is Distinctive

We sequenced the V4 region of 16S rRNA genes from fecal samples of individuals with chronic HIV infection ($n = 22$), recent infection ($n = 3$), and HIV-negative controls ($n = 13$) (Table 1, see Table S1 online). Eleven individuals with chronic infection had not recently undergone ART, and eight had been on ART for ≥ 12 months with viral suppression for ≥ 6 months (see the Experimental Procedures for further cohort details). Individuals were excluded during recruitment if they were pregnant, weighed < 110 pounds, or had received antibiotics within the prior 30 days, resulting in inclusion of only relatively healthy HIV-positive subjects; no chronically HIV-infected individual had a peripheral CD4⁺ T cell count below 204 cells/ μ l (Table 1).

Principal Coordinates Analysis (PCoA) of fecal samples based on 16S rRNA sequences using both unweighted (Figure 1) and weighted (Figure S1) UniFrac showed a strong difference in

Table 1. Description of Cohorts

Cohort	N	Age (Years)	Sex (M, Male; F, Female)	Race					BMI	Duration ART (Months)	CD4 (Cells/ μ L)	Viral Load (Copies/mL)
				W	Wh	AA	AI	?				
1: Recent HIV infection	3	31 \pm 4.36	M, 2; F, 1	3	0	0	0	0	32.5 \pm 8.30	0	614 \pm 521.8 (107–1,342)	164,880 \pm 210,225 (418–449,000)
2: Chronic HIV untreated	11	30 \pm 5.61	M, 11; F, 0	6	2	2	1	0	25.0 \pm 4.83	0	551 \pm 218.2 (270–1095)	139,109 \pm 255,929 (1,350–869,000)
2*: Chronic HIV short-term ART	6 ^a	34 \pm 10.27	M, 5; F, 1	3	2	1	0	0	27.1 \pm 3.67	7.68 \pm 0.78	609 \pm 50.6 (547–681)	83 \pm 105.0 (39–247)
3: Chronic HIV long-term ART	8	45 \pm 8.30	M, 7; F, 1	5	2	1	0	0	24.7 \pm 6.45	33.6 \pm 17.97	483 \pm 258.0 (204–876)	124 \pm 280.8 (<20–819)
4: HIV-seronegative	13	37 \pm 12.26	M, 8; F, 5	10	1	1	0	1	25.6 \pm 3.03	0	NA	0

Race abbreviations are W (white, not Hispanic), Wh (white-Hispanic), AA (African American), AI (American Indian), and ? (unknown). See also Table S1. Values are given as mean \pm SD with ranges in parentheses.

^aThree of the six individuals in cohort 2* are also in cohort 2 because they were sampled both before and after short-term ART treatment.

microbiota of individuals with chronic untreated HIV infection compared to HIV-negative controls. UniFrac evaluates microbiota similarity based on shared evolutionary history of bacterial taxa, and the weighted metric differs from unweighted because it accounts for change in relative abundance in addition to presence or absence (Lozupone and Knight, 2005; Lozupone et al., 2007). Fecal microbiota with chronic untreated HIV infection also had greater overall diversity, with higher evenness as measured by the Shannon diversity index ($p = 0.013$) and richness as measured by phylogenetic diversity (PD) ($p = 0.0003$) (Table 2).

To further determine the degree to which gut microbiota with chronic untreated infection differed from HIV-negative controls, we used a supervised learning technique called Random Forests (Knights et al., 2011a). The purpose of the Random Forests classifier is to learn a function that maps a set of predictive features to a discrete state, in this case healthy versus chronically infected with HIV and without ART. We used as features the relative abundance in each sample of 97% identity (ID) Operational Taxonomic Units (OTUs; clusters in which sequences have $\geq 97\%$ identity over their aligned 16S rRNA genes, which approximates assignment to the same species; Stackebrandt and Goebal, 1994). The measure of the method's success is its ability to classify new samples as coming from an HIV-positive or -negative individual. We only used one sample per individual when multiple time points were collected. The model could classify unknown samples according to HIV status with a $4.33\% \pm 0.821\%$ error rate, which is 10.6 times higher than the baseline error rate for random guessing of 45.83%, indicating that HIV-associated microbiota exhibit highly characteristic differences.

To investigate whether disease severity impacted gut microbiota in HIV infection, we determined whether microbiota diversity correlated with peripheral CD4+ T cell count or plasma HIV-1 RNA viral load in 11 individuals with chronic infection who were not on ART. As an estimate of divergence from healthy, we calculated the average pairwise weighted and unweighted UniFrac distance between each HIV microbiota sample and HIV-negative control samples. The magnitude of divergence from HIV-negative microbiota samples was not correlated with CD4+ T cell count or plasma HIV-1 RNA viral load (all Spearman correlation p values > 0.73).

Individuals Recently Infected with HIV Have Relatively Subtle Microbiota Differences

The microbiota of recently infected untreated individuals differed little from uninfected controls. Samples from two of three recently infected individuals clustered in an intermediate position between healthy and untreated chronic infection in an unweighted UniFrac PCoA plot (Figure 1A). However, these individuals clustered with chronically infected subjects when using weighted UniFrac (Figure S1), suggesting that changes in relative abundance of existing taxa may occur faster than gain or loss of taxa.

Effects of ART Treatment

No chronically infected individual sampled pre and post 6.7–8.5 months of ART exhibited a strong shift toward the HIV-negative individuals (Figure 1A), indicating that short-term ART was insufficient to restore the microbiota. However, the microbiota of two individuals treated for 7.1–8.5 months for whom we did not collect pre-ART samples showed a closer resemblance to HIV-negative individuals than to subjects with untreated chronic HIV-1 infection (Figure 1B). Of the eight HIV-chronic individuals on long-term ART (12.1–57.6 months), five clustered with untreated chronically infected individuals, and three clustered with HIV-negative individuals (Figure 1B). Individuals on ART for as long as 57.6 months could still cluster with HIV-positive subjects with chronic infection, indicating that ART duration alone does not predict restoration to a HIV-negative phenotype. Untreated HIV-infected individuals had significantly higher alpha diversity (PD and evenness) compared to ART-treated and HIV-negative individuals, which were not significantly different from each other (Table 2).

Taxa that Discriminate between HIV-Negative Individuals and Chronic HIV Infection

The fecal microbiota of untreated individuals with chronic infection exhibited significantly higher relative abundances of Prevotellaceae (Prevotella), Erysipelotrichaceae (Catenibacterium and Bulleidia), Veillonellaceae (Dialister and Mitsuoakella), Clostridium cluster XIII, and the genus Desulfovibrio compared to HIV negative (Figure 1, Figure 2, Table S2). HIV-negative individuals had increased Bacteroidaceae (Bacteroides), Rikenellaceae

Table 2. Comparison of Alpha Diversity in HIV-Positive and -Negative Individuals with and without ART Treatment

	Observed Species	Shannon	PD
HIV negative	508 ± 21.6	5.6 ± 0.19	27.9 ± 1.46
Chronic HIV, ART untreated	563 ± 30.5	6.3 ± 0.15	36.4 ± 1.31
Chronic HIV on long-term ART	469 ± 39.6	5.68 ± 0.16	28.69 ± 1.19
Recent HIV	517 ± 64.4	5.76 ± 0.49	31.7 ± 3.06
p value (ART untreated versus negative) t test	0.144	0.013	0.0003
p value (ART untreated versus ART treated) t test	0.070	0.016	0.0006

Values are given as mean ± SD.

with *Bacteroides*- versus *Prevotella*-rich enterotypes (Wu et al., 2011).

PCoA clustering with unweighted UniFrac was largely driven by technical variation between studies (Figure 3A), as might be expected since they varied methodologically (Table S3) (Lozupone et al., 2013). However, by coloring the points by the relative abundance of *Bacteroides* (Figure 3B) and *Prevotella* (Figure 3C), it is clear that the major diversity differences within all three data sets are associated with a tradeoff between these genera along the same axis of variation. Samples from HIV-positive individuals from the U.S. clustered closer to individuals from Malawi and Venezuelan Amerindians than to HIV-negative individuals from the U.S. (Figure 3D). Additionally, the discriminative 97% ID OTUs that differentiated adults from Western and agrarian cultures using Random Forests analysis (Yatsunenkov et al., 2012) were highly related and sometimes identical to those that discriminated between HIV-negative and -positive individuals (Figure S2). This was particularly evident for the *Bacteroidales* order, where Western and HIV-negative associated OTUs clustered together in the *Bacteroides* and *Alistipes* genera and agrarian and HIV-positive associated OTUs clustered together in the *Prevotella* genus (Figure S2A). However, consistent with the notion that *Bacteroides* and *Prevotella* are indicator taxa for more complex community assemblages, parallels between agrarian cultures and HIV-positive individuals were also observed in other taxonomic groups (see Figure S2 for more details).

Since HIV infection induces chronic gut inflammation (Bjarnason et al., 1996; Brenchley and Douek, 2008), we also used meta-analysis to determine whether there is a relationship with a chronic inflammatory disease of the gut using data from fecal samples collected from individuals with Crohn's disease and healthy controls in Willing et al. (2010). Samples from HIV-infected individuals (including recent and chronic untreated and ART treated individuals) aligned with individuals with Ileal Crohn's disease across a distinct axis of variation from that associated with diet/culture (see Figure S3 and the Supplemental Information).

Immune Responses to Selected Bacteria Modulated by HIV Infection

Microbiota changes with HIV infection can have several underlying causes, including (1) a compromised ability of the innate and/

or adaptive immune system to control commensal bacteria, (2) the indirect selection of inflammation-tolerant versus sensitive bacteria resulting from a chronic inflammatory state, or (3) a loss of interaction with CD4+ T cells that produce regulatory responses that promote tolerance of beneficial microbes. Although the immunologic driving factors of microbiota changes in HIV infection are likely to be complex, we began to explore drivers of compositional differences by examining cytokine and CD4+ T cell proliferative response to bacteria highly related to OTUs that differed with HIV infection status using peripheral blood mononuclear cells (PBMCs) from a separate cohort of untreated chronic HIV-positive and HIV-negative individuals. We tested (1) two species from the genus *Bacteroides*, which had a conserved decrease with HIV infection (*Bacteroides dorei* and *Bacteroides fluxus*); (2) two species highly related to OTUs that increased with HIV infection, *P. copri* (Prevotellaceae) and *E. bifforme* (Erysipelotrichaceae) (Table S2, Figure S2); and (3) *E. coli* serotype O55:K59(B5), which did not change.

We first examined proinflammatory (TNF- α) and anti-inflammatory (IL-10) cytokines in six HIV-negative and six HIV-positive subjects after incubating PBMCs for 24 hr with bacterial lysate. HIV-infected subjects tended to have elevated levels of both TNF- α and IL-10 to all bacteria in comparison with HIV-negative controls (Figure S4). Previous studies have shown that both of these cytokines are elevated in plasma of HIV-positive subjects (Dutertre et al., 2012; Said et al., 2010). However, *E. bifforme* had a significantly higher TNF- α :IL-10 ratio (Figure 4A) in both HIV-negative and HIV-infected subjects, indicating that it is more proinflammatory than the other bacteria tested.

Next, we examined CD4+ T cell proliferative responses to these bacteria in 10 subjects with HIV infection and 11 HIV-negative controls. Because responses to gut bacteria in peripheral blood are often below the limit of detection directly ex vivo (Howe et al., 2009), we labeled PBMC with Cell Trace to track cell division and cultured the PBMC with the bacteria lysate for 6 days. This method amplifies bacteria-specific CD4+ T cells to easily detected frequencies. All bacteria, but not LPS, induced CD4+ T cell proliferation in both HIV-infected and HIV-negative subjects. However, responses to *B. fluxus* and *B. dorei* were reduced with HIV infection, while responses to *P. copri*, *E. bifforme*, and *E. coli* were not significantly different (Figure 4B).

DISCUSSION

Because HIV targets central players of innate and adaptive immunity including CD4+ T cells, monocytes, and macrophages (McMichael et al., 2010; Mogensen et al., 2010), changes in gut microbiota with infection support the notion that the immune system plays an important role in shaping composition. That immune dysfunction can cause gut microbiota changes with concomitant health effects has been supported in mice lacking specific components of the immune system, including TLR5, IL-22, or the inflammasome. Compositional shifts in gut microbiota in these models were associated with development of colitis (Carvalho et al., 2012; Elinav et al., 2011; Zenewicz et al., 2013) and metabolic syndrome (Vijay-Kumar et al., 2010), diseases that are increased in prevalence with HIV (Knox et al., 2000; Wanke, 1999).




Figure 2. Taxonomy Bar Charts Showing Dominance of Prevotellaceae in HIV-Infected Subjects

Families significantly different between ART-naïve individuals chronically infected with HIV and HIV-negative controls are marked with a star ($p < 0.05$ ANOVA with FDR correction; only the first time point was used for individuals who had multiple time points). Those enriched in HIV negative have a blue star, and those enriched in HIV (Chronic) are in green. Samples from the same individual at two time points are adjacent and joined with a bridge. See Tables S2A, S2B, and S2C for lists of discriminatory genera, families, and 97% OTUs, respectively, and Figure S2 for a phylogenetic tree relating discriminatory 97% OTUs.

Strong and characteristic change with HIV infection despite matching the control cohort for age, sex, race, and geographic location is surprising, since studies of the macaque microbiota in SIV infection failed to detect significant changes independent of colitis, and yet macaques experience a similar depletion of CD4+ T cells and inflammation in the gut compared to HIV-infected humans (McKenna et al., 2008; Veazey et al., 1998). This difference may be related in part to SIV infection duration (e.g., macaques in one longitudinal study were only followed 8 weeks post infection (McKenna et al., 2008)). Microbiota changes in two of three recently infected HIV-positive humans in our study were subtle compared to changes with chronic infection, indicating that microbiota changes begin during early infection and continue over time. In fact, a recent study, which characterized gut microbial communities of six wild-living chimpanzees from Gombe National Park in Tanzania who were naturally infected with SIV over the course of a 13 year survey, showed that SIV infection was associated with increased temporal variability of microbiota composition within individuals (Moeller et al., 2013). However, the compositional changes observed in chimpanzees with long-term SIV infection were not as consistent across individuals as what we observe here in HIV-infected humans.

The similarity between gut microbiota of HIV-infected individuals in the U.S. and HIV-negative individuals in agrarian cultures suggests that the relatively strong phenotype with HIV infection in humans may also be due to specific elements of our cohort's lifestyle, since Western populations appear to have a stronger deviation of the pre- versus the post-HIV-infection microbiota. Indeed, similar to healthy humans in agrarian cultures, the

healthy gut microbiota of macaques is already Prevotella rich and Bacteroides poor (McKenna et al., 2008). It will be of interest to see if HIV-associated changes in agrarian cultures are less pronounced than in the U.S. or if they are more pronounced in macaques consuming a Western-type diet.

HIV-associated enteropathy is initiated during acute infection, continues through advanced disease, and involves increased intestinal inflammation (Bjarnason et al., 1996; Brenchley and Douek, 2008). We thus had expected an expansion of bacteria that increase with other chronic intestinal inflammatory diseases and a decrease in diversity, which has been observed in several unhealthy or inflammatory states (Chang et al., 2008; Willing et al., 2010). Although a previous study that used 16S rRNA gene quantification of order-level bacterial taxa showed a trend toward a greater proportion with HIV infection of Enterobacteriales (Ellis et al., 2011), a taxon often associated with inflammatory conditions (Koren et al., 2012; Mukhopadhyaya et al., 2012), we did not detect significant changes in Enterobacteriales at either coarse or fine taxonomic levels. Furthermore, microbiota in untreated HIV infection, but not with ART, had increased diversity compared to the control population. This is consistent with the resemblance to the microbiota in agrarian cultures, which also has increased diversity compared to that of healthy U.S. adults (Yatsunenko et al., 2012), but not with other unhealthy states such as inflammatory bowel disease (Willing et al., 2010) or recurrent *Clostridium difficile* infection (Chang et al., 2008). Since we excluded individuals with GI symptoms that warranted antibiotic treatment during the preceding 30 days, we cannot exclude the possibility that Enterobacteriales are enriched or




Figure 3. The Fecal Microbiota of HIV-Positive Individuals in the U.S. Resemble those of HIV-Negative Individuals in Agrarian Cultures at the Community Level

Unweighted UniFrac PCoA plot comparing data from this study with a global survey of fecal microbial community composition conducted in individuals from Malawi, the Amazonas State of Venezuela, and the U.S. (Yatsunenکو et al., 2012) (global gut) and (2) a study linking long-term dietary patterns with gut microbial enterotypes (Wu et al., 2011) (long-term diet), using a reference mapping protocol.

(A) Points colored by study.

(B) Same plot, but with points colored across a red-blue spectrum from low to high relative abundance of the genus *Bacteroides*.

(C) Same as for (B), but for the *Prevotella* genus.

(D) Points colored by country of residence and HIV status.

See Figure S3 for an additional meta-analysis exploring the relationship of HIV infection with Crohn's disease.

diversity decreased in HIV-infected individuals with acute diarrhea or advanced AIDS.

However, our analysis of the pro-/anti-inflammatory properties of cultured isolates related to OTUs that increased with chronic HIV infection indicated that *E. bifforme*, but not *P. copri*, was significantly more proinflammatory than bacteria in the HIV-negative-associated *Bacteroides* genus. *E. bifforme* is enriched in individuals with irritable bowel syndrome (Ponnusamy et al., 2011), and it increased in the active disease stage compared to remission in an individual with ulcerative colitis (Harrington et al., 2008). *D. piger* which increased with HIV infection is significantly higher in IBD patients compared to healthy controls (Jia et al., 2012; Loubinoux et al., 2002). Enrichment of these taxa with HIV infection is thus consistent with the relationship we

observed between HIV and Ileal Crohn's disease in both ART-treated and ART-untreated individuals at the community level.

Despite potential links between certain bacterial lineages and proinflammatory states in HIV infection, overall the predominant pattern of microbiota change with HIV infection was surprisingly along a *Prevotella*-*Bacteroides* gradient. *Prevotella*- versus *Bacteroides*-rich communities correlate with overall diversity differences observed both across cultures and within Western populations, although there is some controversy over whether they represent discrete "enterotypes," as initially proposed (Arumugam et al., 2011), or whether this variation occurs across a gradient (Koren et al., 2013; Wu et al., 2011; Yatsunenکو et al., 2012). Diet has been implicated as a primary driving factor of this gradient, since *Prevotella* were enriched in healthy U.S. adults




Figure 4. Characterization of Immune Responses to Selected Bacteria in Untreated HIV-Positive and HIV-Negative Subjects

(A) TNF- α to IL-10 ratio (as measured using a bead array assay) after a 24 hr stimulation. An asterisk indicates that *E. bifforme* had a significantly higher TNF- α /IL-10 ratio compared to the other bacterial isolates with the Kruskal-Wallis ANOVA and Dunn test. Red signifies HIV-positive and blue HIV-negative subjects.

(B) Percentage of divided (cell trace low) CD4+ T cells enumerated by flow cytometry after a 6 day stimulation. Statistical significance was determined using Mann-Whitney t tests. Bacteria that decreased in relative abundance with chronic, untreated HIV infection are shown in red, those that increased are in blue, and no difference is shown in black. Points in the scatterplots depict each individual, and the bar is the median. See also Figure S4.

who consumed diets poor in animal protein and saturated fats and rich in carbohydrates and simple sugars (Wu et al., 2011), and in healthy individuals in agrarian cultures with diets dominated by maize-, cassava-, and other plant-derived polysaccharides (De Filippo et al., 2010; Yatsunenko et al., 2012). Although we do not have dietary information for our HIV-positive cohort and thus do not know whether they uniformly consume a diet that is particularly carbohydrate rich or animal fat/protein poor, that both HIV infection duration and ART treatment had an effect on the degree/consistency of *Prevotella* richness with HIV infection further supports immunologic and not strictly dietary driving factors. This raises the intriguing possibility that the predominant type of gut microbiota diversity variation that occurs across cultures and within Western populations is driven by an interplay between diet and the immune system.

CD4+ T cells are profoundly depleted in GALT with HIV infection (Brenchley and Douek, 2008). Interactions between CD4+ T cells and commensal gut bacteria and concomitant changes in microbiota composition are not well understood, although recent work has demonstrated that immunoglobulin A (IgA) is important for maintaining balance between gut bacteria and host (Kawamoto et al., 2012). B cell immunity and subsequent IgA production are dependent on CD4+ T follicular helper cells which are impaired in HIV infection (Cubas et al., 2013). Recently it was shown that innate lymphoid cells regulate CD4+ T cell responses to intestinal bacteria (Hepworth et al., 2013). CD4+ T cell-microbe interactions are perhaps best understood for *Bacteroides fragilis*, which is in a genus that consistently and dramatically decreases with HIV infection. This species, and

specifically a zwitterionic polysaccharide on its capsule called polysaccharide A (PSA), restores CD4+ T cell development in germ-free mice by eliciting expansion of antigen-experienced ("educated") CD4+CD45Rb^{low} T cells (Mazmanian et al., 2005). *B. fragilis* PSA can also mediate conversion of CD4+ T cells into anti-inflammatory Foxp3+ Treg cells (Round and Mazmanian, 2010) and can provide protection from inflammatory disease (Mazmanian et al., 2005, 2008). When administered as a probiotic, another *Bacteroides* species, *B. uniformis*, can correct T cell proliferation deficiencies displayed in "Western diet"-fed mice (Gauffin Cano et al., 2012) and induces higher secretion of IL-10, a negative regulator of inflammatory response.

Given these interactions between *Bacteroides* species and CD4+ T cells, it is of interest that *Bacteroides* decrease in HIV-positive individuals when CD4+ T cell populations are compromised, suggesting that CD4+ T cell interactions may be essential for persistence of *Bacteroides* in the gut. Even within the overall compromised T cell populations of HIV-positive subjects, we show that proliferative responses to *Bacteroides* species are preferentially depleted compared to other species tested. One potential explanation is that CD4+ T cell responses are diminished because of reduced levels of *Bacteroides* antigen. However, an increased proliferative response to *P. copri* and *E. bifforme* was not observed despite their increase in relative abundance with HIV infection, indicating that antigen concentration is not the only explaining factor. Another potential explanation, given the known interaction of *Bacteroides* with CD4+ T cells via PSA, is that *Bacteroides*-specific T cells are preferentially activated and thus killed, which is consistent with

enhanced HIV infection of intestinal lamina propria CD4⁺ T cells generally upon exposure to commensal bacteria (Dillon et al., 2012). Conducting immunologic assays on samples for which we have matched gut microbiota compositional data will help to determine the degree to which T cell proliferative response changes may be driven by changes in antigen presence or concentration.

Mouse experiments have supported the notion that a Bacteroides-poor microbiota in the context of a Western diet may have negative health effects (Gauffin Cano et al., 2012). Oral administration of *B. uniformis* to mice consuming a high-fat diet resulted in decreased inflammation, body weight gain, liver steatosis, liver cholesterol, and triglyceride concentrations and increased small adipocyte numbers. In this light, it is interesting that lipodystrophy, a disorder characterized by some of these same features, has been increasingly observed in HIV-positive individuals (Carr et al., 1998; Wanke, 1999). Although lipodystrophy and dyslipidemia are linked to ART, particularly the protease inhibitors (Carr et al., 1998), it also can occur in the absence of ART, leading some to suggest that it is in part due to HIV infection itself (Madge et al., 1999; Nicholaou et al., 2013).

Prevotella-rich microbiota have been linked with high plasma levels of trimethylamine-*N*-oxide (TMAO), a proatherogenic cardiovascular risk-associated compound that can be produced from dietary L-carnitine in red meat by intestinal microbiota (Koeth et al., 2013). Given high rates and early onset of atherosclerosis and coronary heart disease in HIV-infected individuals (Boccaro et al., 2013; Hsue et al., 2009), it will be of interest to see if a Prevotella-rich microbiota in HIV-infected individuals is associated with high TMAO levels in plasma.

The failure of ART to consistently restore the gut microbiota to a state resembling HIV-negative individuals is consistent with persistence of gastrointestinal diseases with ART. HIV-positive subjects on successful ART remain at greater risk of multiple inflammatory diseases, including atherosclerosis (Hsue et al., 2009). Diseases of nutrition ranging from wasting to lipodystrophy persist, or for lipodystrophy can even be exacerbated by ART (Carr et al., 1998; Wanke, 1999; Wanke et al., 2000). Although ART can improve chronic diarrhea in patients with advanced HIV infection and lead to disappearance of parasites such as *Cryptosporidium* sp. from stool (Foudraine et al., 1998), other parasites can persist (Boyles et al., 2012). A persistence of an HIV-associated microbiota in some individuals on long-term ART is also consistent with the observation that ART treatment generally does not completely restore CD4⁺ T cells in GALT (Estes et al., 2008). Discordance between CD4⁺ T cell counts in GALT and the periphery may explain why we did not observe a correlation between peripheral CD4⁺ T cell count and the degree of divergence in fecal microbiota composition from HIV-negative control samples.

It will be important to further study the effects of HIV infection on the gut microbiota in the context of different disease states, cultures, and diets so that the complex interactions between lifestyle, diets, and the immune system in shaping microbiota composition can be determined. Further studies of the immunomodulatory properties and inflammation tolerance of bacterial isolates related to OTUs that change during HIV infection will provide insight into driving factors of gut microbiota changes and their potential impact on health.

EXPERIMENTAL PROCEDURES

Subject Recruitment

Four cohorts were prospectively recruited from the University of Colorado Hospital Infectious Disease Group Practice: (1) recent HIV-1 infection, individuals likely infected within the prior 6 months; (2) chronic HIV-1 infection untreated, individuals infected for >6 months and ART drug naive or off treatment for >6 months; (3) chronic HIV-1 infection on long-term ART, ART treatment for ≥ 12 months with a minimum of three ART drugs prior to study entry and viral suppressed for >6 months; and (4) healthy controls, HIV-negative individuals matched to HIV-infected participants for sex, age, geographic location, and smoking status (see the [Supplemental Experimental Procedures](#) for further cohort characteristics). Individuals from cohort 2 sometimes initiated ART treatment immediately following the initial visit and donated an additional fecal sample 6.7–8.5 months later. Informed consent was obtained from each subject, and the study protocol was approved by the Colorado Multiple Institution Review Board.

DNA Extraction and Sequencing

Stool samples were collected on a sterile swab by the patient either during or 12 hr prior to the visit and stored at -80°C . DNA was extracted using the standard Power Soil Kit protocol (MoBio). Extracted DNA was PCR amplified with barcoded primers targeting the V4 region of 16S rRNA as detailed in Yatsunenkeno et al. (2012). Control water samples that had undergone the same DNA extraction and PCR amplification procedures were also sequenced. Each PCR product was quantified using PicoGreen (Invitrogen), and equal amounts (ng) of DNA from each sample were pooled and cleaned using the UltraClean PCR Clean-Up Kit (MoBio). Sequences were generated on two runs on a MiSeq personal sequencer (Illumina, San Diego, CA).

Sequence Data Analysis

Raw sequences were quality filtered and assigned to samples based on their barcodes using the default parameters of QIIME version 1.5.0 (Caporaso et al., 2010) (see the [Supplemental Experimental Procedures](#)). Sequences were assigned to 97% ID OTUs by comparing them to a nonredundant reference database of near-full length sequences (February 4, 2011, Greengenes database) (DeSantis et al., 2006), and unassigned sequences were clustered into de novo OTUs using UCLUSTref (Edgar, 2010). Since samples contained between 4,694 and 72,828 sequences, analyses were standardized at 4,600 sequences per sample to avoid biases.

UniFrac PCoA analyses were conducted using QIIME as detailed in the [Supplemental Information](#). Random Forests classification on the HIV data was carried out using SourceTracker software (<http://sourceforge.net/projects/sourcetracker/>) (Knights et al., 2011b) and custom code as detailed in the [Supplemental Information](#). Alpha diversity was calculated using three different measures (1) observed species, the number of 97% ID OTUs observed in 4,600 sequences; (2) Shannon Index (Shannon, 1948); and (3) PD (Faith, 1992). Only one sample was used for individuals sampled at two time points.

Bacterial families and genera in each sample were determined using the RDP classifier retrained on the greengenes taxonomy (McDonald et al., 2012) and statistically compared as detailed in the [Supplemental Information](#). Discriminating OTUs were related to each other and to bacterial type strains using the ARB software package (Ludwig et al., 2004) as detailed in the [Supplemental Information](#).

To compare sequences from our samples to other studies, we used the QIIME database (<http://microbio.me/qiime>), which assigns sequences to 97% ID OTUs by comparing them to a reference database of near-full-length sequences (February 4, 2011, Greengenes database) (DeSantis et al., 2006) using UCLUSTref (Edgar, 2010) as described in Lozupone et al. (2013). Unassigned sequences were dropped from the analysis. We removed samples from individuals not between the ages of 10 and 66 and standardized at 4,600 sequences per sample.

Immunologic Assays

Bacterial lysates for *B. dorei* (DSM 17855), *B. fluxus* (DSM 22534), *P. copri* (DSM18205), *E. bifforme* (DSM 3989), and *E. coli* serotype O55:K59(B5) were prepared and quantified as detailed in the [Supplemental Information](#). PBMCs were isolated from blood of ten untreated HIV-positive subjects with chronic

infection (median CD4+ T cells 508 cells/ μ l; range 248–742 and VL 55,495 copies of HIV RNA/ml; range 554–123,000) and 11 HIV-negative subjects as described in the [Supplemental Information](#). For the T cell proliferation assays, PBMCs were labeled with 2 μ M Cell Trace to track cell division (Invitrogen); plated at 1×10^6 cells/ml in a 48-well plate; and cultured with 1 μ g/ml of bacterial lysates, LPS, 2.5 μ g/ml PHA, or medium alone at 37°C in a humidified 5% CO₂ atmosphere. After 6 days, cells were washed with PBS containing 1% BSA and surface stained with anti-CD4 (PE-Cy5-5; BD Biosciences), anti-CD3 (Qdot 605; Invitrogen), anti-CD19 (Alexa Fluor 405; Caltag), CD45RA, and CD27 for 30 min at 4°C, then washed and fixed with 1% formaldehyde. Cells were analyzed using a LSR-II flow cytometer (BD Immunocytometry Systems) as described in the [Supplemental Information](#).

For the cytokine analysis, PBMCs (1×10^6 cells/ml) were cultured in 96-well plates with 1 μ g/ml of bacterial lysates, LPS, or medium alone. Cultures were incubated at 37°C in a humidified 5% CO₂ atmosphere for 24 hr, and culture supernatants were harvested and centrifuged to remove remaining cells and stored at –80°C. IL-10 and TNF- α levels were assayed using a 7-plex Inflammatory Cytokine kit (Meso Scale Development, Gaithersburg MD). Assays were performed per manufacturer's instructions and analyzed using a Sector Imager 2400 (Meso Scale Discovery).

ACCESSION NUMBERS

The sequence data have been deposited in the QIIME database (<http://www.microbio.me/qiime/>) and at EBI (<http://www.ebi.ac.uk/ena/>; accession number ERP003611).

SUPPLEMENTAL INFORMATION

Supplemental Information includes four figures, three tables, Supplemental Experimental Procedures, and Supplemental References and can be found with this article online at <http://dx.doi.org/10.1016/j.chom.2013.08.006>.

ACKNOWLEDGMENTS

This work was supported by a LHMP-sponsored project (UO1HL098996 to A.P.F., T.B.C., S.C.F., and R.K.) and by the Colorado Clinical and Translational Sciences Institute (UL1TR000005). C.A.L. was supported by K01DK090285, and R.K. was supported by the Howard Hughes Medical Institute. We would like to thank Beverly Putnam and Christine Grismer for subject recruitment, Sam MaWhinney and Amanda Allshouse for statistical support, Dan Knights for contributing custom code for Random Forests analysis, and Maria del Carmen Portillo for help with anaerobic culture. We would particularly like to thank the study participants.

Received: May 23, 2013

Revised: July 3, 2013

Accepted: August 16, 2013

Published: September 11, 2013

REFERENCES

- Arumugam, M., Raes, J., Pelletier, E., Le Paslier, D., Yamada, T., Mende, D.R., Fernandes, G.R., Tap, J., Bruls, T., Batto, J.M., et al.; MetaHIT Consortium. (2011). Enterotypes of the human gut microbiome. *Nature* **473**, 174–180.
- Bjarnason, I., Sharpstone, D.R., Francis, N., Marker, A., Taylor, C., Barrett, M., Macpherson, A., Baldwin, C., Menzies, I.S., Crane, R.C., et al. (1996). Intestinal inflammation, ileal structure and function in HIV. *AIDS* **10**, 1385–1391.
- Boccarda, F., Lang, S., Meuleman, C., Ederhy, S., Mary-Krause, M., Costagliola, D., Capeau, J., and Cohen, A. (2013). HIV and coronary heart disease: time for a better understanding. *J. Am. Coll. Cardiol.* **61**, 511–523.
- Boyles, T.H., Black, J., Meintjes, G., and Mendelson, M. (2012). Failure to eradicate *Isospora belli* diarrhoea despite immune reconstitution in adults with HIV—a case series. *PLoS ONE* **7**, e42844. <http://dx.doi.org/10.1371/journal.pone.0042844>.
- Brenchley, J.M., and Douek, D.C. (2008). HIV infection and the gastrointestinal immune system. *Mucosal Immunol.* **1**, 23–30.
- Brenchley, J.M., Schacker, T.W., Ruff, L.E., Price, D.A., Taylor, J.H., Beilman, G.J., Nguyen, P.L., Khoruts, A., Larson, M., Haase, A.T., and Douek, D.C. (2004). CD4+ T cell depletion during all stages of HIV disease occurs predominantly in the gastrointestinal tract. *J. Exp. Med.* **200**, 749–759.
- Brenchley, J.M., Price, D.A., Schacker, T.W., Asher, T.E., Silvestri, G., Rao, S., Kazzaz, Z., Bornstein, E., Lambotte, O., Altmann, D., et al. (2006). Microbial translocation is a cause of systemic immune activation in chronic HIV infection. *Nat. Med.* **12**, 1365–1371.
- Caporaso, J.G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D., Costello, E.K., Fierer, N., Peña, A.G., Goodrich, J.K., Gordon, J.I., et al. (2010). QIIME allows analysis of high-throughput community sequencing data. *Nat. Methods* **7**, 335–336.
- Carr, A., Samaras, K., Burton, S., Law, M., Freund, J., Chisholm, D.J., and Cooper, D.A. (1998). A syndrome of peripheral lipodystrophy, hyperlipidaemia and insulin resistance in patients receiving HIV protease inhibitors. *AIDS* **12**, F51–F58.
- Carvalho, F.A., Koren, O., Goodrich, J.K., Johansson, M.E., Nalbantoglu, I., Aitken, J.D., Su, Y., Chassaing, B., Walters, W.A., González, A., et al. (2012). Transient inability to manage proteobacteria promotes chronic gut inflammation in TLR5-deficient mice. *Cell Host Microbe* **12**, 139–152.
- Chang, J.Y., Antonopoulos, D.A., Kalra, A., Tonelli, A., Khalife, W.T., Schmidt, T.M., and Young, V.B. (2008). Decreased diversity of the fecal Microbiome in recurrent *Clostridium difficile*-associated diarrhea. *J. Infect. Dis.* **197**, 435–438.
- Cubas, R.A., Mudd, J.C., Savoye, A.L., Perreau, M., van Grevenynghe, J., Metcalf, T., Connick, E., Meditz, A., Freeman, G.J., Abesada-Terk, G., Jr., et al. (2013). Inadequate T follicular cell help impairs B cell immunity during HIV infection. *Nat. Med.* **19**, 494–499.
- De Filippo, C., Cavalieri, D., Di Paola, M., Ramazzotti, M., Poullet, J.B., Massart, S., Collini, S., Pieraccini, G., and Lionetti, P. (2010). Impact of diet in shaping gut microbiota revealed by a comparative study in children from Europe and rural Africa. *Proc. Natl. Acad. Sci. USA* **107**, 14691–14696.
- DeSantis, T.Z., Hugenholtz, P., Larsen, N., Rojas, M., Brodie, E.L., Keller, K., Huber, T., Dalevi, D., Hu, P., and Andersen, G.L. (2006). Greengenes, a chimera-checked 16S rRNA gene database and workbench compatible with ARB. *Appl. Environ. Microbiol.* **72**, 5069–5072.
- Dillon, S.M., Manuzak, J.A., Leone, A.K., Lee, E.J., Rogers, L.M., McCarter, M.D., and Wilson, C.C. (2012). HIV-1 infection of human intestinal lamina propria CD4+ T cells in vitro is enhanced by exposure to commensal *Escherichia coli*. *J. Immunol.* **189**, 885–896.
- Dutertre, C.A., Amraoui, S., DeRosa, A., Jourdain, J.P., Vimeux, L., Goguet, M., Degrelle, S., Feuillet, V., Liovat, A.S., Müller-Trutwin, M., et al. (2012). Pivotal role of M-DC8⁺ monocytes from viremic HIV-infected patients in TNF α overproduction in response to microbial products. *Blood* **120**, 2259–2268.
- Edgar, R.C. (2010). Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* **26**, 2460–2461.
- Elinav, E., Strowig, T., Kau, A.L., Henao-Mejia, J., Thaiss, C.A., Booth, C.J., Peaper, D.R., Bertin, J., Eisenbarth, S.C., Gordon, J.I., and Flavell, R.A. (2011). NLRP6 inflammasome regulates colonic microbial ecology and risk for colitis. *Cell* **145**, 745–757.
- Ellis, C.L., Ma, Z.M., Mann, S.K., Li, C.S., Wu, J., Knight, T.H., Yotter, T., Hayes, T.L., Maniar, A.H., Troia-Cancio, P.V., et al. (2011). Molecular characterization of stool microbiota in HIV-infected subjects by panbacterial and order-level 16S ribosomal DNA (rDNA) quantification and correlations with immune activation. *J. Acquir. Immune Defic. Syndr.* **57**, 363–370.
- Estes, J., Baker, J.V., Brenchley, J.M., Khoruts, A., Barthold, J.L., Bantle, A., Reilly, C.S., Beilman, G.J., George, M.E., Douek, D.C., et al. (2008). Collagen deposition limits immune reconstitution in the gut. *J. Infect. Dis.* **198**, 456–464.
- Faith, D.P. (1992). Conservation evaluation and phylogenetic diversity. *Biol. Conserv.* **61**, 1–10.
- Foudraine, N.A., Weverling, G.J., van Gool, T., Roos, M.T., de Wolf, F., Koopmans, P.P., van den Broek, P.J., Meenhorst, P.L., van Leeuwen, R., Lange, J.M., and Reiss, P. (1998). Improvement of chronic diarrhoea in

- patients with advanced HIV-1 infection during potent antiretroviral therapy. *AIDS* 12, 35–41.
- Gauffin Cano, P., Santacruz, A., Moya, A., and Sanz, Y. (2012). *Bacteroides uniformis* CECT 7771 ameliorates metabolic and immunological dysfunction in mice with high-fat-diet induced obesity. *PLoS ONE* 7, e41079, <http://dx.doi.org/10.1371/journal.pone.0041079>.
- Harrington, C.R., Lucchini, S., Ridgway, K.P., Wegmann, U., Eaton, T.J., Hinton, J.C., Gasson, M.J., and Narbad, A. (2008). A short-oligonucleotide microarray that allows improved detection of gastrointestinal tract microbial communities. *BMC Microbiol.* 8, 195, <http://dx.doi.org/10.1186/1471-2180-8-195>.
- Hepworth, M.R., Monticelli, L.A., Fung, T.C., Ziegler, C.G., Grunberg, S., Sinha, R., Mantegazza, A.R., Ma, H.L., Crawford, A., Angelosanto, J.M., et al. (2013). Innate lymphoid cells regulate CD4+ T-cell responses to intestinal commensal bacteria. *Nature* 498, 113–117.
- Howe, R., Dillon, S., Rogers, L., McCarter, M., Kelly, C., Gonzalez, R., Madinger, N., and Wilson, C.C. (2009). Evidence for dendritic cell-dependent CD4(+) T helper-1 type responses to commensal bacteria in normal human intestinal lamina propria. *Clin. Immunol.* 131, 317–332.
- Hsue, P.Y., Hunt, P.W., Schnell, A., Kalapus, S.C., Hoh, R., Ganz, P., Martin, J.N., and Deeks, S.G. (2009). Role of viral replication, antiretroviral therapy, and immunodeficiency in HIV-associated atherosclerosis. *AIDS* 23, 1059–1067.
- Jia, W., Whitehead, R.N., Griffiths, L., Dawson, C., Bai, H., Waring, R.H., Ramsden, D.B., Hunter, J.O., Cauchi, M., Bessant, C., et al. (2012). Diversity and distribution of sulphate-reducing bacteria in human faeces from healthy subjects and patients with inflammatory bowel disease. *FEMS Immunol. Med. Microbiol.* 65, 55–68.
- Kawamoto, S., Tran, T.H., Maruya, M., Suzuki, K., Doi, Y., Tsutsui, Y., Kato, L.M., and Fagarasan, S. (2012). The inhibitory receptor PD-1 regulates IgA selection and bacterial composition in the gut. *Science* 336, 485–489.
- Knights, D., Costello, E.K., and Knight, R. (2011a). Supervised classification of human microbiota. *FEMS Microbiol. Rev.* 35, 343–359.
- Knights, D., Kuczynski, J., Koren, O., Ley, R.E., Field, D., Knight, R., DeSantis, T.Z., and Kelley, S.T. (2011b). Supervised classification of microbiota mitigates mislabeling errors. *ISME J.* 5, 570–573.
- Knox, T.A., Spiegelman, D., Skinner, S.C., and Gorbach, S. (2000). Diarrhea and abnormalities of gastrointestinal function in a cohort of men and women with HIV infection. *Am. J. Gastroenterol.* 95, 3482–3489.
- Koeth, R.A., Wang, Z., Levison, B.S., Buffa, J.A., Org, E., Sheehy, B.T., Britt, E.B., Fu, X., Wu, Y., Li, L., et al. (2013). Intestinal microbiota metabolism of L-carnitine, a nutrient in red meat, promotes atherosclerosis. *Nat. Med.* 19, 576–585.
- Koren, O., Goodrich, J.K., Cullender, T.C., Spor, A., Laitinen, K., Bäckhed, H.K., Gonzalez, A., Werner, J.J., Angenent, L.T., Knight, R., et al. (2012). Host remodeling of the gut microbiome and metabolic changes during pregnancy. *Cell* 150, 470–480.
- Koren, O., Knights, D., Gonzalez, A., Waldron, L., Segata, N., Knight, R., Huttenhower, C., and Ley, R.E. (2013). A guide to enterotypes across the human body: meta-analysis of microbial community structures in human microbiome datasets. *PLoS Comput. Biol.* 9, e1002863, <http://dx.doi.org/10.1371/journal.pcbi.1002863>.
- Loubinoux, J., Bronowicki, J.P., Pereira, I.A., Mouguel, J.L., and Faou, A.E. (2002). Sulfate-reducing bacteria in human feces and their association with inflammatory bowel diseases. *FEMS Microbiol. Ecol.* 40, 107–112.
- Lozupone, C., and Knight, R. (2005). UniFrac: a new phylogenetic method for comparing microbial communities. *Appl. Environ. Microbiol.* 71, 8228–8235.
- Lozupone, C.A., Hamady, M., Kelley, S.T., and Knight, R. (2007). Quantitative and qualitative beta diversity measures lead to different insights into factors that structure microbial communities. *Appl. Environ. Microbiol.* 73, 1576–1585.
- Lozupone, C.A., Stombaugh, J.I., Gordon, J.I., Jansson, J.K., and Knight, R. (2012). Diversity, stability and resilience of the human gut microbiota. *Nature* 489, 220–230.
- Lozupone, C.A., Stombaugh, J., Gonzalez, A., Ackermann, G., Wendel, D., Vazquez-Baeza, Y., Jansson, J.K., Gordon, J.I., and Knight, R. (2013). Meta-analyses of studies of the human microbiota. *Genome Res.* Published online July 16, 2013. <http://dx.doi.org/10.1101/gr.151803.112>.
- Ludwig, W., Strunk, O., Westram, R., Richter, L., Meier, H., Yadukumar, Buchner, A., Lai, T., Steppi, S., Jobb, G., et al. (2004). ARB: a software environment for sequence data. *Nucleic Acids Res.* 32, 1363–1371.
- Madge, S., Kinloch-de-Loes, S., Mercey, D., Johnson, M.A., and Weller, I.V.D. (1999). Lipodystrophy in patients naive to HIV protease inhibitors. *AIDS* 13, 735–737.
- Mazmanian, S.K., Liu, C.H., Tzianabos, A.O., and Kasper, D.L. (2005). An immunomodulatory molecule of symbiotic bacteria directs maturation of the host immune system. *Cell* 122, 107–118.
- Mazmanian, S.K., Round, J.L., and Kasper, D.L. (2008). A microbial symbiosis factor prevents intestinal inflammatory disease. *Nature* 453, 620–625.
- McDonald, D., Price, M.N., Goodrich, J., Nawrocki, E.P., DeSantis, T.Z., Probst, A., Andersen, G.L., Knight, R., and Hugenholz, P. (2012). An improved Greengenes taxonomy with explicit ranks for ecological and evolutionary analyses of bacteria and archaea. *ISME J.* 6, 610–618.
- McKenna, P., Hoffmann, C., Minkah, N., Aye, P.P., Lackner, A., Liu, Z., Lozupone, C.A., Hamady, M., Knight, R., and Bushman, F.D. (2008). The macaque gut microbiome in health, lentiviral infection, and chronic enterocolitis. *PLoS Pathog.* 4, e20, <http://dx.doi.org/10.1371/journal.ppat.0040020>.
- McMichael, A.J., Borrow, P., Tomaras, G.D., Goonetilleke, N., and Haynes, B.F. (2010). The immune response during acute HIV-1 infection: clues for vaccine development. *Nat. Rev. Immunol.* 10, 11–23.
- Moeller, A.H., Shiits, M., Li, Y., Rudicell, R.S., Lonsdorf, E.V., Pusey, A.E., Wilson, M.L., Hahn, B.H., and Ochman, H. (2013). SIV-induced instability of the chimpanzee gut microbiome. *Cell Host Microbe* 14, this issue, 340–345.
- Mogensen, T.H., Melchjorsen, J., Larsen, C.S., and Paludan, S.R. (2010). Innate immune recognition and activation during HIV infection. *Retrovirology* 7, 54, <http://dx.doi.org/10.1186/1742-4690-7-54>.
- Mukhopadhyay, I., Hansen, R., El-Omar, E.M., and Hold, G.L. (2012). IBD—what role do Proteobacteria play? *Nat. Rev. Gastroenterol. Hepatol.* 9, 219–230.
- Nicholaou, M.J., Martinson, J.J., Abraham, A.G., Brown, T.T., Hussain, S.K., Wolinsky, S.M., and Kingsley, L.A. (2013). HAART-associated dyslipidemia varies by biogeographical ancestry in the multicenter AIDS cohort study. *AIDS Res. Hum. Retroviruses* 29, 871–879.
- Ponnusamy, K., Choi, J.N., Kim, J., Lee, S.Y., and Lee, C.H. (2011). Microbial community and metabolomic comparison of irritable bowel syndrome faeces. *J. Med. Microbiol.* 60, 817–827.
- Round, J.L., and Mazmanian, S.K. (2010). Inducible Foxp3+ regulatory T-cell development by a commensal bacterium of the intestinal microbiota. *Proc. Natl. Acad. Sci. USA* 107, 12204–12209.
- Said, E.A., Dupuy, F.P., Trautmann, L., Zhang, Y., Shi, Y., El-Far, M., Hill, B.J., Noto, A., Ancuta, P., Peretz, Y., et al. (2010). Programmed death-1-induced interleukin-10 production by monocytes impairs CD4+ T cell activation during HIV infection. *Nat. Med.* 16, 452–459.
- Shannon, C.E. (1948). A mathematical theory of communication. *Bell Syst. Tech. J.* 27, 379–423.
- Smith, M.I., Yatsunenkov, T., Manary, M.J., Trehan, I., Mkakosya, R., Cheng, J., Kau, A.L., Rich, S.S., Concannon, P., Mychaleckyj, J.C., et al. (2013). Gut microbiomes of Malawian twin pairs discordant for kwashiorkor. *Science* 339, 548–554.
- Stackebrandt, E., and Goebel, B.M. (1994). Taxonomic note: a place for DNA-DNA reassociation and 16S rRNA sequence analysis in the present species definition in bacteriology. *Int. Syst. Bacteriol.* 44, 846–849.
- Veazey, R.S., DeMaria, M., Chalifoux, L.V., Shvetz, D.E., Pauley, D.R., Knight, H.L., Rosenzweig, M., Johnson, R.P., Desrosiers, R.C., and Lackner, A.A. (1998). Gastrointestinal tract as a major site of CD4+ T cell depletion and viral replication in SIV infection. *Science* 280, 427–431.
- Vijay-Kumar, M., Aitken, J.D., Carvalho, F.A., Cullender, T.C., Mwangi, S., Srinivasan, S., Sitaraman, S.V., Knight, R., Ley, R.E., and Gewirtz, A.T.

- (2010). Metabolic syndrome and altered gut microbiota in mice lacking Toll-like receptor 5. *Science* 328, 228–231.
- Wanke, C.A. (1999). Epidemiological and clinical aspects of the metabolic complications of HIV infection the fat redistribution syndrome. *AIDS* 13, 1287–1293.
- Wanke, C.A., Silva, M., Knox, T.A., Forrester, J., Speigelman, D., and Gorbach, S.L. (2000). Weight loss and wasting remain common complications in individuals infected with human immunodeficiency virus in the era of highly active antiretroviral therapy. *Clin. Infect. Dis.* 31, 803–805.
- Willing, B.P., Dicksved, J., Halfvarson, J., Andersson, A.F., Lucio, M., Zheng, Z., Järnerot, G., Tysk, C., Jansson, J.K., and Engstrand, L. (2010). A pyrosequencing study in twins shows that gastrointestinal microbial profiles vary with inflammatory bowel disease phenotypes. *Gastroenterology* 139, 1844–1854.
- Wu, G.D., Chen, J., Hoffmann, C., Bittinger, K., Chen, Y.Y., Keilbaugh, S.A., Bewtra, M., Knights, D., Walters, W.A., Knight, R., et al. (2011). Linking long-term dietary patterns with gut microbial enterotypes. *Science* 334, 105–108.
- Yatsunencko, T., Rey, F.E., Manary, M.J., Trehan, I., Dominguez-Bello, M.G., Contreras, M., Magris, M., Hidalgo, G., Baldassano, R.N., Anokhin, A.P., et al. (2012). Human gut microbiome viewed across age and geography. *Nature* 486, 222–227.
- Zenewicz, L.A., Yin, X., Wang, G., Elinav, E., Hao, L., Zhao, L., and Flavell, R.A. (2013). IL-22 deficiency alters colonic microbiota to be transmissible and colitogenic. *J. Immunol.* 190, 5306–5312.