

Contents lists available at [ScienceDirect](http://www.sciencedirect.com)

# Bioorganic & Medicinal Chemistry Letters

journal homepage: [www.elsevier.com/locate/bmcl](http://www.elsevier.com/locate/bmcl)

## Development of photo-controllable hydrogen sulfide donor applicable in live cells



Naoki Fukushima<sup>a</sup>, Naoya Ieda<sup>a</sup>, Mitsuyasu Kawaguchi<sup>a</sup>, Kiyoshi Sasakura<sup>b</sup>, Tetsuo Nagano<sup>b</sup>, Kenjiro Hanaoka<sup>b</sup>, Naoki Miyata<sup>a</sup>, Hidehiko Nakagawa<sup>a,\*</sup>

<sup>a</sup> Graduate School of Pharmaceutical Sciences, Nagoya City University, 3-1, Tanabe-dori, Mizuho-ku, Nagoya-shi, Aichi 467-8603, Japan

<sup>b</sup> Graduate School of Pharmaceutical Sciences, The University of Tokyo, 7-3-1, Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

### ARTICLE INFO

#### Article history:

Received 7 November 2014

Revised 27 November 2014

Accepted 29 November 2014

Available online 8 December 2014

#### Keywords:

Hydrogen sulfide

Caged H<sub>2</sub>S

Photolabile protecting group

Xanthone photocage

### ABSTRACT


Hydrogen sulfide (H<sub>2</sub>S) has multiple physiological roles, for example, in vasodilation and inflammation. It is a highly reactive gas under ambient conditions, so controllable H<sub>2</sub>S donors are required for studying its biological functions. Here, we describe the design, synthesis and application of a H<sub>2</sub>S donor (SPD-2) that utilizes xanthone photochemistry to control H<sub>2</sub>S release. H<sub>2</sub>S generation from SPD-2 was completely dependent on UVA-irradiation (325–385 nm), as confirmed by methylene blue assay and by the use of a H<sub>2</sub>S-selective fluorescent probe. SPD-2 was confirmed to provide controlled H<sub>2</sub>S delivery in live cells, and should be suitable for various biological applications.

© 2014 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/3.0/>).

Hydrogen sulfide (H<sub>2</sub>S), like nitric oxide (NO) and carbon monoxide (CO), is a gaseous physiological signal transmitter<sup>1</sup> that has roles in multiple processes, including vascular smooth muscle relaxation,<sup>2</sup> neurotransmission<sup>3</sup> and regulation of inflammation.<sup>4</sup> It is biologically synthesized from L-cysteine and/or L-homocysteine by cystathionine-β-synthase (CBS),<sup>5</sup> cystathionine-γ-lyase (CSE)<sup>6</sup> and 3-mercaptopyruvate sulfur transferase (3MST) coupled with cysteine aminotransferase (CAT).<sup>7</sup> Recently, a novel H<sub>2</sub>S biosynthetic pathway from D-cysteine, involving 3-MST and D-amino acid oxidase (DAO), has also been reported.<sup>8</sup> These enzymes have been demonstrated to influence a wide range of physiological and pathological processes.<sup>9,10</sup>


In biological research on H<sub>2</sub>S, inorganic sulfide salts, such as Na<sub>2</sub>S and NaSH have been widely used as H<sub>2</sub>S sources. However, these sources have substantial disadvantages for examination of the physiological functions of H<sub>2</sub>S. For example, H<sub>2</sub>S generation is very fast, and precise control of the release rate and dosage is not feasible. These compounds cannot mimic biological H<sub>2</sub>S release, which is thought to be relatively slow and continuous. Therefore, controllable H<sub>2</sub>S donors are required for detailed investigation of the physiological functions of H<sub>2</sub>S and for potential therapeutic applications. We focused on photocontrollable H<sub>2</sub>S donors, since photolysis-induced H<sub>2</sub>S release has the potential to allow precise control of the location, timing and dosage of release.

Some photolysis-inducing H<sub>2</sub>S donors, based on geminal-dithiol structure, have already been reported.<sup>11</sup> However, it is difficult to control the rate of H<sub>2</sub>S release from these donors because the formation of H<sub>2</sub>S depends on hydrolysis of geminal dithiol in aqueous media. Recently, we have reported a H<sub>2</sub>S derivative doubly protected with a photolabile protecting group (PPG) as a photo-activatable H<sub>2</sub>S donor,<sup>12</sup> based on ketoprofenate PPG<sup>13</sup> which has good light-responsiveness (SPD-1, Scheme 1a). Ketoprofenate




Scheme 1. Structures and photoreactions of SPD-1 and SPD-2.

\* Corresponding author.




**Scheme 2.** Synthesis of SPD-2. Reagents and conditions: (a) 2-iodobenzoic acid,  $\text{Cs}_2\text{CO}_3$ , CuCl, TDA-1, dioxane; (b)  $\text{H}_2\text{SO}_4$ , 85 °C, 2 steps 52%; (c)  $\text{H}_2\text{SO}_4$ , MeOH, 96%; (d) paraformaldehyde,  $\text{K}_2\text{CO}_3$ , DMSO, 56%; (e) MsCl,  $\text{NEt}_3$ ,  $\text{CH}_2\text{Cl}_2$ , quant.; (f) LiBr, MeCN, 36%; (g) sodium sulfide nonahydrate, toluene,  $\text{H}_2\text{O}$ ,  $\text{C}_{16}\text{H}_{33}(\text{C}_4\text{H}_9)_3\text{P}^+\text{Br}^-$ , 46%; (h) LiOH, THF,  $\text{H}_2\text{O}$ , MeOH, 84%. TDA-1 = tris[2-(2-methoxyethoxy)ethyl]amine, DMSO = dimethylsulfoxide, MsCl = methanesulfonyl chloride, THF = tetrahydrofuran.




**Figure 1.** Release of  $\text{H}_2\text{S}$  from SPD-2 (100  $\mu\text{M}$ ) in PBS, pH 7.4, with 1% DMSO as detected by the methylene blue method. Formation of methylene blue after irradiation of SPD-2 for various times was determined from the absorbance at 671 nm. Photoirradiation was performed at 325–385 nm and the light intensity was set at 2.0  $\text{mW}/\text{cm}^2$ . The data represent the average of three independent experiments with standard deviations.

PPG has many advantageous features, such as good water solubility, formation of benign photo-products and a higher release rate than *o*-nitrobenzyl PPG,<sup>14</sup> the most widely used PPG. However, SPD-1 shows maximum absorbance at 260 nm (Fig. S1b), and consequently the efficiency of  $\text{H}_2\text{S}$  release from SPD-1 is insufficient for application in living cells when UV-A (330–380 nm), widely equipped in some microscopies, is utilized as an uncaging light source. Therefore, we set out to design a new photo-induced  $\text{H}_2\text{S}$  donor suitable for cellular application. Recently, xanthone PPGs have been reported as next-generation carbanion-mediated PPGs<sup>15</sup> with excellent absorption in the UVA region (>320 nm). We designed a novel photo-controllable  $\text{H}_2\text{S}$  donor, SPD-2, by utilizing xanthone PPG to protect  $\text{H}_2\text{S}$  directly (Scheme 1b). SPD-2 showed superior photochemical properties to the ketopropionate-type  $\text{H}_2\text{S}$  donor, and we confirmed that it is available for photo-controlled, site-specific  $\text{H}_2\text{S}$  release in live cells.



**Figure 2.** Fluorescence measurement for detection of  $\text{H}_2\text{S}$  generation using HSip-1. The concentration of SPD-2 was 100  $\mu\text{M}$  and that of HSip-1 was 1  $\mu\text{M}$  in phosphate buffer, pH 7.4, containing 1% DMSO. Fluorescence emission was measured at 516 nm with excitation at 491 nm. Measurements were made after incubation of HSip-1: (a) without SPD-2 or irradiation; (b) with SPD-2 but without irradiation; (c) without SPD-2 after irradiation for 300 s; (d) with SPD-2 after irradiation for 300 seconds. Photoirradiation was performed at 325–385 nm and the light intensity was set at 2.0  $\text{mW}/\text{cm}^2$ . The data represent the average of three independent experiments with standard deviations.

SPD-2 was synthesized as shown in Scheme 2. Preparation of the synthetic intermediate was performed as described previously.<sup>15</sup> Xanthone derivative **5** was prepared by Ullmann condensation of *o*-iodobenzoic acid and 2-(4-hydroxyphenyl)propanoic acid (**3**), followed by acid-catalyzed Friedel–Crafts type reaction. After esterification of **5**, hydroxymethylation by treatment with paraformaldehyde and  $\text{K}_2\text{CO}_3$  gave the desired alcohol **7**. This was mesylated to obtain **8**, which was converted to the brominated derivative **9**. Then, sulfide **10** was synthesized by using sodium sulfide nonahydrate. In this reaction, 2-(prop-1-en-2-yl)-9*H*-xanthene-9-one (Scheme 1b, the proposed photoproduct **2**) was obtained as a by-product, and this was isolated for structural confirmation of the photo-product of SPD-2. Finally, hydrolysis gave the desired product SPD-2. The structure and purity of SPD-2 were confirmed by  $^1\text{H}$  NMR,  $^{13}\text{C}$  NMR, mass spectrometry and elemental analysis.



**Figure 3.** Fluorescence images of cells treated with HSip-1 DA (30  $\mu\text{M}$ ) in the presence or absence of SPD-2 AM with UVA photoirradiation. The treated cells were photoirradiated with UVA (330–380 nm) for 5 min. Representative images are shown in panels a–d: (a and b) Fluorescence images of cells treated with SPD-2 AM (a) before and (b) after photoirradiation; (c and d) images of cells without SPD-2 AM (c) before and (d) after photoirradiation. Photoirradiation was performed at the area indicated by the circle. All cells were treated with HSip-1 DA. Scale bar = 20  $\mu\text{m}$ . (e) Ratio of average fluorescence intensities inside the indicated circle at 10 and 0 min ( $F_{\text{after}}/F_{\text{before}}$ ) after 5 min UV-irradiation (330–380 nm) in D-PBS. Data represent means and standard deviation ( $n = 3$ ). \* $P < 0.05$  versus control.

To verify  $\text{H}_2\text{S}$  release from SPD-2 upon photoirradiation, we utilized methylene blue assay, a widely used colorimetric method for  $\text{H}_2\text{S}$  detection.<sup>16</sup> In this study, a 100  $\mu\text{M}$  solution of SPD-2 in pH 7.4 phosphate buffer (containing 1% DMSO as a co-solvent) was prepared. When a solution of SPD-2 was subjected to UV irradiation at 325–385 nm for various durations,  $\text{H}_2\text{S}$  release was clearly irradiation-dependent (Figs. 1, S2). When 100  $\mu\text{M}$  SPD-2 was irradiated for 400 s, the concentration of released  $\text{H}_2\text{S}$  was determined as 47  $\mu\text{M}$ , while SPD-1 released only 15  $\mu\text{M}$   $\text{H}_2\text{S}$  under the same conditions.<sup>10</sup> This result indicated that SPD-2 is a more efficient  $\text{H}_2\text{S}$  donor than SPD-1, presumably because the maximum absorption range of SPD-2 is well matched to the wavelength of the irradiating light, whereas that of SPD-1 is not (Fig. S1).

For further confirmation of photo-dependent  $\text{H}_2\text{S}$  release, we employed HSip-1, a  $\text{H}_2\text{S}$ -selective fluorescent probe that utilizes azamacrocyclic copper (II) ion complex chemistry to control the fluorescence.<sup>17</sup> As shown in Figure 2, the fluorescence intensity of HSip-1 was dramatically increased upon irradiation in the presence of 100  $\mu\text{M}$  SPD-2 in 100 mM PBS, pH 7.4 containing 1% DMSO as a co-solvent (an aliquot was taken after irradiation for 300 s). Almost no fluorescence increment was observed in the controls (Fig. 2a–c).

We confirmed the formation of the photoproduct **2**, illustrated in Scheme 1b. Photodecomposition of SPD-2 was monitored by HPLC. As shown in Figure S3, the peak of SPD-2 decreased and a new peak around 14 min was observed in the irradiated solution. The absorbance and retention time of the newly formed peak was identical to those of authentic **2**. This result confirmed that photolysis-induced  $\text{H}_2\text{S}$  generation proceeded in the expected manner.

Next, we examined the stability of SPD-2. A solution of 100  $\mu\text{M}$  SPD-2 in pH 7.4 phosphate buffer was incubated for 1 month at 37  $^\circ\text{C}$  in the dark. However, no decomposition was detected by means of HPLC (data not shown). This result suggested that SPD-2 is remarkably stable in the dark and has good light-responsiveness as a  $\text{H}_2\text{S}$  photo-donor.

Next, we applied this donor to living cells (Fig. 3). However, SPD-2 probably has poor membrane permeability due to the hydrophilic carboxyl moieties. Therefore, these moieties were acetoxymethylated to improve the membrane permeability. Acetoxymethylated SPD-2 (SPD-2 AM) should permeate cell membrane and be intracellularly hydrolyzed to SPD-2. HEK293 cells were incubated with 100  $\mu\text{M}$  SPD-2 AM and 30  $\mu\text{M}$  HSip-1 DA (membrane-permeable

HSip-1) in Dulbecco's modified Eagle's medium (DMEM) for 30 min, then washed with Dulbecco's phosphate-buffered saline (D-PBS), and photo-irradiated for 5 min at the site indicated by the circle in Figure 3a. Upon photoirradiation, fluorescence enhancement was observed (Fig. 3b and e), while little fluorescence increment was observed in cells outside the photoirradiated area. Additionally, SPD-2 did not show strong toxicity at low micromolar level which is anticipated as the active dose of  $\text{H}_2\text{S}$  donors.<sup>18</sup> (Fig. S4). Under the same conditions, SPD-1 showed no fluorescence enhancement (data not shown), probably because its efficiency of  $\text{H}_2\text{S}$  release on irradiation at this wavelength was much lower than that of SPD-2. These results indicated that SPD-2 is suitable for use in live cells.

In conclusion, we have developed a novel photo-controllable  $\text{H}_2\text{S}$  donor, SPD-2, using xanthone PPG. Generation of  $\text{H}_2\text{S}$  from SPD-2 can be precisely controlled by irradiation at 325–385 nm (UVA). This donor is available both in aqueous solution and in live cells. We expect that SPD-2 will be a useful tool for elucidating a wide range of biological functions of  $\text{H}_2\text{S}$ .

#### Acknowledgments

This work was supported by the JST PRESTO program (H.N.) of the Japan Science and Technology Agency, as well as by a JSPS KAKENHI Grant Number 25293028 (H.N.), a Grant from Hoansha Foundation (H.N.), and a Sasakawa Scientific Research Grant from The Japan Scientific Society (No. 25-311, N.F.).

#### Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.bmcl.2014.11.084>.

#### References and notes

- (a) Paul, B. D.; Snyder, S. H. *Nat. Rev. Mol. Cell Biol.* **2012**, *13*, 499; (b) Mustafa, A. K.; Gadalla, M. M.; Snyder, S. H. *Sci. Signal.* **2009**, *2*, 2; (c) Szabo, C. *Nat. Rev. Drug Disc.* **2007**, *6*, 917.
- (a) Zhao, W.; Zhang, J.; Lu, Y.; Wang, R. *EMBO J.* **2001**, *20*, 6008; (b) Yang, G.; Wu, L.; Jiang, B.; Qi, J.; Meng, Q.; Mustafa, A. K.; Mu, W.; Zhang, S.; Snyder, S. H.; Wang, R. *Science* **2008**, *322*, 587.
- Abe, K.; Kimura, H. *J. Neurosci.* **1996**, *16*, 1066.
- Zanardo, R. C. O.; Brancaleone, V.; Distrutti, E.; Fiorucci, S.; Cirino, G.; Wallace, J. L. *FASEB J.* **2006**, *20*, 2118.

5. Chen, X.; Jhee, K.; Kruger, W. D. *J. Biol. Chem.* **2004**, *279*, 52082.
6. Chiku, T.; Padovani, D.; Zhu, W.; Singh, S.; Vitvitsky, V.; Banerjee, R. *J. Biol. Chem.* **2002**, *277*, 42680.
7. Shibuya, N.; Tanaka, M.; Yoshida, M.; Ogasawara, Y.; Togawa, T.; Ishii, K.; Kimura, H. *Antioxid. Redox Signal.* **2009**, *11*, 703.
8. Shibuya, N.; Koike, S.; Tanaka, M.; Ishigami-Yuasa, M.; Kimura, Y.; Ogasawara, Y.; Fukui, K.; Nagahara, N.; Kimura, H. *Nat. Commun.* **2013**, *4*, 1366.
9. Ida, T.; Sawa, T.; Ihara, H.; Tsuchiya, Y.; Watanabe, Y.; Kumagai, Y.; Suematsu, M.; Motohashi, H.; Fujii, S.; Matsunaga, T.; Yamamoto, M.; Ono, K.; Devarie-Baez, N. O.; Xian, M.; Fukuto, J. M.; Akaike, T. *Proc. Natl. Acad. Sci. U.S.A.* **2014**, *111*, 7606.
10. Ono, K.; Akaike, T.; Sawa, T.; Kumagai, Y.; Wink, D. A.; Tantillo, D. J.; Hobbs, A. J.; Nagy, P.; Xian, M.; Lin, J.; Fukuto, J. M. *Free Radic. Biol. Med.* **2014**, *77*, 82.
11. Devarie-Baez, N. O.; Bagdon, P. E.; Peng, B.; Zhao, Y.; Park, C.; Xian, M. *Org. Lett.* **2013**, *15*, 2786.
12. Fukushima, N.; Ieda, N.; Sasakura, K.; Nagano, T.; Hanaoka, K.; Suzuki, T.; Miyata, N.; Nakagawa, H. *Chem. Commun.* **2014**, 587.
13. Lukeman, M. C.; Scaiano, J. J. *Am. Chem. Soc.* **2005**, *127*, 7698.
14. Il'ichev, Y. V.; Schwörer, M. A.; Wirz, J. *J. Am. Chem. Soc.* **2004**, *126*, 4581.
15. (a) Blake, J. A.; Lukeman, M.; Scaiano, J. C. *J. Am. Chem. Soc.* **2009**, *131*, 4127; (b) Blake, J. A.; Gagnon, E.; Lukeman, M.; Scaiano, J. C. *Org. Lett.* **2013**, *15*, 2786.
16. (a) Siegel, L. M. *Anal. Biochem.* **1965**, *11*, 126; (b) Shen, X.; Pattillo, C. B.; Pardue, S.; Bir, S. C.; Wang, R.; Kevil, C. G. *Free Radic. Biol. Med.* **2011**, *50*, 1021.
17. Sasakura, K.; Hanaoka, K.; Shibuya, N.; Mikami, Y.; Kimura, Y.; Komatsu, T.; Ueno, T.; Terai, T.; Kimura, H.; Nagano, T. *J. Am. Chem. Soc.* **2011**, *133*, 18003.
18. Zhao, Y.; Biggs, T. D.; Xian, M. *Chem. Commun.* **2014**, 11788.