


Epithelial IL-18 Equilibrium Controls Barrier Function in Colitis

Graphical Abstract



Authors

Roni Nowarski, Ruaidhrí Jackson, Nicola Gagliani, ..., Morven Graham, Eran Elinav, Richard A. Flavell

Correspondence

richard.flavell@yale.edu

In Brief

Production of the cytokine IL-18 by epithelial cells, previously thought to protect the mucosal barrier from the effect of inflammation, is critical to drive the pathologic breakdown of intestinal barrier integrity, directly inhibiting goblet cell maturation prior to the onset of colitis.

Highlights

- IL-18/IL-18R signaling in intestinal epithelial cells promotes DSS-induced colitis
- Hyperactive epithelial IL-18 signaling drives goblet cell depletion during colitis
- Epithelial IL-18 signaling prevents goblet cell maturation prior to colitis
- IL-18 intercepts the transcriptional program controlling goblet cell development



Epithelial IL-18 Equilibrium Controls Barrier Function in Colitis

Roni Nowarski,^{1,5} Ruaidhrí Jackson,^{1,5} Nicola Gagliani,¹ Marcel R. de Zoete,¹ Noah W. Palm,¹ Will Bailis,¹ Jun Siong Low,¹ Christian C.D. Harman,¹ Morven Graham,² Eran Elinav,³ and Richard A. Flavell^{1,4,*}

¹Department of Immunobiology, Yale University School of Medicine, New Haven, CT 06520, USA

²Center for Cellular and Molecular Imaging, Yale School of Medicine, New Haven, CT 06520, USA

³Immunology Department, Weizmann Institute of Science, Rehovot 76100, Israel

⁴Howard Hughes Medical Institute, Yale University, New Haven, CT 06520, USA

⁵Co-first author

*Correspondence: richard.flavell@yale.edu

<http://dx.doi.org/10.1016/j.cell.2015.10.072>

SUMMARY

The intestinal mucosal barrier controlling the resident microbiome is dependent on a protective mucus layer generated by goblet cells, impairment of which is a hallmark of the inflammatory bowel disease, ulcerative colitis. Here, we show that IL-18 is critical in driving the pathologic breakdown of barrier integrity in a model of colitis. Deletion of *Il18* or its receptor *Il18r1* in intestinal epithelial cells (Δ/EC) conferred protection from colitis and mucosal damage in mice. In contrast, deletion of the IL-18 negative regulator *Il18bp* resulted in severe colitis associated with loss of mature goblet cells. Colitis and goblet cell loss were rescued in *Il18bp*^{-/-};*Il18r* ^{Δ/EC} mice, demonstrating that colitis severity is controlled at the level of IL-18 signaling in intestinal epithelial cells. IL-18 inhibited goblet cell maturation by regulating the transcriptional program instructing goblet cell development. These results inform on the mechanism of goblet cell dysfunction that underlies the pathology of ulcerative colitis.

INTRODUCTION

Inflammatory bowel disease (IBD) is a complex and debilitating disorder that can be sub-classified into the distinct multifactorial disorders Crohn's disease (CD) and ulcerative colitis (UC) (Kaser et al., 2010; Maloy and Powrie, 2011). While both are characterized by chronic relapsing pathogenic inflammation and intestinal epithelial cell injury, they differ substantially in their clinical manifestations. CD patients exhibit discontinuous lesions throughout the entirety of the intestinal tract and disease pathology is closely associated with a dysregulation of the antimicrobial peptide (AMP) response (Fellermann et al., 2003; Neurath, 2014). A genetic basis for CD susceptibility has been linked to genes involved in autophagy and ER stress (e.g., *Atg16l1* and *Xbp1*), as well as microbial recognition (e.g., *Nod2*), in AMP-producing Paneth cells (Adolph et al., 2013; Cadwell et al., 2008; Hugot et al., 2001; Ogura et al., 2001). Interestingly, however, no major

defects in AMP production have been observed in UC patients (Nuding et al., 2007; Wehkamp et al., 2007), indicating distinct mechanistic differences in disease etiology. Despite UC having greater worldwide prevalence than CD (Danese and Fiocchi, 2011), surprisingly little is known about the specific underlying host factors that drive susceptibility to disease. One unique and defining feature of human UC pathology is major depletion of mucin-producing goblet cells and the mucus layer, which correlates with increased microbiota-induced colonic inflammation and disease pathology (McCormick et al., 1990; Pullan et al., 1994; Strugala et al., 2008; Trabucchi et al., 1986). Intriguingly, the in vivo mechanisms responsible for this important clinical observation during inflammation remain obscure.

Members of the IL-1 family of cytokines play critical roles in intestinal homeostasis and inflammation (Lopetuso et al., 2013; Neurath, 2014; Saleh and Trinchieri, 2011). In particular, IL-18 has emerged as an indispensable factor in governing host-microorganism homeostasis and has been postulated to be a key determining factor in IBD (Dinarelo et al., 2013; Elinav et al., 2011; Nakamura et al., 1989). IL-18 is initially synthesized as an inactive precursor molecule that requires coordinated inflammatory activation of the cysteine protease caspase-1 to cleave proIL-18 into a functional mature bioactive cytokine (Fantuzzi et al., 1999; Martinon et al., 2002). Upon activation and release, IL-18 is free to bind the IL-18 receptor alpha chain (IL-18R1) and in cells co-expressing the IL-18R accessory protein (IL-18Rap), ligand binding triggers receptor heterodimerization and the formation of an intracellular Myd88 signaling platform (Adachi et al., 1998; Born et al., 1998; Hoshino et al., 1999). This elicits the recruitment of IRAK and TRAF6, facilitating activation of the inhibitor of κ B ($\text{I}\kappa\text{B}$) kinases (IKKs), IKK α and IKK β (Medzhitov et al., 1998; Mercurio et al., 1997). In turn, IKK β can phosphorylate $\text{I}\kappa\text{B}\alpha$, targeting the protein for proteasomal degradation and allowing the NF- κ B subunit p65 to translocate to the nucleus to initiate diverse gene expression programs such as proinflammatory cytokine production and NOD-Like Receptor (NLR) upregulation (Bauernfeind et al., 2009). As such, IL-18 signaling requires tight regulation to prevent autoimmunity and this is thought to be directly accomplished by the soluble decoy receptor IL-18 binding protein (IL-18BP), as its transgene overexpression has been shown to neutralize IL-18 activity in vivo to prevent hyper NF- κ B activation and inflammation (Fantuzzi et al., 2003).

The use of IL-18- and IL-18R1-deficient mice identified IL-18 as a putative host molecule required to protect intestinal epithelial cells from intestinal inflammation and colitis (Salcedo et al., 2010). In support of a role for IL-18 in promoting intestinal epithelial integrity and protection from acute experimental colitis, mice deficient in the key processing subunits of IL-18, caspase 1, and the NLRP3 inflammasome are also highly susceptible to disease pathology (Dupaul-Chicoine et al., 2010; Zaki et al., 2010). Administration of exogenous recombinant IL-18 rescues colitis in the aforementioned and other inflammasome-deficient mice, further supporting a protective role for IL-18 in colitis (Oficjalska et al., 2015). In contrast, however, inhibition of IL-18 has also been shown to instigate protection in experimental colitis, supporting a pro-colitogenic role for IL-18 (Kanai et al., 2001; Siegmund et al., 2001; Ten Hove et al., 2001). Such conflicting findings have led to much controversy and discussion in the field, and the true role of IL-18 in intestinal homeostasis and inflammation is still unresolved (Asquith and Powrie, 2010; Dinarello et al., 2013; Gagliani et al., 2014; Siegmund, 2010).

Underlying this discourse is the fact that most previous work studying the complete IL-18 deletion in mice is confounded by IL-18 effect on colitogenic microbiota (Elinav et al., 2011; Henao-Mejia et al., 2012), while equally important roles of IL-18 during inflammation are masked by dysbiosis. Compound-associated phenotypic alterations in *Il18*^{-/-} mice such as metabolic syndrome may further obscure the direct contribution of IL-18 to intestinal function (Netea et al., 2006). At present, no genetic models exist to specifically dissect the role of IL-18 in colitis risk, and the need for new genetic tools is therefore paramount. To this end, we generated conditional knockout mice for both IL-18 and IL-18R1 to delineate the direct involvement of IL-18 in epithelial and hematopoietic cells to homeostasis and colitis. Here, we show that IL-18 production, irrespective of its cellular source, exacerbated colitis severity after administration of the colitis-inducing agent dextran sodium sulfate (DSS). Deletion of IL-18R in epithelial cells (*Il18r*^{Δ/EC}) protected mice from developing colitis, suggesting that IL-18 directly disrupts epithelial cell integrity during colitis. By deleting IL-18 binding protein (*Il18bp*), the IL-18 negative regulator, we asked if increased bioavailability of IL-18 would promote barrier function or rather drive colitis. Remarkably, *Il18bp*^{-/-} mice developed severe colitis associated with progressive loss of mature goblet cells, which could be reversed by specifically deleting the epithelial IL-18R in these mice. Finally, we show that IL-18-mediated goblet cell dysfunction precedes clinical disease manifestation and is caused by a defect in terminal goblet cell maturation through transcriptional regulation of goblet cell differentiation factors. Taken together, these results uncover the direct role of IL-18 in promoting goblet cell dysfunction during colitis, leading to breakdown of the mucosal barrier. This study may therefore offer a genetic understanding to the pathology of human ulcerative colitis.

RESULTS

Epithelial IL-18/IL-18R Signaling Promotes DSS-Induced Colitis

IL-18 is a key mediator of intestinal homeostasis and inflammation, yet the cellular partners and molecular mechanisms driving

these effects remain poorly understood. To delineate the compound role of IL-18 in intestinal inflammation, we conditionally deleted *Il18* or *Il18r1* in intestinal epithelial cells by generating *Villin-cre*⁺;*Il18*^{fl/fl} (hereafter called *Il18*^{Δ/EC}) and *Villin-cre*⁺;*Il18r*^{fl/fl} (*Il18r*^{Δ/EC}) mice (Figures S1A–S1D). To enable mechanistic evaluation of IL-18's microbiota-independent roles, throughout this study knockout mice were compared to their cohoused floxed (fl/fl) wild-type littermates. Indeed, bacterial 16S rRNA sequencing confirmed equalized bacterial composition in both *Il18*^{Δ/EC} and *Il18r*^{fl/fl} littermates (Figure S2A). IL-18 production in *Il18*^{Δ/EC} total colon explants was markedly reduced (Figure S1B), confirming IECs as the major source of IL-18 under physiological conditions (Takeuchi et al., 1997). Steady-state colon sections did not show gross structural or cellular irregularities in *Il18*^{Δ/EC} or *Il18r*^{Δ/EC} mice, including goblet cell maturation and tight junction formation, as determined by MUC2, β-catenin, and ZO-1 staining (Figure S3). Nevertheless, *Il18*^{Δ/EC} mice were surprisingly resistant to colonic inflammation following administration of DSS, as reflected by reduced weight loss compared to *Il18*^{fl/fl} littermates (Figure 1A). Colonoscopy performed on day 7 post DSS showed increased tissue damage in control *Il18*^{fl/fl} mice, measured by the degree of bleeding, colon wall granularity and translucency, as well as stool consistency (Figure 1B). Similarly to *Il18*^{Δ/EC} mice, DSS-treated *Il18r*^{Δ/EC} mice were protected against weight loss, as compared to *Il18r*^{fl/fl} littermates (Figure 1C). To more rigorously assess these effects in the presence of a "colitogenic" microbiota, *Il18*^{Δ/EC} and *Il18r*^{fl/fl} were cohoused for 8 weeks with dysbiotic *Il18*^{-/-} mice in order to introduce transmissible dominantly colitogenic bacteria (Elinav et al., 2011) (Figure S2B). Despite an overall higher degree of inflammation, *Il18r*^{Δ/EC} mice had reduced weight loss and lower colonoscopy score than control *Il18r*^{fl/fl} mice (Figures 1D and 1E). Severe colitis and deterioration of tissue integrity in *Il18r*^{fl/fl} mice, but not in *Il18r*^{Δ/EC} mice, was corroborated by histological examination of distal colon sections performed on day 8 post DSS (Figure 1F). These results suggest that IL-18 promotes the pathology of DSS-induced colitis through a mechanism dependent on its action on intestinal epithelial cells.

Hematopoietic/Endothelial IL-18, but Not IL-18R, Promotes DSS-Induced Colitis

In addition to epithelial cells, IL-18 and IL-18R are also expressed by various hematopoietic and endothelial cells, in particular under inflammatory conditions (Siegmund, 2010). To address the role of the IL-18 axis in these cells during colitis, we generated *Flk1-cre*⁺;*Il18*^{fl/fl} (*Il18*^{Δ/H_E}) and *Flk1-cre*⁺;*Il18r*^{fl/fl} (*Il18r*^{Δ/H_E}) mice in which *Il18* or *Il18r* are specifically deleted in all hematopoietic and endothelial cells (Figure S1B). As above, knockout mice were compared to their cohoused floxed (fl/fl) wild-type littermates, with both featuring similar microbiome configurations (including the colitogenic *Prevotellaceae* species), thus enabling us to study in detail the microbiome-independent contribution of hematopoietic IL-18 to the intestinal pathology in these mice (Figures S2C and S2D). Consistent with deletion of IL-18 in epithelial cells, *Il18*^{Δ/H_E} mice were highly protected in DSS-induced colitis, as indicated by reduced weight loss and colonoscopy scores compared to *Il18*^{fl/fl} littermates (Figures 2A and 2B). In contrast, *Il18r*^{Δ/H_E} mice were susceptible to extensive




Figure 1. Epithelial IL-18/IL-18R Signaling Promotes DSS-Induced Colitis

To induce colitis, mice were administered 2% DSS in drinking water for 7 days.

(A) Weight loss of cohoused *Il18^{fl/fl}* and *Il18^{Δ/EC}* littermates (n = 11–14).

(B) Colonoscopy severity score of *Il18^{fl/fl}* and *Il18^{Δ/EC}* mice on day 7 and 11 post DSS (left) and representative endoscopic view of the mouse colon on day 7 post DSS (right).

(C) Weight loss of cohoused *Il18^{fl/fl}* and *Il18^{Δ/EC}* littermates (n = 8).

(D) Littermate *Il18^{fl/fl}* and *Il18^{Δ/EC}* mice were cohoused with *Il18^{-/-}* mice for 8 weeks, after which DSS was administered and weight loss recorded (n = 7–13).

(E) Colonoscopy severity score of cohoused *Il18^{fl/fl}*, *Il18^{Δ/EC}*, and *Il18^{-/-}* on day 7.

(F) Representative H&E staining of distal colon sections obtained from cohoused *Il18^{fl/fl}*, *Il18^{Δ/EC}*, and *Il18^{-/-}* mice given water (top) or DSS (bottom) and assessed on day 8. Note the disruption of crypt structure and mucosal immune cell infiltration in *Il18^{fl/fl}* and *Il18^{-/-}* mice. Scale bar, 25 μm. Data are represented as mean ± SEM. *p < 0.05; **p < 0.01; ***p < 0.001; ****p < 0.0001 by unpaired Student's t test.

See also [Figures S1, S2, and S3](#).

weight loss and tissue damage, to a comparable degree as their *Il18^{fl/fl}* littermates ([Figures 2C and 2D](#)). Histology performed on day 8 post DSS confirmed similar extent of colitis in both *Il18^{fl/fl}* and *Il18^{Δ/EC}* mice ([Figure 2E](#)). These results further demonstrate

that irrespective of its cellular source, IL-18 production during colitis drives disease progression. Colitis severity, however, is not exacerbated by IL-18R signaling in hematopoietic and/or endothelial cells, in contrast to what is observed in epithelial




Figure 2. Hematopoietic/Endothelial IL-18, but Not IL-18R, Promotes DSS-Induced Colitis

(A) Weight loss of cohoused *Il18^{fl/fl}* and *Il18^{Δ/HE}* littermates treated with 2% DSS for 7 days ($n = 4$). (B) Colonoscopy severity score of *Il18^{fl/fl}* and *Il18^{Δ/HE}* mice on day 7 and 11.

(C) Weight loss of cohoused *Il18^{r/fl}* and *Il18^{r/Δ/HE}* littermates ($n = 6$).

(D) Colonoscopy severity score of *Il18^{r/fl}* and *Il18^{r/Δ/HE}* mice on day 7.

(E) Representative H&E staining of distal colon sections obtained from cohoused *Il18^{fl/fl}* and *Il18^{Δ/HE}* mice given water (top) or DSS (bottom) and assessed on day 8. Scale bar, 25 μm . Data are represented as mean \pm SEM. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; by unpaired Student's *t* test. See also Figures S1, S2, and S3.

sion levels of *Il18bp* in the steady-state colon were low, it was highly induced during the course of colitis, returning to baseline levels following recovery (Figure 3A). To better understand the mechanism by which IL-18 enhances susceptibility to colitis, we generated mice with hyperactive IL-18 signaling by deleting *Il18bp* (Figure S1E). *Il18bp* expression was undetectable in *Il18bp^{-/-}* mice, whereas the expression of neighboring genes was unaffected (Figure S1F). Furthermore, in the steady-state *Il18bp^{-/-}* mice had equalized flora compared to their wild-type (WT) littermates (Figure S2E) and displayed normal goblet cell development and tight junction structure (Figure S3). Although *Il18* mRNA expression was comparable in WT and *Il18bp^{-/-}* mice, the active secreted form of IL-18 was elevated in *Il18bp^{-/-}* colon explant supernatants, both in the steady state and following DSS treatment (Figure 3B). During DSS colitis, *Il18bp^{-/-}* mice developed rapid and severe morbidity associated with extensive bleeding and tissue damage (Figures 3C and 3D). Extensive tissue deterioration and colitis were also evident in histological sections of *Il18bp^{-/-}* mice but not of their WT littermate controls (Figure 3E). Remarkably, *Il18bp^{-/-}* mice suffered an overwhelming loss of mucus-producing goblet cells (Figure 3E). The absence of mature goblet cells and associated mucus layer in *Il18bp^{-/-}* mice was verified by AB/PAS staining (Figure 3E). Goblet cell enumeration in histological sections showed a dramatic decrease in goblet cell counts in *Il18bp^{-/-}* mice compared to WT mice at day 8 post DSS (Figure 3E). Decreased goblet cell counts were also noted in *Il18^{fl/fl}* and *Il18^{r/fl}* mice compared to *Il18^{Δ/EC}* or *Il18^{r/Δ/EC}* littermates

cells. Together, these data show that the target of IL-18-mediated pathology is the epithelium.

Hyperactive IL-18 Signaling Drives Colitis and Goblet Cell Depletion in *Il18bp^{-/-}* Mice

IL-18 is negatively regulated by the IL-18 binding protein (IL-18BP), which serves as a decoy receptor and prevents IL-18 association with IL-18R (Novick et al., 1999). While basal expres-

sion levels of *Il18bp* in the steady-state colon were low, it was highly induced during the course of colitis, returning to baseline levels following recovery (Figure 3A). To better understand the mechanism by which IL-18 enhances susceptibility to colitis, we generated mice with hyperactive IL-18 signaling by deleting *Il18bp* (Figure S1E). *Il18bp* expression was undetectable in *Il18bp^{-/-}* mice, whereas the expression of neighboring genes was unaffected (Figure S1F). Furthermore, in the steady-state *Il18bp^{-/-}* mice had equalized flora compared to their wild-type (WT) littermates (Figure S2E) and displayed normal goblet cell development and tight junction structure (Figure S3). Although *Il18* mRNA expression was comparable in WT and *Il18bp^{-/-}* mice, the active secreted form of IL-18 was elevated in *Il18bp^{-/-}* colon explant supernatants, both in the steady state and following DSS treatment (Figure 3B). During DSS colitis, *Il18bp^{-/-}* mice developed rapid and severe morbidity associated with extensive bleeding and tissue damage (Figures 3C and 3D). Extensive tissue deterioration and colitis were also evident in histological sections of *Il18bp^{-/-}* mice but not of their WT littermate controls (Figure 3E). Remarkably, *Il18bp^{-/-}* mice suffered an overwhelming loss of mucus-producing goblet cells (Figure 3E). The absence of mature goblet cells and associated mucus layer in *Il18bp^{-/-}* mice was verified by AB/PAS staining (Figure 3E). Goblet cell enumeration in histological sections showed a dramatic decrease in goblet cell counts in *Il18bp^{-/-}* mice compared to WT mice at day 8 post DSS (Figure 3E). Decreased goblet cell counts were also noted in *Il18^{fl/fl}* and *Il18^{r/fl}* mice compared to *Il18^{Δ/EC}* or *Il18^{r/Δ/EC}* littermates




Figure 3. Hyperactive IL-18 Signaling Drives Colitis and Goblet Cell Loss in *Il18bp*^{-/-} Mice

(A) Wild-type (WT) mice were treated with 2% DSS in drinking water and *Il18bp* mRNA expression in the distal colon was measured over 14 days.

(B) IL-18 mRNA expression in distal colon (left) and protein secretion in colonic explants (right) of *Il18bp*^{-/-} and WT littermates.

(C) Weight loss following DSS treatment in cohoused WT and *Il18bp*^{-/-} littermates (n = 10).

(D) Colonoscopy severity score of WT and *Il18bp*^{-/-} mice.

(E) Representative H&E and AB/PAS staining of distal colon sections obtained from cohoused WT (top) and *Il18bp*^{-/-} (bottom) littermates on day 8 post DSS treatment. Asterisks mark example goblet cells. Right, enumeration of goblet cells in histological sections from cohoused WT and *Il18bp*^{-/-} littermates. Scale bar, 25 μ m. Data are represented as mean \pm SEM. *p < 0.05; **p < 0.01; ***p < 0.001; ****p < 0.0001 by unpaired Student's t test.

See also Figures S1, S2, S3, and S4.

following DSS (Figure S4A). These results suggest that hyper-IL-18 signaling leads to exacerbated colitis associated with mature goblet cell depletion.

Deletion of Epithelial IL-18R Rescues Goblet Cell Loss and Colitis in *Il18bp*^{-/-} Mice

We next asked whether goblet cell loss and the increased colitis severity observed in *Il18bp*^{-/-} mice arise as a general consequence of immune cell activation and inflammation due to hyperactive IL-18, or is rather directly dependent on IL-18 signaling mediated perturbation of epithelial cell function. To address this question in vivo, we generated *Il18bp*^{-/-};*Il18r* ^{Δ EC} double knockout mice, with the hypothesis that if increased colitis in *Il18bp*^{-/-} mice is a result of hyper immune activation, loss of IL-18R signaling in epithelial cells will not affect the course of disease. However, if increased colitis in *Il18bp*^{-/-} mice is a result of epithelial cell dysregulation, loss of IL-18R signaling in epithelial

cells may rescue the pathology in *Il18bp*^{-/-} mice. Following DSS treatment, *Il18bp*^{-/-};*Il18r* ^{Δ EC} mice were dramatically protected from weight loss in comparison to *Il18bp*^{-/-} littermates, to the same level as *Il18bp*^{+/+} WT littermates (Figure 4A). This correlated with decreased intestinal bleeding and tissue damage in *Il18bp*^{-/-};*Il18r* ^{Δ EC} mice, assessed by colonoscopy at day 7 post DSS (Figure 4B). Upon dissection at day 8 post DSS, *Il18bp*^{-/-} mice exhibited macroscopic indications of severe intestinal inflammation with extensive bleeding throughout the intestine and significant shortening of the colon, while the intestines of *Il18bp*^{-/-};*Il18r* ^{Δ EC} mice appeared normal (Figure 4C). Colon tissue sections obtained at day 8 post DSS revealed reduced immune cell infiltration and maintenance of structural integrity in *Il18bp*^{-/-};*Il18r* ^{Δ EC} mice, while *Il18bp*^{-/-} mice showed extensive tissue pathology (Figure 4D). Importantly, goblet cells were completely recovered in *Il18bp*^{-/-};*Il18r* ^{Δ EC} mice, representing the baseline levels observed in WT mice, as




Figure 4. Deletion of Epithelial IL-18R Rescues Goblet Cell Loss and Colitis in $Il18bp^{-/-}$ Mice

(A) Weight loss of cohoused $Il18bp^{-/-}$, $Il18bp^{-/-}; Il18r^{\Delta/EC}$, and WT littermates treated with 2% DSS for 7 days (n = 10–14).

(B) Colonoscopy severity score of $Il18bp^{-/-}$ and $Il18bp^{-/-}; Il18r^{\Delta/EC}$ littermates on day 7.

(C) Gross pathology of colons on day 8 post DSS. Note the extensive shortening, bleeding, and diarrhea in $Il18bp^{-/-}$ mice.





(D) Representative H&E and AB/PAS staining of distal colon sections obtained from cohoused $Il18bp^{-/-}$ (top) and $Il18bp^{-/-}; Il18r^{\Delta/EC}$ mice (bottom) on day 8 post DSS treatment. Asterisks mark example goblet cells. Scale bar, 25 μ m. Data are represented as mean \pm SEM. **p < 0.01; ***p < 0.001; ****p < 0.0001 by unpaired Student's t test.

demonstrated by both H&E and AB/PAS staining (Figure 4D; compare to Figure 3E). These results indicate that goblet cell loss and increased susceptibility to colitis are a direct consequence of increased IL-18 signaling in intestinal epithelial cells.

Hyperactive IL-18 Signaling Prevents Goblet Cell Maturation Prior to Colitis

To determine whether the observed goblet cell dysfunction is a cause rather than a consequence of inflammation, we examined histological colon sections from $Il18bp^{-/-}$ mice at day 4 post DSS, preceding significant weight loss and clinical symptoms of colitis. Although goblet cells were readily identified by H&E staining in $Il18bp^{-/-}$ mice, AB/PAS staining indicated decreased abundance of mature PAS⁺ goblet cells in $Il18bp^{-/-}$ mice compared to WT or $Il18bp^{-/-}; Il18r^{\Delta/EC}$ littermates (Figure 5A).

Goblet cell maturation was further assessed by staining for fucosylated glycoproteins with the lectin Ulex europaeus agglutinin-1 (UEA-1) and the abundant goblet cell mucin MUC2, both of which accumulate as goblet cells mature. $Il18bp^{-/-}$ mice exhibited an increase of immature goblet cells, determined by low area MUC2 staining (<10 μ m in diameter) in UEA-1^{lo/-} cells and decrease in large mature MUC2⁺UEA-1^{bright} goblet cells compared to $Il18bp^{-/-}; Il18r^{\Delta/EC}$ mice (Figure 5B). The mature/immature goblet cell ratio on day 4 post DSS decreased to 0.58 in $Il18bp^{-/-}$ mice compared to 1.39 in $Il18bp^{-/-}; Il18r^{\Delta/EC}$ mice and 1.84 in $Il18bp^{+/+}$ (WT) mice (Figures 5C, S4B, and S4C). As noted above, mature goblet cells were markedly depleted in $Il18bp^{-/-}$ mice on day 8 post DSS; however, small MUC2⁺UEA-1^{+/+} cells were still highly represented, notably at the lower half of the crypt (Figure S4D). To determine whether



(legend on next page)

dysregulation of goblet cell maturation reflects a transcriptional imbalance, we measured expression of transcription factors involved in goblet cell differentiation and maturation. Whereas no change was noted in the secretory lineage differentiation factors *Math1* (*Hath1*; *Atoh1*) and *Hes1*, expression of the goblet cell differentiation/maturation factors *Gfi1*, *Spdef*, and *Klf4* was markedly inhibited in *Il18bp^{-/-}* mice (Figure 5D). These results suggest that IL-18 promotes colitis by preventing functional goblet cell maturation through regulation of the goblet cell transcriptional maturation program.

IL-18 Directly Controls Goblet Cell Maturation and Colitis

We finally assessed the direct role of IL-18 in goblet cell dysfunction leading to colitis, by injecting recombinant IL-18 protein to WT mice during the course of DSS administration. Disease severity was increased in mice receiving daily IL-18 injections, as determined by weight loss and macroscopic examination of the colon at day 8 post DSS (Figures 6A and 6B). In line with our observations in *Il18bp^{-/-}* mice, AB/PAS staining showed gradual decrease in the abundance of mature PAS⁺ goblet cells in mice receiving IL-18 compared to PBS (Figure 6C). The state of goblet cell maturation was corroborated in colon sections obtained following 5 daily injections prior to weight loss and clinical symptoms of colitis, demonstrating an IL-18-mediated block in goblet cell maturation (Figures 6D and 6E). The ratio of mature/immature goblet cell decreased further in IL-18-injected mice on day 8 (Figures S4D and S4E). IL-18 injection was sufficient to reduce *Gfi1*, *Spdef*, and *Klf4* gene expression in isolated IECs, further supporting direct regulation of goblet cell maturation by IL-18 (Figure 6F). These results suggest that elevated IL-18 production during inflammation is responsible for dysregulated goblet cell maturation.

DISCUSSION

Despite great strides in our understanding of IL-18 over the past 15 years, its precise contributions to host homeostasis, intestinal inflammation, and its overall relevance to IBD still remain controversial and elusive. On one hand, complete loss of IL-18 (or IL-18R) predisposes mice to increased intestinal epithelial damage and fosters an altered inflammatory environment that potentiates intestinal tumor formation (Salcedo et al., 2010; Takagi et al., 2003). This could be explained, at least in part, by the recently identified role of IL-18 in controlling the outgrowth of colitogenic bacterial species (Elinav et al., 2011). On the other hand, IL-18 is a potent proinflammatory cytokine with the ability to promote colitis through the induction of inflammatory mediators such TNF α and chemokines (Sivakumar et al., 2002; Ten Hove et al., 2001).

The role of IL-18 in intestinal homeostasis and inflammation and its mechanistic segregation from microbiota-dependent functions therefore remained unresolved.

Previous interpretations of IL-18 functionality have been limited by the lack of precise genetic models required to systematically determine its roles in intestinal biology. Therefore, IL-18 function has been inferred from complete deletion of IL-18, inflammasomes, caspase 1/11, or the multifunctional adaptor protein ASC. Such studies have led to the conclusion that epithelial-derived IL-18 is required to promote barrier integrity during early inflammation, as acute treatment with recombinant IL-18 during early colitis promotes epithelial proliferation in inflammasome-deficient mice, rescuing intestinal pathology (Dupaul-Chicoine et al., 2010; Zaki et al., 2010). However, extrapolation of direct IL-18 functionality from these models should be approached with caution. First, deficiency of NLRP3, which is highly expressed in the myeloid compartment, results in numerous phenotypic alterations beyond IL-18 processing. Most obvious is an inherent defect in processing the closely related and equally important cytokine IL-1 β . Like IL-18, IL-1 β is also thought to mediate a dichotomous role in intestinal homeostasis and inflammation (Bamias et al., 2012; Lopetuso et al., 2013). Notably, bone marrow chimera experiments have shown that hematopoietic-derived IL-1 β is also sufficient to rescue epithelial cell damage and promote epithelial restitution during experimental colitis (Bersudsky et al., 2014). Therefore, in NLRP3-deficient mice, which harbor defects in IL-1 family member maturation, IL-18 may compensate for the lack of IL-1 β ; however, whether this occurs physiologically (or at physiologically relevant levels of IL-18) remains unclear. In addition, caspase 1 plays a key role in the clearance of intracellular intestinal pathogens through the regulated cell death process of pyroptosis (Miao et al., 2010). Although the role of pyroptosis in colitis is still under investigation, the use of pyroptosis-defective mice to examine the specific IL-18 functionality in the intestine proves problematic. The study of direct functions of IL-18 in the intestine is further complicated by NLRP6 regulation of dysbiosis and the outgrowth of pathogenic intestinal microbial communities (Elinav et al., 2011). As demonstrated by Levy et al. (2015) in this issue of *Cell*, IL-18 processing by the NLRP6 inflammasome shapes the steady-state host-microbiome interface by regulating the downstream anti-microbial peptide (AMP) landscape, thereby maintaining intestinal homeostasis. Normally, this axis is controlled by indigenous microbiota-modulated metabolites. However, it can also be directly subverted by inflammasome suppressing metabolites derived from a disease-causing microbiota, which hijacks this pathway, thereby facilitating dysbiosis development and persistence in an invaded host. This highlights the

Figure 5. Hyperactive IL-18 Signaling Prevents Goblet Cell Maturation Prior to Colitis

(A) Representative H&E and AB/PAS staining of distal colon sections obtained from WT, *Il18bp^{-/-}*, and *Il18bp^{-/-};Il18 ^{Δ IEC}* littermates on day 4 post DSS treatment prior to onset of colitis. Note reduction in PAS⁺ mature goblet cells in *Il18bp^{-/-}* mice. Scale bar, 25 μ m.

(B) Epifluorescence images (left, top right panels) and confocal stacks (bottom right panels) of distal colon sections obtained on day 4 post DSS and stained with the fucose-binding lectin UEA-1 and anti-MUC2. Scale bar, 150 μ m.

(C) Enumeration of immature and mature goblet cells in distal colon sections stained as in (B). Immature goblet cells were scored as UEA-1^{low} cells containing low area (<10 μ m in diameter) MUC2 staining, and mature goblet cells were scored as cells containing large area (>10 μ m in diameter) MUC2⁺UEA-1^{bright} staining.

(D) Distal colon samples were obtained on day 8 post DSS and used for gene expression analysis by qPCR. Data are represented as mean \pm SEM. * p < 0.05; **** p < 0.0001 by unpaired Student's *t* test.

See also Figure S4.




Figure 6. IL-18 Directly Controls Goblet Cell Maturation and Colitis

(A) WT mice received daily intraperitoneal (i.p.) injections of 1 μ g recombinant IL-18 or PBS during the course of 7 day 2% DSS administration and weight loss was recorded (n = 5).

(B) Gross pathology of colons on day 8 post DSS. Note reduction in colon length following IL-18 treatment.

(C) Representative AB/PAS staining of distal colon sections obtained on day 5 or 8 post DSS treatment from WT littermates receiving recombinant IL-18 or PBS. Note reduction in PAS⁺ mature goblet cells in mice receiving recombinant IL-18. Scale bar, 25 μ m.

(legend continued on next page)

importance of using cohoused littermate control mice, as in the present study, as they harbor near identical bacterial species enabling distinction of the genetic contribution of IL-18 from that of flora driven inflammation.

In this study, we show that during inflammation, not only is IL-18 production in intestinal epithelial and hematopoietic cells not required for protection against experimental colitis, IL-18 signaling in epithelial cells amplifies intestinal damage. This pathogenic role of IL-18 correlates with clinical observations whereby an increase in both epithelial and hematopoietic IL-18 expression and cytokine bioactivity have been demonstrated in patients with increased severity of IBD (Monteleone et al., 1999; Pizarro et al., 1999). However, the mechanism through which this upregulation of IL-18 in the intestine may contribute to increased disease severity was unknown. An emerging realization in the complexity of IBD is that pathology is not wholly shaped by a dysregulated immune response but highly dependent on an intact mucosal barrier and coordinated cross talk between the intestinal epithelial and immune cells with the microbiota (Kaser et al., 2011; Schreiber et al., 2005; Xavier and Podolsky, 2007). One possible mechanism to explain this association is that increased IL-18 release from epithelial cells acts on resident immune cell to upregulate IL-18 and other proinflammatory mediators, which induce endothelial VCAM-1 expression to enhance immune cell infiltration into the mucosa and together trigger severe auto-inflammation. In support of this model, we show that deletion of IL-18 production in the hematopoietic compartment results in significant amelioration of intestinal damage during colitis. However, deletion of IL-18R signaling in the hematopoietic compartment fails to rescue mice from DSS-induced inflammation. This suggests that the pathology driven by IL-18 does not occur via signaling in hematopoietic cells, in line with previous reports (Dupaul-Chicoine et al., 2010; Malvin et al., 2012; Saleh and Trinchieri, 2011; Zaki et al., 2010). Rather, we found that deletion of the IL-18R from intestinal epithelial cells dramatically protects mice from DSS-induced colitis, suggesting that elevated IL-18 expression during colitis is directly pathogenic to the epithelial cell barrier.

Ulcerative colitis is characterized by mucosal barrier dysfunction, most notably in epithelial goblet cells and mucus production (Danese and Focchi, 2011; Gersemann et al., 2009; McCormick et al., 1990; Pullan et al., 1994; Trabucchi et al., 1986). As goblet cell secretion of protective mucins, trefoil factors, and other proteins is essential for barrier integrity and for preventing microflora-driven intestinal inflammation, such dysregulation underlies the pathology exhibited in UC patients. In order to investigate how IL-18 may specifically contribute to intestinal barrier breakdown during DSS colitis, we deleted its decoy receptor inhibitor, IL-18BP. Interestingly, *Il18bp*^{-/-} mice were characterized by increased colitis severity and lethality associated with major depletion of mature goblet cells, which was reversed in

Il18bp^{-/-}; *Il18r*^{Δ/EC} double knockout mice. Thus, excessive IL-18 signaling on the epithelium leads to progressive depletion of goblet cells and may represent a major risk factor for intestinal inflammation and UC. As severe intestinal inflammation has previously been suggested to result in goblet cell depletion (Bergstrom et al., 2008), we analyzed mice during preclinical manifestation of colitis in order to explore mechanistically if IL-18 was the key determining factor governing goblet cell loss and risk for colitis. Whereas we observed no discernible differences in goblet cell numbers at preclinical time points, we instead discovered that IL-18 promotes disequilibrium in the state of goblet cell development and maturation, decreasing the pool of functional mature goblet cells.

Studies on the transcriptional regulation of intestinal lineage determination have identified the factors controlling goblet cell differentiation and maturation (McCauley and Guasch, 2015). Notch1 signaling drives epithelial progenitor cell specification to absorptive enterocytes and inhibits secretory cell differentiation via the Hes1 transcription factor (Fre et al., 2005; Jensen et al., 2000), whereas Notch1 inhibition results in robust conversion of proliferating crypt progenitors into terminally differentiated goblet cells (van Es et al., 2005). Below a threshold of Notch1 signal, upregulation of Math1 (also known as Hath1, Atoh1) governs commitment of progenitor cells to the epithelial secretory cell lineages, including goblet cells, Paneth cells, and neuroendocrine cells (Yang et al., 2001). The transcriptional repressor Gfi1 functions downstream of Math1 to control intestinal secretory cell differentiation, specification, and maintenance and is required for goblet and Paneth cell development (Shroyer et al., 2005). The transcription factors Spdef and Klf4 in turn control goblet cell development and maturation downstream to Gfi1 and are essential for mucosal barrier protection (Gregorieff et al., 2009; Katz et al., 2002; Noah et al., 2010). Analysis of the lineage commitment transcription factors Hes1 and Math1 in the intestinal epithelium of mice with hyperactive IL-18 did not indicate dysregulation at the lineage specification level. Conversely however, the downstream secretory lineage transcription factor Gfi1 and the terminal goblet cell maturation effectors Spdef and Klf4 were markedly downregulated by IL-18. Importantly, all three transcription factors are required to be constitutively expressed in mature goblet cells and loss of either Gfi1, Spdef, or Klf4 expression was shown to result in loss of mature goblet cells (Gregorieff et al., 2009; Katz et al., 2002; Shroyer et al., 2005). These data highlight an unexpected role for IL-18 signaling in directly modulating goblet cell maturation and fate downstream to secretory lineage commitment during intestinal inflammation.

In conclusion, we propose that a strict equilibrium of epithelial IL-18 signaling must be maintained in order to preserve its homeostatic functions while preventing progressive loss of mature goblet cells and increased risk of colitis. This work suggests that IL-18 targeting or specific targeting of epithelial IL-18R may

(D) Epifluorescence images (left, top right panels) and confocal stacks (bottom right panels) of distal colon sections obtained on day 5 post DSS and stained with UEA-1 and anti-MUC2. Scale bar, 150 μ m.

(E) Enumeration of immature and mature goblet cells in distal colon sections stained as in (D). Immature and mature goblet cells were scored as in Figure 5C.

(F) Distal colon samples were obtained on day 8 post DSS and used for gene expression analysis by qPCR. Data are represented as mean \pm SEM. * $p < 0.05$; ** $p < 0.01$; **** $p < 0.0001$ by unpaired Student's t test.

See also Figure S4.

represent a novel strategy to prevent the pathologic breakdown of the mucosal barrier in human ulcerative colitis.

EXPERIMENTAL PROCEDURES

Mice

Conditional knockout mice with floxed IL-18 alleles or floxed IL-18R alleles were generated using standard homologous gene recombination techniques (Rongvaux et al., 2014). *Il18*^{-/-} mice have been previously reported (Takeda et al., 1998). *Il18bp* deletion was generated using Cas9/CrispR technology as described (Wang et al., 2013). All experiments were performed using co-housed littermate control mice. For transmissible dysbiosis cohousing experiments, littermate *Il18*^{fl/fl} and *Il18*^{Δ/EC} mice and age and gender matched *Il18*^{-/-} mice were cohoused at ratios of 1:1:1 for 8 weeks. All animal experimentation was performed in compliance with Yale Institutional Animal Care and Use Committee protocols.

Experimental Colitis

For acute experimental colitis induction, mice were administered 2% DSS (M.W. = 36,000–50,000 Da; MP Biomedicals) in their drinking water ad libitum for 7 days, followed by regular drinking water. According to the animal protocol, mice were sacrificed if they lost more than 30% of their initial body weight.

Colonoscopy

Colonoscopy was performed using a high resolution mouse video endoscopic system ("Coloview," Carl Storz). The severity of colitis was blindly scored using MEICS (Murine Endoscopic Index of Colitis Severity) based on four parameters: granularity of mucosal surface, vascular pattern, translucency of the colon mucosa, and stool consistency (Becker et al., 2006).

Histology

Colons were fixed in Bouin's medium and embedded in paraffin. Blocks were serially sectioned along the cephalocaudal axis of the gut to the level of the lumen; the next 5 mm-thick section was stained with H&E. For goblet cell and mucus layer preservation *ex vivo*, immediately after excision, colons were submerged in Ethanol-Carnoy's fixative at 4°C for 2 hr and then placed into 100% ethanol. Fixed colon tissues were embedded in paraffin and cut into 5-μm sections. Tissues were stained with Alcian blue/PAS. Images were acquired with Leica DMI6000B inverted microscope, and data were analyzed using the LAS-AF software.

SUPPLEMENTAL INFORMATION

Supplemental Information includes Supplemental Experimental Procedures and four figures and can be found with this article online at <http://dx.doi.org/10.1016/j.cell.2015.10.072>.

AUTHOR CONTRIBUTIONS

R.N. and R.J. conceived, performed, and analyzed the experiments and wrote the manuscript. N.G., M.R.d.Z., N.W.P., W.B., J.S.L., C.C.D.H., M.G., and E.E. provided technical assistance in various experiments. R.A.F. supervised the project and participated in interpreting the results and writing the manuscript.

ACKNOWLEDGMENTS

We would like to thank Caroline Lieber, Judith Stein, Jon Alderman, Cindy Hughes, and Elizabeth Hughes-Picard for technical assistance. N.G. is supported by the Dr. Keith Landesman Memorial Fellowship of the Cancer Research Institute, M.R.d.Z. is supported by a Rubicon Fellowship from the Netherlands Organization of Scientific Research, N.W.P. and W.B. are supported by the Cancer Research Institute Irvington Fellowship Program, C.C.D.H. is supported by a Howard Hughes Medical Institute International Student Research Fellowship, and R.N. is supported by the Jane Coffin Childs Fund Postdoctoral Fellowship. This work was supported in part by the Howard Hughes Medical Institute and the Blavatnik Family Foundation (R.A.F.).

Received: June 30, 2015

Revised: September 3, 2015

Accepted: October 23, 2015

Published: December 3, 2015

REFERENCES

- Adachi, O., Kawai, T., Takeda, K., Matsumoto, M., Tsutsui, H., Sakagami, M., Nakanishi, K., and Akira, S. (1998). Targeted disruption of the MyD88 gene results in loss of IL-1- and IL-18-mediated function. *Immunity* 9, 143–150.
- Adolph, T.E., Tomczak, M.F., Niederreiter, L., Ko, H.J., Böck, J., Martinez-Naves, E., Glickman, J.N., Tschurtschenthaler, M., Hartwig, J., Hosomi, S., et al. (2013). Paneth cells as a site of origin for intestinal inflammation. *Nature* 503, 272–276.
- Asquith, M., and Powrie, F. (2010). An innately dangerous balancing act: intestinal homeostasis, inflammation, and colitis-associated cancer. *J. Exp. Med.* 207, 1573–1577.
- Bamias, G., Corridoni, D., Pizarro, T.T., and Cominelli, F. (2012). New insights into the dichotomous role of innate cytokines in gut homeostasis and inflammation. *Cytokine* 59, 451–459.
- Bauernfeind, F.G., Horvath, G., Stutz, A., Alnemri, E.S., MacDonald, K., Speert, D., Fernandes-Alnemri, T., Wu, J., Monks, B.G., Fitzgerald, K.A., et al. (2009). Cutting edge: NF-κappaB activating pattern recognition and cytokine receptors license NLRP3 inflammasome activation by regulating NLRP3 expression. *J. Immunol.* 183, 787–791.
- Becker, C., Fantini, M.C., and Neurath, M.F. (2006). High resolution colonoscopy in live mice. *Nat. Protoc.* 1, 2900–2904.
- Bergstrom, K.S.B., Guttman, J.A., Rumi, M., Ma, C., Bouzari, S., Khan, M.A., Gibson, D.L., Vogl, A.W., and Vallance, B.A. (2008). Modulation of intestinal goblet cell function during infection by an attaching and effacing bacterial pathogen. *Infect. Immun.* 76, 796–811.
- Bersudsky, M., Luski, L., Fishman, D., White, R.M., Ziv-Sokolovskaya, N., Dotan, S., Rider, P., Kaplanov, I., Aychek, T., Dinarello, C.A., et al. (2014). Non-redundant properties of IL-1α and IL-1β during acute colon inflammation in mice. *Gut* 63, 598–609.
- Born, T.L., Thomassen, E., Bird, T.A., and Sims, J.E. (1998). Cloning of a novel receptor subunit, AcPL, required for interleukin-18 signaling. *J. Biol. Chem.* 273, 29445–29450.
- Cadwell, K., Liu, J.Y., Brown, S.L., Miyoshi, H., Loh, J., Lennerz, J.K., Kishi, C., Kc, W., Carrero, J.A., Hunt, S., et al. (2008). A key role for autophagy and the autophagy gene Atg16l1 in mouse and human intestinal Paneth cells. *Nature* 456, 259–263.
- Danese, S., and Fiocchi, C. (2011). Ulcerative colitis. *N. Engl. J. Med.* 365, 1713–1725.
- Dinarello, C.A., Novick, D., Kim, S., and Kaplanski, G. (2013). Interleukin-18 and IL-18 binding protein. *Front. Immunol.* 4, 289.
- Dupaul-Chicoine, J., Yeretssian, G., Doiron, K., Bergstrom, K.S., McIntire, C.R., LeBlanc, P.M., Meunier, C., Turbide, C., Gros, P., Beauchemin, N., et al. (2010). Control of intestinal homeostasis, colitis, and colitis-associated colorectal cancer by the inflammatory caspases. *Immunity* 32, 367–378.
- Elinav, E., Strowig, T., Kau, A.L., Henao-Mejia, J., Thaiss, C.A., Booth, C.J., Peaper, D.R., Bertin, J., Eisenbarth, S.C., Gordon, J.I., and Flavell, R.A. (2011). NLRP6 inflammasome regulates colonic microbial ecology and risk for colitis. *Cell* 145, 745–757.
- Fantuzzi, G., Reed, D.A., and Dinarello, C.A. (1999). IL-12-induced IFN-γ is dependent on caspase-1 processing of the IL-18 precursor. *J. Clin. Invest.* 104, 761–767.
- Fantuzzi, G., Banda, N.K., Guthridge, C., Vondracek, A., Kim, S.-H., Siegmund, B., Azam, T., Sennello, J.A., Dinarello, C.A., and Arend, W.P. (2003). Generation and characterization of mice transgenic for human IL-18-binding protein isoform a. *J. Leukoc. Biol.* 74, 889–896.

- Fellermann, K., Wehkamp, J., Herrlinger, K.R., and Stange, E.F. (2003). Crohn's disease: a defensin deficiency syndrome? *Eur. J. Gastroenterol. Hepatol.* *15*, 627–634.
- Fre, S., Huyghe, M., Mourikis, P., Robine, S., Louvard, D., and Artavanis-Tsakonas, S. (2005). Notch signals control the fate of immature progenitor cells in the intestine. *Nature* *435*, 964–968.
- Gagliani, N., Palm, N.W., de Zoete, M.R., and Flavell, R.A. (2014). Inflammasomes and intestinal homeostasis: regulating and connecting infection, inflammation and the microbiota. *Int. Immunol.* *26*, 495–499.
- Gersemann, M., Becker, S., Kübler, I., Koslowski, M., Wang, G., Herrlinger, K.R., Griger, J., Fritz, P., Fellermann, K., Schwab, M., et al. (2009). Differences in goblet cell differentiation between Crohn's disease and ulcerative colitis. *Differentiation* *77*, 84–94.
- Gregorieff, A., Stange, D.E., Kujala, P., Begthel, H., van den Born, M., Korving, J., Peters, P.J., and Clevers, H. (2009). The ets-domain transcription factor Spdef promotes maturation of goblet and paneth cells in the intestinal epithelium. *Gastroenterology* *137*, 1333–1345, e1–3.
- Henao-Mejia, J., Elinav, E., Jin, C., Hao, L., Mehal, W.Z., Strowig, T., Thaiss, C.A., Kau, A.L., Eisenbarth, S.C., Jurczak, M.J., et al. (2012). Inflammation-mediated dysbiosis regulates progression of NAFLD and obesity. *Nature* *482*, 179–185.
- Hoshino, K., Tsutsui, H., Kawai, T., Takeda, K., Nakanishi, K., Takeda, Y., and Akira, S. (1999). Cutting edge: generation of IL-18 receptor-deficient mice: evidence for IL-1 receptor-related protein as an essential IL-18 binding receptor. *J. Immunol.* *162*, 5041–5044.
- Hugot, J.P., Chamaillard, M., Zouali, H., Lesage, S., Cézard, J.P., Belaiche, J., Almer, S., Tysk, C., O'Morain, C.A., Gassull, M., et al. (2001). Association of NOD2 leucine-rich repeat variants with susceptibility to Crohn's disease. *Nature* *411*, 599–603.
- Jensen, J., Pedersen, E.E., Galante, P., Hald, J., Heller, R.S., Ishibashi, M., Kageyama, R., Guillemot, F., Serup, P., and Madsen, O.D. (2000). Control of endodermal endocrine development by Hes-1. *Nat. Genet.* *24*, 36–44.
- Kanai, T., Watanabe, M., Okazawa, A., Sato, T., Yamazaki, M., Okamoto, S., Ishii, H., Totsuka, T., Iiyama, R., Okamoto, R., et al. (2001). Macrophage-derived IL-18-mediated intestinal inflammation in the murine model of Crohn's disease. *Gastroenterology* *121*, 875–888.
- Kaser, A., Zeissig, S., and Blumberg, R.S. (2010). Inflammatory bowel disease. *Annu. Rev. Immunol.* *28*, 573–621.
- Kaser, A., Niederreiter, L., and Blumberg, R.S. (2011). Genetically determined epithelial dysfunction and its consequences for microflora-host interactions. *Cell. Mol. Life Sci.* *68*, 3643–3649.
- Katz, J.P., Perreault, N., Goldstein, B.G., Lee, C.S., Labosky, P.A., Yang, V.W., and Kaestner, K.H. (2002). The zinc-finger transcription factor Klf4 is required for terminal differentiation of goblet cells in the colon. *Development* *129*, 2619–2628.
- Levy, M., Thaiss, C.A., Zeevi, D., Dohnalová, L., Zilberman-Schapira, G., Ali Mahdi, J., David, E., Savidor, A., Korem, T., Herzig, Y., et al. (2015). Microbiota-modulated metabolites shape the intestinal microenvironment by regulating NLRP6 inflammasome signaling. *Cell* *163*, this issue, 1428–1443.
- Lopetuso, L.R., Chowdhry, S., and Pizarro, T.T. (2013). Opposing functions of classic and novel IL-1 family members in gut health and disease. *Front. Immunol.* *4*, 181.
- Maloy, K.J., and Powrie, F. (2011). Intestinal homeostasis and its breakdown in inflammatory bowel disease. *Nature* *474*, 298–306.
- Malvin, N.P., Seno, H., and Stappenbeck, T.S. (2012). Colonic epithelial response to injury requires Myd88 signaling in myeloid cells. *Mucosal Immunol.* *5*, 194–206.
- Martinon, F., Burns, K., and Tschopp, J. (2002). The inflammasome: a molecular platform triggering activation of inflammatory caspases and processing of proIL- β . *Mol. Cell* *10*, 417–426.
- McCauley, H.A., and Guasch, G. (2015). Three cheers for the goblet cell: maintaining homeostasis in mucosal epithelia. *Trends Mol. Med.* *21*, 492–503.
- McCormick, D.A., Horton, L.W., and Mee, A.S. (1990). Mucin depletion in inflammatory bowel disease. *J. Clin. Pathol.* *43*, 143–146.
- Medzhitov, R., Preston-Hurlburt, P., Kopp, E., Stadlen, A., Chen, C., Ghosh, S., and Janeway, C.A., Jr. (1998). MyD88 is an adaptor protein in the hToll/IL-1 receptor family signaling pathways. *Mol. Cell* *2*, 253–258.
- Mercurio, F., Zhu, H., Murray, B.W., Shevchenko, A., Bennett, B.L., Li, J., Young, D.B., Barbosa, M., Mann, M., Manning, A., and Rao, A. (1997). IKK-1 and IKK-2: cytokine-activated I κ B kinases essential for NF- κ B activation. *Science* *278*, 860–866.
- Miao, E.A., Leaf, I.A., Treuting, P.M., Mao, D.P., Dors, M., Sarkar, A., Warren, S.E., Wewers, M.D., and Aderem, A. (2010). Caspase-1-induced pyroptosis is an innate immune effector mechanism against intracellular bacteria. *Nat. Immunol.* *11*, 1136–1142.
- Monteleone, G., Trapasso, F., Parrello, T., Biancone, L., Stella, A., Iuliano, R., Luzza, F., Fusco, A., and Pallone, F. (1999). Bioactive IL-18 expression is up-regulated in Crohn's disease. *J. Immunol.* *163*, 143–147.
- Nakamura, K., Okamura, H., Wada, M., Nagata, K., and Tamura, T. (1989). Endotoxin-induced serum factor that stimulates gamma interferon production. *Infect. Immun.* *57*, 590–595.
- Netea, M.G., Joosten, L.A.B., Lewis, E., Jensen, D.R., Voshol, P.J., Kullberg, B.J., Tack, C.J., van Krieken, H., Kim, S.-H., Stalenhoef, A.F., et al. (2006). Deficiency of interleukin-18 in mice leads to hyperphagia, obesity and insulin resistance. *Nat. Med.* *12*, 650–656.
- Neurath, M.F. (2014). Cytokines in inflammatory bowel disease. *Nat. Rev. Immunol.* *14*, 329–342.
- Noah, T.K., Kazanjian, A., Whitsett, J., and Shroyer, N.F. (2010). SAM pointed domain ETS factor (SPDEF) regulates terminal differentiation and maturation of intestinal goblet cells. *Exp. Cell Res.* *316*, 452–465.
- Novick, D., Kim, S.H., Fantuzzi, G., Reznikov, L.L., Dinarello, C.A., and Rubinstein, M. (1999). Interleukin-18 binding protein: a novel modulator of the Th1 cytokine response. *Immunity* *10*, 127–136.
- Nuding, S., Fellermann, K., Wehkamp, J., and Stange, E.F. (2007). Reduced mucosal antimicrobial activity in Crohn's disease of the colon. *Gut* *56*, 1240–1247.
- Oficjalska, K., Raverdeau, M., Aviello, G., Wade, S.C., Hickey, A., Sheehan, K.M., Corr, S.C., Kay, E.W., O'Neill, L.A., Mills, K.H., and Creagh, E.M. (2015). Protective role for caspase-11 during acute experimental murine colitis. *J. Immunol.* *194*, 1252–1260.
- Ogura, Y., Bonen, D.K., Inohara, N., Nicolae, D.L., Chen, F.F., Ramos, R., Britton, H., Moran, T., Karaliuskas, R., Duerr, R.H., et al. (2001). A frameshift mutation in NOD2 associated with susceptibility to Crohn's disease. *Nature* *411*, 603–606.
- Pizarro, T.T., Michie, M.H., Bentz, M., Woratanadham, J., Smith, M.F., Jr., Foley, E., Moskaluk, C.A., Bickston, S.J., and Cominelli, F. (1999). IL-18, a novel immunoregulatory cytokine, is up-regulated in Crohn's disease: expression and localization in intestinal mucosal cells. *J. Immunol.* *162*, 6829–6835.
- Pullan, R.D., Thomas, G.A., Rhodes, M., Newcombe, R.G., Williams, G.T., Allen, A., and Rhodes, J. (1994). Thickness of adherent mucus gel on colonic mucosa in humans and its relevance to colitis. *Gut* *35*, 353–359.
- Rongvaux, A., Jackson, R., Harman, C.C., Li, T., West, A.P., de Zoete, M.R., Wu, Y., Yordy, B., Lakhani, S.A., Kuan, C.Y., et al. (2014). Apoptotic caspases prevent the induction of type I interferons by mitochondrial DNA. *Cell* *159*, 1563–1577.
- Salcedo, R., Worschech, A., Cardone, M., Jones, Y., Gyulai, Z., Dai, R.-M., Wang, E., Ma, W., Haines, D., O'Huigin, C., et al. (2010). MyD88-mediated signaling prevents development of adenocarcinomas of the colon: role of interleukin 18. *J. Exp. Med.* *207*, 1625–1636.
- Saleh, M., and Trinchieri, G. (2011). Innate immune mechanisms of colitis and colitis-associated colorectal cancer. *Nat. Rev. Immunol.* *11*, 9–20.
- Schreiber, S., Rosenstiel, P., Albrecht, M., Hampe, J., and Krawczak, M. (2005). Genetics of Crohn disease, an archetypal inflammatory barrier disease. *Nat. Rev. Genet.* *6*, 376–388.

- Shroyer, N.F., Wallis, D., Venken, K.J., Bellen, H.J., and Zoghbi, H.Y. (2005). Gfi1 functions downstream of Math1 to control intestinal secretory cell subtype allocation and differentiation. *Genes Dev.* *19*, 2412–2417.
- Siegmund, B. (2010). Interleukin-18 in intestinal inflammation: friend and foe? *Immunity* *32*, 300–302.
- Siegmund, B., Fantuzzi, G., Rieder, F., Gamboni-Robertson, F., Lehr, H.A., Hartmann, G., Dinarello, C.A., Endres, S., and Eigler, A. (2001). Neutralization of interleukin-18 reduces severity in murine colitis and intestinal IFN-gamma and TNF-alpha production. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* *281*, R1264–R1273.
- Sivakumar, P.V., Westrich, G.M., Kanaly, S., Garka, K., Born, T.L., Derry, J.M., and Viney, J.L. (2002). Interleukin 18 is a primary mediator of the inflammation associated with dextran sulphate sodium induced colitis: blocking interleukin 18 attenuates intestinal damage. *Gut* *50*, 812–820.
- Strugala, V., Dettmar, P.W., and Pearson, J.P. (2008). Thickness and continuity of the adherent colonic mucus barrier in active and quiescent ulcerative colitis and Crohn's disease. *Int. J. Clin. Pract.* *62*, 762–769.
- Takagi, H., Kanai, T., Okazawa, A., Kishi, Y., Sato, T., Takaishi, H., Inoue, N., Ogata, H., Iwao, Y., Hoshino, K., et al. (2003). Contrasting action of IL-12 and IL-18 in the development of dextran sodium sulphate colitis in mice. *Scand. J. Gastroenterol.* *38*, 837–844.
- Takeda, K., Tsutsui, H., Yoshimoto, T., Adachi, O., Yoshida, N., Kishimoto, T., Okamura, H., Nakanishi, K., and Akira, S. (1998). Defective NK cell activity and Th1 response in IL-18-deficient mice. *Immunity* *8*, 383–390.
- Takeuchi, M., Nishizaki, Y., Sano, O., Ohta, T., Ikeda, M., and Kurimoto, M. (1997). Immunohistochemical and immuno-electron-microscopic detection of interferon-gamma-inducing factor ("interleukin-18") in mouse intestinal epithelial cells. *Cell Tissue Res.* *289*, 499–503.
- Ten Hove, T., Corbaz, A., Amitai, H., Aloni, S., Belzer, I., Graber, P., Drillenburger, P., van Deventer, S.J., Chvatchko, Y., and Te Velde, A.A. (2001). Blockade of endogenous IL-18 ameliorates TNBS-induced colitis by decreasing local TNF-alpha production in mice. *Gastroenterology* *121*, 1372–1379.
- Trabucchi, E., Mukenge, S., Baratti, C., Colombo, R., Fregoni, F., and Montorsi, W. (1986). Differential diagnosis of Crohn's disease of the colon from ulcerative colitis: ultrastructure study with the scanning electron microscope. *Int. J. Tissue React.* *8*, 79–84.
- van Es, J.H., van Gijn, M.E., Riccio, O., van den Born, M., Vooijs, M., Begthel, H., Cozijnsen, M., Robine, S., Winton, D.J., Radtke, F., and Clevers, H. (2005). Notch/gamma-secretase inhibition turns proliferative cells in intestinal crypts and adenomas into goblet cells. *Nature* *435*, 959–963.
- Wang, H., Yang, H., Shivalila, C.S., Dawlaty, M.M., Cheng, A.W., Zhang, F., and Jaenisch, R. (2013). One-step generation of mice carrying mutations in multiple genes by CRISPR/Cas-mediated genome engineering. *Cell* *153*, 910–918.
- Wehkamp, J., Schmid, M., and Stange, E.F. (2007). Defensins and other antimicrobial peptides in inflammatory bowel disease. *Curr. Opin. Gastroenterol.* *23*, 370–378.
- Xavier, R.J., and Podolsky, D.K. (2007). Unravelling the pathogenesis of inflammatory bowel disease. *Nature* *448*, 427–434.
- Yang, Q., Bermingham, N.A., Finegold, M.J., and Zoghbi, H.Y. (2001). Requirement of Math1 for secretory cell lineage commitment in the mouse intestine. *Science* *294*, 2155–2158.
- Zaki, M.H., Boyd, K.L., Vogel, P., Kastan, M.B., Lamkanfi, M., and Kanneganti, T.D. (2010). The NLRP3 inflammasome protects against loss of epithelial integrity and mortality during experimental colitis. *Immunity* *32*, 379–391.