

Some Characterizations of the Multivariate t Distribution*

PI-ERH LIN

*Department of Statistics, Florida State University, Tallahassee, Florida 32306**Communicated by E. Lukacs*

A multivariate t vector \mathbf{X} is represented in two different forms, one associated with a normal vector and an independent chi-squared variable, and the other with a normal vector and an independent Wishart matrix. We show that \mathbf{X} is multivariate t with mean $\boldsymbol{\mu}$, covariance matrix $\nu(\nu - 2)^{-1}\Sigma$, $\nu > 2$ and degrees of freedom ν if and only if for any $\mathbf{a} \neq \mathbf{0}$, $(\mathbf{a}'\Sigma\mathbf{a})^{-1/2} \mathbf{a}'(\mathbf{X} - \boldsymbol{\mu})$ has the Student's t distribution with ν degrees of freedom under both representations. Some other characterizations are also obtained.

1. INTRODUCTION AND SUMMARY

The multivariate t distribution was first derived independently by Cornish [4] and Dunnett and Sobel [8]. It arises in multiple decision problems concerned with the selection and ranking of population means of several normal populations having a common unknown variance (see Bechhofer, Dunnett and Sobel [3]). It can be used in setting up simultaneous confidence bounds for the means of correlated normal variables, for parameters in a linear model and for future observations from a multivariate normal distribution (see John [11]). The multivariate t distribution also appears in the Bayesian multivariate analysis of variance and regression, treated by Tiao and Zellner [16], Geisser and Cornfield [9], Raiffa and Schlaifer [13], and Ando and Kaufmann [2], where the normal-Wishart distribution is considered to be the conjugate prior distribution (in the sense of Raiffa and Schlaifer) of the mean vector and covariance matrix

Received March 4, 1971; revised March 8, 1972.

AMS 1970 subject classification: Primary 62H05; Secondary 62H10.

Key words: Symmetric square root of a positive definite matrix; conditionally independent; conditional marginal probability density; almost surely with respect to a σ -finite measure.

* Research supported in part by the Florida State University Grant No. 011204-043 and in part by the National Institute of General Medical Sciences through Training Grant 5T1 GM-913.

of a multivariate normal distribution. However, the representation of the multivariate t vector used in the Bayesian multivariate analysis is somewhat different from that employed by Cornish [4], and Dunnett and Sobel [8].

The multivariate t can be considered as a generalization of the univariate t . Siddiqui [14] generalizes the univariate t to a bivariate t in a different direction and Dickey [7] generalizes the multivariate t to a matricivariate t . The probability integrals and the percentage points of a multivariate t have been studied by Gupta [10] and Krishnaiah and Armitage [12].

In Section 2 of this note we consider the distribution of a multivariate t vector which may be represented in two different forms, one associated with a normal vector and an independent chi-squared variable, and the other with a normal vector and an independent Wishart matrix. In Section 3, the characterization properties of a t vector are developed through multivariate normal theory.

As usual, we denote by $N(\boldsymbol{\mu}, \Sigma)$ a multivariate normal distribution with mean $\boldsymbol{\mu}$ and covariance matrix Σ ; by $W(\Sigma, n)$ a Wishart distribution with parameters n and Σ ; by χ_ν^2 a chi-squared distribution with ν degrees of freedom; and by $F'_{p,\nu}(\tau^2)$ a noncentral F distribution with p and ν degrees of freedom and non-centrality τ^2 . We write $F_{p,\nu}$ for $F'_{p,\nu}(0)$.

2. THE MULTIVARIATE t DISTRIBUTION

A p -variate random vector $\mathbf{X} = (X_1, \dots, X_p)'$ is said to have a (nonsingular) multivariate t distribution with mean vector $\boldsymbol{\mu} = (\mu_1, \dots, \mu_p)'$ and covariance matrix $\nu(\nu - 2)^{-1}\Sigma$, $\nu > 2$, denoted by $T_\nu(\boldsymbol{\mu}, \Sigma, p)$, if it has the probability density function (pdf) given by

$$f(\mathbf{x}) = \frac{\Gamma[\frac{1}{2}(\nu + p)]}{(\pi\nu)^{\frac{1}{2}\nu} \Gamma(\frac{1}{2}\nu) |\Sigma|^{\frac{1}{2}}} \left\{ 1 + \frac{(\mathbf{x} - \boldsymbol{\mu})' \Sigma^{-1} (\mathbf{x} - \boldsymbol{\mu})}{\nu} \right\}^{-\frac{1}{2}(\nu + p)}, \quad \nu > 2. \quad (1)$$

It is noted that $T_\nu(0, 1, 1) = t_\nu$ is the Student's t distribution with ν degrees of freedom.

Unless otherwise mentioned, it is assumed that \mathbf{X} , $\boldsymbol{\mu}$, \mathbf{a} , and $\mathbf{0}$ are $p \times 1$ vectors, V and Σ are $p \times p$ positive definite symmetric matrices, and I is the identity matrix of order p .

For the purpose of establishing the characterizations and the distribution theory in Section 3 it is convenient to represent a multivariate t vector in the following forms:

Representation A. Let $\mathbf{X} \sim T_\nu(\boldsymbol{\mu}, \Sigma, p)$. Then \mathbf{X} may be written as

$$\mathbf{X} = S^{-1}\mathbf{Y} + \boldsymbol{\mu}, \quad (2)$$

where $\mathbf{Y} \sim N(\mathbf{0}, \Sigma)$ and $\nu S^2 \sim \chi_\nu^2$, independent of \mathbf{Y} . This implies that $\mathbf{X} | S^2 = s^2 \sim N(\boldsymbol{\mu}, s^{-2}\Sigma)$, where $\nu S^2 \sim \chi_\nu^2$. The Student's t random variable is then represented as $X | S^2 = s^2 \sim N(0, s^{-2})$, where $\nu S^2 \sim \chi_\nu^2$.

Representation B. Let $\mathbf{X} \sim T_\nu(\boldsymbol{\mu}, \Sigma, p)$. Then \mathbf{X} may be written as

$$\mathbf{X} = (V^{1/2})^{-1}\mathbf{Y} + \boldsymbol{\mu}, \tag{3}$$

where $V^{1/2}$ is the symmetric square root of V , i.e.,

$$V^{1/2}V^{1/2} = V \sim W(\Sigma^{-1}, \nu + p - 1)$$

and $\mathbf{Y} \sim N(\mathbf{0}, \nu I)$, independent of V . This implies that $\mathbf{X} | V \sim N(\boldsymbol{\mu}, \nu V^{-1})$, where $V \sim W(\Sigma^{-1}, \nu + p - 1)$. The Student's t random variable is then represented as $X | V \sim N(0, \nu \mathbf{a}' V^{-1} \mathbf{a} / \mathbf{a}' \Sigma \mathbf{a})$ for any $\mathbf{a} \neq \mathbf{0}$, where

$$V \sim W(\Sigma^{-1}, \nu + p - 1).$$

It can be shown that the pdf of \mathbf{X} , under both Representations A and B, is given by (1).

3. CHARACTERIZATIONS OF THE MULTIVARIATE t

Cornish [6] gives a characterization of the multivariate t using Representation A. His proof is lengthy and complicated. We present here the same result under Representation B and also give a simpler proof under Representation A, utilizing multivariate normal distribution theory. Two other characterizations are also obtained under Representation A.

THEOREM 1. $\mathbf{X} \sim T_\nu(\boldsymbol{\mu}, \Sigma, p)$ if and only if, for any $\mathbf{a} \neq \mathbf{0}$,

$$(\mathbf{a}' \Sigma \mathbf{a})^{-1/2} \mathbf{a}'(\mathbf{X} - \boldsymbol{\mu}) \sim t_\nu.$$

Proof. (i) Assume Representation A.

$$\begin{aligned} & \mathbf{X} | S^2 = s^2 \sim N(\boldsymbol{\mu}, s^{-2}\Sigma), \\ \Leftrightarrow & (\mathbf{a}' \Sigma \mathbf{a})^{-1/2} \mathbf{a}'(\mathbf{X} - \boldsymbol{\mu}) | S^2 = s^2 \sim N(0, s^{-2}), \text{ for any } \mathbf{a} \neq \mathbf{0} \\ \Leftrightarrow & (\mathbf{a}' \Sigma \mathbf{a})^{-1/2} \mathbf{a}'(\mathbf{X} - \boldsymbol{\mu}) \sim t_\nu. \end{aligned}$$

(ii) Assume Representation B.

$$\begin{aligned} & \mathbf{X} | V \sim N(\boldsymbol{\mu}, \nu V^{-1}) \\ \Leftrightarrow & (\mathbf{a}' \Sigma \mathbf{a})^{-1/2} \mathbf{a}'(\mathbf{X} - \boldsymbol{\mu}) | V \sim N[0, \nu(\mathbf{a}' V^{-1} \mathbf{a}) / (\mathbf{a}' \Sigma \mathbf{a})], \text{ for any } \mathbf{a} \neq \mathbf{0} \\ \Leftrightarrow & (\mathbf{a}' \Sigma \mathbf{a})^{-1/2} \mathbf{a}'(\mathbf{X} - \boldsymbol{\mu}) \sim t_\nu, \end{aligned}$$

since $\mathbf{a}'\Sigma\mathbf{a}/\mathbf{a}'V^{-1}\mathbf{a} \sim \chi_\nu^2$. (See, for example, Stein [15, pp. 32-33].) This completes the proof of the theorem.

In order to establish some further characterizations, we need the following lemmas. The proofs are straightforward and are omitted.

LEMMA 1. Let $\mathbf{X} \sim T_\nu(\boldsymbol{\mu}, \Sigma, p)$, then for any nonsingular $p \times p$ scalar matrix C and any \mathbf{a} , $C\mathbf{X} + \mathbf{a} \sim T_\nu(C\boldsymbol{\mu} + \mathbf{a}, C\Sigma C', p)$.

LEMMA 2. Let $\mathbf{X} \sim T_\nu(\boldsymbol{\mu}, \Sigma, p)$, then $\mathbf{X}'\Sigma^{-1}\mathbf{X}/p \sim F'_{p,\nu}(\boldsymbol{\mu}'\Sigma^{-1}\boldsymbol{\mu}/p)$. When $\boldsymbol{\mu} = \mathbf{0}$, the distribution is central F .

LEMMA 3. Let Y be a symmetrically distributed random variable. Then $Y^2 \sim \chi_1^2$ if, and only if $Y \sim N(0, 1)$.

LEMMA 4. Let Y_1, \dots, Y_p be independent random variables with finite variances. Assume that Y_k , ($k = 1, \dots, p$), has pdf $h_k(y)$ such that $h_k(y) > 0$ and differentiable for all y , $-\infty < y < \infty$. Then the joint pdf of Y_1, \dots, Y_p is a function only of $y_1^2 + \dots + y_p^2$ if, and only if $(Y_1, \dots, Y_p)' \sim N(\mathbf{0}, \sigma^2 I)$, for some $\sigma^2 < \infty$.

For the remainder of this section, we will consider a multivariate t vector using Representation A.

THEOREM 2. Let $\nu S^2 \sim \chi_\nu^2$ and let X_1, \dots, X_p be continuous random variables. Assume that X_1, \dots, X_p are conditionally independent, and that X_k is symmetrically distributed with mean μ_k and finite variance, given $S^2 = s^2$. Then

$$\sum_{k=1}^p (X_k - \mu_k)^2 / (p\sigma_k^2) \sim F_{p,\nu}, \quad (4)$$

if and only if $(X_1, \dots, X_p)' \sim T_\nu(\boldsymbol{\mu}, D, p)$, where $\boldsymbol{\mu} = (\mu_1, \dots, \mu_p)'$ and D is a $p \times p$ diagonal matrix with its k -th diagonal element equal to σ_k^2 .

Proof. The sufficiency is a special case of Lemma 2. We shall show the necessity. Let $Y_k = (X_k - \mu_k)/\sigma_k$, $k = 1, \dots, p$. Then the Y 's also satisfy the conditions of the theorem with mean 0 and finite variance, given $S^2 = s^2$. It is well known that an $F_{p,\nu}$ -distributed random variable may be represented as the ratio of two independent chi-squared random variables with p and ν degrees of freedom, respectively, multiplied by a suitable constant. Thus, letting $S_1^2 \sim \chi_p^2$ independent of S^2 , we have

$$\sum_{k=1}^p Y_k^2 \sim S_1^2/S^2, \quad (5)$$

or

$$\sum_{k=1}^p (SY_k)^2 | S^2 \sim S_1^2 | S^2 \sim \chi_p^2, \tag{6}$$

which is independent of S^2 . This implies that

$$(SY_k)^2 | S^2 \sim \chi_1^2, \quad k = 1, \dots, p, \tag{7}$$

and by Lemma 3, we have $SY | S^2 = s^2 \sim N(\mathbf{0}, I)$, where $\mathbf{Y} = (Y_1, \dots, Y_p)'$. Thus, by Representation A, $\mathbf{Y} \sim T,(\mathbf{0}, I, p)$ and by Lemma 1, $\mathbf{X} \sim T,(\boldsymbol{\mu}, D, p)$.

THEOREM 3. *Let $\nu S^2 \sim \chi_p^2$ and let $X_k, k = 1, \dots, p$, be conditionally independent random variables, given $S^2 = s^2$, with $\text{Var}(X_k | s^2) = \sigma^2/s^2, \sigma^2 < \infty$. Assume that $X_k (k = 1, \dots, p)$, given $S^2 = s^2$, has conditional pdf $f_k(x | s^2)$ which is positive and differentiable for all $x, -\infty < x < \infty$. Then the joint pdf of X_1, \dots, X_p is a function only of $x_1^2 + \dots + x_p^2$ if, and only if $(X_1, \dots, X_p)' \sim T,(\mathbf{0}, \sigma^2 I, p)$.*

Proof. The sufficiency is easily seen from the joint pdf of X_1, \dots, X_p . We shall prove the necessity. Since the joint pdf of X_1, \dots, X_p is a function of $x_1^2 + \dots + x_p^2$, it follows that the conditional joint pdf of X_1, \dots, X_p , given $S^2 = s^2$, is a function of $x_1^2 + \dots + x_p^2$ and s^2 almost surely with respect to a σ -finite measure m , i.e.,

$$\prod_{k=1}^p f_k(x_k | s^2) = g_0 \left(\sum_{k=1}^p x_k^2, s^2 \right), \quad \text{a.s. } (m). \tag{8}$$

Similarly, the conditional marginal pdf of $X_k (k = 1, \dots, p)$, given $S^2 = s^2$, is

$$f_k(x_k | s^2) = g_k(x_k^2, s^2), \quad \text{a.s. } (m), \tag{9}$$

where $g_k(t, s^2) (k = 0, \dots, p)$ is a positive and differentiable function of $t \geq 0$. Now, the result of Lemma 4 and the fact that $\text{Var}(X_k | s^2) = \sigma^2/s^2$, for all $k = 1, \dots, p$, show that $(X_1, \dots, X_p)' | S^2 = s^2 \sim N(\mathbf{0}, s^{-2}\sigma^2 I)$. Therefore, $(X_1, \dots, X_p)' \sim T,(\mathbf{0}, \sigma^2 I, p)$.

REFERENCES

[1] ANDERSON, T. W. (1958). *An Introduction to Multivariate Statistical Analysis*. Wiley, New York.
 [2] ANDO, A. AND KAUFMANN, G. W. (1965). Bayesian Analysis of the independent multinormal process—neither mean nor precision known. *J. Amer. Statist. Assoc.* **60** 347–358.

- [3] BECHHOFFER, R. E., DUNNETT, C. W., AND SOBEL, M. (1954). A two-sample multiple decision procedure for ranking means of normal populations with a common unknown variance. *Biometrika* **41** 170–176.
- [4] CORNISH, E. A. (1954a). The multivariate t -distribution associated with a set of normal sample deviates. *Austral. J. Phys.* **7** 531–542.
- [5] CORNISH, E. A. (1954b). The sampling distributions of statistics derived from the multivariate t -distribution. *Austral. J. Phys.* **8** 193–199.
- [6] CORNISH, E. A. (1962). The multivariate t -distribution associated with the general multivariate normal distribution. CSIRO Tech. paper No. 13. CSIRO Div. Math. Stat., Adelaide.
- [7] DICKEY, J. M. (1967). Matricvariate generalizations of the multivariate t distribution and the inverted multivariate t distribution. *Ann. Math. Stat.* **38** 511–518.
- [8] DUNNETT, C. W. AND SOBEL, M. (1954). A bivariate generalization of Student t -distribution with tables for certain special cases. *Biometrika* **41** 153–169.
- [9] GEISSER, S. AND CORNFIELD, J. (1963). Posterior distributions for multivariate normal parameters. *J. Roy. Statist. Soc. Ser. B* **25** 368–376.
- [10] GUPTA, S. S. (1963). Probability integrals of multivariate normal and multivariate t . *Ann. Math. Stat.* **34** 792–828.
- [11] JOHN, S. (1961). On the evaluation of the probability integral of the multivariate t -distribution. *Biometrika* **48** 409–417.
- [12] KRISHNAIAH, P. R. AND ARMITAGE, J. V. (1965). Percentage points of the multivariate t distribution. ARL Tech. Rep. No. 65-199. Aerospace Res. Labs. Wright-Patterson AFB, Ohio.
- [13] RAIFFA, H. AND SCHLAIFER, R. (1961). *Applied Statistical Decision Theory*. Division of Research, Harvard Business School.
- [14] SIDDIQUI, M. M. (1967). A bivariate t distribution. *Ann. Math. Stat.* **38** 162–166.
- [15] STEIN, C. M. (1969). *Multivariate Analysis I*. (Notes prepared by M. L. Eaton). Tech. Rep. No. 42, Dept. of Statistics, Stanford University.
- [16] TIAO, G. C. AND ZELLNER, A. (1964). On the Bayesian estimation of multivariate regression. *J. Roy. Statist. Soc. Ser. B* **26** 277–285.