

chondrogenic phenotype of the MSC was assessed following short-term (1 week) culture.

Results: Addition of rAC to the culture media of rat articular chondrocytes led to elevated expression of Sox-9, collagen IIA1, FGF2, TGF beta-1 and several other chondrogenic markers. Similar results were obtained for cells grown in monolayer cultures or 3-D collagen scaffolds. Elevated Sox-9 and collagen IIA1 expression also was observed following rAC treatment of human chondrocytes from a patient with osteoarthritis. No changes in the total cell numbers were observed despite these positive changes in the chondrogenic phenotype, indicating that the primary effect of rAC was on the differentiation capacity of the cells, rather than on cell survival. We also evaluated the effect of rAC on rat bone marrow grown for 1 week, and found that the number of MSC in these cultures was increased ~2–3-fold as assessed by colony forming units (CFU-F) and flow cytometry, and that their chondrogenic phenotype was positively influenced as well.

Conclusions: Based on these results we suggest that AC is an important enzyme required to maintain the differentiated phenotype of primary chondrocytes, and that supplementation of media with rAC results in improved chondrocyte quality following ex vivo expansion. rAC also improves the yield of bone marrow-derived MSC from short-term cultures, and their chondrogenic phenotype as well. rAC may therefore be an important reagent that can be used to improve the production of chondrocytes for cartilage repair, either ex vivo or through direct administration to damaged cartilage sites.

235

DETAILED EVALUATION OF CHONDRAL DEFECT REPAIR AND AUTOLOGOUS BONE MARROW DERIVED MESENCHYMAL CELLS TRANSPLANTATION. A NONHUMAN PRIMATE MODEL

S. Araki, S. Imai, M. Kubo, T. Mimura, K. Nishizawa, H. Ueba. *Shiga Univ of Med Sci., Otsu, Japan*

Purpose: Articular cartilage injury remains one of the major concerns in orthopaedic surgery. Although the repair process of articular cartilage defects has been studied in many species. The extent and time course of articular cartilage defect healing in humans are not well described. Mesenchymal stem cells (MSCs) are an important cell source for cartilage regeneration. A number of successful results in transplantation MSCs into cartilage defects have been reported in animal models. However, Clinical studies of MSCs transplantation for cartilage repair were very few. Physicians not much know efficacy and safety of MSCs transplantation for cartilage repair in clinical.

Evaluation of the status of articular cartilage repair and MSCs transplantation for articular cartilage repair at different time points in a primate model may provide a healing process, efficacy, safety and postoperative activity recommendations in clinical.

Methods: Full-thickness osteochondral defects were created on the medial femoral condyles (5 mm in diameters and 5 mm in depth) and trochlea (3 mm in diameters and 5 mm in depth) of 18 cynomolgus macaques, and the animals were divided into three groups: MSC, Gel, and Defect. In the MSC group, the defects were filled with bone marrow-derived MSCs embedded in collagen gel. In the Gel group, the defects were filled with collagen gel without cells. In the Defect group, the defects were left empty and additional defect were created on the lateral femoral condyles (2 mm in diameters and 5 mm in depth). All groups were evaluated by gross and histologic examination at 6-, 12- and 24-weeks.

Results: In the defect group, Cartilaginous repair responses failed to occur in the larger 3 mm defects, which was covered only by fibrous scar tissue. In contrast, hyaline-like articular cartilage was regenerated by 24 weeks in 2 mm defects. In the MSC group, the border between bone and cartilage moved upwards, integrations between native cartilage and regenerated tissue were improved. After 24 weeks, histological scores of the MSC group improved and were better than those of two other control groups.

Conclusions: 2 mm diameter full-thickness cartilage defect whose size were critical for primate knees. In the primate animal model, significant improvements in the extent and quality of cartilage repair were observed from the 12- to 24-week time points after transplantation of MSCs.

236

PROLIFERATIVE AND DIFFERENTIATION POTENTIAL OF STEM CELLS DERIVED FROM THREE MESENCHYMAL TISSUES IN LATE STAGE OSTEOARTHRITIC PATIENTS

L. Simion¹, G. Shaw², K. Mashayekhi², P. Botez¹, M. Murphy², F. Barry².
¹Univ. Med. and Pharmacy, Iasi, Romania; ²REMEDI NUI, Galway, Ireland

Purpose: To assess the presence and biological potential of stem cells in three mesenchymal tissues (subchondral bone, synovial layer, periarticular adipose tissue) in late stages osteoarthritic patients, subjects of total knee replacement (TKR)

Methods: Samples were collected from patients undergoing TKR, plastic adherent cells cultured in complete α MEM with FGF β for successive passages: cell morphology and growth potential was recorded. Flow cytometric surface cell markers detection for stemness antibodies was performed. Differentiation assays for three mesenchymal lineages (osteogenesis adipogenesis chondrogenesis) was assessed by qualitative and quantitative method. Time lapse life cell imaging of nondifferentiated cells over 24 hours period was used to determine cell kinetics.

Results: Mesenchymal cells derived from all donors and tissue types displayed morphology and growth potential of MSCs, were positive for stemness related antibodies (CD 105, CD 73, CD 90), underwent differentiation toward three lineages with significant differences between tissue of origin and not between donors. Cell kinetics, recorded for one donor, was different for adipose derived, synovial derived and trabecular bone derived MSCs, a parallel could be made between growth kinetics and recorded cell speed.

Conclusions: Late stage OA derived human tissues (subchondral trabecular bone, extraarticular adipose tissue and synovial layer) contain phenotypically different MSCs which have different growth and differentiation potential which could be related with the pathogeny and progression of OA, further to be investigated; these tissues can be considered as cell source for regenerative therapies.

237

TGF- β TYPE II RECEPTOR AS A NOVEL INDICATOR OF CHONDROGENESIS IN EQUINE BONE MARROW-DERIVED MESENCHYMAL STEM CELLS

K.K. Suter¹, K.A. Payne¹, V. Yao¹, L.A. Fortier², C.R. Chu¹. ¹Univ. of Pittsburgh, Pittsburgh, PA, USA; ²Coll. of Vet. Med., Cornell Univ., Ithaca, NY, USA

Purpose: Articular cartilage lacks the essential components for self-repair, often resulting in inadequate healing of isolated cartilage injury, which can lead to degenerative osteoarthritis (OA). Designing techniques to improve cartilage repair after injury may prevent or delay development of OA. Bone marrow-derived mesenchymal stem cells (MSC) are a promising cell source for articular cartilage regeneration. *In vitro*, MSC have been shown to undergo chondrogenic differentiation in the presence of transforming growth factor β 1 (TGF- β 1), via a signaling pathway known to play a key regulatory role in chondrogenesis. Various knockout studies suggest that the TGF- β type II receptor (T β R-II) is specifically important in skeletal muscle formation and bone morphogenesis, but its role has not been clearly defined in chondrogenesis. The purpose of this study is to test the hypothesis that levels of T β R-II can predict the cellular responsiveness of equine bone marrow-derived MSC to TGF- β 1 and are related to subsequent chondrogenic potential.

Methods: Bone marrow aspirate was collected from the sternum of 9 horses and MSC were isolated and expanded in culture until passage two. To evaluate chondrogenesis, equine MSC were cultured as pellets in chondrogenic medium with or without addition of TGF- β 1 (10 ng/ml). At Day 21, pellets were collected and pellet area was measured using a stereomicroscope. Pellets were paraffin-embedded, sectioned and stained with safranin-O/fast green for sulfated glycosaminoglycans (GAG). The Bern score was used for histological grading with a higher score indicating higher chondrogenesis. Presence of T β R-II was analyzed by immunofluorescence (K105, Cell Signaling Technology) and five to seven regions of each pellet were imaged with fluorescence microscopy at 40X. The T β R-II intensity of each region was quantified with Visiopharm image analysis software and normalized to the number of cells. Comparisons between groups were performed using paired t-tests and correlations were evaluated by Pearson's correlation. Statistical analysis was performed using SPSS and $p < 0.05$ was considered significant.