

DIRECT original article

Trans-fatty acid content of food products in Spain in 2015



CrossMark

Napoleón Pérez-Farinós*, María Ángeles Dal Re Saavedra,
Carmen Villar Villalba, Teresa Robledo de Dios

Spanish Agency for Consumer Affairs, Food Safety and Nutrition, Ministry of Health, Social Services and Equality, Madrid, Spain

ARTICLE INFO

Article history:

Received 26 February 2016

Accepted 5 April 2016

Available online 4 June 2016

Keywords:

Trans-fatty acids

Diet

Food and nutrition

Nutrition policy

Reformulation

Food quality

Cardiovascular diseases

ABSTRACT

Objective: To ascertain the content of trans-fatty acids (TFA) in food products in Spain in 2015 and assess trends in TFA content since 2010.

Methods: We analysed the fat content of 277 food products purchased in Spanish supermarkets in 2015 and calculated both the total fat and TFA content and the proportion of TFA to total fats. The results obtained in 2015 were compared to those yielded by a similar study in 2010.

Results: In 2015, the majority of food products studied had a TFA content of less than 0.2 g/100 g product, and a TFA/total fat ratio of less than 2%. No significant increases were found compared to 2010. Food groups with a higher TFA content were dairy products of possible natural origin.

Conclusions: TFA content in Spain is low and has significantly fallen since 2010.

© 2016 SESPAS. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Contenido de ácidos grasos *trans* en los alimentos en España en 2015

RESUMEN

Palabras clave:

Ácidos grasos *trans*

Dieta

Alimentos y nutrición

Políticas de nutrición

Reformulación

Calidad de los alimentos

Enfermedades cardiovasculares

Objetivo: Conocer el contenido de ácidos grasos *trans* en los alimentos en España en 2015 y evaluar la tendencia desde 2010.

Métodos: Se analizó la composición grasa de 277 alimentos adquiridos en supermercados en España en 2015. Se calcularon el contenido en ácidos grasos *trans* y en grasas totales, y el porcentaje de ácidos grasos *trans* respecto de la grasa total. Se compararon los resultados con los de un estudio similar realizado en 2010.

Resultados: En 2015 la mayoría de los alimentos estudiados tenían menos de 0,2 g de ácidos grasos *trans* por 100 g de producto, y menos del 2% de ácidos grasos *trans* respecto a las grasas totales. No se hallaron aumentos significativos desde 2010. Los grupos con contenidos más altos de ácidos grasos *trans* eran derivados lácteos, con posible origen natural.

Conclusiones: El contenido de ácidos grasos *trans* en España es bajo y se ha reducido significativamente desde 2010.

© 2016 SESPAS. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Trans fatty acids (TFA) are unsaturated fatty acids with at least one double bond in the transconfiguration.¹ This configuration can come about naturally from ruminant-based meat and dairy products, or artificially through partial hydrogenation of vegetable oils in the food industry and manufacturing processes.^{1,2}

There is scientific evidence to show that TFA intake is associated with cardiovascular diseases in different ways.^{3,4} In addition, associations between TFA intake and other diseases, such as diabetes mellitus,^{1,3–5} cancer⁶ and asthma,⁷ have also been proposed.

Due to those health risks, it is clear that TFA intake should be as low as possible, and that TFA content in food products should

be reduced or eliminated. Some countries have regulated TFA content, such as Denmark, by means of an Order which came into force in 2003.⁸ In Spain, the 2011 Food Safety and Nutrition Act included a specific provision targeted at reducing artificial TFA in food products.⁹

Although an important decrease in TFA content and intake has been detected in some countries,^{10,11} as a result of statutory regulation, health promotion actions and voluntary food-industry initiatives, more efforts are nevertheless called for.

The aim of this study was thus to ascertain the average TFA content of food products in Spain in 2015, and assess time trends in TFA content since 2010 when an earlier study was undertaken.¹⁰

Methods

To assess the TFA content in Spain, food products were acquired from hypermarkets and supermarkets countrywide.

* Corresponding author.

E-mail address: nperez@mssi.es (N. Pérez-Farinós).

Table 1

Trans-fatty-acids and total fat content (g/100 g of product) of food products in Spain.

Food groups	N	TFA (g/100 g of product)						Total fats (g/100 g of product)					
		2010		2015		p ^a	2010		2015		Mean	SD	p ^a
		Mean	SD	Mean	SD		Mean	SD	Mean	SD			
Salty snacks	16	0.057	0.057	0.003	0.007	0.005	22.213	5.278	22.525	6.204	0.856		
Confectionary and pastries	21	0.210	0.381	0.008	0.014	<0.001	27.629	12.935	24.276	3.953	0.079		
Processed meats	29	0.056	0.037	0.051	0.029	0.332	21.572	7.463	20.472	9.312	0.381		
Breakfast cereals and cereal bars	7	<0.001	0.000	<0.001	0.000	0.998	3.200	0.924	2.857	0.776	0.104		
Chocolates and cocoa products	24	0.099	0.086	0.038	0.056	0.003	21.746	12.206	22.388	12.440	0.796		
Biscuits	24	0.049	0.035	<0.001	0.000	<0.001	15.267	5.085	14.004	4.839	0.025		
Ice creams	19	<0.001	0.000	0.025	0.058	0.067	11.937	3.852	12.305	3.513	0.888		
Butter	7	1.927	0.228	2.011	0.209	0.176	81.814	2.078	82.029	0.706	0.735		
Margarine	10	0.254	0.130	0.136	0.205	0.093	59.640	9.582	59.480	9.243	0.681		
Industrial bread	11	<0.001	0.000	<0.001	0.000	0.997	3.509	1.127	3.045	0.962	0.091		
Chips	16	0.071	0.056	0.030	0.089	0.049	33.975	2.884	34.531	2.402	0.660		
Ready meals	23	0.013	0.023	0.023	0.043	0.218	8.352	2.502	7.752	1.984	0.236		
Products of fast-food restaurants	13	0.196	0.200	0.127	0.220	0.241	10.538	4.822	7.862	4.074	0.236		
Spreadable cheeses	12	0.522	0.095	0.536	0.181	0.790	20.417	4.784	21.358	4.728	0.238		
Tomato sauce	13	<0.001	0.000	<0.001	0.000	0.999	3.531	0.588	3.692	0.621	0.181		
Yoghurts and dairy products	32	0.055	0.055	0.073	0.039	0.074	3.544	2.194	3.178	1.585	0.046		

SD: Standard deviation; TFA: trans fatty acids.

^a p for differences between means from 2010 to 2015.

The most sold food items in each group were selected for a previous study conducted by the Spanish Agency for Consumer Affairs, Food Safety and Nutrition (*Agencia Española de Consumo, Seguridad Alimentaria y Nutrición/AECOSAN*) in 2010.¹⁰ We selected the same products for the purposes of our study, in order to compare TFA content between 2010 and 2015. The methodology used in both the 2010 and 2015 studies was similar.

We measured total fat and TFA content in the selected food items (g/100 g of product). Analytical tests were performed by an independent laboratory accredited by the Spanish National Accreditation Board (*Entidad Nacional de Acreditación/ENAC*), using gas chromatography with a flame ionization detector for food samples.

Mean and standard deviation were computed for TFA and total fat content in each group of food items. The TFA/total fat ratio was also calculated for each food product and group.

To assess the differences between 2010 and 2015, we compared the mean TFA values (g/100 g of product) and TFA/total

fat percentage ratios for those years using the U Mann-Whitney test.

In any case where there was no exact match between 2010 and 2015, the product concerned was excluded from the analysis to ensure the comparability of all groups.

Results

Based on the 2010 study, a set of 443 food items was initially selected.¹⁰ After the excluding all items that failed to show a match between 2010 and 2015, a total of 277 products was finally analyzed and grouped (**Table 1**).

In 2015, the average TFA content was less than 0.2 g/100 g of product across all food groups, with only two exceptions (butter: 2.011 g; spreadable cheeses: 0.536 g). TFA were undetectable in several groups, such as breakfast cereals, biscuits, industrial bread and tomato sauce. Comparison against the 2010 results showed reductions in TFA content in most groups, with some

Table 2

Average proportion of trans-fatty-acids/total fats in food products in Spain.

Food groups	N	2010		2015		p ^a
		TFA/total fats	SD	TFA/total fats	SD	
Salty snacks	16	0.231	0.216	0.011	0.023	0.001
Confectionary and pastries	21	0.657	0.836	0.034	0.055	<0.001
Processed meats	29	0.248	0.147	0.248	0.068	0.673
Breakfast cereals and cereal bars	7	0.034	0.010	0.037	0.011	0.345
Chocolates and cocoa products	24	0.721	0.707	0.140	0.208	<0.001
Biscuits	24	0.311	0.238	0.008	0.003	<0.001
Ice creams	19	0.010	0.005	0.271	0.715	0.061
Butter	7	2.356	0.282	2.452	0.252	0.176
Margarine	10	0.413	0.183	0.274	0.466	0.093
Industrial bread	11	0.031	0.008	0.035	0.008	0.091
Chips	16	0.210	0.168	0.088	0.267	0.020
Ready meals	23	0.131	0.231	0.307	0.583	0.638
Products of fast-food restaurants	13	1.609	1.870	1.959	4.477	0.382
Spreadable cheeses	12	2.646	0.540	2.524	0.709	0.814
Tomato sauce	13	0.029	0.005	0.028	0.004	0.182
Yoghurts and dairy products	32	1.656	1.297	2.297	0.465	0.064

SD: standard deviation; TFA: trans fatty acids.

^a p for differences between means from 2010 to 2015.

of these decreases proving statistically significant (salty snacks, confectionery and pastries, chocolates, biscuits and chips), and indicated only a few non-significant increases (**Table 1**).

The average proportion of TFA to total fats in 2015 was lower than 2% across all food-product groups, except yoghurts and dairy products (2.30%), butter (2.45%) and spreadable cheeses (2.52%), as shown in **Table 2**. Non-significant increases were found and there were some significant reductions in salty snacks, confectionery and pastries, chocolates, biscuits, and chips.

Discussion

There is solid scientific evidence to show an association between TFA intake and cardiovascular diseases. A reduction in TFA content in food products and TFA intake should thus be recommended. While TFA content in Spain was reported to be high by a number of studies,^{2,12} one study suggested that in 2010 TFA content had been reduced in the majority of the food groups studied.¹⁰

This study confirms the situation found five years previously. The comparison between TFA content in 2010 and 2015 showed that there had been no significant changes. In some groups, such as breakfast cereals, industrial bread and tomato sauce, TFA were undetectable, and in other groups, significant reductions were even found (salty snacks, confectionery and pastries, chocolates, biscuits, chips). Some food groups displayed slight, non-significant increases but the magnitude of the content was so low as to be of no concern.

It is also interesting to establish the proportion of TFA to total fats because the majority of regulations use this ratio to set the permitted limit. Our study shows that only in three groups (yoghurts and dairy products, butter, and spreadable cheeses) is this proportion higher than 2%. Even so, TFA content in yoghurts and dairy products is very low (0.073% of the total product). Moreover, TFA of natural origin can be present in meat and dairy products from ruminants (which is the case in point). Another issue lies in the fact that in some groups, such as yoghurts and dairy products, total fat content has been significantly reduced, and this could go some way to explain the rise in the TFA/total fat ratio.

Insofar as TFA of natural origin is concerned, some studies have postulated that such TFA are not associated with cardiovascular diseases and, indeed, might even have some health benefits.¹³ More studies are needed to address this issue.

Following the Danish initiative, some laws in Europe (including the Spanish Food Safety and Nutrition Act) and the USA have sought to reinforce the reduction in TFA content in the case of food products.^{8,11,14} In June 2015, the US Food and Drug Administration (FDA) finalized its determination whereby food manufacturers were ordered to remove TFA from their products in 2018. A combination of statutory regulations and voluntary actions would appear to be the most advisable strategy.¹⁵

The social responsibility of food manufacturers in Spain warrants a special mention. The food industry has made remarkable efforts to reduce TFA since the consequences to health became known.

The main strength of this study is its strict methodology which successfully ensured comparability of the results obtained in 2010 and 2015.

One limitation might lie in the fact that the selected food-product sample belonged to leading consumer brands enjoying widespread distribution, whereas there are also other locally distributed, lower-impact brands which may have been overlooked by the study. Subsequent assessments are called for.

The main conclusion of our study is that the average TFA content of food products in Spain is low, and that in the context of a balanced diet, exposure to TFA should thus be minimal. Although it cannot be stated with certainty in the absence of data on TFA intake, it is safe

to assume that, where exposure is low, the average intake in Spain would likewise be low. Nonetheless, regular surveillance of the TFA content of food products should be maintained. By way of a public health goal, reformulation of food products must be reinforced in order to ensure that the nutritional quality of food is improved.

What is known about the topic?

Trans fatty acids (TFA) are related to cardiovascular diseases. TFA content of food products in Spain was high at the beginning of 21st century but there were indications that it was being reduced. It was necessary to assess whether TFA content was really being reduced.

What does this study add to the literature?

This study confirms that the TFA content of food products in Spain is currently low and does not pose a public health problem. This fact is important in the current context of food-product reformulation policies in Spain, Europe and the rest of the world.

Editor in charge

Carlos Álvarez-Dardet.

Transparency declaration

The corresponding author on behalf of the other authors guarantee the accuracy, transparency and honesty of the data and information contained in the study, that no relevant information has been omitted and that all discrepancies between authors have been adequately resolved and described.

Funding

The study was entirely funded by the Spanish Agency for Consumer Affairs, Food Safety and Nutrition.

Authorship contributions

All authors conceived and designed the study, collaborated in the analysis and interpretation of results, and reviewed and approved the final version of the manuscript. N. Pérez-Farinós wrote the article and was chiefly responsible for the methodological aspects and analysis of the data.

Conflicts of interest

None.

References

1. Mozaffarian D, Katan MB, Ascherio A, et al. Trans fatty acids and cardiovascular disease. *N Engl J Med.* 2006;354:1601–13.
2. Riobó P, Breton I. Informe FESNAD sobre la ingesta de grasas trans. Situación en España. Madrid: Federación Española de Sociedades Científicas de Alimentación, Nutrición y Dietética; 2013. p. 20.
3. Nishida C, Uauy R. WHO Scientific Update on health consequences of trans fatty acids: introduction. *Eur J Clin Nutr.* 2009;63:S1–4.
4. Hammad S, Pu S, Jones PJ. Current evidence supporting the link between dietary fatty acids and cardiovascular disease. *Lipids.* 2016;51:507–17.
5. De Souza RJ, Mente A, Maroleanu A, et al. Intake of saturated and trans unsaturated fatty acids and risk of all cause mortality, cardiovascular disease, and type 2 diabetes: systematic review and meta-analysis of observational studies. *BMJ.* 2015;351:1–16.

6. Chajès V, Thiébaut ACM, Rotival M, et al. Association between serum trans-monounsaturated fatty acids and breast cancer risk in the E3N-EPIC study. *Am J Epidemiol.* 2008;167:1312–20.
7. Weiland SK, von Mutius E, Husing A, et al. Intake of trans fatty acids and prevalence of childhood asthma and allergies in Europe. *Lancet.* 1999;353:2040–1.
8. Leth T, Jensen HG, Mikkelsen AE, et al. The effect of the regulation on trans fatty acid content in Danish food. *Atheroscler Suppl.* 2006;7:53–6.
9. Ley 17/2011, de 5 de julio, de seguridad alimentaria y nutrición. España: Boletín Oficial del Estado 160, 6 de julio de 2011: 71283–319.
10. Robledo de Dios T, Dal Re Saavedra MA, Villar Villalba C, et al. Contenido de ácidos grasos trans en los alimentos en España. *Nutr Hosp.* 2015;32:1329–33.
11. Temme EHM, Millenaar IL, Donkersgoed GVAN, et al. Impact of fatty acid food reformulations on intake of Dutch young adults. *Acta Cardiol.* 2011;66:721–8.
12. Fernández-San Juan PM. Trans fatty acids (tFA): sources and intake levels, biological effects and content in commercial Spanish food. *Nutr Hosp.* 2009;24:515–20.
13. Chardigny JM, Destaillets F, Malpuech-Brugère C, et al. Do trans fatty acids from industrially produced sources and from natural sources have the same effect on cardiovascular disease risk factors in healthy subjects? Results of the trans Fatty Acids Collaboration (TRANSFACT) study. *Am J Clin Nutr.* 2008;87:558–66.
14. Monge-Rojas R, Colón-Ramos U, Jacoby E, et al. Voluntary reduction of trans-fatty acids in Latin America and the Caribbean: current situation. *Rev Panam Salud Pública.* 2011;29:126–9.
15. Downs SM, Thow M, Leeder SR. The effectiveness of policies for reducing dietary trans fat: a systematic review of the evidence. *Bull World Health Organ.* 2013;91:262–9.