

109 MESENCHYMAL STEM CELLS MIGRATE INTO OSTEOARTHRITIS JOINT FROM SYSTEMIC CIRCULATION: AN ANIMAL MODEL

M.C. Arufe^{1,2}, A. De la Fuente², I.A. Lesende-Rodríguez², I. Fuentes¹, F.J. De Toro¹, F.J. Blanco^{1,2}. ¹Dept. of Med. Faculty of Hlth.Sci. INIBIC-Univ. of A Coruña, A Coruña, Spain; ²Osteoarticular and Aging Res. Lab. INIBIC-CHUAC, A Coruña, Spain

Purpose: There are not clear evidences about the migration of mesenchymal stem cells from a systemic circulation to another niche as an injury place in the organism. The current study treats to answer this question.

Methods: Mesenchymal stem cells (MSCs) have been injected via intravenous and directly into the monkey joint, these MSCs were previously characterized by flow cytometry revealing that were positive for CD90 and CD44 and negatives for CD34 and CD45 indicating that presence of MSCs markers. CD105 enriched sub-population of MSCs was labelled with oxycarbocyanine – DiO (green) when they were injected via intravenous (IV), with octadecyl (C18) indocarbocyanine – Dil (red) when they were injected directly into the knee, intra-articular (IA) and another group of MSCs were modified to express constitutively the green fluorescence protein (GFP) to follow their evolutions through the animals injected by different ways and their localization. The animal models used were adult monkeys *Macaca Fascicularis* which had been injured into the left knee to create an Osteoarthritis (OA) animal model and the right knee was used as control. First step: one group of three animals was injected with 1×10^6 GFP-MSCs/KG of animal in the vein twice and these animals were sacrificed only one week after the second injection because the purpose of this experiment was following the GFP-MSCs into the organism. Second step: one group of two animals was injected twice in the vein with 1×10^6 CD105⁺-MSCs labelled with DiO only and another two animals were injected directly in the knee twice with 1×10^6 CD105⁺-MSCs labelled with Dil at the same time these animals were injected twice in the vein with 1×10^6 CD105⁺-MSCs labelled with DiO. The injections were realized into the animals with an interval of one week between them. The animals were sacrificed one month after treatment.

Results: Immunohistochemistry analysis of the tissues from the animals revealed that and GFP-MSCs and CD105⁺-MSCs migrated towards the injured knee joint. Neither labelled cells nor teratomes were found in other vital organs into the animals. The CD105⁺-MSCs labelled with Dil (red) and with DiO (green) were only located in the injured knee (0.01% from the total injected) and they were not found in the normal one. Furthermore CD105⁺-MSCs labelled with Dil (red) and with DiO (green) were found in crossed ligaments and knee muscle (vastus medialis), synovial membrane, both meniscus and cartilage of patello femoral groove all of them only in the injured joint and not in the normal one. An exhaustive immunofluorescence analysis of DiO labelled cells founded in the injured knee revealed that they were negatives for CD68, which is a macrophage antibody marker, and 4% of CD105⁺-MSCs labelled with DiO (green) were positives for SDF-1, which is a marker of osteoclastogenesis and produces liberation of IL-6, in the patello femoral groove and crossed ligaments of injured joint and not in the normal one.

Conclusions: These results seem indicate that:

1. The MSCs injected in a peripheral vein go to injured tissue, in this case injury knee.
2. The MSCs injected directly in an injured place keep in the same place.
3. The MSCs could be recruitment by SDF-1.

110 SHORT TERM HIGH-FAT DIET INDUCES OSTEOPHYTE FORMATION, APOPTOSIS AND DEGENERATION OF ARTICULAR CHONDROCYTE IN MURINE KNEE JOINT

M. Iwata¹, H. Ochi², Y. Hara¹, D. Koga³, M. Tagawa¹, A. Okawa³, Y. Asou³. ¹Div. of Vet. Surgery, Sch. of Vet. Med., Nippon Vet. and Life Sci. Univ., 1-7-1, Kyonan-cho, Musashino-shi, Tokyo, Japan; ²Section of Nephrology, Endocrinology and Metabolism, Dept. of Internal Med., Sch. of Med., Keio Univ., 35 Shinanomachi Shinjuku Tokyo, Japan; ³Dept. of Orthopaedics Surgery, Tokyo Med. and Dental Univ., 1-5-45, Yushima Bunkyo-ku, Tokyo, Japan

Purpose: Obesity and high body mass index are associated with a higher incidence risk of OA. To elucidate the relationship between obesity and osteoarthritis, several attempts have been made to observe the effect

of high fat diet (HFD) on murine articular degeneration. Although HFD have succeeded in inducing osteoarthritis after long term treatment (for more than 6months), initial response of articular tissues to HFD is still unknown. The aim of this study is to evaluate sub-acute effect of HFD on histological features of murine knee joint.

Materials and Methods: C57BL/6J male mice were fed a diet containing 32% fat for HFD group and normal diet for control group from 7 weeks of age for less than 12 weeks. Mice feeding a high fat diet and control have been divided into three groups, such as 4 weeks, 8 weeks and 12 weeks of treatment. Whole knee joints were removed by dissection, fixed in 4% paraformaldehyde, and decalcified in EDTA. After dehydration and paraffin embedding, serial 5- μ m sagittal sections were made from the whole medial compartment of the knee joint. Three sections were picked up at 100- μ m intervals (from lane 1; medial to lane 3; lateral) from the weight-bearing region of each medial plateau of tibia, and were stained with Safranin O-fast green and HE. OA severity in the tibial plateau was evaluated according to Mankin's histologic grading system. Osteophyte area was quantified in each slide using Image-Pro Plus 4.1 software. ColX expression was examined by immunohistochemistry. TUNEL assay was performed and the number of TUNEL positive cells were compared between HFD group and control.

Results: Body weight was increased 10% in 4weeks, 20% in 8 weeks, 50% in 12 weeks of HFD group compared to controls respectively. As soon as 4weeks of HFD treatment, the area of intrapatellar fat pad (IPFP) was increased in HFD group prior to cartilage degeneration. Infiltration of inflammatory cells was observed at synovium and IPFP. Prior to osteophyte formation, increased synovial cells were abundantly observed at anterior edge of tibial plateau. Simultaneously, angiogenesis was activated in IPFP. At 8 weeks of HFD treatment, TUNEL positive cells were abundantly observed at the site of osteophyte formation and superficial or middle zone of articular cartilage. The number of TUNEL positive cells was significantly elevated in HFD group. Osteophyte formation was detected at the medio-anterior edge of tibial plateau. Mankin score was significantly elevated in HFD group from this period. At 12 weeks of treatment, osteophyte formation was apparently advanced in HFD group as a result of activated endochondral ossification. Type X collagen expression in articular cartilage was enhanced at weight bearing region in HFD group.

Conclusion: We demonstrated that osteoarthritic change, such as osteophyte formation, apoptosis and degeneration of articular chondrocyte, could be induced by HFD within three month of treatment. Prior to OA progression, thickening of IPFP accompanied by mild inflammation was observed, suggesting involvement of IPFP in the initiation of OA change. In vivo model described herein offers opportunities to noninvasively induce osteoarthritis without surgery.

111 ANTI-INFLAMMATORY EFFECT OF ROSUVASTATIN AND ROSIGLITAZONE SUPPRESSES DEVELOPMENT OF DIET-INDUCED OSTEOARTHRITIS IN HUMAN CRP TRANSGENIC MICE

L.M. Gierman¹, F. van der Ham¹, A. Koudijs¹, P.Y. Wielinga¹, R. Kleemann¹, T. Kooistra¹, R. Stoop¹, G.J. van Osch², V. Stojanovic-Susulic³, M. Kloppenburg⁴, T. Huizinga⁴, A-M. Zuurmond¹. ¹TNO, Leiden, Netherlands; ²Erasmus MC, Rotterdam, Netherlands; ³Centocor, Radnor, PA, USA; ⁴LUMC, Leiden, Netherlands

Purpose: Obesity is associated with systemic inflammation and a risk factor for osteoarthritis (OA) development. We hypothesized that metabolic stress-induced inflammation is responsible for the development of high fat diet (HFD)-induced OA in mice and not mechanical overload.

Methods: To test this hypothesis, human C-reactive protein (huCRP) transgenic mice received HFD without or with 0.005% w/w rosuvastatin or 0.018% w/w rosiglitazone, two different types of drugs with anti-inflammatory capacities. Control mice fed chow were also included. After 42 weeks mice were sacrificed and histological grading of the knees was performed to assess OA development. To monitor the overall inflammatory state, systemic huCRP levels were determined. In addition, the adipokines leptin and resistin, for which a role in OA has been suggested, were measured.

Results: Male mice on HFD had significantly higher OA grades than mice on chow and showed no correlation between severity of OA and body weight. HFD-induced OA was significantly inhibited by rosuvastatin or rosiglitazone to OA grades observed in age-matched control mice on