



SCRAPPER-Dependent Ubiquitination of Active Zone Protein RIM1 Regulates Synaptic Vesicle Release

Ikuko Yao, Hiroshi Takagi, Hiroshi Ageta, Tomoaki Kahyo, Showbu Sato, Ken Hatanaka, Yoshiyuki Fukuda, Tomoki Chiba, Nobuhiro Morone, Shigeki Yuasa, Kaoru Inokuchi, Toshihisa Ohtsuka, Grant R. MacGregor, Keiji Tanaka, and Mitsutoshi Setou*

*Correspondence: setou@nips.ac.jp DOI 10.1016/j.cell.2007.09.015

(Cell 130, 943-957; September 7, 2007)

In the above article, the accession number for SCRAPPER was not properly updated due to a production error. The correct NCBI CoreNucleotide accession number is EF649694. We apologize for this error.

The gene locus of the gene coding SCRAPPER is 17q21.2 in human and 11D in mice. SCRAPPER shows similarities to the FBXL family of proteins (Jin, J., Cardozo, T., Lovering, R.C., Elledge, S.J., Pagano, M., and Harper, J.W. [2004]. Systematic analysis and nomenclature of mammalian F-box proteins. Genes Dev. 18, 2573-2580).

SCRAPPER protein is 76.8% similar throughout to FBXL2, a protein encoded by a different gene locus. In addition, FBXL20 (NM_028149) mRNA sequence is identical to SCRAPPER from nucleotide 302 on, but has a different N-terminal sequence, suggesting that FBXL20 and SCRAPPER are splicing variants encoded by the same gene.

Exonuclease-1 Deletion Impairs DNA Damage Signaling and Prolongs Lifespan of Telomere-Dysfunctional Mice

Sonja Schaetzlein, N.R. Kodandaramireddy, Zhenyu Ju, Andre Lechel, Anna Stepczynska, Dana R. Lilli, Alan B. Clark, Cornelia Rudolph, Florian Kuhnel, Kaichun Wei, Brigitte Schlegelberger, Peter Schirmacher, Thomas A. Kunkel, Roger A. Greenberg, Winfried Edelmann, and K. Lenhard Rudolph* *Correspondence: lenhard.rudolph@uni-ulm.de

DOI 10.1016/j.cell.2007.09.033

(Cell 130, 863-877; September 7, 2007)

When the above article was originally published, Florian Kuhnel was accidentally omitted from the author list. The error has been corrected online, and the complete author list appears here.