



ELSEVIER

journal homepage

View metadata, citation and similar papers at core.ac.uk

Formation of MgF_3^- -dependent complexes between an AAA^+ ATPase and σ^{54}

Nan Zhang¹, Martin Buck^{*,1}

Division of Cell and Molecular Biology, Sir Alexander Fleming Building, Imperial College London, Exhibition Road, London SW7 2AZ, UK

ARTICLE INFO

Article history:

Received 7 March 2012

Revised 29 March 2012

Accepted 10 April 2012

Keywords:

AAA^+ ATPase
Nucleotide analogue
Transcription
bEBP
PspF
 σ^{54}

ABSTRACT

The widely distributed bacterial σ^{54} -dependent transcription regulates pathogenicity and numerous adaptive responses in diverse bacteria. Formation of the σ^{54} -dependent open promoter complex is a multi-step process driven by AAA^+ ATPases. Non-hydrolysable nucleotide analogues are particularly suitable for studying such complexity by capturing various intermediate states along the energy coupling pathway. Here we report a novel ATP analogue, ADP-MgF_3^- , which traps an AAA^+ ATPase with its target σ^{54} . The MgF_3^- -dependent complex is highly homogeneous and functional assays suggest it may represent an early transcription intermediate state valuable for structural studies.

© 2012 Federation of European Biochemical Societies. Published by Elsevier B.V.

Open access under [CC BY-NC-ND license](http://creativecommons.org/licenses/by-nc-nd/4.0/).

1. Introduction

The high-energy phosphoryl transfer reaction is a principle mechanism exploited by mechanochemical enzymes such as the AAA^+ ATPases (ATPases associated with various cellular activities). The AAA^+ ATPases convert the chemical energy released from ATP hydrolysis to the remodelling of a diverse array of their substrates, to achieve for example protein unfolding, membrane fusion, DNA repair and transcription activation [1,2]. However, this energy transfer process is transient. In order to capture the AAA^+ ATPases and their substrates for kinetic and structural studies, nucleotide analogues are widely used. These analogues, in many cases, consist of an ADP and a metallo-halide (e.g., AlF_x , which occupies the γ -phosphate position within the ATP catalytic site) and are reported to represent different ATP states (AMP-AlF_x and ADP-BeF_3^- for the ATP ground state and ADP-AlF_x for the ATP transition state). Being the most frequently used γ -phosphate analogue, the AlF_x moiety shows complexity in binding to the catalytic site. Schlichting et al. [3] surveyed the majority of the AlF_x -containing crystal structures and demonstrated that the pH of the crystallographic experiment determined whether AlF_3 or AlF_4^- was present in the crystal (thus abbreviated as AlF_x in this paper). The AlF_3 moiety adopts a trigonal bipyramidal arrangement with the axial coordination sites being occupied by oxygens from the β -phosphate and hydrolytic

water, which is thought to closely mimic the γ -phosphate at the point of hydrolysis in geometry [4]. The AlF_4^- moiety adopts an octahedral arrangement with a net negative charge, complementary to the transition state γ -phosphate [4]. The fact that both AlF_x species are found in the crystal structures suggests the catalytic site has enough flexibility to accommodate either without much reconfiguration and energy loss [3].

Vincent et al. reported a high-affinity dimeric complex formation between the RhoA GTPase and the p190 RhoGAP (Rho GTPase-activating protein) in a fluoride- and magnesium-dependent manner but an aluminium- and GDP-independent manner [5]. To investigate whether the same effect can be observed on AAA^+ ATPases and their substrates, we employed a bacterial enhancer binding protein (bEBP) called the *Escherichia coli* phage shock protein F (PspF) – a Clade 6 hexameric AAA^+ ATPase for this study [6]. PspF or its AAA^+ domain alone (residues 1–275, PspF_{1–275}) can activate the *psp* operon (*pspABCDE* and *pspG*) [7] by reorganising the $\text{E}\sigma^{54}$ -DNA complex through PspF surface-exposed loops in a nucleotide-dependent manner. Recently, the Cryo-EM contour structure of a PspF_{1–275}• $\text{E}\sigma^{54}$ • ADP-AlF_x complex has been resolved [8]. However, the high-resolution hexameric crystal structure of PspF is yet to be obtained, partially due to the interference from precipitation arising from high concentrations of AlCl_3 used.

Here, we report an MgF_3^- -dependent complex formation between the ADP-bound PspF_{1–275} and σ^{54} . We demonstrated that this novel MgF_3^- -dependent complex was more homogeneous than the previously described complexes with AlF_x and may represent an intermediate state early along the activation pathway. We propose that MgF_3^- will serve as a new reagent to obtain high-resolution

* Corresponding author. Fax: +44 (0) 2075045419.

E-mail address: m.buck@imperial.ac.uk (M. Buck).¹ N.Z. and M.B. designed the experiments. N.Z. performed the experiments. N.Z. and M.B. contributed to the writing.

structural information on co-complexes of some AAA⁺ ATPases with their remodelling targets.

2. Materials and methods

2.1. Protein expression

E. coli PspF_{1–275} was purified as previously described [9]. *Klebsiella pneumoniae* σ^{54} was purified as previously described [10].

2.2. Native gel mobility shift assay

Reactions were performed in 10 μ l volumes containing 10 μ M PspF_{1–275}, 2.35 μ M σ^{54} , \pm AlCl₃, \pm MgCl₂, \pm ADP and \pm NaF in STA buffer (2.5 mM Tris–acetate pH 8.0, \pm 8 mM Mg–acetate, \pm 8 mM K–acetate, 10 mM KCl, 1 mM DTT, 3.5% (w/v) PEG 8000) at 37 °C for 15 min. Complexes were analysed on 4% native gels.

2.3. Gel filtration

The trapped complexes with 20 μ M PspF_{1–275} and 4.7 μ M σ^{54} were formed after 15 min incubation with reagents at 37 °C and run with gel filtration buffer (20 mM Tri–HCl pH 8.0, 50 mM NaCl, 15 mM MgCl₂) in a Superdex 200 Column (10/30, 24 ml, GE Healthcare) at room temperature.

2.4. In vitro RP₀ formation assay

The RP₀ formation assay was conducted as previously described [9]. Typically in 10 μ l volumes, 4 μ M PspF_{1–275}, 100 nM holoenzyme (1:4 ratio of E: σ^{54}), 4 mM dATP and 20 nM linear *Sinorhizobium meliloti* *nifH* promoter probes (Sigma–Aldrich) were incubated at 37 °C for 15 min before the elongation mixture (0.5 mM dinucleotide primer UpG, 0.2 μ Ci/ μ l [α -³²P GTP] (3000 Ci/mmol) and 0.2 mg/ml heparin) was added for another 10 min incubation. Reactions were quenched by addition of 4 μ l formamide stop dye and run on a sequencing gel.

3. Results

3.1. The Mg²⁺-promoted PspF_{1–275}– σ^{54} complex requires ADP but not Al³⁺

In the presence of the ATP transition state analogue ADP–AlF_x, the PspF surface-exposed L1 loops extend to stably engage σ^{54} [11]. The resulting PspF_{1–275}• σ^{54} •ADP–AlF_x trapped species represents a sub-complex of one of the intermediate states en route to open complex formation (RP₀) [12] to support transcription initiation by making the start site available [13]. However, heterogeneity is often observed in the population of ADP–AlF_x trapped complexes (Fig. 1 lane 4), which can lead to potential complications in mass spectroscopic analyses and crystallography. Higashijima et al. have shown that at high F[–] concentrations, Mg²⁺ can replace Al³⁺ in transforming the G protein α subunit into a more active state, possibly by associating with three F[–] ions to mimic the γ -phosphate of GTP [14]. In an attempt to obtain a more homogeneous population of PspF_{1–275}– σ^{54} trapped complexes, possibly with new geometrical and functional features, we performed the trapping experiment by in situ formation of MgF₃[–] in the absence and presence of nucleotides.

The trapping reaction buffer (STA buffer), which has routinely been used in various binding and transcription activation assays [9], contains 8 mM Mg²⁺–acetate. We initially assessed whether the intrinsic Mg²⁺ concentration from the reaction buffer was sufficient to support the formation of MgF₃[–] moieties. Indeed, without




Fig. 1. PspF_{1–275} forms nucleotide- and MgF₃[–]-dependent trapped complexes with σ^{54} . The STA (+8 mM Mg²⁺) buffer has been routinely used in various native gel shift assays and transcription activation assays and contains 2.5 mM Tris–acetate pH 8.0, 8 mM Mg²⁺–acetate, 10 mM KCl, 1 mM DTT, and 3.5% PEG 8000. To assess whether the trapped complex formation was dependent on the intrinsic Mg²⁺ ions, the STA (–8 mM Mg²⁺) buffer was used in which the 8 mM Mg²⁺–acetate was replaced by 8 mM K⁺–acetate.

any added Al³⁺, a more homogeneous population of Mg²⁺-promoted complexes was observed, whose formation was absolutely dependent on the presence of ADP (Fig. 1 lanes 5 and 6) and NaF (Fig. 1 lane 11). Addition of further Mg²⁺ ions to the PspF_{1–275}– σ^{54} interaction assay did not seem to increase the yield of complexes, even though the concentrations of PspF_{1–275} and σ^{54} were not limiting (Fig. 1 compare lane 8 with lane 6). When the Mg²⁺ ions were removed from the STA buffer, the PspF_{1–275}– σ^{54} complex formation was completely abolished (Fig. 1 lane 12) but restored once the Mg²⁺ ions were added back (Fig. 1 lane 10), further confirming the Mg²⁺-dependent nature of this newly trapped complex. The gel filtration data (Fig. 2) demonstrated that the ADP–MgF₃[–]-dependent complex eluted as a single homogenous peak (at 10.06 ml) before the doubly peaked ADP–AlF_x-dependent complexes (at 10.16/10.91 ml), suggesting a different intermediate state is likely to be represented by the ADP–MgF₃[–]-dependent complex.

The above observations suggest that the AlF_x-dependent trapped complexes formed in the presence of Mg²⁺ ions are likely to be a mixture of PspF_{1–275}• σ^{54} •ADP–AlF_x/–MgF₃[–] with ADP–AlF_x species dominating.

3.2. ADP–AlF_x stabilises the PspF_{1–275}– σ^{54} complex more strongly than does ADP–MgF₃[–]

Since trapped complexes formed in STA buffer contain a mixture of PspF_{1–275}• σ^{54} •ADP–AlF_x/–MgF₃[–] due to the presence of both Mg²⁺ and Al³⁺ ions in the reaction, we examined whether or not the AlF_x-dependent complexes could form in the absence of Mg²⁺ ions. As shown in Fig. 3A, adding NaF and Al³⁺ ions to the Mg²⁺–acetate free STA buffer shifted nearly all the σ^{54} into the trapped complex (Fig. 3A lane 6). The addition of 0.4mM Mg²⁺ ions (same concentration as Al³⁺ ions) or a 20-fold higher concentration of Mg²⁺ ions yielded 16% and 37% ADP–MgF₃[–] trapped complexes compared to the Al³⁺-dependent assays (Fig. 3A compare lanes 4 and 5 with 6). Furthermore, a titration experiment revealed that




Fig. 2. Gel filtration of the ADP-AIF_x/⁻MgF₃⁻ dependent trapped complexes at room temperature.

a relatively low concentration of the Al³⁺ ions (0.04 mM) was required to form the AIF_x-dependent complexes, much lower than




Fig. 3. (A) Al³⁺ ions are more efficient at promoting the trapped complex formation. (B) Titration of the amount of Al³⁺ ions required for trapped complex formation.




Fig. 4. Comparison of the ADP-AIF_x and ADP-MgF₃⁻ dependent complexes in the RP₀ formation assay. The amount of abortive tetra-nucleotides (UpGpGpG) generated from this assay directly correlates with the amount of open complex formation. Three DNA sequences with mismatches on the non-template strand (thick curves in cartoons) were used: the homoduplex WT/WT, the early-melted -12-11/WT (mimicking the DNA conformation in the closed promoter complex), and the late-melted -10-1/WT (mimicking the DNA conformation in the open promoter complex).

the 0.4 mM routinely used (Fig. 3B). The above observations suggest that although both Al³⁺ and Mg²⁺ ions can form the ADP-dependent trapped complexes independently of one another's presence, the Al³⁺ ions are far more efficient at promoting the complex formation.

3.3. ADP-MgF₃⁻ is a functional analogue of ADP-BeF₃⁻ in RP₀ formation

Burrows et al. [13] devised a short primed RNA (sprRNA) synthesis assay and demonstrated that the putative transition state ADP-AIF_x complex could reorganise Eσ⁵⁴ to a near open complex state on a pre-opened linear DNA probe (the *S. meliloti nifH* promoter with the non-template 'melted' from -10 to -1). However, the ground state ADP-BeF₃⁻ complex failed to do so [13]. Here we employed the sprRNA assay to assess whether the MgF₃⁻-dependent complexes are transcriptionally active and/or carry any functional

similarities to either ADP- AlF_x or ADP- BeF_3^- dependent complexes.

In the presence of a dinucleotide primer UpG and radio-labelled GTP, abortive tetra-nucleotides UpGpGpG are generated on the linear *nifH* promoter by preventing $\text{E}\sigma^{54}$ from transcribing beyond the +3 site. The amount of UpGpGpG formed reflects the amount of RP_O -like activity in the presence of ADP-metal fluorides. Consistent with the previous data, the ADP- AlF_x dependent complex was able to generate an RP_O -like activity from the pre-melted DNA probe (Fig. 4). However, the ADP- MgF_3^- -dependent complex failed to yield RP_O -like activity from any linear DNA probes used in this experiment (Fig. 4) – a similar functional phenotype was exhibited by the ADP- BeF_3^- dependent complexes [13]. We propose that MgF_3^- and BeF_3^- may (i) similarly change the organisation of the ATPase catalytic site and (ii) represent functional intermediate states of the AAA^+ domain in ATP hydrolysis which form prior to the state created by the ADP- AlF_x .

4. Discussion

We have identified a new method dependent on the formation of MgF_3^- moieties, to stably trap an AAA^+ ATPase PspF_{1–275} with its target substrate σ^{54} . The MgF_3^- -dependent complexes can co-exist in solution with the AlF_x -dependent complexes when both metal ions are present – a condition under which most of the previous biochemical, mass spectroscopic and crystallographic experiments were performed. This potential heterogeneity of complex formation with AlCl_3 and NaF in the presence of Mg^{2+} is not readily detected given the AlF_x functions more efficiently in trapping conditions, and so could have been easily overlooked. As a potential source of heterogeneity in protein conformation, the presence of MgF_3^- and AlF_x may interfere with protein crystallisation. Based on the pH effect [3], we reason that the AlF_x moiety under the trapping conditions in this work (pH 8.0) is more likely to assume a trigonal bipyromidal AlCl_3 configuration than an octahedral AlF_4^- configuration. However, Xiaoxia et al. [15] argue that there is a dominant role of charge in selection of the best bound ATP analogues and thus the AlF_4^- moiety might be considered the better binding candidate species compared to AlCl_3 , as has been observed in other classes of ATP hydrolysing enzymes. Clearly, further detailed analyses and high resolution structural information are required to determine the precise ATP analogue species bound and roles of charge/geometry relationships in their binding to the bEBP class of ATPases.

Vincent et al. suggested that additional mechanistic roles could be assigned to the MgF_3^- moiety. Their observation of MgF_3^- -dependent GTPase-GAP complex formation in the absence of GDP challenges the widely held γ -phosphate mimicking role for MgF_3^- [5]. Our MgF_3^- -dependent trapping data revealed an absolute requirement for ADP for PspF_{1–275} to interact with σ^{54} , suggesting that MgF_3^- in AAA^+ ATPases is confined to function solely as a γ -phosphate mimick. We reason that in contrast to the relatively 'simple' GTP catalytic site between the GTPase-GAP heterodimer, the ATP catalytic sites at the hexameric interfaces of an AAA^+ ATPase need to be precisely organised and selective for nucleotide analogues in order to productively coordinate the energy relay across subunits [16].

The MgF_3^- and BeF_3^- moieties as trapping reagents displayed similar phenotypic traits at the level of the PspF_{1–275} engaging its target. Both moieties are less efficient at promoting the PspF_{1–275}- σ^{54} complex formation than is the AlF_x moiety (14% by BeF_3^- and 16% by MgF_3^- in comparison to 100% by AlF_x), and are

unable to productively reorganise RP_C to yield an RP_O -like complex on a pre-opened DNA probe ([13] and this work). Graham et al. suggest that the geometry of these two moieties is different at the catalytic site, as BeF_3^- adopts a tetrahedral arrangement and MgF_3^- adopts a trigonal bipyromidal arrangement [4]. Thus, MgF_3^- and BeF_3^- in combination with ADP may represent slightly different intermediate early states of bound ATP prior to ATP hydrolysis. Clearly the MgF_3^- - and ADP-dependent PspF_{1–275}- σ^{54} complex has novelty and is the first such complex reported for an AAA^+ ATPase, with the potential to advance high-resolution structural studies between nucleotide-bound AAA^+ ATPases and their targets in pre-hydrolysis state.

Acknowledgements

This work was supported by a Biotechnology and Biological Sciences Research Council Project Grant (BB/J002828/1) to M.B. We thank Prof. G. Dodson and Dr. N. Joly for stimulating discussion and members of M.B. lab for their constructive comments and support.

References

- [1] Snider, J. and Houry, W.A. (2008) AAA^+ proteins: diversity in function, similarity in structure. *Biochem. Soc. Trans.* 36, 72–77.
- [2] Tucker, P.A. and Sallai, L. (2007) The AAA^+ superfamily – a myriad of motions. *Curr. Opin. Struct. Biol.* 17, 641–652.
- [3] Schlichting, I. and Reinstein, J. (1999) PH influences fluoride coordination number of the AlF_x phosphoryl transfer transition state analog. *Nat. Struct. Biol.* 6, 721–723.
- [4] Graham, D.L., Lowe, P.N., Grime, G.W., Marsh, M., Rittinger, K., Smerdon, S.J., Gamblin, S.J. and Eccleston, J.F. (2002) MgF_3^- as a transition state analog of phosphoryl transfer. *Chem. Biol.* 9, 375–381.
- [5] Vincent, S., Brouns, M., Hart, M.J. and Settleman, J. (1998) Evidence for distinct mechanisms of transition state stabilization of GTPases by fluoride. *Proc. Natl. Acad. Sci. USA* 95, 2210–2215.
- [6] Erzberger, J.P. and Berger, J.M. (2006) Evolutionary relationships and structural mechanisms of AAA^+ proteins. *Annu. Rev. Biophys. Biomol. Struct.* 35, 93–114.
- [7] Joly, N., Engl, C., Jovanovic, G., Huvet, M., Toni, T., Sheng, X., Stumpf, M.P. and Buck, M. (2010) Managing membrane stress: the phage shock protein (Psp) response, from molecular mechanisms to physiology. *FEMS Microbiol. Rev.* 34, 797–827.
- [8] Bose, D., Pape, T., Burrows, P.C., Rappas, M., Wigneshweraraj, S.R., Buck, M. and Zhang, X. (2008) Organization of an activator-bound RNA polymerase holoenzyme. *Mol. Cell* 32, 337–346.
- [9] Zhang, N., Joly, N., Burrows, P.C., Jovanovic, M., Wigneshweraraj, S.R. and Buck, M. (2009) The role of the conserved phenylalanine in the sigma54-interacting GAFTGA motif of bacterial enhancer binding proteins. *Nucleic Acids Res.* 37, 5981–5992.
- [10] Cannon, W.V., Gallegos, M.T. and Buck, M. (2000) Isomerization of a binary sigma-promoter DNA complex by transcription activators. *Nat. Struct. Biol.* 7, 594–601.
- [11] Rappas, M., Schumacher, J., Niwa, H., Buck, M. and Zhang, X. (2006) Structural basis of the nucleotide driven conformational changes in the AAA^+ domain of transcription activator PspF. *J. Mol. Biol.* 357, 481–492.
- [12] Leach, R.N., Gell, C., Wigneshweraraj, S., Buck, M., Smith, A. and Stockley, P.G. (2006) Mapping ATP-dependent activation at a sigma54 promoter. *J. Biol. Chem.* 281, 33717–33726.
- [13] Burrows, P.C., Joly, N. and Buck, M. (2010) A prehydrolysis state of an AAA^+ ATPase supports transcription activation of an enhancer-dependent RNA polymerase. *Proc. Natl. Acad. Sci. USA* 107, 9376–9381.
- [14] Higashijima, T., Ferguson, K.M., Sternweis, P.C., Ross, E.M., Smigel, M.D. and Gilman, A.G. (1987) The effect of activating ligands on the intrinsic fluorescence of guanine nucleotide-binding regulatory proteins. *J. Biol. Chem.* 262, 752–756.
- [15] Xiaoxia, L., Marston, J.P., Baxter, N.J., Hounslow, A.M., Yufen, Z., Blackburn, G.M., Cliff, M.J. and Waltho, J.P. (2011) Prioritization of charge over geometry in transition state analogues of a dual specificity protein kinase. *J. Am. Chem. Soc.* 133, 3989–3994.
- [16] Joly, N. and Buck, M. (2010) Engineered interfaces of an AAA^+ ATPase reveal a new nucleotide-dependent coordination mechanism. *J. Biol. Chem.* 285, 15178–15186.