



ELSEVIER

Available online at www.sciencedirect.com ScienceDirect

Journal of Algebra 312 (2007) 228–256

**JOURNAL OF
Algebra**

www.elsevier.com/locate/jalgebra

Bicrossproduct approach to the Connes–Moscovici Hopf algebra

T. Hadfield ^{*,1}, S. Majid ²*School of Mathematical Sciences, Queen Mary, University of London 327 Mile End Road, London E1 4NS, England, UK*

Received 26 January 2006

Available online 1 December 2006

Communicated by J.T. Stafford

Abstract

We give a rigorous proof that the (codimension one) Connes–Moscovici Hopf algebra \mathcal{H}_{CM} is isomorphic to a bicrossproduct Hopf algebra linked to a group factorisation of the diffeomorphism group $\text{Diff}^+(\mathbb{R})$. We construct a second bicrossproduct U_{CM} equipped with a nondegenerate dual pairing with \mathcal{H}_{CM} . We give a natural quotient Hopf algebra $k_\lambda[\text{Heis}]$ of \mathcal{H}_{CM} and Hopf subalgebra $U_\lambda(\mathbf{heis})$ of U_{CM} which again are in duality. All these Hopf algebras arise as deformations of commutative or cocommutative Hopf algebras that we describe in each case. Finally we develop the noncommutative differential geometry of $k_\lambda[\text{Heis}]$ by studying first order differential calculi of small dimension.

© 2006 Elsevier Inc. All rights reserved.

Keywords: Quantum groups; Noncommutative geometry; Group factorisation; Heisenberg group

1. The Connes–Moscovici Hopf algebra \mathcal{H}_{CM}

The Connes–Moscovici Hopf algebras originally appeared in [5], arising from a longstanding internal problem of noncommutative geometry, the computation of the index of transversally elliptic operators on foliations. This family of Hopf algebras (one for each positive integer) was found to reduce transverse geometry to a universal geometry of affine nature, and provided the initial impetus for the development of Hopf-cyclic cohomology. The cyclic cohomology of these

* Corresponding author.

E-mail addresses: t.hadfield@qmul.ac.uk (T. Hadfield), s.majid@qmul.ac.uk (S. Majid).¹ Supported by an EPSRC postdoctoral fellowship.² Supported during completion of the work by the Perimeter Institute, Waterloo, Ontario.

Hopf algebras was shown by Connes and Moscovici to serve as an organising principle for the computation of the cocycles in their local index formula [4]. They are also closely related to the Connes–Kreimer Hopf algebras of rooted trees arising from renormalisation of quantum field theories [2]. More recently these Hopf algebras have appeared in number theory, in the context of operations on spaces of modular forms and modular Hecke algebras [6] and spaces of \mathbb{Q} -lattices [3]. They appear to play a near-ubiquitous role as symmetries in noncommutative geometry. There is also an algebraic approach to diffeomorphism groups [14], which we link to Connes and Moscovici’s work.

In this paper we focus on the simplest example, the codimension one Connes–Moscovici Hopf algebra. We work with a right-handed version of this algebra, which we denote \mathcal{H}_{CM} . The algebras in [5] were implicitly defined over \mathbb{R} or \mathbb{C} , but throughout this paper we will work over an arbitrary field k of characteristic zero.

Definition 1. We define \mathcal{H}_{CM} to be the Hopf algebra (over k) generated by elements X, Y, δ_n ($n \geq 1$), with

$$\begin{aligned}
 [Y, X] &= X, & [X, \delta_n] &= \delta_{n+1}, & [Y, \delta_n] &= n\delta_n, & [\delta_m, \delta_n] &= 0 \quad \forall m, n, \\
 \Delta(X) &= X \otimes 1 + 1 \otimes X + Y \otimes \delta_1, \\
 \Delta(Y) &= Y \otimes 1 + 1 \otimes Y, & \Delta(\delta_1) &= \delta_1 \otimes 1 + 1 \otimes \delta_1, \\
 \varepsilon(X) &= 0 = \varepsilon(Y), & \varepsilon(\delta_n) &= 0 \quad \forall n, \\
 S(Y) &= -Y, & S(X) &= Y\delta_1 - X, & S(\delta_1) &= -\delta_1
 \end{aligned} \tag{1}$$

with $\Delta(\delta_{n+1}), S(\delta_{n+1})$ defined inductively from the relation $[X, \delta_n] = \delta_{n+1}$.

This differs from the Hopf algebra defined in [5, p. 206], in that Connes and Moscovici take $\Delta(X) = X \otimes 1 + 1 \otimes X + \delta_1 \otimes Y$. We will denote this original left-handed version by $\mathcal{H}_{CM}^{\text{left}}$.

The first part of this paper gives a rigorous proof that \mathcal{H}_{CM} is isomorphic to a bicrossproduct Hopf algebra linked to a factorisation of the group $\text{Diff}^+(\mathbb{R})$ of positively-oriented diffeomorphisms of the real line. Recall that a group X is said to factorise into subgroups G and M if group multiplication gives a set bijection $G \times M \rightarrow X$. We write $X = G \bowtie M$. As remarked in [5], the group

$$\text{Diff}^+(\mathbb{R}) = \{ \varphi \in \text{Diff}(\mathbb{R}) : \varphi'(x) > 0 \forall x \in \mathbb{R} \} \tag{2}$$

factorises into the two subgroups

$$D_0 = \text{Diff}_0^+(\mathbb{R}) = \{ \phi \in \text{Diff}^+(\mathbb{R}) : \phi(0) = 0, \phi'(0) = 1 \} \tag{3}$$

and $B_+ = \{(a, b) : x \mapsto ax + b, a, b \in \mathbb{R}, a > 0\}$ the subgroup of affine diffeomorphisms, which we identify with its faithful matrix representation

$$B_+ = \left\{ (a, b) = \begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix} : a, b \in \mathbb{R}, a > 0 \right\}. \tag{4}$$

Given a group factorisation $X = G \bowtie M$ of a finite group X , there is a natural construction of dually-paired finite-dimensional bicrossproduct Hopf algebras denoted $k[M] \blacktriangleright kG$,

$kM \bowtie k[G]$ [10,12,19]. In our case, the group $X = \text{Diff}^+(\mathbb{R})$ is very far from finite, so it remains a challenge to construct an analogous pair of infinite-dimensional bicrossproduct Hopf algebras. Fortunately the bicrossproduct construction is more general than the group factorisation case. Given Hopf algebras \mathcal{A} and \mathcal{H} , with \mathcal{A} a left \mathcal{H} -module algebra and \mathcal{H} a right \mathcal{A} -comodule coalgebra with action and coaction compatible in an appropriate sense, then it is possible to equip the vector space $\mathcal{A} \otimes \mathcal{H}$ with the structure of a Hopf algebra, the (left–right) bicrossproduct denoted $\mathcal{A} \bowtie \mathcal{H}$ [12,15]. Similarly, we can construct a (right–left) bicrossproduct $\mathcal{H} \bowtie \mathcal{A}$ from a right \mathcal{H} -module algebra \mathcal{A} and left \mathcal{A} -comodule coalgebra \mathcal{H} . Thus in our case we define Hopf algebras $k[D_0]$, $U(\mathfrak{b}_+)$, together with a left action of $U(\mathfrak{b}_+)$ on $k[D_0]$ and right coaction of $k[D_0]$ on $U(\mathfrak{b}_+)$ which we prove are compatible in the sense necessary for the construction of a bicrossproduct Hopf algebra $k[D_0] \bowtie U(\mathfrak{b}_+)$ (Theorem 18). We prove that $k[D_0] \bowtie U(\mathfrak{b}_+)$ is isomorphic to \mathcal{H}_{CM} . We then construct a second bicrossproduct Hopf algebra $U_{\text{CM}} = U(\mathfrak{d}_0) \bowtie k[B_+]$ (Proposition 22) equipped with a nondegenerate dual pairing with \mathcal{H}_{CM} .

We explain carefully how the actions and coactions giving rise to these bicrossproducts can be derived from the factorisation $\text{Diff}^+(\mathbb{R}) = B_+ \bowtie D_0$. This serves as motivation and is not part of our proof. However, if we simply presented compatible actions and coactions without indicating how they arose, although no rigour would be lost this would leave things very opaque.

We note that the original Connes–Moscovici Hopf algebra $\mathcal{H}_{\text{CM}}^{\text{left}}$, defined as in (1), but with $\delta_1 \otimes Y$ rather than $Y \otimes \delta_1$ appearing in $\Delta(X)$, can also be shown to be a bicrossproduct linked to this group factorisation. The construction is given in Appendix A. However, as we explain \mathcal{H}_{CM} rather than $\mathcal{H}_{\text{CM}}^{\text{left}}$ is in an appropriate sense the natural bicrossproduct associated to this factorisation.

In the second part of the paper, we define two families of Hopf algebras, denoted $\mathcal{H}_{\text{CM}}^\lambda$, U_{CM}^λ , parameterised by $\lambda \in k$. For $\lambda \neq 0$, the corresponding element of each family is isomorphic to the bicrossproduct \mathcal{H}_{CM} respectively U_{CM} . For $\lambda = 0$ (the so-called classical limit) the Hopf algebra $\mathcal{H}_{\text{CM}}^\lambda$ is commutative, and can be realised as functions on the semidirect product $\mathbb{R}^2 \rtimes D_0$. We construct a natural quotient Hopf algebra $k_\lambda[\text{Heis}]$ of $\mathcal{H}_{\text{CM}}^\lambda$, which for $\lambda = 0$ similarly corresponds to the coordinate algebra of the Heisenberg group. For $\lambda \neq 0$ $k_\lambda[\text{Heis}]$ pairs with a Hopf subalgebra $U_\lambda(\mathfrak{heis})$ of U_{CM}^λ . By passing to an extended bicrossproduct $U(\mathfrak{d}_0) \bowtie F[B_+]_\lambda$ we give the correct classical limits of U_{CM}^λ and $U_\lambda(\mathfrak{heis})$. Finally we show that $k_\lambda[\text{Heis}]$ and $U_\lambda(\mathfrak{heis})$ are linked to a local factorisation of the group $SL_2(\mathbb{R})$, in the same way \mathcal{H}_{CM} and U_{CM} are linked to the factorisation of $\text{Diff}^+(\mathbb{R})$. We remark that locally compact quantum groups (in the von Neumann algebra setting) similar to $k_\lambda[\text{Heis}]$ and $U_\lambda(\mathfrak{heis})$ were previously constructed by Vaes [20], linked to a factorisation of the continuous Heisenberg group rather than $SL_2(\mathbb{R})$.

Finally, a bicrossproduct coacts canonically on one of its factors (the Schrödinger coaction) hence a corollary of our results is that $\mathcal{H}_{\text{CM}}^\lambda$ and $k_\lambda[\text{Heis}]$ coact canonically on $U_\lambda(\mathfrak{b}_+)$. The latter is $U(\mathfrak{b}_+)$ viewed as a noncommutative space, i.e. with scaling parameter λ introduced in such a way as to be commutative when $\lambda = 0$. This puts $\mathcal{H}_{\text{CM}}^\lambda$ and $k_\lambda[\text{Heis}]$ in the same family as the coordinate algebras of the Euclidean quantum group of [13] and the κ -Poincaré quantum group [11,17] coacting on algebras $U_\lambda(\mathfrak{b}_+^n)$ of various dimensions. In such models one is also interested in the covariant noncommutative differential geometry of the coordinate algebras of both the noncommutative space and the coacting quantum group. Thus in the final part of the paper we study low-dimensional covariant first order differential calculi over $U_\lambda(\mathfrak{b}_+)$ and $k_\lambda[\text{Heis}]$.

2. Preliminaries

In this section we recall from [15] the construction of the (left–right) bicrossproduct Hopf algebra $\mathcal{A} \blacktriangleright \triangleleft \mathcal{H}$ from Hopf algebras \mathcal{A} and \mathcal{H} , with \mathcal{A} a left \mathcal{H} -module algebra and \mathcal{H} a right \mathcal{A} -comodule coalgebra. For completeness we also give the definition of a factorisation of a group X into subgroups G and M , and the construction of a dual pair of finite-dimensional bicrossproduct Hopf algebras associated to a factorisation of a finite group (this is not used directly in our constructions of infinite-dimensional bicrossproducts, but is an important part of the motivation). We then define the Hopf algebras $k[\mathbf{D}_0]$, $U(\mathbf{d}_0)$, $U(\mathbf{b}_+)$, $k[\mathbf{B}_+]$ which we use to construct bicrossproducts. As shown by Figueroa and Gracia-Bondia [8], $k[\mathbf{D}_0]$ is isomorphic to two other well-known Hopf algebras, the comeasuring Hopf algebra of the real line \mathcal{C} [14] and the Faà di Bruno Hopf algebra \mathcal{F} . We use this to give a more convenient alternative presentation (9) of \mathcal{H}_{CM} using the generators t_n of \mathcal{C} instead of the δ_n .

2.1. Bicrossproduct Hopf algebras

Throughout this paper we work over a field k assumed to be of characteristic zero. For a Hopf algebra \mathcal{H} , we use the Sweedler notation $\Delta(x) = \sum x_{(1)} \otimes x_{(2)}$ for the coproduct. We denote a right coaction $\Delta_R: \mathcal{M} \rightarrow \mathcal{M} \otimes \mathcal{H}$ of \mathcal{H} on a k -vector space \mathcal{M} by $\Delta_R(m) = \sum m^{(1)} \otimes m^{(2)}$. Now let \mathcal{A} be an algebra and \mathcal{C} a coalgebra (over k).

Definition 2. \mathcal{A} is a left \mathcal{H} -module algebra if there exists a left action $\triangleright: \mathcal{H} \otimes \mathcal{A} \rightarrow \mathcal{A}$ such that $h \triangleright (ab) = \sum (h_{(1)} \triangleright a)(h_{(2)} \triangleright b)$ and $h \triangleright 1 = \varepsilon(h)1$, for all $h \in \mathcal{H}$ and $a, b \in \mathcal{A}$.

Definition 3. \mathcal{H} is a right \mathcal{C} -comodule coalgebra if there exists a right coaction $\Delta_R: \mathcal{H} \rightarrow \mathcal{H} \otimes \mathcal{C}$ such that for all $h \in \mathcal{H}$, $\sum \varepsilon(h^{(1)})h^{(2)} = \varepsilon(h)1$ and

$$\sum h^{(1)} \otimes h^{(2)} \otimes h^{(3)} = \sum h_{(1)} \otimes h_{(2)} \otimes h_{(1)} h_{(2)}.$$

Definition 4. We say that Hopf algebras \mathcal{H}, \mathcal{K} are dually paired (in duality) if there exists a bilinear form $\langle \cdot, \cdot \rangle: \mathcal{H} \times \mathcal{K} \rightarrow k$ such that

$$\begin{aligned} \langle a, xy \rangle &= \sum \langle a_{(1)}, x \rangle \langle a_{(2)}, y \rangle, & \langle ab, x \rangle &= \sum \langle a, x_{(1)} \rangle \langle b, x_{(2)} \rangle, \\ \langle S(a), x \rangle &= \langle a, S(x) \rangle, & \langle a, 1 \rangle &= \varepsilon(a), & \langle 1, x \rangle &= \varepsilon(x) \end{aligned}$$

for all $a, b \in \mathcal{H}$, $x, y \in \mathcal{K}$. We say that the pairing is nondegenerate if for every nonzero $a \in \mathcal{H}$, there exists some $x \in \mathcal{K}$ so that $\langle a, x \rangle \neq 0$, and for every nonzero $y \in \mathcal{K}$, there exists some $b \in \mathcal{H}$ so that $\langle b, y \rangle \neq 0$.

If \mathcal{A} and \mathcal{H} are bialgebras, with \mathcal{H} acting on \mathcal{A} , and \mathcal{A} coacting on \mathcal{H} (with action and coaction compatible in a suitable sense) then the bicrossproduct construction [12, 15] manufactures a larger bialgebra, the bicrossproduct of \mathcal{A} and \mathcal{H} , containing both \mathcal{A} and \mathcal{H} as sub-bialgebras. If \mathcal{A} and \mathcal{H} are Hopf algebras, then so is the bicrossproduct. Explicitly:

Theorem 5. [15, Theorem 6.2.2] *Let \mathcal{A} and \mathcal{H} be Hopf algebras, with \mathcal{A} a left \mathcal{H} -module algebra, and \mathcal{H} a right \mathcal{A} -comodule coalgebra, such that:*

- (1) $\varepsilon(h \triangleright a) = \varepsilon(h)\varepsilon(a)$, $\Delta(h \triangleright a) = \sum h_{(1)} \overline{(1)} \triangleright a_{(1)} \otimes h_{(1)}^{(2)}(h_{(2)} \triangleright a_{(2)})$,
- (2) $\Delta_R(1) = 1 \otimes 1$, $\Delta_R(gh) = \sum g_{(1)} \overline{(1)} h^{(1)} \otimes g_{(1)}^{(2)}(g_{(2)} \triangleright h^{(2)})$,
- (3) $\sum h_{(2)} \overline{(1)} \otimes (h_{(1)} \triangleright a)h_{(2)}^{(2)} = \sum h_{(1)} \overline{(1)} \otimes h_{(1)}^{(2)}(h_{(2)} \triangleright a)$

for all $a, b \in \mathcal{A}$, $g, h \in \mathcal{H}$. Then the vector space $\mathcal{A} \otimes \mathcal{H}$ can be given the structure of a Hopf algebra, the left–right bicrossproduct denoted $\mathcal{A} \blacktriangleleft \mathcal{H}$, via:

$$\begin{aligned} (a \otimes h)(b \otimes g) &= \sum a(h_{(1)} \triangleright b) \otimes h_{(2)}g, & S(a \otimes h) &= \sum (1 \otimes Sh \overline{(1)})(S(ah^{(2)}) \otimes 1), \\ S(a \otimes h) &= \sum (1 \otimes Sh \overline{(1)})(S(ah^{(2)}) \otimes 1). \end{aligned} \tag{5}$$

The left–right reversed result, constructing a Hopf algebra $\mathcal{H} \blacktriangleleft \mathcal{A}$ from a right \mathcal{H} -module algebra \mathcal{A} and left \mathcal{A} -comodule coalgebra \mathcal{H} is [15, Theorem 6.2.3].

2.2. Group factorisations and finite-dimensional bicrossproducts

A group X is said to factorise into subgroups G and M if group multiplication gives a set bijection $G \times M \rightarrow X$. That is, given $x \in X$, there are unique $g \in G$, $m \in M$ such that $gm = x$. We write $X = G \bowtie M$. Here for any $m \in M$, $g \in G$ there exist unique $g' \in G$, $m' \in M$ such that $mg = g'm'$. Writing $g' = m \triangleright g$, $m' = m \triangleleft g$, it is straightforward to check that this defines a natural left action \triangleright of M on G , and a natural right action \triangleleft of G on M .

For any group Γ denote by $k\Gamma$ the group algebra (over k) of Γ , with coproduct $\Delta(g) = g \otimes g$, for all $g \in \Gamma$, and (if Γ is finite) by $k[\Gamma]$ the commutative algebra of k -valued functions on Γ , with basis the delta-functions $\{\delta_g\}_{g \in \Gamma}$ and coproduct $\Delta(\delta_g) = \sum_{xy=g} \delta_x \otimes \delta_y$. Then if a finite group X factorises as $X = G \bowtie M$, we can construct two dually-paired bicrossproducts:

- (1) $k[M] \blacktriangleleft kG$. We have a compatible left action and right coaction

$$g \triangleright \delta_m := \delta_{m \triangleleft g^{-1}}, \quad g \mapsto \sum_{m \in M} (m \triangleright g) \otimes \delta_m \tag{6}$$

so we can form the bicrossproduct $k[M] \blacktriangleleft kG$, with relations

$$g\delta_m = \delta_{(m \triangleleft g^{-1})}g, \quad \Delta(\delta_m) = \sum_{xy=m} \delta_x \otimes \delta_y, \quad \Delta(g) = \sum_{m \in M} (m \triangleright g) \otimes \delta_m g.$$

- (2) $kM \blacktriangleleft k[G]$. We have a compatible right action and left coaction

$$\delta_g \triangleleft m := \delta_{m^{-1} \triangleright g}, \quad m \mapsto \sum_{g \in G} \delta_g \otimes (m \triangleleft g).$$

The bicrossproduct $kM \blacktriangleleft k[G]$ has relations

$$\delta_g m = m\delta_{m^{-1} \triangleright g}, \quad \Delta(\delta_g) = \sum_{xy=g} \delta_x \otimes \delta_y, \quad \Delta(m) = \sum_{g \in G} m\delta_x \otimes (m \triangleleft g).$$

When G and M are infinite, both these constructions fail. The challenge for factorisations of infinite groups is to construct bicrossproduct Hopf algebras analogous to those above. In Section 3 we solve this problem for the group factorisation $\text{Diff}^+(\mathbb{R}) = \mathbf{B}_+ \bowtie \mathbf{D}_0$.

2.3. The commutative Hopf subalgebra $k[\mathbf{D}_0]$

We define $k[\mathbf{D}_0]$ to be the unital commutative subalgebra of \mathcal{H}_{CM} generated by $\{\delta_n: n = 1, 2, \dots\}$, which, as shown in [5], is a Hopf subalgebra of \mathcal{H}_{CM} with $\Delta(\delta_n) = \sum_{k=1}^n D_{n,k} \otimes \delta_k$ for some $D_{n,k} \in k[\mathbf{D}_0]$. It was shown in [8] that $k[\mathbf{D}_0]$ is isomorphic to both the comeasuring Hopf algebra \mathcal{C} of the real line and the Faà di Bruno Hopf algebra \mathcal{F} , whose definitions we now recall.

Definition 6. [14] The comeasuring Hopf algebra \mathcal{C} of the real line is the commutative Hopf algebra over k generated by indeterminates $\{t_n: n = 1, 2, \dots\}$ with $t_1 = 1$, counit $\varepsilon(t_n) = \delta_{n,1}$, and coproduct

$$\Delta(t_n) = \sum_{k=1}^n \left(\sum_{i_1+\dots+i_k=n} t_{i_1} \cdots t_{i_k} \right) \otimes t_k. \tag{7}$$

Adapting results of [8], the antipode on \mathcal{C} is given by

$$S(t_{n+1}) = \sum_{\mathbf{c} \in \mathcal{S}} (-1)^{n-c_1} \frac{(2n - c_1)! c_1! t_1^{c_1} t_2^{c_2} \cdots t_{n+1}^{c_{n+1}}}{(n + 1)! c_1! c_2! \cdots c_{n+1}!}, \tag{8}$$

where $\mathcal{S} = \{(c_1, \dots, c_{n+1}): \sum_{j=1}^{n+1} c_j = n, \sum_{j=1}^{n+1} j c_j = 2n\}$.

If we rewrite Definition 6 in terms of generators $a_n = n!t_n$, this gives the usual presentation of the Faà di Bruno Hopf algebra \mathcal{F} .

Proposition 7. [8] $k[\mathbf{D}_0]$, the Faà di Bruno Hopf algebra \mathcal{F} and the comeasuring Hopf algebra of the real line \mathcal{C} are all isomorphic, via

$$\delta_n \mapsto n! \sum_{\mathbf{c} \in \mathcal{S}} (-1)^{n-c_1} \frac{(n - c_1)!}{c_2! \cdots c_{n+1}!} (t_1)^{c_1} (2t_2)^{c_2} \cdots ((n + 1)t_{n+1})^{c_{n+1}},$$

where $\mathcal{S} = \{(c_1, \dots, c_{n+1}): \sum_{j=1}^{n+1} c_j = n + 1, \sum_{j=1}^{n+1} j c_j = 2n + 1\}$, and

$$(n + 1)t_{n+1} \mapsto \sum_{c_1+2c_2+\dots+nc_n=n} \frac{\delta_1^{c_1} \cdots \delta_n^{c_n}}{c_1! \cdots c_n! (1!)^{c_1} \cdots (n!)^{c_n}}.$$

In the sequel, we will use the presentation of $k[\mathbf{D}_0]$ as the commutative Hopf algebra with generators $\{t_n\}_{n \geq 1}$, and coproduct and antipode given by (7), (8).

Lemma 8. $k[\mathbf{D}_0]$ is an \mathbb{N} -graded Hopf algebra, via the grading defined on monomials by $|t_{n_1} \cdots t_{n_A}| = n_1 + \cdots + n_A - A$.

Proof. Let $k[\mathbf{D}_0]_N$ be the linear span of monomials $\mathbf{t} = t_{n_1} \cdots t_{n_A}$ with $|\mathbf{t}| = N$. Then $k[\mathbf{D}_0]_m k[\mathbf{D}_0]_n \subseteq k[\mathbf{D}_0]_{m+n}$ for all m, n . From (8) $S(k[\mathbf{D}_0]_N) \subseteq k[\mathbf{D}_0]_N$. It also follows from (7) that $\Delta(k[\mathbf{D}_0]_N) \subseteq \bigoplus_{n=0}^N k[\mathbf{D}_0]_n \oplus k[\mathbf{D}_0]_{N-n}$. \square

As a corollary of Proposition 7, we have:

Corollary 9. *In terms of the t_n , the presentation (1) of \mathcal{H}_{CM} becomes:*

$$\begin{aligned} [Y, X] &= X, & [X, t_n] &= (n + 1)t_{n+1} - 2t_2 t_n, & [Y, t_n] &= (n - 1)t_n, \\ \Delta(X) &= X \otimes 1 + 1 \otimes X + Y \otimes 2t_2, & \Delta(Y) &= Y \otimes 1 + 1 \otimes Y, & \varepsilon(t_n) &= \delta_{n,1}, \\ S(X) &= -X + 2Yt_2, & S(Y) &= -Y, & \varepsilon(X) &= 0 = \varepsilon(Y) \end{aligned} \tag{9}$$

with $\Delta(t_n), S(t_n)$ given by (7), (8).

Taking $k = \mathbb{R}$ or \mathbb{C} , the generators δ_n, t_n can be realised as functions on \mathbf{D}_0 :

$$\delta_n(f) = [\log f']^{(n)}(0), \quad t_n(f) = \frac{1}{n!} f^{(n)}(0), \quad f \in \mathbf{D}_0. \tag{10}$$

2.4. *The Hopf algebra $U(\mathbf{d}_0)$*

Definition 10. We define \mathbf{d}_+ to be the Lie algebra (over k) with countably many generators $\{z_n\}_{n \in \mathbb{N}}$ and relations $[z_m, z_n] = (n - m)z_{m+n-1}$ for all m, n . Define \mathbf{d}_0 to be the Lie subalgebra generated by $\{z_n\}_{n \geq 2}$. Then $U(\mathbf{d}_0)$ is the universal enveloping algebra of \mathbf{d}_0 with canonical Hopf structure.

Note that \mathbf{d}_+ has a natural representation as differential operators $z_n = x^n \frac{d}{dx}$ acting on the unital algebra $k[x]$ of polynomials in a single indeterminate x .

Lemma 11. *$U(\mathbf{d}_0)$ is an \mathbb{N} -graded Hopf algebra, via the grading defined on monomials by $|1| = 0, |z_{m_1} \cdots z_{m_p}| = m_1 + \cdots + m_p - p$.*

Proof. Denote by $U(\mathbf{d}_0)_N$ the linear span of monomials of degree N . Since $|z_m z_n| = m + n - 2 = |z_n z_m| = |z_{m+n-1}|$, then $U(\mathbf{d}_0)_m U(\mathbf{d}_0)_n \subseteq U(\mathbf{d}_0)_{m+n}$ for all $m, n \in \mathbb{N}$. Further, as $\Delta(z_n) = z_n \otimes 1 + 1 \otimes z_n$, it follows that $\Delta(U(\mathbf{d}_0)_N) \subseteq \bigoplus_{n=0}^N U(\mathbf{d}_0)_n \otimes U(\mathbf{d}_0)_{N-n}$. Finally $S(U(\mathbf{d}_0)_N) \subseteq U(\mathbf{d}_0)_N$. \square

Proposition 12. *There is a nondegenerate dual pairing (in the sense of Definition 4) of the Hopf algebras $U(\mathbf{d}_0)$ and $k[\mathbf{D}_0]$, defined on generators by*

$$\langle z_m, t_n \rangle = \delta_{m,n} \quad \forall m \geq 2, n \geq 1, \tag{11}$$

equivalently by $\langle z_m, \delta_n \rangle = m! \delta_{m,n+1}$. The pairing satisfies

$$\langle z_{m_1} \cdots z_{m_p}, t_n \rangle = \begin{cases} \prod_{j=1}^{p-1} (n + j - 1 - \sum_{l=1}^j m_l), & \sum_{j=1}^p m_j = n + p - 1, \\ 0, & \text{otherwise,} \end{cases}$$

$$\langle z_m, t_{n_1} \cdots t_{n_A} \rangle = \begin{cases} 1, & \{n_1, \dots, n_A\} = \{m, 1, \dots, 1\} \text{ as sets,} \\ 0, & \text{otherwise.} \end{cases} \tag{12}$$

Proof. Assuming the pairing is well-defined, the identities follow by a straightforward induction. For example,

$$\langle z_m, t_{n_1} \cdots t_{n_A} \rangle = \langle \Delta^{A-1}(z_m), t_{n_1} \otimes \cdots \otimes t_{n_A} \rangle = \sum_{l=1}^A \varepsilon(t_{n_l}) \cdots \langle z_m, t_{n_l} \rangle \cdots \varepsilon(t_{n_A}).$$

To check well-defined, as $\langle z_m z_n, t_{n_1} \cdots t_{n_A} \rangle = \langle \Delta^{A-1}(z_m z_n), t_{n_1} \otimes \cdots \otimes t_{n_A} \rangle$ it suffices to check $\langle z_m z_n, t_p \rangle$. By the above, $\langle z_m z_n, t_p \rangle = n \delta_{m+n, p+1}$. So $\langle z_m z_n - z_n z_m, t_p \rangle = (n - m) \delta_{m+n-1, p} = (n - m) \langle z_{m+n-1}, t_p \rangle$.

It follows from (8), (12) that $\langle S(z_m), t_n \rangle = \langle z_m, S(t_n) \rangle$ for all m, n , so $\langle S(z_m), \mathbf{t} \rangle = \langle z_m, S(\mathbf{t}) \rangle$ for $\mathbf{t} = t_{n_1} \cdots t_{n_A}$ hence for all $\mathbf{t} \in k[\mathbb{B}_+]$. Then

$$\begin{aligned} \langle S(z_{m_1} \cdots z_{m_p}), \mathbf{t} \rangle &= \langle S(z_{m_p}) \otimes \cdots \otimes S(z_{m_1}), \Delta^{p-1}(\mathbf{t}) \rangle \\ &= \langle z_{m_1} \otimes \cdots \otimes z_{m_p}, \Delta^{p-1}(S(\mathbf{t})) \rangle = \langle z_{m_1} \cdots z_{m_p}, S(\mathbf{t}) \rangle. \end{aligned}$$

To show nondegeneracy, we need the \mathbb{N} -gradings of Lemmas 8 and 11.

Lemma 13. For $\mathbf{z} = z_{m_1}^{a_1} \cdots z_{m_p}^{a_p}$ and $\mathbf{t} = t_{n_1} \cdots t_{n_A}$, with $2 \leq m_1 < m_2 < \cdots < m_p$, $2 \leq n_1 \leq n_2 \leq \cdots \leq n_A$, and $a_1, \dots, a_p, A \geq 1$, then:

- (1) $\langle \mathbf{z}, \mathbf{t} \rangle = 0$ if $A > a_1 + \cdots + a_p$.
- (2) $\langle \mathbf{z}, \mathbf{t} \rangle = 0$ unless $|\mathbf{z}| = |\mathbf{t}|$, i.e. unless $a_1 m_1 + \cdots + a_p m_p = n_1 + \cdots + n_A$.
- (3) If $A = a_1 + \cdots + a_p$, $\langle \mathbf{z}, \mathbf{t} \rangle = a_1! \cdots a_p! \delta_{m_1, n_1} \cdots \delta_{m_1, n_{a_1}} \delta_{m_2, n_{a_1+1}} \cdots \delta_{m_p, n_A}$.

Proof. For part (1), using $\langle \mathbf{z}, \mathbf{t} \rangle = \langle \Delta^{A-1}(\mathbf{z}), t_{n_1} \otimes \cdots \otimes t_{n_A} \rangle$, if $A > a_1 + \cdots + a_p$ then every term in $\Delta^{A-1}(\mathbf{z})$ will contain at least one component $- \otimes 1 \otimes -$, which pairs to zero with the corresponding t_{n_k} . Hence $\langle \mathbf{z}, \mathbf{t} \rangle = 0$.

As $k[\mathbb{D}_0]$ and $U(\mathfrak{d}_0)$ are both \mathbb{N} -graded, part (2) follows using (12).

We prove part (3) by induction. It holds for $p = A = 1$. Suppose it holds for \mathbf{z}, \mathbf{t} . Then for $m_p \leq m_{p+1}, n_A \leq n_{A+1}$, we have

$$\begin{aligned} \langle \mathbf{z} z_{m_{p+1}}, \mathbf{t} t_{n_{A+1}} \rangle &= \langle \mathbf{z} \otimes z_{m_{p+1}}, \Delta(\mathbf{t} t_{n_{A+1}}) \rangle = \langle \mathbf{z} \otimes z_{m_{p+1}}, \sum \cdots \otimes t_{i_1} \cdots t_{i_{A+1}} \rangle \\ &= \sum_{k=1}^A \langle \mathbf{z}, t_{n_1} \cdots \hat{t}_{n_k} \cdots t_{n_{A+1}} \rangle \delta_{m_{p+1}, n_k} + \langle \mathbf{z}, \mathbf{t} \rangle \delta_{m_{p+1}, n_{A+1}} \end{aligned} \tag{13}$$

using (12). The first A terms contain $\delta_{m_p, n_{A+1}} \delta_{m_{p+1}, n_k}$. If $m_p < m_{p+1}$ this is zero, so $\langle \mathbf{z} z_{m_{p+1}}, \mathbf{t} t_{n_{A+1}} \rangle = \langle \mathbf{z}, \mathbf{t} \rangle \delta_{m_{p+1}, n_{A+1}}$. If $m_p = m_{p+1}$, then (13) becomes

$$\sum_{k=A-a_p+1}^A \langle \mathbf{z}, t_{n_1} \cdots \hat{t}_{n_k} \cdots t_{n_{A+1}} \rangle \delta_{m_p, n_k} + \langle \mathbf{z}, \mathbf{t} \rangle \delta_{m_p, n_{A+1}} = (a_p + 1) \langle \mathbf{z}, \mathbf{t} \rangle \delta_{m_p, n_{A+1}},$$

which completes the inductive step. \square

We can now prove nondegeneracy. Given $\mathbf{z} = \sum \lambda(\mathbf{m}, \mathbf{a}) z_{m_1}^{a_1} \cdots z_{m_p}^{a_p}$ such that $\langle \mathbf{z}, \mathbf{t} \rangle = 0$ for all $\mathbf{t} \in k[\mathbf{D}_0]$, by Lemma 13, part (2) we can restrict to $\mathbf{z} \in U(\mathbf{d}_0)_N$ for some N . Define $A = \max\{a_1 + \cdots + a_p\}$ taken over monomials occurring in \mathbf{z} . By Lemma 13, part (1),

$$\langle \mathbf{z}, \mathbf{t} \rangle = \left\langle \sum_{\mathbf{m}, \mathbf{a}: a_1 + \cdots + a_p = A} \lambda(\mathbf{m}, \mathbf{a}) z_{m_1}^{a_1} \cdots z_{m_p}^{a_p}, t_{n_1} \cdots t_{n_A} \right\rangle.$$

So by Lemma 13, part (3), choosing \mathbf{t} appropriately gives $\lambda(\mathbf{m}, \mathbf{a}) = 0$ for all such \mathbf{m}, \mathbf{a} . Nondegeneracy follows, completing the proof of Proposition 12. \square

2.5. The Hopf algebras $U(\mathfrak{b}_+)$ and $k[\mathbf{B}_+]$

The group \mathbf{B}_+ was defined in (4). Let \mathfrak{b}_+ be the Lie algebra (over k) generated by X, Y satisfying the relation $[Y, X] = X$. We now define a commutative Hopf algebra $k[\mathbf{B}_+]$ and a nondegenerate dual pairing of $k[\mathbf{B}_+]$ and $U(\mathfrak{b}_+)$.

Definition 14. $k[\mathbf{B}_+]$ is the commutative Hopf algebra (over k) generated by elements $\alpha^{\pm 1}, \beta$ satisfying

$$\begin{aligned} \Delta(\alpha) &= \alpha \otimes \alpha, & \Delta(\beta) &= \alpha \otimes \beta + \beta \otimes 1, \\ \varepsilon(\alpha) &= 1, & \varepsilon(\beta) &= 0, & S(\alpha) &= \alpha^{-1}, & S(\beta) &= -\alpha^{-1}\beta. \end{aligned}$$

Lemma 15. There is a unique nondegenerate dual pairing of the Hopf algebras $U(\mathfrak{b}_+)$ and $k[\mathbf{B}_+]$, defined on generators by

$$\langle X, \alpha \rangle = 0, \quad \langle X, \beta \rangle = 1 = \langle Y, \alpha \rangle, \quad \langle Y, \beta \rangle = 0.$$

This satisfies

$$\langle X^j Y^k, \alpha^t \beta^r \rangle = j! \delta_{j,r} t^k \quad \forall j, k, r \in \mathbb{N}, t \in \mathbb{Z}, \tag{14}$$

where we use the convention $X^0 = 1 = Y^0 = \alpha^0 = \beta^0, 0! = 1$, and $0^0 = 1$.

Proof. This is more straightforward than the proof of Proposition 12, and is in fact well known. We give the details for completeness. For $t \geq 1, \langle Y, \alpha^t \rangle = \sum_{i=1}^t \langle 1 \otimes \cdots \otimes Y \cdots \otimes 1, \alpha^{\otimes t} \rangle = t$. We also have $\langle Y, \alpha^{-1} \rangle = \langle Y, S(\alpha) \rangle = \langle S(Y), \alpha \rangle = -\langle Y, \alpha \rangle = -1$, and in fact $\langle Y, \alpha^t \rangle = t$ for all $t \in \mathbb{Z}$. Then $\langle Y^k, \alpha^t \rangle = \langle Y^{\otimes k}, \Delta^{k-1}(\alpha^t) \rangle = \langle Y, \alpha^t \rangle^k = t^k$. Suitably interpreted, this holds also for $k = 0$. In the same way, $\langle Y^k, \beta^r \rangle = \delta_{k,0} \delta_{r,0}$, for all $k, r \geq 0$. So $\langle Y^k, \alpha^t \beta^r \rangle = \delta_{r,0} t^k$. Further, $\langle X^j, \alpha^t \rangle = \delta_{j,0}$ and $\langle X^j \beta^r \rangle = j! \delta_{j,r}$ for all $j, r, t \geq 0$, hence $\langle X^j, \alpha^t \beta^r \rangle = j! \delta_{j,r}$. Using $\langle X^j Y^k, \alpha^t \beta^r \rangle = \langle X^j \otimes Y^k, \Delta(\alpha^t \beta^r) \rangle$ the result follows. It is also straightforward to check that $\langle S(X^j Y^k), \alpha^t \beta^r \rangle = \langle X^j Y^k, S(\alpha^t \beta^r) \rangle$.

To check well-defined, we have $\langle XY, \alpha^t \beta^r \rangle = \delta_{1,r} t$, and

$$\begin{aligned} \langle YX, \alpha^t \beta^r \rangle &= \left\langle Y \otimes X, (\alpha^t \otimes \alpha^t) \left(\sum_{s=0}^r \binom{r}{s} \alpha^s \beta^{r-s} \otimes \beta^s \right) \right\rangle \\ &= \sum_{s=0}^r \binom{r}{s} \delta_{0,r-s} (s+t) \delta_{1,s} = \delta_{1,r} (t+1). \end{aligned}$$

Hence $\langle YX - XY, \alpha^t \beta^r \rangle = \delta_{1,r} = \langle X, \alpha^t \beta^r \rangle$. Finally, given (14), nondegeneracy of the pairing is immediate. \square

3. Bicrossproduct structure of the Connes–Moscovici Hopf algebra

We prove that $k[D_0]$ can be given the structure of a left $U(\mathfrak{b}_+)$ -module algebra, and $U(\mathfrak{b}_+)$ the structure of a right $k[D_0]$ -comodule coalgebra, with action and coaction compatible in the sense of Theorem 5. This enables the construction of a bicrossproduct Hopf algebra $k[D_0] \blacktriangleright U(\mathfrak{b}_+)$, which we prove is isomorphic to the Connes–Moscovici Hopf algebra \mathcal{H}_{CM} . We also construct a second bicrossproduct $U_{CM} := U(\mathfrak{d}_0) \blacktriangleleft k[B_+]$, equipped with a nondegenerate dual pairing with \mathcal{H}_{CM} . We explain how these bicrossproducts are linked to the factorisation of the group $\text{Diff}^+(\mathbb{R})$ into the subgroups B_+ and D_0 . The factorisation argument is not part of our proof, but rather serves as motivation.

3.1. The bicrossproduct $k[D_0] \blacktriangleright U(\mathfrak{b}_+)$

Lemma 16. $k[D_0]$ is a left $U(\mathfrak{b}_+)$ -module algebra via the action defined by

$$X \triangleright t_n = (n + 1)t_{n+1} - 2t_2 t_n, \quad Y \triangleright t_n = (n - 1)t_n \tag{15}$$

equivalently defined by $X \triangleright \delta_n = \delta_{n+1}$, $Y \triangleright \delta_n = n\delta_n$.

Proof. As $k[D_0]$ is commutative and $U(\mathfrak{b}_+)$ cocommutative, it is easy to check that (15) extends to a well-defined left action of $U(\mathfrak{b}_+)$ on $k[D_0]$. For example,

$$\begin{aligned} Y \triangleright (X \triangleright t_n) &= Y \triangleright [(n + 1)t_{n+1} - 2t_2 t_n] = (n^2 + n)t_{n+1} - 2nt_2 t_n, \\ X \triangleright (Y \triangleright t_n) &= (n - 1)X \triangleright t_n = (n^2 - n)t_{n+1} - 2(n - 1)t_2 t_n. \end{aligned}$$

Hence $(YX - XY) \triangleright t_n = (n + 1)t_{n+1} - 2t_2 t_n = X \triangleright t_n$. The action on the δ_n follows from Proposition 7. \square

Lemma 17. $U(\mathfrak{b}_+)$ is a right $k[D_0]$ -comodule coalgebra, via the coaction defined on generators by

$$\Delta_R(X) = X \otimes 1 + Y \otimes 2t_2, \quad \Delta_R(Y) = Y \otimes 1, \tag{16}$$

and extended by $\Delta_R(gh) = \sum g_{(1)} \overline{(1)} h \overline{(1)} \otimes g_{(1)}^{(2)} (g_{(2)} \triangleright h^{(2)})$.

Proof. We check that these formulae define a coaction. Suppose $g \in U(\mathfrak{b}_+)$ satisfies $(\text{id} \otimes \Delta) \Delta_R(g) = (\Delta_R \otimes \text{id}) \Delta_R(g)$ (this holds for $g = X, Y$). Then

$$\begin{aligned} \Delta_R(gY) &= \sum g_{(1)} \overline{(1)} Y \overline{(1)} \otimes g_{(1)}^{(2)} (g_{(2)} \triangleright Y^{(2)}) = \sum g \overline{(1)} Y \otimes g^{(2)} \\ \Rightarrow (\Delta_R \otimes \text{id}) \Delta_R(gY) &= \sum g \overline{(1)} \overline{(1)} Y \otimes g^{(1)(2)} \otimes g^{(2)} \\ &= \sum g \overline{(1)} Y \otimes g^{(2)}_{(1)} \otimes g^{(2)}_{(2)} = (\text{id} \otimes \Delta) \Delta_R(gY) \end{aligned}$$

and in the same way $(\Delta_R \otimes \text{id})\Delta_R(Xg) = (\text{id} \otimes \Delta)\Delta_R(Xg)$. We check that Δ_R is well-defined. We have

$$\begin{aligned} \Delta_R(YX) &= \sum Y_{(1)} \overline{(1)} X \overline{(1)} \otimes Y_{(1)}^{(2)} (Y_{(2)} \triangleright X^{(2)}) = YX \otimes 1 + (Y^2 + Y) \otimes 2t_2, \\ \Delta_R(XY) &= XY \otimes 1 + Y^2 \otimes 2t_2. \end{aligned}$$

Hence $\Delta_R(YX - XY) = (YX - XY) \otimes 1 + Y \otimes 2t_2 = \Delta_R(X)$. \square

Theorem 18. *The left action (15) and right coaction (16) are compatible in the sense of Theorem 5.*

Proof. We check conditions (1)–(3) of Theorem 5. For (1), to show $\varepsilon(h \triangleright a) = \varepsilon(h)\varepsilon(a)$, it is enough to show $\varepsilon(X \triangleright a) = 0 = \varepsilon(Y \triangleright a)$ for all a . Using the \mathbb{N} -grading of $k[\mathbb{D}_0]$ (Lemma 8) we see that $X \triangleright k[\mathbb{D}_0]_N \subseteq k[\mathbb{D}_0]_{N+1}$, $Y \triangleright k[\mathbb{D}_0]_N \subseteq k[\mathbb{D}_0]_N$, and $\varepsilon(k[\mathbb{D}_0]_N) = 0$ for $N \geq 1$. We also need to check that

$$\Delta(h \triangleright a) = \sum h_{(1)} \overline{(1)} \triangleright a_{(1)} \otimes h_{(1)}^{(2)} (h_{(2)} \triangleright a_{(2)}) \tag{17}$$

for all $h \in U(\mathfrak{b}_+)$, $a \in k[\mathbb{D}_0]$. Suppose for fixed h, k (17) holds for all a . Then

$$\begin{aligned} \Delta(hk \triangleright a) &= \Delta(h \triangleright (k \triangleright a)) = \sum h_{(1)} \overline{(1)} \triangleright (k \triangleright a)_{(1)} \otimes h_{(1)}^{(2)} (h_{(2)} \triangleright (k \triangleright a)_{(2)}) \\ &= \sum (h_{(1)} \overline{(1)} k_{(1)} \overline{(1)}) \triangleright a_{(1)} \otimes h_{(1)}^{(2)} (h_{(2)(1)} \triangleright k_{(1)}^{(2)}) (h_{(2)(2)} k_{(2)} \triangleright a_{(2)}). \end{aligned}$$

Now,

$$\begin{aligned} (\Delta_R \otimes \text{id})\Delta(hk) &= \sum (hk)_{(1)} \overline{(1)} \otimes (hk)_{(1)}^{(2)} \otimes (hk)_{(2)} \\ &= \sum h_{(1)(1)} \overline{(1)} k_{(1)} \overline{(1)} \otimes h_{(1)(1)}^{(2)} (h_{(1)(2)} \triangleright k_{(1)}^{(2)}) \otimes h_{(2)} k_{(2)} \\ &= \sum h_{(1)} \overline{(1)} k_{(1)} \overline{(1)} \otimes h_{(1)}^{(2)} (h_{(2)(1)} \triangleright k_{(1)}^{(2)}) \otimes h_{(2)(2)} k_{(2)} \\ \Rightarrow \Delta(hk \triangleright a) &= \sum (hk)_{(1)} \overline{(1)} \triangleright a_{(1)} \otimes (hk)_{(1)}^{(2)} ((hk)_{(2)} \triangleright a_{(2)}). \end{aligned}$$

So it will be enough to check $h = X, Y$ only. It is straightforward to check that if (17) holds for $\Delta(X \triangleright a), \Delta(X \triangleright b)$, then it holds for $\Delta(X \triangleright ab)$, and similarly for Y . Hence we need only check $X \triangleright t_n, Y \triangleright t_n$. Now, $\Delta(Y \triangleright t_n) = (n - 1)\Delta(t_n)$, whereas the right-hand side of (17) is

$$\begin{aligned} &\sum Y_{(1)} \overline{(1)} \triangleright t_{n(1)} \otimes Y_{(1)}^{(2)} (Y_{(2)} \triangleright t_{n(2)}) \\ &= \sum [Y \triangleright (t_n)_{(1)} \otimes (t_n)_{(2)} + (t_n)_{(1)} \otimes Y \triangleright (t_n)_{(2)}] \\ &= \sum_{k=1}^n \sum_{i_1 + \dots + i_k = n} [Y \triangleright (t_{i_1} \dots t_{i_k}) \otimes t_k + t_{i_1} \dots t_{i_k} \otimes Y \triangleright t_k] \\ &= \sum_{k=1}^n \sum_{i_1 + \dots + i_k = n} (i_1 + \dots + i_k - 1) t_{i_1} \dots t_{i_k} \otimes t_k = (n - 1)\Delta(t_n) = \Delta(Y \triangleright t_n). \end{aligned}$$

In the same way, we check that (17) holds for $\Delta(X \triangleright t_2)$, with the general case $\Delta(X \triangleright t_n)$ following by induction. Hence condition (1) of Theorem 5 holds.

Condition (2) is automatic from the definition of Δ_R . Finally, since $k[D_0]$ is commutative and $U(\mathfrak{b}_+)$ cocommutative, condition (3) is immediate. \square

All conditions of Theorem 5 hold, so we can construct the left–right bicrossproduct Hopf algebra $k[D_0] \blacktriangleright U(\mathfrak{b}_+)$, which we can think of as an analogue of the finite-dimensional bicrossproduct $k[M] \blacktriangleright kG$ defined in Section 2.2.

Theorem 19. \mathcal{H}_{CM} and $k[D_0] \blacktriangleright U(\mathfrak{b}_+)$ are isomorphic Hopf algebras.

Proof. Using (5), it follows that $k[D_0] \blacktriangleright U(\mathfrak{b}_+)$ has generators X, Y, t_n with relations coinciding exactly with the presentation (9) of \mathcal{H}_{CM} . \square

Finally, we remark that the original codimension one Connes–Moscovici Hopf algebra $\mathcal{H}_{CM}^{\text{left}}$, which differs from \mathcal{H}_{CM} only in that $Y \otimes \delta_1$ is replaced by $\delta_1 \otimes Y$ in (1), is isomorphic to a right–left bicrossproduct $U(\mathfrak{b}_+) \blacktriangleleft k[D_0]$ which is also linked to the factorisation of $\text{Diff}^+(\mathbb{R})$, but in a less natural way than \mathcal{H}_{CM} . For completeness, this is outlined (without proofs) in Appendix A.

3.2. Relation to group factorisation I

We motivate the above bicrossproduct constructions using a factorisation of the group $\text{Diff}^+(\mathbb{R})$ of orientation preserving diffeomorphisms of the real line (2). As shown in [5], the factorisation $\text{Diff}^+(\mathbb{R}) = B_+ \triangleright D_0$ is as follows. Given $\varphi \in \text{Diff}^+(\mathbb{R})$, we have $\varphi = (a, b) \circ \phi$ for unique $(a, b) \in B_+, \phi \in D_0$, with

$$(a, b) = (\varphi'(0), \varphi(0)), \quad \phi(x) = \frac{\varphi(x) - \varphi(0)}{\varphi'(0)} \quad \forall x \in \mathbb{R}. \tag{18}$$

Since $(\phi \circ (a, b))(x) = \phi(ax + b)$, the corresponding left action of D_0 on B_+ and right action of B_+ on D_0 are given by

$$\phi \triangleright (a, b) = (a\phi'(b), \phi(b)), \quad (\phi \triangleleft (a, b))(x) = \frac{\phi(ax + b) - \phi(b)}{a\phi'(b)}. \tag{19}$$

We identify $X, Y \in \mathfrak{b}_+$ with the matrices $\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$. By slight abuse of notation, for any $s \in \mathbb{R}$ denote by e^{sX}, e^{sY} the elements $(1, s), (e^s, 0)$ of B_+ .

To understand the origin of (15), consider the factorisation (18) of $\text{Diff}^+(\mathbb{R})$. For any function $\xi : D_0 \rightarrow k$, using (19) we define a left action of B_+ via $((a, b) \triangleright \xi)(\phi) = \xi(\phi \triangleleft (a, b))$, and (taking $k = \mathbb{R}$ or \mathbb{C}) by differentiation a left action of $U(\mathfrak{b}_+)$ on $k[D_0]$. So $(\phi \triangleleft e^{sX})(x) = \frac{\phi(x+s) - \phi(s)}{\phi'(s)}$, then (10) gives

$$\begin{aligned} t_n(\phi \triangleleft e^{sX}) &= \frac{\phi^{(n)}(s)}{n!\phi'(s)} = \frac{[\phi^{(n)}(0) + s\phi^{(n+1)}(0) + O(s^2)]}{n![\phi'(0) + s\phi''(0) + O(s^2)]} \\ &= \frac{1}{n!}[\phi^{(n)}(0) - s\phi''(0)\phi^{(n)}(0) + s\phi^{(n+1)}(0)] + O(s^2) \end{aligned}$$

$$\begin{aligned} \Rightarrow (X \triangleright t_n)(\phi) &= \left. \frac{d}{ds} \right|_{s=0} t_n(\phi \triangleleft e^{sX}) = \frac{1}{n!} [-\phi''(0)\phi^{(n)}(0) + \phi^{(n+1)}(0)] \\ &= [(n+1)t_{n+1} - 2t_2t_n](\phi) \end{aligned}$$

giving $X \triangleright t_n = (n+1)t_{n+1} - 2t_2t_n$. Similarly, $\delta_n(\phi \triangleleft e^{sX}) = g^{(n)}(0)$, where

$$g(x) = \log(\phi \triangleleft e^{sX})'(x) = \log \phi'(x+s) - \log \phi'(s).$$

Hence $g^{(n)}(x) = h^{(n)}(x+s)$, where $h(x) = \log \phi'(x)$, so $h^{(n)}(0) = \delta_n(\phi)$ for all $n \geq 1$. So $g^{(n)}(0) = h^{(n)}(s) = h^{(n)}(0) + sh^{(n+1)}(0) + O(s^2)$. Thus,

$$(X \triangleright \delta_n)(\phi) = \left. \frac{d}{ds} \right|_{s=0} \delta_n(\phi \triangleleft e^{sX}) = h^{(n+1)}(0) = \delta_{n+1}(\phi).$$

So $X \triangleright \delta_n = \delta_{n+1}$. The formulae for Y follow similarly. So using the group factorisation we recover (15), which we already showed to be a left action.

Next we explain how the coaction (16) can be recovered from the factorisation (18). For any group factorisation $X = G \bowtie M$ we define a k -linear map

$$\tilde{\Delta}_R : kG \rightarrow k[M, kG], \quad \tilde{\Delta}_R(g)(m) = m \triangleright g,$$

where $k[M, kG]$ is the k -vector space of maps $M \rightarrow kG$. If X is finite then $k[M, kG] \cong kG \otimes k[M]$ as vector spaces, and $\tilde{\Delta}_R$ is the right coaction (6). In our situation, taking $k = \mathbb{R}$ then for any $s \in \mathbb{R}, \phi \in D_0$,

$$\tilde{\Delta}_R(e^{sX})(\phi) = \phi \triangleright e^{sX} = \phi \triangleright (1, s) = (\phi'(s), \phi(s)).$$

We induce a linear map $\Delta_R : U(\mathfrak{b}_+) \rightarrow \mathbb{R}[D_0, U(\mathfrak{b}_+)]$ by differentiation:

$$\Delta_R(X)(\phi) = \left. \frac{d}{ds} \right|_{s=0} \tilde{\Delta}_R(e^{sX})(\phi) = (\phi'(0), \phi'(0)) + (\phi''(0), \phi(0)) = Xt_1(\phi) + Y2t_2(\phi).$$

So we can identify $\Delta_R(X)$ with $X \otimes t_1 + Y \otimes 2t_2 \in U(\mathfrak{b}_+) \otimes \mathbb{R}[D_0]$. As $t_1 = 1$ we retrieve (16). Further, $\tilde{\Delta}_R(e^{sY})(\phi) = \phi \triangleright (e^s, 0) = (e^s \phi'(0), \phi(0))$. So

$$\Delta_R(Y)(\phi) = \left. \frac{d}{ds} \right|_{s=0} \tilde{\Delta}_R(e^{sY})(\phi) = (\phi'(0), \phi(0)) = Yt_1(\phi).$$

We identify $\Delta_R(Y)$ with $Y \otimes t_1 = Y \otimes 1$. So (working with $k = \mathbb{R}$) we recover (16), which as we already showed defines a right coaction (for general k).

3.3. The bicrossproduct U_{CM}

We now manufacture a second bicrossproduct $U_{CM} := U(\mathfrak{d}_0) \blacktriangleleft k[B_+]$, which we equip with a nondegenerate dual pairing with \mathcal{H}_{CM} . The action and coaction used to construct the bicrossproduct can again be motivated by considering the group factorisation (18) of $\text{Diff}^+(\mathbb{R})$. First of all:

Lemma 20. $k[\mathbf{B}_+]$ is a right $U(\mathbf{d}_0)$ -module algebra via the action defined by

$$\alpha \triangleleft z_n = n\alpha\beta^{n-1}, \quad \beta \triangleleft z_n = -\beta^n \quad (n \geq 2). \tag{20}$$

Proof. It is enough to check that the action (20) defined on generators is compatible with the algebra relations. For example,

$$(\alpha \triangleleft z_m) \triangleleft z_n = m(\alpha\beta^{m-1}) \triangleleft z_n = m(n - m + 1)\alpha\beta^{m+n-2}.$$

Hence $\alpha \triangleleft (z_m z_n - z_n z_m) = (n - m)(m + n - 1)\alpha\beta^{m+n-2} = \alpha \triangleleft [z_m, z_n]$. \square

Lemma 21. $U(\mathbf{d}_0)$ is a left $k[\mathbf{B}_+]$ -comodule coalgebra via the coaction defined on generators by

$$\Delta_L(z_n) = \sum_{j=2}^n \binom{n}{j} \alpha^{j-1} \beta^{n-j} \otimes z_j \tag{21}$$

and extended to all of $U(\mathbf{d}_0)$ via $\Delta_L(hg) = \sum (h^{(0)} \triangleleft g_{(1)}) g_{(2)}^{(0)} \otimes h^{(\overline{1})} g_{(2)}^{(\overline{1})}$.

Proof. In the same way as Lemma 17 one can check that these formulae extend to a left coaction. In particular, it is straightforward that $(\Delta \otimes \text{id})\Delta_L(z_n) = (\text{id} \otimes \Delta_L)\Delta_L(z_n)$, and $\Delta_L(z_m z_n) - \Delta_L(z_n z_m) = (n - m)\Delta_L(z_{m+n-1})$. \square

As in Theorem 18 it can be checked that the right action (20) and left coaction (21) are compatible in the sense of [15, Theorem 6.2.3]. Then:

Proposition 22. The bicrossproduct Hopf algebra $U_{\text{CM}} := U(\mathbf{d}_0) \bowtie k[\mathbf{B}_+]$ has generators z_n ($n \geq 2$), $\alpha^{\pm 1}$, β with relations $[\alpha, \beta] = 0$,

$$\begin{aligned} [z_m, z_n] &= (n - m)z_{m+n-1}, & [z_n, \alpha] &= -n\alpha\beta^{n-1}, & [z_n, \beta] &= \beta^n, & \Delta(\alpha) &= \alpha \otimes \alpha, \\ \Delta(\beta) &= \alpha \otimes \beta + \beta \otimes 1, & \Delta(z_n) &= z_n \otimes 1 + \sum_{j=2}^n \binom{n}{j} \alpha^{j-1} \beta^{n-j} \otimes z_j \end{aligned} \tag{22}$$

with antipode and counit defined accordingly.

U_{CM} is an analogue of the finite-dimensional bicrossproduct $kM \bowtie k[G]$ of Section 2.2. By the general theory of bicrossproduct Hopf algebras [15]:

Theorem 23. There is a nondegenerate dual pairing of U_{CM} , \mathcal{H}_{CM} , given by

$$\langle \mathbf{z}\xi, \mathbf{t}x \rangle := \langle \mathbf{z}, \mathbf{t} \rangle \langle x, \xi \rangle \tag{23}$$

for all $\mathbf{z} = z_{m_1} \cdots z_{m_p} \in U(\mathbf{d}_0)$, $\xi = \alpha^i \beta^j \in k[\mathbf{B}_+]$, $\mathbf{t} = t_{n_1} \cdots t_{n_A} \in k[\mathbf{D}_0]$, $x = X^r Y^s \in U(\mathbf{b}_+)$, where on the right-hand side we use the pairings (11), (14).

3.4. Relation to group factorisation II

As in Section 3.2, we explain how the right action (20) and left coaction (21) used in the construction of U_{CM} can be recovered from the factorisation (18). Again this is background and not part of the proof.

Recall that the Lie algebra \mathfrak{d}_+ can be represented as differential operators $z_n = x^n \frac{d}{dx}$. Taking $k = \mathbb{R}$, each z_n gives a flow on \mathbb{R} by solving the ODE $x'(t) = x(t)^n$. For z_0 , $x(t) = x(0) + t$, for z_1 , $x(t) = x(0)e^t$, and for z_{n+1} , with $n \geq 1$, $x(t) = x(0)[1 - nx(0)^n t]^{-1/n}$. The flows defined by $f_t(x(0)) = x(t)$ are:

$$\begin{aligned} z_0: \quad f_t(x) &= x + t, & z_1: \quad f_t(x) &= xe^t, \\ z_{n+1}, \quad n \geq 1: \quad f_t(x) &= x[1 - nx^n t]^{-1/n} = x[1 + x^n t] + O(t^2). \end{aligned} \tag{24}$$

Obviously these are not defined for all t . From (19) $\phi \triangleright (a, b) = (a\phi'(b), \phi(b))$, for all $\phi \in D_0$, $(a, b) \in B_+$. We use this to recover the right action of $U(\mathfrak{d}_0)$ on $k[B_+]$. The flow (24) corresponding to z_{n+1} ($n \geq 1$) is $f_\varepsilon(x) = x[1 + x^n \varepsilon] + O(\varepsilon^2)$, hence $\frac{df_\varepsilon}{dx}(x) = 1 + (n + 1)x^n \varepsilon + O(\varepsilon^2)$. So for z_{n+1} , formally we have

$$f_\varepsilon \triangleright (a, b) = (a + (n + 1)ab^n \varepsilon + O(\varepsilon^2), b + b^{n+1} \varepsilon + O(\varepsilon^2)).$$

For $\xi \in k[B_+]$, define $(\xi \triangleleft z_{n+1})(a, b) := \frac{d}{d\varepsilon} \Big|_{\varepsilon=0} \xi(f_\varepsilon \triangleright (a, b))$, as formally $f_\varepsilon = e^{\varepsilon z_{n+1}}$. Hence

$$(\alpha \triangleleft z_{n+1})(a, b) = \frac{d}{d\varepsilon} \Big|_{\varepsilon=0} a(1 + (n + 1)b^n \varepsilon + O(\varepsilon^2)) = (n + 1)ab^n.$$

So $\alpha \triangleleft z_{n+1} = (n + 1)\alpha\beta^n$ for all $n \geq 1$, and the formulae for β follow in the same way. This explains the motivation for (20), and in Lemma 20 we already proved that this is an action as claimed.

Next, for any group factorisation $X = G \bowtie M$, define a k -linear map

$$\tilde{\Delta}_L : kM \rightarrow k[G, kM], \quad \tilde{\Delta}_L(m)(g) = m \triangleleft g,$$

where $k[G, kM]$ is the k -vector space of maps $G \rightarrow kM$. We now take $k = \mathbb{R}$. For z_{n+1} , the flow (24) is $f_\varepsilon(x) = x[1 + x^n \varepsilon] + O(\varepsilon^2)$. Hence

$$\begin{aligned} (\tilde{\Delta}_L(e^{\varepsilon z_{n+1}})(a, b))(x) &= (f_\varepsilon \triangleright (a, b))(x) = \frac{f_\varepsilon(ax + b) - f_\varepsilon(b)}{af'_\varepsilon(b)} \\ &= \frac{(ax + b)[1 + (ax + b)^n \varepsilon] - b[1 + b^n \varepsilon]}{a[1 + (n + 1)b^n \varepsilon]} + O(\varepsilon^2) \\ &= x + \varepsilon \sum_{k=2}^{n+1} \binom{n+1}{k} a^{k-1} b^{n+1-k} x^k + O(\varepsilon^2). \end{aligned}$$

Differentiating with respect to ε and evaluating at $\varepsilon = 0$ gives a map $\Delta_L(z_{n+1}) : B_+ \rightarrow U(\mathfrak{d}_0)$ which we can identify with (21).

3.5. Schrödinger action

Starting with a bicrossproduct Hopf algebra $\mathcal{H} \bowtie \mathcal{A}$, suppose we have a Hopf algebra \mathcal{A}' equipped with a nondegenerate dual pairing with \mathcal{A} . Then:

Lemma 24. [15] \mathcal{A}' is a left $\mathcal{H} \bowtie \mathcal{A}$ -module algebra, via the Schrödinger action

$$(h \otimes a) \triangleright \phi = \sum h \triangleright \phi_{(1)} \langle \phi_{(2)}, a \rangle, \tag{25}$$

where the left action of \mathcal{H} on \mathcal{A}' is defined by $(h \triangleright \phi)(a) = \phi(a \triangleleft h)$.

Corollary 25. $U(\mathfrak{b}_+)$ is a left U_{CM} -module algebra, via the Schrödinger action

$$\begin{aligned} z_n \triangleright X &= 2Y\delta_{n,2}, & \alpha \triangleright X &= X, & \beta \triangleright X &= 1, \\ z_n \triangleright Y &= 0, & \alpha \triangleright Y &= Y + 1, & \beta \triangleright Y &= 0. \end{aligned}$$

Proof. We have $(z \otimes \xi) \triangleright x = \sum z \triangleright x_{(1)} \langle x_{(2)}, \xi \rangle$, for all $z \in U(\mathfrak{d}_0)$, $\xi \in k[\mathbf{B}_+]$, $x \in U(\mathfrak{b}_+)$. So $(z \otimes 1) \triangleright X = z \triangleright X$, defined via $\langle z \triangleright X, h \rangle = \langle X, h \triangleleft z \rangle$. By (14), $\langle z_n \triangleright X, \alpha^t \beta^r \rangle = \langle X, (\alpha^t \triangleleft z_n) \beta^r + r \alpha^t \beta^{r-1} (\beta \triangleleft z_n) \rangle = (nt - r) \langle X, \alpha^t \beta^{n+r-1} \rangle = \delta_{1, n+r-1} (nt - r) \delta_{s,0} = 2\delta_{n,2} \delta_{r,0} t$ (since $n \geq 2$) whereas $\langle Y, \alpha^t \beta^r \rangle = t \delta_{r,0}$. So $z_n \triangleright X = 2Y\delta_{n,2}$ as claimed. The other results follow in the same way. \square

There is also a corresponding dual Schrödinger coaction. First of all, we note that bicrossproducts behave well with respect to dual pairings:

Lemma 26. [15] Suppose we are given a bicrossproduct $\mathcal{H} \bowtie \mathcal{A}$, together with Hopf algebras \mathcal{A}' , \mathcal{H}' equipped with nondegenerate dual pairings with \mathcal{A} , \mathcal{H} , respectively. Suppose further that \mathcal{A}' is a right \mathcal{H}' -comodule coalgebra via a coaction $\phi \mapsto \sum \phi^{(1)} \otimes \phi^{(2)}$ which is dual to $\triangleleft : \mathcal{A} \otimes \mathcal{H} \rightarrow \mathcal{A}$ in the sense that

$$\langle \phi, a \triangleleft h \rangle = \left\langle \sum \phi^{(1)} \otimes \phi^{(2)}, a \otimes h \right\rangle.$$

Then \mathcal{H}' is a left \mathcal{A}' -module algebra via the left action defined by

$$\triangleright : \mathcal{A}' \otimes \mathcal{H}' \rightarrow \mathcal{H}', \quad (\phi \triangleright z)(h) := \sum \langle \phi, h^{(0)} \rangle \langle z, h^{(1)} \rangle.$$

Furthermore, this left action and right coaction are compatible in the sense of Theorem 5, enabling us to form the bicrossproduct $\mathcal{H}' \bowtie \mathcal{A}'$.

Now consider the linear map

$$\Delta_R : \mathcal{A}' \rightarrow \mathcal{A}' \otimes \mathcal{H}' \otimes \mathcal{A}', \quad \phi \mapsto \sum \phi_{(1)}^{(1)} \otimes \phi_{(1)}^{(2)} \otimes \phi_{(2)}. \tag{26}$$

It is straightforward to check that \mathcal{A}' is a right $\mathcal{H}' \bowtie \mathcal{A}'$ -comodule coalgebra via Δ_R . The Schrödinger action (25) and coaction (26) are dual in the sense:

$$\triangleright : \mathcal{H} \bowtie \mathcal{A} \otimes \mathcal{A}' \rightarrow \mathcal{A}' = (\langle \cdot, \cdot \rangle \otimes \text{id}_{\mathcal{A}'}) (\text{id}_{\mathcal{H} \bowtie \mathcal{A}} \otimes \tau \circ \Delta_R),$$

where $\tau : \mathcal{A}' \otimes \mathcal{H}' \bowtie \mathcal{A}' \rightarrow \mathcal{H}' \bowtie \mathcal{A}' \otimes \mathcal{A}'$ is the flip map. We have:

Lemma 27. *The Schrödinger coaction (26) of \mathcal{H}_{CM} on $U(\mathfrak{b}_+)$ is*

$$\Delta_R(X) = X \otimes 1 + 1 \otimes X + Y \otimes 2t_2, \quad \Delta_R(Y) = Y \otimes 1 + 1 \otimes Y.$$

4. Scalings and deformations

In this section we introduce a natural scale parameter $\lambda \in k$ into the Hopf algebras $\mathcal{H}_{\text{CM}}, U_{\text{CM}}$ of the previous section. We first define a family of bicrossproducts $\{\mathcal{H}_{\text{CM}}^\lambda\}_{\lambda \in k}$, with $\mathcal{H}_{\text{CM}}^\lambda \cong \mathcal{H}_{\text{CM}}$ for each $\lambda \neq 0$, while for $\lambda = 0$ (the so-called classical limit) we obtain a commutative Hopf algebra which can be realised as functions on the semidirect product $\mathbb{R}^2 \rtimes D_0$. We construct a natural quotient Hopf algebra $k_\lambda[\text{Heis}]$ of $\mathcal{H}_{\text{CM}}^\lambda$, which for $\lambda = 0$ corresponds to the coordinate algebra of the Heisenberg group. We define a second family $\{U_{\text{CM}}^\lambda\}_{\lambda \in k}$, again all isomorphic to U_{CM} for $\lambda \neq 0$, and find a Hopf subalgebra $U_\lambda(\text{heis})$ with a nondegenerate dual pairing with $k_\lambda[\text{Heis}]$. Finally, by passing to an extended bicrossproduct $U(\mathfrak{d}_0) \blacktriangleright F[\mathbb{B}_+]_\lambda$ we identify the expected classical limits of U_{CM}^λ and $U_\lambda(\text{heis})$.

4.1. The deformed Heisenberg bicrossproducts $\mathcal{H}_{\text{CM}}^\lambda$ and $k_\lambda[\text{Heis}]$

Definition 28. For each $\lambda \in k$, we define $\mathcal{H}_{\text{CM}}^\lambda$ to be the Hopf algebra with generators $X, Y, \{t_n : n = 1, 2, \dots\}$, with $t_1 = 1$ and relations

$$[Y, X] = \lambda X, \quad [Y, t_n] = \lambda(n - 1)t_n, \quad [X, t_n] = \lambda((n + 1)t_{n+1} - 2t_2 t_n) \quad (27)$$

with coproduct and antipode defined by (7)–(9).

For $\lambda \neq 0$, the map $\mathcal{H}_{\text{CM}} \rightarrow \mathcal{H}_{\text{CM}}^\lambda$ given by

$$X \mapsto \lambda^{-2}X, \quad Y \mapsto \lambda^{-1}Y, \quad t_n \mapsto \lambda^{1-n}t_n \quad (28)$$

is a Hopf algebra isomorphism. For $\lambda = 0$, (27) reduces to a commutative Hopf algebra, denoted $k[\mathbb{R}^2 \rtimes D_0]$, with the generators X, Y, t_n realisable as functions on the semidirect product $\mathbb{R}^2 \rtimes D_0$.

Lemma 29. *Let \mathcal{I} be the two-sided ideal of $\mathcal{H}_{\text{CM}}^\lambda$ generated by $\{\tilde{t}_n := t_n - t_2^{n-1}\}_{n \geq 3}$. Then $\Delta(\mathcal{I}) \subseteq \mathcal{H}_{\text{CM}}^\lambda \otimes \mathcal{I} + \mathcal{I} \otimes \mathcal{H}_{\text{CM}}^\lambda$ and $\varepsilon(\mathcal{I}) = 0$.*

Proof. First, $\varepsilon(\tilde{t}_n) = 0 \forall n$, so $\varepsilon(\mathcal{I}) = 0$. We have

$$\begin{aligned} \Delta(t_n) &= \sum_{k=1}^n \sum_{i_1+\dots+i_k=n} t_{i_1} \cdots t_{i_k} \otimes t_k \\ &= \sum_{k=1}^n \sum_{i_1+\dots+i_k=n} (\tilde{t}_{i_1} + t_2^{i_1-1}) \cdots (\tilde{t}_{i_k} + t_2^{i_k-1}) \otimes (\tilde{t}_k + t_2^{k-1}) \\ &= \sum_{k=1}^n \sum_{i_1+\dots+i_k=n} t_2^{i_1+\dots+i_k-k} \otimes t_2^{k-1} \quad \text{modulo } \mathcal{H}_{\text{CM}}^\lambda \otimes \mathcal{I} + \mathcal{I} \otimes \mathcal{H}_{\text{CM}}^\lambda \end{aligned}$$

$$\begin{aligned} &= \sum_{k=1}^n \sum_{i_1+\dots+i_k=n} t_2^{n-k} \otimes t_2^{k-1} = \sum_{k=1}^n \binom{n-1}{k-1} t_2^{n-k} \otimes t_2^{k-1} \\ &= \sum_{k=0}^{n-1} \binom{n-1}{k} t_2^{n-k-1} \otimes t_2^k = (t_2 \otimes 1 + 1 \otimes t_2)^{n-1} = \Delta(t_2^{n-1}). \end{aligned}$$

So $\Delta(\tilde{t}_n) \in \mathcal{H}_{\text{CM}}^\lambda \otimes \mathcal{I} + \mathcal{I} \otimes \mathcal{H}_{\text{CM}}^\lambda$. Since the \tilde{t}_n generate \mathcal{I} , the result follows. \square

Corollary 30. *The quotient bialgebra $k_\lambda[\text{Heis}] := \mathcal{H}_{\text{CM}}^\lambda/\mathcal{I}$ is in fact a Hopf algebra, generated by $X, Y, t = 2t_2$ satisfying*

$$\begin{aligned} [Y, X] &= \lambda X, & [X, t] &= \frac{1}{2}\lambda t^2, & \Delta(X) &= 1 \otimes X + X \otimes 1 + Y \otimes t, \\ [Y, t] &= \lambda t, & \Delta(Y) &= Y \otimes 1 + 1 \otimes Y, & \Delta(t) &= t \otimes 1 + 1 \otimes t, & S(Y) &= -Y, \\ S(X) &= -X + Yt, & S(t) &= -t, & \varepsilon(X) &= 0 = \varepsilon(Y) = \varepsilon(t). \end{aligned} \tag{29}$$

Proof. By Lemma 29 the bialgebra structure of $\mathcal{H}_{\text{CM}}^\lambda$ descends to the quotient, and it is straightforward to check that there is a unique antipode S (defined as shown) that gives $k_\lambda[\text{Heis}]$ the structure of a Hopf algebra. \square

We denote $U(\mathbf{b}_+)$ with the scaled relation $[Y, X] = \lambda X$ by $U_\lambda(\mathbf{b}_+)$. Obviously $U_\lambda(\mathbf{b}_+)$ and $U(\mathbf{b}_+)$ are isomorphic Hopf algebras for $\lambda \neq 0$. Finally, $k[t]$ is the commutative unital algebra of polynomials in t .

Proposition 31. $k_\lambda[\text{Heis}]$ is a bicrossproduct $k[t] \blacktriangleright U_\lambda(\mathbf{b}_+)$, via the action

$$\triangleright : U_\lambda(\mathbf{b}_+) \otimes k[t] \rightarrow k[t], \quad X \triangleright t = \frac{1}{2}\lambda t^2, \quad Y \triangleright t = \lambda t \tag{30}$$

and coaction $\Delta_R : U_\lambda(\mathbf{b}_+) \rightarrow U_\lambda(\mathbf{b}_+) \otimes k[t]$ defined on generators by

$$\Delta_R(X) = X \otimes 1 + Y \otimes t, \quad \Delta_R(Y) = Y \otimes 1 \tag{31}$$

and extended by $\Delta_R(gh) = \sum g_{(1)} \overline{(1)} h \overline{(1)} \otimes g_{(1)}^{(2)} (g_{(2)} \triangleright h^{(2)})$.

Proof. It is straightforward to check that the given coaction is well-defined, and $k[t]$ is a left $U_\lambda(\mathbf{b}_+)$ -module algebra, $U_\lambda(\mathbf{b}_+)$ a right $k[t]$ -comodule coalgebra. As in Theorem 18 it can be checked that action and coaction are compatible in the sense of Theorem 5. Hence we can construct the bicrossproduct $k[t] \blacktriangleright U_\lambda(\mathbf{b}_+)$, whose presentation using (5) coincides with (29). \square

The three-dimensional Heisenberg group $\mathbb{H}_3(k)$ is the matrix group

$$\mathbb{H}_3(k) = \left\{ (a, b, c) := \begin{pmatrix} 1 & a & b \\ 0 & 1 & c \\ 0 & 0 & 1 \end{pmatrix} : a, b, c \in k \right\}.$$

If we write the coproduct (29) of $k_\lambda[\text{Heis}]$ using the matrix notation

$$\Delta \begin{pmatrix} 1 & Y & X \\ 0 & 1 & t \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & Y & X \\ 0 & 1 & t \\ 0 & 0 & 1 \end{pmatrix} \otimes \begin{pmatrix} 1 & Y & X \\ 0 & 1 & t \\ 0 & 0 & 1 \end{pmatrix}$$

we see that for $\lambda = 0$, $k_\lambda[\text{Heis}]$ is isomorphic to the commutative Hopf algebra generated by the coordinate functions $Y(a, b, c) = a$, $X(a, b, c) = b$, $t(a, b, c) = c$ on $\mathbb{H}_3(k)$. We therefore consider $k_\lambda[\text{Heis}]$ to be a deformation of the Heisenberg group coordinate algebra.

4.2. The deformed Heisenberg bicrossproducts U_{CM}^λ and $U_\lambda(\mathbf{heis})$

Definition 32. For each $\lambda \in k$, we define U_{CM}^λ to be the Hopf algebra with generators z_n ($n \geq 2$), α , β and relations

$$\begin{aligned} [\alpha, \beta] &= 0, & [z_m, z_n] &= (n - m)z_{m+n-1}, \\ [z_n, \alpha] &= -\lambda^{n-1}n\alpha\beta^{n-1}, & [z_n, \beta] &= \lambda^{n-1}\beta^n \end{aligned}$$

with coproduct and antipode given by (22).

For $\lambda \neq 0$, there is an isomorphism of Hopf algebras $U_{\text{CM}} \rightarrow U_{\text{CM}}^\lambda$ defined by

$$z_n \mapsto \lambda^{n-1}z_n, \quad \alpha \mapsto \alpha, \quad \beta \mapsto \lambda^2\beta. \tag{32}$$

Definition 33. We define $U_\lambda(\mathbf{heis})$ to be the Hopf subalgebra of U_{CM}^λ generated by $z := z_2$, α , β . The presentation of $U_\lambda(\mathbf{heis})$ is then:

$$\begin{aligned} [z, \alpha] &= -2\lambda\alpha\beta, & [z, \beta] &= \lambda\beta^2, & [\alpha, \beta] &= 0, \\ \Delta(\alpha) &= \alpha \otimes \alpha, & \Delta(\beta) &= \beta \otimes 1 + \alpha \otimes \beta, & \Delta(z) &= z \otimes 1 + \alpha \otimes z. \end{aligned} \tag{33}$$

$U_\lambda(\mathbf{heis})$ corresponds to a Heisenberg version of the Planck scale Hopf algebra [16]. As before, the $U_\lambda(\mathbf{heis})$ are all isomorphic for $\lambda \neq 0$. Now let $U(z)$ be the unital commutative Hopf algebra generated by z , with $\Delta(z) = z \otimes 1 + 1 \otimes z$.

Proposition 34. $U_\lambda(\mathbf{heis})$ is a bicrossproduct $U(z) \bowtie k[\mathbf{B}_+]$, via the right action $\triangleleft : k[\mathbf{B}_+] \otimes U(z) \rightarrow k[\mathbf{B}_+]$ defined by $\alpha \triangleleft z = 2\lambda\alpha\beta$, $\beta \triangleleft z = -\lambda\beta^2$, and left coaction $\Delta_L : U(z) \rightarrow k[\mathbf{B}_+] \otimes U(z)$, $h \mapsto \sum h^{(0)} \otimes h^{(1)}$ defined by

$$\Delta_L(z) = \alpha \otimes z, \quad \Delta_L(hg) = \sum (h^{(0)} \triangleleft g_{(1)})g_{(2)}^{(0)} \otimes h^{(1)}g_{(2)}^{(1)} \quad \forall h, g \in U(z).$$

Proof. It is easily checked that [15, Theorem 6.2.3] applies. \square

Lemma 35. For $\lambda \neq 0$, there is a nondegenerate dual pairing of $k_\lambda[\text{Heis}]$ and $U_\lambda(\mathbf{heis})$ given by $\langle t^i X^j Y^k, z^p \alpha^q \beta^r \rangle = j!(\lambda q)^k \delta_{i,p} \delta_{j,r}$.

Proof. This follows from (23) together with (28) and (32). \square

Obtaining the “correct” classical limit $\lambda = 0$ of $U_\lambda(\mathbf{heis})$ (in the sense of duality with $k[\mathbf{Heis}]$) is more subtle, since we would like to obtain the universal enveloping algebra $U(\mathbf{heis})$ of the Heisenberg Lie algebra. From the geometric point of view, consider the \mathbb{R} -valued functions $A, \{\alpha_t\}_{t \in \mathbb{R}}, \beta$ on \mathbf{B}_+ :

$$A(a, b) = \log a, \quad \alpha_t(a, b) = a^t, \quad \beta(a, b) = b. \tag{34}$$

We have $\alpha_{t_1} \alpha_{t_2} = \alpha_{t_1+t_2}, \alpha_0 = 1$ and formally, $\alpha_t = e^{tA}$. To treat the case $\lambda = 0$ we wish to work with A as a generator of $U_\lambda(\mathbf{heis})$ rather than $\alpha = \alpha_1$. This can be formulated rigorously in two different ways. One well-known approach is by working over the ring of formal power series $k[[\lambda]]$. A second approach which we now sketch is as follows. Motivated by (34), for any $\lambda \in k$ define $F[\mathbf{B}_+]_\lambda$ as the commutative Hopf algebra (over k) generated by $\{\alpha_t\}_{t \in k}, A, \beta$ with $\alpha_0 = 1, \alpha_{t_1} \alpha_{t_2} = \alpha_{t_1+t_2}$, and

$$\Delta(\alpha_t) = \alpha_t \otimes \alpha_t, \quad \Delta(A) = A \otimes 1 + 1 \otimes A, \quad \Delta(\beta) = \alpha_\lambda \otimes \beta + \beta \otimes 1.$$

Then there is a right action of $U(\mathbf{d}_0)$ on $F[\mathbf{B}_+]_\lambda$, and left coaction of $F[\mathbf{B}_+]_\lambda$ on $U(\mathbf{d}_0)$ defined by

$$\begin{aligned} \alpha_t \triangleleft z_n &= \lambda^{n-2} n t \alpha_t \beta^{n-1}, & A \triangleleft z_n &= n \lambda^{n-2} \beta^{n-1}, & \beta \triangleleft z_n &= -\lambda^{n-1} \beta^n, \\ \Delta_L(z_n) &= \sum_{j=2}^n \lambda^{n-j} \binom{n}{j} \alpha_{\lambda(j-1)} \beta^{n-j} \otimes z_j. \end{aligned}$$

This action and coaction are compatible in the sense of [15, Theorem 6.2.3], hence for each $\lambda \in k$ there is a bicrossproduct $U(\mathbf{d}_0) \blacktriangleright F[\mathbf{B}_+]_\lambda$ containing U_{CM}^λ as a Hopf subalgebra. Define an extended version of $U_\lambda(\mathbf{heis})$, denoted $\tilde{U}_\lambda(\mathbf{heis})$, to be the Hopf subalgebra generated by $z = z_2, \alpha = \alpha_\lambda, A$ and β . For $\lambda \neq 0$ this corresponds to $U_\lambda(\mathbf{heis})$ adjoined the primitive element A , with $[z, A] = -2\beta$, while for $\lambda = 0$ we have $\alpha = \alpha_0 = 1$, so z, A and β are primitive with relations $[z, A] = -2\beta, [z, \beta] = 0 = [A, \beta]$. So for $\lambda = 0$ then $\tilde{U}_\lambda(\mathbf{heis})$ is isomorphic to $U(\mathbf{heis})$. Similarly, the correct classical limit of U_{CM}^λ is the cocommutative Hopf algebra generated by primitive elements $\{z_n\}_{n \geq 2}, A, \beta$, with $[z_m, z_n] = (n - m)z_{m+n-1}, [z_n, A] = -2\beta \delta_{n,2}$, and $[z_n, \beta] = 0 = [A, \beta]$. These remarks are for clarification purposes. We do not use this approach in the sequel.

Next, the Schrödinger action of Corollary 25 is compatible with the scaling:

Lemma 36. For $\lambda \neq 0, U_\lambda(\mathbf{b}_+)$ is a left $U_\lambda(\mathbf{heis})$ -module algebra via

$$\begin{aligned} z \triangleright X &= 2\lambda Y, & z \triangleright Y &= 0, & \alpha \triangleright X &= X, \\ \alpha \triangleright Y &= Y + \lambda, & \beta \triangleright X &= \lambda, & \beta \triangleright Y &= 0. \end{aligned}$$

When $\lambda = 0$ there is the known action of the Heisenberg algebra on $k[X, Y]$ by $z = 2Y \frac{\partial}{\partial X}, A = \frac{\partial}{\partial Y}, \beta = \frac{\partial}{\partial X}$, and the Schrödinger action of Lemma 36 should be thought of as a deformation of this.

We note also that it is immediate that the right Schrödinger coaction of Lemma 27 restricts to a right coaction of $k_\lambda[\mathbf{Heis}]$ on $U_\lambda(\mathfrak{b}_+)$.

Finally, a typical feature of bicrossproduct Hopf algebras associated to group factorisations where neither factor group is compact is that the actions have singularities [12,13]. These singularities do not appear at the algebraic bicrossproduct level, which is why we have not encountered them in constructing $k_\lambda[\mathbf{Heis}]$ and $U_\lambda(\mathbf{heis})$, but rather when one tries to pass to C^* - or von Neumann completions. We note that a pair of locally compact quantum groups corresponding to $U_\lambda(\mathbf{heis})$ and $k_\lambda[\mathbf{Heis}]$ was previously constructed by Vaes [20, Example 3.4], applying the techniques of [1] to group factorisations $X \cong G \bowtie M$, where both G and M are locally compact. Explicitly, the correspondence between the generators of the Hopf algebras $U_\lambda(\mathbf{heis})$ and $k_\lambda[\mathbf{Heis}]$, and the (unbounded) operators A_i, B_i, C_i ($i = 1, 2$) generating these von Neumann algebras is:

$$\begin{aligned}
 U_\lambda(\mathbf{heis}) : \alpha &\mapsto A_1^2, & \beta &\mapsto A_1 B_1, & z &\mapsto \lambda C_1 A_1^2, \\
 k_\lambda[\mathbf{Heis}] : X &\mapsto \lambda B_2, & Y &\mapsto \frac{1}{2} \lambda A_2, & t &\mapsto 2C_2.
 \end{aligned}$$

It is natural to ask whether this could be extended to give faithful representations of \mathcal{H}_{CM} and U_{CM} as (unbounded) operators on some Hilbert space, affiliated to a locally compact quantum group. An obstacle is the fact that $\text{Diff}^+(\mathbb{R})$ is not locally compact, nor does it have any interesting locally compact subgroups [9]. This question will be pursued elsewhere.

4.3. Local factorisation of $SL_2(\mathbb{R})$

It is natural to ask for a geometrical picture in terms of a group factorisation linked to the $k_\lambda[\mathbf{Heis}]$ and $U_\lambda(\mathbf{heis})$ bicrossproducts, in the same way as \mathcal{H}_{CM} and U_{CM} were shown to be linked to the factorisation of $\text{Diff}^+(\mathbb{R})$. In this section we show that the relevant group is $SL_2(\mathbb{R})$, which locally (but not globally) factorises as $SL_2(\mathbb{R}) \approx \mathbf{B}_+ \bowtie \mathbb{R}$.

Using the coaction (31) define a linear map $f : U(\mathfrak{b}_+) \rightarrow U(\mathfrak{b}_+)$ by $f(x) = \sum x^{(1)} \langle x^{(2)}, z \rangle$ where the pairing $\langle \cdot, \cdot \rangle : k[t] \otimes U(z) \rightarrow k$ is $\langle t^m, z^n \rangle = 2^m \delta_{m,n}$. Then $f(X) = 2Y$, $f(Y) = 0$, and further $f(YX) = 2Y(Y + 1)$, $f(XY) = 2Y^2$, hence $f(YX - XY) = 2Y = f(X)$. Identifying X, Y with the generators of \mathfrak{b}_+ , and z with the generator of the Lie algebra \mathfrak{r} of \mathbb{R} , we have a well-defined left action of \mathfrak{r} on \mathfrak{b}_+ , given by $z \triangleright x = f(x)$, satisfying

$$z \triangleright X = 2Y, \quad z \triangleright Y = 0. \tag{35}$$

The adjoint of the left action (30) defines a right action of $U(\mathfrak{b}_+)$ on $U(z)$:

$$\begin{aligned}
 \langle t^n, z \triangleleft Y \rangle &:= \langle Y \triangleright t^n, z \rangle = \langle nt^n, z \rangle = 2n \delta_{n,1}, \\
 \langle t^n, z \triangleleft X \rangle &:= \langle X \triangleright t^n, z \rangle = \left\langle \frac{1}{2} nt^{n+1}, z \right\rangle = \frac{1}{2} n \delta_{n+1,1} = 0.
 \end{aligned}$$

This gives a right action of \mathfrak{b}_+ on \mathfrak{r} , satisfying

$$z \triangleleft X = 0, \quad z \triangleleft Y = z. \tag{36}$$

It is straightforward to check that \mathfrak{b}_+ and \mathfrak{r} equipped with the actions (35), (36) are a matched pair of Lie algebras in the sense of [15, Definition 8.3.1]. From this point we take $k = \mathbb{R}$. Since we have a matched pair, the \mathbb{R} -vector space $\mathfrak{g} := \mathfrak{b}_+ \oplus \mathfrak{r}$ can be given the structure of a Lie algebra, with Lie bracket

$$[x, y]_{\mathfrak{g}} := [x, y]_{\mathfrak{b}_+}, \quad [z, x]_{\mathfrak{g}} := z \triangleleft x + z \triangleright x, \quad \forall x, y \in \mathfrak{b}_+.$$

Then $[z, X]_{\mathfrak{g}} = z \triangleleft X + z \triangleright X = 2Y$, $[z, Y]_{\mathfrak{g}} = z \triangleleft Y + z \triangleright Y = z$, so $\mathfrak{g} \cong \mathfrak{sl}_2(\mathbb{R})$. Since both B_+ and \mathbb{R} are simply connected Lie groups, we conclude that $B_+ \bowtie \mathbb{R} \approx SL_2(\mathbb{R})$ in so far as the actions on the left-hand side exponentiate. We embed B_+ and \mathbb{R} as subgroups of $SL_2(\mathbb{R})$ by

$$(a, b) = \begin{pmatrix} a^{\frac{1}{2}} & a^{-\frac{1}{2}}b \\ 0 & a^{-\frac{1}{2}} \end{pmatrix}, \quad (c) = \begin{pmatrix} 1 & 0 \\ -c & 1 \end{pmatrix}. \tag{37}$$

Lemma 37. *There is a local factorisation $SL_2(\mathbb{R}) \approx B_+ \bowtie \mathbb{R}$ given by*

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \mapsto (d^{-2}, d^{-1}b)(-d^{-1}c), \quad d \neq 0.$$

Proof. For $d \neq 0$, $\begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} d^{-1} & b \\ 0 & d \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -d^{-1}c & 1 \end{pmatrix}$. Then apply (37). \square

The resulting left action of \mathbb{R} on B_+ and right action of B_+ on \mathbb{R} are given by:

$$c \triangleright (a, b) = \left(\frac{a}{(1-bc)^2}, \frac{b}{1-bc} \right), \quad c \triangleleft (a, b) = \frac{ac}{1-bc}.$$

As in Sections 3.2 and 3.4 we can use this factorisation to rederive the actions and coactions used to construct the bicrossproducts $k_\lambda[\text{Heis}]$, $U_\lambda(\mathbf{heis})$.

5. Differential calculi over $U_\lambda(\mathfrak{b}_+)$ and $k_\lambda[\text{Heis}]$

The model obtained above can be seen as a variant of one of a family of previously-studied bicrossproducts, which act on noncommutative algebras denoted $U_\lambda(\mathfrak{b}_+^n)$ of varying dimension n . First of all, in [12,13], a bicrossproduct $U(\mathfrak{so}_3) \bowtie \mathbb{C}[B_+^3]$ associated to a local factorisation of $SO(3, 1)$ was constructed. This bicrossproduct can be regarded as corresponding to a deformation of the Euclidean group of motions, and acts naturally on the algebra $U_\lambda(\mathfrak{b}_+^3)$. Similarly, in [17], a bicrossproduct $U(\mathfrak{so}_{3,1}) \bowtie \mathbb{C}[B_+^{3,1}]$ associated to a factorisation of $SO(3, 2)$ was constructed, and interpreted as corresponding to a deformation of the Poincaré group. This bicrossproduct acts naturally on the algebra $U_\lambda(\mathfrak{b}_+^{3,1})$. These and other examples [16] have been widely studied in the mathematical physics literature. Our new example coming from the Connes–Moscovici algebra has an analogous geometrical picture.

From this point of view it is natural to extend the theory to include noncommutative differential geometry both on $U_\lambda(\mathfrak{b}_+)$ and on $k_\lambda[\text{Heis}]$ as ‘coordinate algebras.’ Recall [21] that a first order differential calculus (FODC) over an algebra \mathcal{A} is an \mathcal{A} -bimodule Ω^1 with a linear map $d : \mathcal{A} \rightarrow \Omega^1$ such that

- (1) d obeys the Leibniz rule $d(ab) = (da)b + adb$, for all $a, b \in \mathcal{A}$.
- (2) Ω^1 is the linear span of elements adb .

If in addition \mathcal{A} is a right \mathcal{H} -comodule algebra for some Hopf algebra \mathcal{H} , then Ω^1 is said to be right covariant if there exists a linear map $\Delta_R: \Omega^1 \rightarrow \Omega^1 \otimes \mathcal{H}$, extending the coaction Δ_R of \mathcal{H} on \mathcal{A} , in the sense that

$$\Delta_R(awb) = \Delta_R(a)\Delta_R(\omega)\Delta_R(b), \quad \Delta_R(da) = (d \otimes \text{id})\Delta_R(a)$$

for all $a, b \in \mathcal{H}$, $\omega \in \Omega^1$. Left covariant and bicovariant are defined similarly. Covariant FODC over finite-dimensional bicrossproducts have been intensively studied, in particular, see [7], which deals with the case of bicrossproducts arising from finite group factorisations.

5.1. Differential calculi over $U_\lambda(\mathbf{b}_+)$

There is a standard calculus on $U_\lambda(\mathbf{b}_+)$, the so-called Oeckl calculus [18] generated as a left $U_\lambda(\mathbf{b}_+)$ -module Ω^1 by dX, dY , with

$$[dX, X] = 0 = [dX, Y], \quad [dY, X] = \lambda dX, \quad [dY, Y] = \lambda dY. \tag{38}$$

The right Schrödinger coaction (26) gives $U_\lambda(\mathbf{b}_+)$ the structure of a $k_\lambda[\text{Heis}]$ -comodule algebra. This right coaction is dual to the left Schrödinger action (Lemma 36) in that $a \triangleright x = \sum x^{(1)} \langle a, x^{(2)} \rangle$ for all $x \in U_\lambda(\mathbf{b}_+)$, $a \in U_\lambda(\mathbf{heis})$. This extends to a right coaction of $k_\lambda[\text{Heis}]$ on Ω^1 , if

$$\Delta_R(dX) = dX \otimes 1 + dY \otimes t, \quad \Delta_R(dY) = dY \otimes 1 \tag{39}$$

is well defined.

Lemma 38. *The Oeckl calculus (38) is covariant with the right coaction (39) if and only if $\lambda = 0$.*

Proof. $\Delta_R((dX)X) - \Delta_R(X dX) = \lambda(dX \otimes t + \frac{1}{2}dY \otimes t^2)$, but $[dX, X] = 0$. \square

This parallels what is found for the higher-dimensional bicrossproducts mentioned above, where it is known that the natural translation-invariant calculus on $U_\lambda(\mathbf{b}_+^n)$ is not covariant under the bicrossproduct symmetry group.

Theorem 39. *There exists a unique FODC Ω^1 over $U_\lambda(\mathbf{b}_+)$ such that:*

- (1) Ω^1 has basis (as a left $U_\lambda(\mathbf{b}_+)$ -module) $\{dX, dY\}$.
- (2) Ω^1 is covariant under the right coaction (39) of $k_\lambda[\text{Heis}]$.

The explicit presentation of Ω^1 is then

$$\begin{aligned} (dX)X &= X dX, & (dX)Y &= (Y - \lambda/2) dX, \\ (dY)X &= (\lambda/2) dX + X dY, & (dY)Y &= (Y + \lambda/2) dY. \end{aligned}$$

Proof. It follows from our assumptions that

$$\begin{aligned} (dX)X &= a_1 dX + a_2 dY, & (dX)Y &= b_1 dX + b_2 dY, \\ (dY)X &= c_1 dX + c_2 dY, & (dY)Y &= e_1 dX + e_2 dY, \end{aligned}$$

for some $a_1, \dots, e_2 \in U_\lambda(\mathbf{b}_+)$. Then from $d(YX) - d(XY) = \lambda dX$, we have $c_1 = b_1 - Y + \lambda$, $c_2 = X + b_2$. Next,

$$\begin{aligned} \Delta_R((dY)Y) &= (dY)Y \otimes 1 + dY \otimes Y = e_1 dX \otimes 1 + (e_2 \otimes 1 + 1 \otimes Y)(dY \otimes 1), \\ \Delta_R(e_1 dX + e_2 dY) &= \Delta(e_1)(dX \otimes 1) + [\Delta(e_1)(1 \otimes t) + \Delta(e_2)](dY \otimes 1). \end{aligned}$$

Hence $\Delta(e_1) = e_1 \otimes 1$, $\Delta(e_2) + \Delta(e_1)(1 \otimes t) = e_2 \otimes 1 + 1 \otimes Y$, so $e_1 = e'_1$, $e_2 = Y - e'_1 t + e'_2$, where e'_1, e'_2 are constants. Applying Δ_R to the other expressions gives

$$\begin{aligned} (dX)X &= (X + a'_1) dX + a'_2 dY, & (dX)Y &= (Y - \lambda/2) dX + (a'_1/2) dY, \\ (dY)X &= (\lambda/2) dX + (X + a'_1/2) dY, & (dY)Y &= (Y + \lambda/2) dY, \end{aligned}$$

where a'_1, a'_2 are constants. Then the constraint $(dX)YX - (dX)XY = \lambda(dX)X$, and similarly for dY gives the result. \square

We note further that the left coaction $U_\lambda(\mathbf{b}_+) \rightarrow k_\lambda[\text{Heis}] \otimes U_\lambda(\mathbf{b}_+)$ given by $X \mapsto 1 \otimes X$, $Y \mapsto 1 \otimes Y + Y \otimes 1$ extends to a left coaction on this Ω^1 making it into a left covariant FODC. The left and right coactions are compatible, hence Ω^1 is in fact bicovariant under $k_\lambda[\text{Heis}]$.

5.2. Differential calculi over $k_\lambda[\text{Heis}]$

Similarly, one would like a calculus on the bicrossproduct quantum group itself. In previously-studied higher-dimensional cases it has been found that any bicovariant calculus needs to have extra nonclassical generators. In our case we would expect a four-dimensional calculus on $k_\lambda[\text{Heis}]$. We are therefore interested to find FODC over $k_\lambda[\text{Heis}]$ which are bicovariant with respect to the coactions induced from the coproduct. This implies

$$\Delta_L(dX) = 1 \otimes dX + Y \otimes dt, \quad \Delta_R(dX) = dX \otimes 1 + dY \otimes t, \tag{40}$$

while both dY and dt are left- and right-invariant.

As shown by Woronowicz [21], covariant FODC can be classified in terms of one-sided ideals of the dual Hopf algebra invariant under the (left or right) adjoint coaction of the dual on itself. Using the presentation (33) of $U_\lambda(\mathbf{heis})$ it is straightforward to give a complete list of right-covariant FODC over $k_\lambda[\text{Heis}]$ of dimension at most 4. Then one could check by hand for bicovariance. This would be very laborious and we prefer to proceed directly. First of all:

Theorem 40. *Let Ω^1 be the $k_\lambda[\text{Heis}]$ -bimodule with left module basis $\{dX, dY, dt\}$ and relations*

$$\begin{aligned} (dX)X &= X dX + \lambda X dt, & (dX)Y &= Y dX, & (dX)t &= t dX + \lambda t dt, \\ (dY)X &= X dY + \lambda dX, & (dY)Y &= (Y + \lambda) dY, & (dY)t &= t dY + \lambda dt, \\ (dt)X &= X dt, & (dt)Y &= Y dt, & (dt)t &= t dt. \end{aligned}$$

Then Ω^1 is a left-covariant FODC for the left coaction (40).

Theorem 41. For $\lambda \neq 0$, there exist two nonisomorphic right-covariant three-dimensional FODC over $k_\lambda[\text{Heis}]$, with basis $\{dX, dY, dt\}$, extending the two-dimensional FODC of Theorem 39. Explicitly, these are

$$\begin{aligned} (dX)X &= X dX, & (dX)Y &= (Y - \lambda/2) dX, & (dX)t &= t dX + g t dt, \\ (dY)X &= (\lambda/2) dY + X dY, & (dY)Y &= (Y + \lambda/2) dY, & (dY)t &= t dY + g dt, \\ (dt)X &= (X + (g - \lambda)t) dt, & (dt)Y &= (Y + g - \lambda) dt, & (dt)t &= t dt, \end{aligned}$$

with $g = 0$ or $\lambda/2$.

Proof. We start in the same way as Theorem 39, with the given relations $(dX)X = X dX, \dots, (dY)Y = (Y + \lambda/2) dY$ together with

$$\begin{aligned} (dX)t &= c_1 dX + c_2 dY + c_3 dt, & (dY)t &= g_1 dX + \dots, \\ (dt)X &= h_1 dX + \dots, & (dt)Y &= j_1 dX + \dots, & (dt)t &= k_1 dX + k_2 dY + k_3 dt, \end{aligned}$$

for some $c_1, \dots, k_3 \in k_\lambda[\text{Heis}]$. Applying Δ_R to both sides of $(dY)Y, (dY)t, (dt)Y, (dt)t$, and using the relation $d(Yt) - d(tY) = \lambda dt$, we have

$$\begin{aligned} (dY)Y &= f'_1 dX + (Y - f'_1 t + f'_2) dY + f'_3 dt, \\ (dY)t &= g'_1 dX + (g'_2 + (1 - g'_1)t) dY + g'_3 dt, \\ (dt)t &= k'_1 dX + (k'_2 - k'_1 t) dY + (t + k'_3) dt, \\ (dt)Y &= g'_1 dX + (g'_2 - g'_1 t) dY + (Y + g'_3 - \lambda) dt, \end{aligned}$$

where f'_1, \dots, k'_3 are scalars. Doing the same for $(dX)t$ and $(dt)X$ gives

$$\begin{aligned} (dX)t &= [(1 + g'_1)t + \lambda k'_1/2] dX + [-g'_1 t^2 + (g'_2 - \lambda k'_1/2)t + (h'_2 + \lambda k'_2/2)] dY \\ &\quad + [g'_3 t + (h'_3 + \lambda k'_3/2)] dt, \\ (dt)X &= (g'_1 t) dX + (\lambda/2)(k'_2 - k'_1 t) dY + (\lambda/2)(2t + k'_3) dt. \end{aligned}$$

Demanding consistency of all possible relations $(dt)(YX - XY) = \lambda(dt)X, \dots, (dY)(Xt - tX) = (\lambda/2)(dY)t^2$ gives the result. \square

It is straightforward to check that none of the covariant FODC of Theorems 40 and 41 are bicovariant. In fact:

Theorem 42. Let Ω^1 be a three-dimensional FODC over $k_\lambda[\text{Heis}]$, with basis $\{dX, dY, dt\}$. Then Ω^1 cannot be bicovariant.

Proof. This follows in exactly the same way as Theorem 39. \square

We now look for four-dimensional covariant FODC Ω^1 over $k_\lambda[\text{Heis}]$ of a typical form making the calculus ‘inner.’

Theorem 43. *Suppose that Ω^1 is a four-dimensional right-covariant FODC Ω^1 over $k_\lambda[\text{Heis}]$, with basis (as a left $k_\lambda[\text{Heis}]$ -module) $\{dX, dY, dt, \theta\}$, with $\theta a - a\theta = da$ for all $a \in k_\lambda[\text{Heis}]$, and $\Delta_R(\theta) = \theta \otimes 1$. Such an Ω^1 cannot contain as a sub-bimodule the two-dimensional calculus of Theorem 39.*

Proof. As before, write

$$(dX)X = a_1 dX + a_2 dY + a_3 dt + a_4\theta, \quad \dots, \quad (dt)t = k_1 dX + k_2 dY + k_3 dt + k_4\theta$$

for some $a_1, \dots, k_4 \in k_\lambda[\text{Heis}]$. Applying right-covariance and the relations $[Y, X] = \lambda X$ and so on gives

$$\begin{aligned} (dX)X &= [X + f_1't^2 + (\lambda - 2e_1')t + a_1']dX \\ &\quad + [-f_1't^3 + (f_2' - \lambda/2)t^2 + (2e_1' - a_1')t + a_2']dY \\ &\quad + [f_3't^2 + 2e_3't + a_3']dt + [f_4't^2 + a_4']\theta, \\ (dX)Y &= [Y + f_1't + (e_1' - \lambda)]dX + [-f_1't^2 + (f_2' - e_1')t + e_2']dY \\ &\quad + [f_3't + e_3']dt + [f_4't + e_4']\theta, \\ (dX)t &= [(1 + g_1')t + (c_1' + (\lambda/2)k_1')]dX \\ &\quad + [-g_1't^2 + (g_2' - c_1' - (\lambda/2)k_1')t + (c_2' + (\lambda/2)k_2')]dY \\ &\quad + [g_3't + (c_3' + (\lambda/2)k_3')]dt + [g_4't + (c_4' + (\lambda/2)k_4')] \theta, \\ (dY)X &= [f_1't + e_1']dX + [X - f_1't^2 + (f_2' - e_1')t + e_2']dY \\ &\quad + [f_3't + e_3']dt + [f_4't + e_4']\theta, \\ (dY)Y &= f_1'dX + [Y - f_1't + f_2']dY + f_3'dt + f_4'\theta, \\ (dY)t &= g_1'dX + [g_2' + (1 - g_1')t]dY + g_3'dt + g_4'\theta, \\ (dt)X &= [g_1't + c_1']dX + [-g_1't^2 + (g_2' - c_1')t + c_2']dY \\ &\quad + [X + (g_3' - \lambda)t + c_3']dt + [g_4't + c_4']\theta, \\ (dt)Y &= g_1'dX + [g_2' - g_1't]dY + [Y + g_3' - \lambda]dt + g_4'\theta, \\ (dt)t &= k_1'dX + [k_2' - k_1't]dY + [t + k_3']dt + k_4'\theta, \end{aligned} \tag{41}$$

for scalars a_1', \dots, k_4' . There are many constraints imposed by $(dX)[Y, X] = \lambda(dX)X$ and so on, we do not list these. For Ω^1 to contain as a sub-bimodule the calculus of Theorem 39, we need $(dX)X = X dX$ in (41), which implies

$$X + f_1't^2 + (\lambda - 2e_1')t + a_1' = X, \quad -f_1't^3 + (f_2' - \lambda/2)t^2 + (2e_1' - a_1')t + a_2' = 0.$$

Hence $\lambda = 2e_1', a_1' = 0$ and $2e_1' = a_1'$, which has no solution for $\lambda \neq 0$. \square

Theorem 44. *Suppose that Ω^1 is a four-dimensional bicovariant FODC over $k_\lambda[\text{Heis}]$ with basis $\{dX, dY, dt, \theta\}$, with $\theta a - a\theta = da$ for all $a \in k_\lambda[\text{Heis}]$, $\Delta_L(\theta) = 1 \otimes \theta$ and $\Delta_R(\theta) = \theta \otimes 1$. Then no such Ω^1 can exist.*

Proof. Starting with the relations (41) and applying Δ_L to $(dt)t, (dt)Y, \dots, (dX)t$ gives $c'_1 = 0, e'_1 = \lambda, e'_2 = e'_3 = e'_4 = 0, f'_2 = \lambda, f'_1 = f'_3 = f'_4 = 0, g'_3 = \lambda, g'_1 = g'_2 = g'_4 = 0, k'_i = 0$ for all i . Hence

$$(dX)X = [X - \lambda t + a'_1]dX + [(\lambda/2)t^2 + (2\lambda - a'_1)t + a'_2]dY + a'_3 dt + a'_4 \theta.$$

Applying Δ_L to both sides of this gives on the left-hand side:

$$\begin{aligned} \Delta_L((dX)X) &= X \otimes dX + 1 \otimes (dX)X + Y \otimes (dt)X + YX \otimes dt \\ &= [X \otimes 1 + 1 \otimes X - 1 \otimes \lambda t + a'_1(1 \otimes 1)](1 \otimes dX) + \text{other,} \end{aligned}$$

whereas $\Delta_L(\text{right-hand side}) = [\Delta(X) - \lambda\Delta(t) + a'_1(1 \otimes 1)](1 \otimes dX) + \text{other}$ (where “other” denotes terms in dY, dt, θ), and these are inconsistent. \square

Acknowledgments

We thank Gerald Goldin for very useful discussions about diffeomorphism groups. We are also grateful for the hospitality of the Perimeter Institute and the Isaac Newton Institute during the time this work was completed. Finally we thank the referee for their helpful comments.

Appendix A. Bicrossproduct description of the left-handed Connes–Moscovici Hopf algebra $\mathcal{H}_{\text{CM}}^{\text{left}}$

We now describe how the original codimension one Connes–Moscovici Hopf algebra $\mathcal{H}_{\text{CM}}^{\text{left}}$, defined in [5, p. 206], is isomorphic to a right–left bicrossproduct $U(\mathfrak{b}_+) \bowtie k[\mathbf{D}_0]$ which we now construct. The presentation of $\mathcal{H}_{\text{CM}}^{\text{left}}$ differs from that of \mathcal{H}_{CM} given in (1) only in that $Y \otimes \delta_1$ is replaced by $\delta_1 \otimes Y$ in the definition of $\Delta(X)$. The new bicrossproduct is also associated to the factorisation $\text{Diff}^+(\mathbb{R}) = \mathbf{B}_+ \bowtie \mathbf{D}_0$, but in a less natural way than \mathcal{H}_{CM} . In particular, the corresponding dual bicrossproduct $k[\mathbf{B}_+] \blacktriangleright U(\mathfrak{d}_0)$ is much more complicated. This is why we chose to work with \mathcal{H}_{CM} throughout this paper. Proofs of the assertions in this section are completely analogous to those given in Section 3.1, and we omit the details.

Lemma 45. *$k[\mathbf{D}_0]$ is a right $U(\mathfrak{b}_+)$ -module algebra via the action*

$$t_n \triangleleft X = -(n + 1)t_{n+1} + 2t_2 t_n, \quad t_n \triangleleft Y = (1 - n)t_n \tag{42}$$

equivalently defined by $\delta_n \triangleleft X = -\delta_{n+1}, \delta_n \triangleleft Y = -n\delta_n$.

Lemma 46. *$U(\mathfrak{b}_+)$ is a left $k[\mathbf{D}_0]$ -comodule coalgebra via the coaction*

$$\Delta_L(X) = 1 \otimes X + 2t_2 \otimes Y, \quad \Delta_L(Y) = 1 \otimes Y \tag{43}$$

(equivalently, $\Delta_L(X) = 1 \otimes X + \delta_1 \otimes Y$, $\Delta_L(Y) = 1 \otimes Y$) extended to all of $U(\mathfrak{b}_+)$ via $\Delta_L(hg) = \sum (h^{(1)} \triangleleft g_{(1)}) g_{(2)}^{(1)} \otimes h^{(2)} g_{(2)}^{(2)}$.

This action and coaction can be derived from (19) as follows. Define a right action of B_+ on $k[D_0]$ via

$$(\xi \triangleleft (a, b))(\phi) := \xi(\phi \triangleleft (a, b)^{-1}).$$

In the same way as Section 3.1 the formulae (42), (43) can be recovered. Then:

Proposition 47. *The right action (42) and left coaction (43) are compatible in the sense of [15, Theorem 6.2.3].*

This means that there is a well-defined right–left bicrossproduct Hopf algebra $U(\mathfrak{b}_+) \bowtie k[D_0]$. Using [15, Theorem 6.2.3] to write out its presentation, this turns out to coincide with the presentation of $\mathcal{H}_{\text{CM}}^{\text{left}}$. Hence:

Theorem 48. *The bicrossproduct $U(\mathfrak{b}_+) \bowtie k[D_0]$ is isomorphic to the Connes–Moscovici Hopf algebra $\mathcal{H}_{\text{CM}}^{\text{left}}$.*

References

- [1] S. Baaj, G. Skandalis, C^* -algèbres de Hopf et théorie de Kasparov équivariante, *K-Theory* 2 (1989) 683–721.
- [2] A. Connes, D. Kreimer, Renormalization in quantum field theory and the Riemann–Hilbert problem. I. The Hopf algebra structure of graphs and the main theorem, *Comm. Math. Phys.* 210 (1) (2000) 249–273.
- [3] A. Connes, M. Marcolli, \mathbb{Q} -lattices: Quantum statistical mechanics and Galois theory, *J. Geom. Phys.* 56 (1) (2006) 2–23.
- [4] A. Connes, H. Moscovici, The local index formula in noncommutative geometry, *Geom. Funct. Anal.* 5 (2) (1995) 174–243.
- [5] A. Connes, H. Moscovici, Hopf algebras, cyclic cohomology and the transverse index theorem, *Comm. Math. Phys.* 198 (1998) 199–246.
- [6] A. Connes, H. Moscovici, Modular Hecke algebras and their Hopf symmetry, *Mosc. Math. J.* 4 (1) (2004) 67–109, 310.
- [7] J.-P. Ezin, S. Majid, F. Ngakeu, Classification of differentials and Cartan calculus on bicrossproducts, *Acta Appl. Math.* 84 (2004) 193–236.
- [8] H. Figueroa, J. Gracia-Bondia, Combinatorial Hopf algebras in quantum field theory, I, *Rev. Math. Phys.* 17 (8) (2005) 881–976.
- [9] G. Goldin, Lectures on diffeomorphism groups in quantum physics, in: J. Govaerts, M.N. Hounkonnou, A.Z. Msezane (Eds.), *Contemporary Problems in Mathematical Physics, Proceedings of the Third International Workshop*, Cotonou, Republic of Benin, 2003, World Scientific, 2004.
- [10] G.I. Kac, V.G. Paljutkin, Finite ring groups, *Trans. Amer. Math. Soc.* 15 (1966) 251–294.
- [11] J. Lukierski, H. Ruegg, Quantum κ -Poincaré in any dimension, *Phys. Lett. B* 329 (2–3) (1994) 189–194.
- [12] S. Majid, Non-commutative-geometric groups by a bicrossproduct construction: Hopf algebras at the Planck scale, PhD thesis, Harvard University Archives, 1988.
- [13] S. Majid, Hopf–von Neumann algebra bicrossproducts, Kac algebra bicrossproducts, and the classical Yang–Baxter equations, *J. Funct. Anal.* 95 (1991) 291–319.
- [14] S. Majid, Quantum and braided diffeomorphism groups, *J. Geom. Phys.* 28 (1998) 94–128.
- [15] S. Majid, *Foundations of Quantum Group Theory*, Cambridge Univ. Press, 2000.
- [16] S. Majid, R. Oeckl, Twisting of quantum differentials and the Planck scale Hopf algebra, *Comm. Math. Phys.* 205 (1999) 617–655.
- [17] S. Majid, H. Ruegg, Bicrossproduct structure of the κ -Poincaré group and non-commutative geometry, *Phys. Lett. B* 334 (1994) 348–354.

- [18] R. Oeckl, Classification of differential calculi on $U_q(\mathfrak{b}_+)$, classical limits, and duality, *J. Math. Phys.* 40 (7) (1999) 3588–3603.
- [19] M. Takeuchi, Matched pairs of groups and bismash products, *Comm. Algebra* 9 (1981) 841–882.
- [20] S. Vaes, Examples of locally compact quantum groups through the bicrossed product construction, in: XIIIth International Congress on Mathematical Physics, London, 2000, Int. Press, Boston, MA, 2001, pp. 341–348.
- [21] S.L. Woronowicz, Differential calculus on compact matrix pseudogroups (quantum groups), *Comm. Math. Phys.* 122 (1989) 125–170.