

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

Appl. Comput. Harmon. Anal. 20 (2006) 26–40

---



---

**Applied and  
Computational  
Harmonic Analysis**


---



---

[www.elsevier.com/locate/acha](http://www.elsevier.com/locate/acha)

## Frame expansions for Gabor multipliers

John J. Benedetto<sup>a,\*</sup>, Götz E. Pfander<sup>b,2</sup><sup>a</sup> *Department of Mathematics, University of Maryland, College Park, MD 20742, USA*<sup>b</sup> *School of Engineering and Science, International University Bremen, 28759 Bremen, Germany*

Received 1 November 2004; accepted 12 March 2005

Available online 27 July 2005

Communicated by Akram Aldroubi

---

### Abstract

Discrete Gabor multipliers are composed of rank one operators. We shall prove, in the case of rank one projection operators, that the generating operators for such multipliers are either Riesz bases (exact frames) or not frames for their closed linear spans. The same dichotomy conclusion is valid for general rank one operators under mild and natural conditions. This is relevant since discrete Gabor multipliers have an emerging role in communications, radar, and waveform design, where redundant frame decompositions are increasingly applicable.

© 2005 Elsevier Inc. All rights reserved.

*Keywords:* Gabor multipliers; Gabor frames; Symplectic Fourier transforms

---

### 1. Introduction

Inspired and initiated by von Neumann in quantum mechanics [1, pp. 405 ff.], and Gabor in communications and acoustics [2], decompositions of functions  $f \in L^2(\mathbb{R}^d)$ , such as

$$f = \sum_{(x,\xi) \in \Lambda} \langle f, M_\xi T_x h \rangle M_\xi T_x g, \quad (1)$$

---

\* Corresponding author.

*E-mail address:* [jjb@math.umd.edu](mailto:jjb@math.umd.edu) (J.J. Benedetto).

<sup>1</sup> John J. Benedetto was supported in part by NSF DMS Grant 0139759.

<sup>2</sup> Götz E. Pfander was supported by the Max-Kade Foundation.

have become a fundamental tool in time–frequency analysis and applications dealing with time-varying spectra, e.g., [3–8]. In (1),  $g, h \in L^2(\mathbb{R}^d)$  are given square integrable functions on Euclidean space  $\mathbb{R}^d$ ,  $\Lambda \subset \mathbb{R}^d \times \mathbb{R}^d$  is a full rank lattice (such as  $\mathbb{Z}^{2d}$ ) where  $\hat{\mathbb{R}}^d$  is  $\mathbb{R}^d$  considered as a spectral domain,  $T_x$  is the translation operator  $T_x k(y) = k(y-x)$ ,  $M_\xi$  is the modulation operator  $M_\xi k(y) = e^{2\pi i y \cdot \xi} k(y)$ ,  $\langle \cdot, \cdot \rangle$  is the inner product in  $L^2(\mathbb{R}^d)$ , and convergence is in  $L^2(\mathbb{R}^d)$ .

The expansion (1) can be written operator theoretically in terms of the resolution of the identity  $Id_{L^2} : L^2(\mathbb{R}^d) \rightarrow L^2(\mathbb{R}^d)$  as

$$Id_{L^2} = \sum_{\lambda \in \Lambda} \rho(\lambda) P_{g,h}, \tag{2}$$

where the rank one operator  $P_{g,h} : L^2(\mathbb{R}^d) \rightarrow L^2(\mathbb{R}^d)$  is defined by  $f \mapsto \langle f, h \rangle g$ , and where  $\rho(\lambda) P_{g,h}$  is the composition (conjugation)

$$\rho(\lambda) P_{g,h} = \pi(\lambda) \circ P_{g,h} \circ \pi(\lambda)^*, \tag{3}$$

for  $\pi(\lambda)k(y) = M_\xi T_x k(y)$ ,  $\lambda = (x, \xi) \in \Lambda$ , and for the adjoint  $U^*$  of the unitary operator  $U$ . The equivalence of (1) and (2) follows from the elementary calculation

$$\pi(\lambda) \circ P_{g,h} \circ \pi(\lambda)^* f(y) = \langle f, \pi(\lambda)h \rangle \pi(\lambda)g(y). \tag{4}$$

Further, the right side of (4) is

$$\int_{\mathbb{R}^d} f(z) e^{-2\pi i(z-y) \cdot \xi} g(y-x) \overline{h(z-x)} dz;$$

and so, from (3) and (4),  $\rho(\lambda) P_{g,h}$  is a Hilbert–Schmidt operator with kernel  $k_\lambda(y, z) = e^{-2\pi i(z-y) \cdot \xi} g(y-x) \overline{h(z-x)}$ , where  $\lambda = (x, \xi)$ . We denote  $P_{g,g}$  by  $P_g$ .

In this context, the “dichotomy” theorem we shall prove, under mild necessary conditions on  $g$  and  $h$ , is that  $\{\rho(\lambda) P_{g,h}\}_{\lambda \in \Lambda}$  is a Riesz basis for its closed linear span in the space  $HS(\mathbb{R}^d)$  of Hilbert–Schmidt operators, or it is not a frame for this span, see Theorem 3.1.

The reason for the abstraction to the setting of  $HS(\mathbb{R}^d)$  for our theorem is the emerging importance of Gabor multipliers  $G_m$ , which are formally defined by a weighted version of (2), namely

$$G_m = \sum_{\lambda \in \Lambda} m_\lambda \rho(\lambda) P_{g,h}, \quad m_\lambda \in \mathbb{C} \text{ for } \lambda \in \Lambda, \tag{5}$$

e.g., [9–11], and the revitalization of underspread operators in the mathematical community, e.g., [12].

As a concluding application of our dichotomy result and the inherent characterization of Riesz bases of the form  $\{\rho(\lambda) P_{g,h}\}_{\lambda \in \Lambda}$ , we shall describe the role of the volume of the lattice  $\Lambda$  in (5) in terms of operator identification. In fact, we shall show that with natural hypotheses the Gabor multiplier class spanned by  $\{\rho(\lambda) P_{g,h}\}_{\lambda \in \Lambda}$  is identifiable if the volume of  $\Lambda$  is greater than one and not identifiable if the volume of  $\Lambda$  is less than one, see Theorem 5.2.

We begin in Section 2 with mathematical preliminaries concerning Gabor analysis, Hilbert–Schmidt operators, and shift invariant spaces. Section 3 contains a precise statement and proof of our dichotomy result, mentioned above, as well as some related results. Section 4 is devoted to relevant examples, and Section 5 to Gabor multipliers and identification.

## 2. Preliminaries

Throughout this paper we shall use standard notation from harmonic analysis and in particular Gabor analysis as found in [7]. For example, we shall use the unitary Fourier transformation  $\mathcal{F}$  on  $L^2(\mathbb{R}^d)$  which is normalized to satisfy  $\mathcal{F}g(\gamma) = \hat{g}(\gamma) = \int_{\mathbb{R}^d} g(y)e^{-2\pi iy \cdot \gamma} dy$ ,  $\gamma \in \hat{\mathbb{R}}^d$ , for  $g \in L^1(\mathbb{R}^d) \cap L^2(\mathbb{R}^d)$ .

In addition, the notation  $P \asymp R$  on  $X$  is used to abbreviate the following statement: there exist  $A, B > 0$  such that for all  $x \in X$  we have  $AP(x) \leq R(x) \leq BP(x)$ .

### 2.1. Gabor analysis

The short time Fourier transform of  $f \in L^2(\mathbb{R}^d)$  with respect to a window function  $g \in L^2(\mathbb{R}^d) \setminus \{0\}$  is given by

$$V_g f(\lambda) = \langle f, \pi(\lambda)g \rangle = \int_{\mathbb{R}^d} f(y) \overline{g(y-x)} e^{-2\pi iy \cdot \xi} dy, \quad \lambda = (x, \xi) \in \mathbb{R}^d \times \hat{\mathbb{R}}^d.$$

We have  $V_g f \in L^2(\mathbb{R}^d \times \hat{\mathbb{R}}^d)$  and  $\|V_g f\|_{L^2(\mathbb{R}^d \times \hat{\mathbb{R}}^d)} = \|f\|_{L^2(\mathbb{R}^d)} \|g\|_{L^2(\mathbb{R}^d)}$ . Further, we can *synthesize*  $f \in L^2(\mathbb{R}^d)$ , using translates and modulates of any  $h \in L^2(\mathbb{R}^d)$  with  $\langle h, g \rangle = 1$ , in the sense that the integral  $\int_{\lambda} V_g f(\lambda) \pi(\lambda) h d\lambda$  converges weakly to  $f$ .

A central goal in Gabor analysis is to find  $g, h \in L^2(\mathbb{R}^d)$  and full rank lattices  $\Lambda \subset \mathbb{R}^d \times \hat{\mathbb{R}}^d$  which allow a *discretization* of the reconstruction formula  $f \equiv \int_{\lambda} V_g f(\lambda) \pi(\lambda) h d\lambda$  of the form

$$f = \sum_{\lambda \in \Lambda} V_g f(\lambda) \pi(\lambda) h, \quad f \in L^2(\mathbb{R}^d), \quad (6)$$

with convergence in  $L^2(\mathbb{R}^d)$  and where  $g$  and  $h$  are independent of  $f$ .

A discussion of the validity of (6) entails Bessel sequences, Gabor frames, and Riesz bases, notions we now define.

Let  $g \in L^2(\mathbb{R}^d)$  and let  $\Lambda \subset \mathbb{R}^d \times \hat{\mathbb{R}}^d$  be a full rank lattice. Formally, consider the discrete analysis operator  $C_g$  defined by

$$C_g : L^2(\mathbb{R}^d) \rightarrow l^2(\Lambda), \quad f \mapsto \{V_g f(\lambda)\}_{\lambda \in \Lambda},$$

and the discrete synthesis operator  $T_g = C_g^*$  defined by

$$T_g : l^2(\Lambda) \rightarrow L^2(\mathbb{R}^d), \quad \{c_\lambda\}_{\lambda \in \Lambda} \mapsto \sum_{\lambda \in \Lambda} c_\lambda \pi(\lambda) g.$$

The set  $(g, \Lambda) = \{\pi(\lambda)g\}_{\lambda \in \Lambda}$  is called a *Gabor system*; and a Gabor system is a *Bessel sequence* if  $C_g$  is a well-defined linear operator in which case both  $C_g$  and  $T_g$  are bounded. A Bessel sequence  $(g, \Lambda)$  is a *frame* for  $L^2(\mathbb{R}^d)$  if  $C_g$  is also stable, i.e., if  $\|f\|_{L^2(\mathbb{R}^d)} \asymp \|C_g f\|_{l^2(\Lambda)}$  for  $f \in L^2(\mathbb{R}^d)$ , and it is a *Riesz basis (bounded unconditional basis)* for its closed linear span in  $L^2(\mathbb{R}^d)$  if  $T_g$  is stable in addition to being bounded, i.e., if  $\|\{c_\lambda\}\|_{l^2(\Lambda)} \asymp \|T_g \{c_\lambda\}\|_{L^2(\mathbb{R}^d)}$ , for  $\{c_\lambda\} \in l^2(\Lambda)$ .

The right-hand side of (6) is well defined if the Gabor systems  $(g, \Lambda)$  and  $(h, \Lambda)$  are Bessel sequences. Further, if  $(g, \Lambda)$  and  $(h, \Lambda)$  are frames, then the operator

$$S_{g,h} : L^2(\mathbb{R}^d) \rightarrow L^2(\mathbb{R}^d), \quad f \mapsto \sum_{\lambda \in \Lambda} V_g f(\lambda) \pi(\lambda) h$$

is an algebraic and topological isomorphism. If  $(g, \Lambda)$  and  $(h, \Lambda)$  are frames, and  $S_{g,h} = I : f \mapsto f$ , we say that  $(h, \Lambda)$  is a *dual frame* of  $(g, \Lambda)$  [7].

Fundamental to the analysis of Gabor systems  $(g, \Lambda)$  is the *volume*  $|\Lambda|$  of the full rank lattice  $\Lambda$ , which is given by  $|\Lambda| = |\det A|$ , where  $A$  is chosen such that  $A\mathbb{Z}^{2d} = \Lambda$ . In fact, if  $(g, \Lambda)$  is a Riesz basis for its closed linear span in  $L^2(\mathbb{R}^d)$ , then  $|\Lambda| \geq 1$ ; and if  $(g, \Lambda)$  is a frame for  $L^2(\mathbb{R}^d)$ , and therefore complete in  $L^2(\mathbb{R}^d)$ , then  $|\Lambda| \leq 1$ .

In the case that  $\Lambda$  has *critical density*, i.e., if  $|\Lambda| = 1$ , and  $(g, \Lambda)$  is a frame for  $L^2(\mathbb{R}^d)$ , then  $(g, \Lambda)$  is automatically a Riesz basis for  $L^2(\mathbb{R}^d)$ , or, equivalently, an *exact frame* for  $L^2(\mathbb{R}^d)$ , i.e.,  $(g, \Lambda)$  ceases to be a frame if any one of its elements is removed. In case  $|\Lambda| < 1$ , any frame  $(g, \Lambda)$  for  $L^2(\mathbb{R}^d)$  is *non-exact (overcomplete)*, and one can remove any finite number of elements from  $(g, \Lambda)$  and the resulting family remains a frame for  $L^2(\mathbb{R}^d)$ . Further, if  $|\Lambda| < 1$ , then, for any  $g \in L^2(\mathbb{R}^d)$ , there exists non-trivial  $\{c_\lambda\}_{\lambda \in \Lambda} \in l^2(\Lambda) \setminus \{0\}$  for which  $0 = \sum_{\lambda \in \Lambda} c_\lambda \pi(\lambda)g$  in  $L^2(\mathbb{R}^d)$ .

The uncertainty principle provides insight into any decomposition such as (6) [13–18]. For example, in the case of Gabor systems one manifestation of the uncertainty principle is the Balian–Low theorem [19–21], which asserts that if  $(g, \Lambda)$  is a Riesz basis for  $L^2(\mathbb{R}^d)$ , in which case we have  $|\Lambda| = 1$ , then  $g$  cannot be well localized in time and frequency, in the sense that

$$\left( \int |yg(y)|^2 dy \right) \cdot \left( \int |\eta\hat{g}(\eta)|^2 d\eta \right) = \infty \tag{7}$$

must occur.

We shall sometimes use the Feichtinger algebra  $S_0(\mathbb{R}^d)$  in place of  $L^2(\mathbb{R}^d)$ .  $S_0(\mathbb{R}^d)$  is the Banach algebra composed of those functions  $f \in L^2(\mathbb{R}^d)$  with the property that  $V_{g_0}f \in L^1(\mathbb{R}^d \times \hat{\mathbb{R}}^d)$  for the Gaussian  $g_0(x) = e^{-\|x\|^2}$ ,  $x \in \mathbb{R}^d$ . The norm  $\|f\|_{S_0(\mathbb{R}^d)} = \|V_{g_0}f\|_{L^1(\mathbb{R}^d \times \hat{\mathbb{R}}^d)}$  gives  $S_0(\mathbb{R}^d)$  a Banach algebra structure under pointwise multiplication and/or convolution. For equivalent definitions of  $S_0(\mathbb{R}^d)$ , as well as basic theory, see [22].

### 2.2. Hilbert–Schmidt operators

A Hilbert–Schmidt operator  $H \in HS(\mathbb{R}^d)$  is a compact integral operator on  $L^2(\mathbb{R}^d)$ , i.e.,  $H$  is defined by

$$Hf(x) = \int \kappa_H(x, t)f(t) dt = \int \kappa_H(x, x - t)f(x - t) dt \quad \text{a.e., } f \in L^2(\mathbb{R}^d),$$

with kernel  $\kappa_H \in L^2(\mathbb{R}^{2d})$ . The space of Hilbert–Schmidt operators is a Hilbert space with inner product  $\langle H_1, H_2 \rangle_{HS} = \langle \kappa_{H_1}, \kappa_{H_2} \rangle_{L^2}$  [23,24]. For any orthonormal basis  $\{e_i\}_{i \in I}$  of  $L^2(\mathbb{R}^d)$  we have

$$\|H\|_{HS}^2 = \langle H, H \rangle_{HS} = \sum_{i \in I} \|He_i\|_{L^2(\mathbb{R}^d)}^2,$$

and therefore  $\|H\|_{HS} \geq \|H\|_{\mathcal{L}}$  where  $\|\cdot\|_{\mathcal{L}}$  denotes the operator norm of  $H \in \mathcal{L}(L^2(\mathbb{R}^d), L^2(\mathbb{R}^d))$ .

Our use of families of Hilbert–Schmidt operators is carried out on a symbolic level. For any Hilbert–Schmidt operator  $H$  with kernel  $\kappa_H \in L^2(\mathbb{R}^{2d})$ , the *Kohn–Nirenberg symbol*  $\sigma_H$  of  $H$  is defined as

$$\sigma_H(\lambda) = \sigma_H(x, \xi) = \int_{\mathbb{R}^d} \kappa_H(x, x - y)e^{-2\pi iy \cdot \xi} dy \quad \text{a.e.}$$

[25]. The operator  $H$  can then be expressed using the Kohn–Nirenberg symbol by means of the formula

$$Hf(x) = \int_{\hat{\mathbb{R}}^d} \sigma_H(x, \xi) \hat{f}(\xi) e^{2\pi i x \cdot \xi} d\xi \quad \text{a.e.}$$

Critical to our analysis is the fact that the linear operator  $K : \kappa_H \mapsto \sigma_H$  is the composition of a partial Fourier transformation and a volume preserving axis transformation. Hence,  $K$  is unitary and, consequently,

$$\langle H_1, H_2 \rangle_{HS} = \langle \kappa_{H_1}, \kappa_{H_2} \rangle_{L^2(\mathbb{R}^{2d})} = \langle \sigma_{H_1}, \sigma_{H_2} \rangle_{L^2(\mathbb{R}^d \times \hat{\mathbb{R}}^d)}. \tag{8}$$

Since, in addition to (8), the Kohn–Nirenberg symbol of  $\rho(\lambda)H = \pi(\lambda) \circ H \circ \pi(\lambda)^*$  for  $H \in HS(\mathbb{R}^d)$  and  $\lambda \in \mathbb{R}^d \times \hat{\mathbb{R}}^d$  satisfies  $\sigma_{\rho(\lambda)H} = T_\lambda \sigma_H$ , we obtain that  $\{\rho(\lambda)P_{g,h}\}_{\lambda \in \Lambda}$  is a frame or Riesz basis for its closed linear span in  $HS(\mathbb{R}^d)$  if and only if  $\{T_\lambda \sigma_{P_{g,h}}\}_{\lambda \in \Lambda}$  is a frame or Riesz basis for its closed linear span in  $L^2(\mathbb{R}^d \times \hat{\mathbb{R}}^d)$ . The question of asking if  $\{T_\lambda \sigma_{P_{g,h}}\}_{\lambda \in \Lambda}$  is a frame or Riesz basis for the closed shift invariant space generated by  $\{T_\lambda \sigma_{P_{g,h}}\}_{\lambda \in \Lambda}$  in  $L^2(\mathbb{R}^d \times \hat{\mathbb{R}}^d)$  can be answered using zero set criteria for spectral periodizations, e.g., [17,26–28] and Theorem 2.1.

### 2.3. Shift-invariance of functions defined on phase space

We have reduced the analysis of sequences  $\{\rho(\lambda)P_{g,h}\}_{\lambda \in \Lambda}$  of Hilbert–Schmidt operators in  $HS(\mathbb{R}^d)$  to the analysis of function sequences  $\{T_\lambda \sigma_{P_{g,h}}\}_{\lambda \in \Lambda}$  in  $L^2(\mathbb{R}^d \times \hat{\mathbb{R}}^d)$ . Since the sequences  $\{\rho(\lambda)P_{g,h}\}_{\lambda \in \Lambda}$  are defined on phase space, we shall state a symplectic version of Theorem 1.4.1 in [17] as Theorem 2.1, part (b). The Fourier version of Theorem 2.1, part (a), is well-known and elementary to prove; and so the proof of Theorem 2.1, part (a), is also straightforward.

The symplectic Fourier transformation,  $\mathcal{F}_s$ , of functions defined on the phase space  $\mathbb{R}^d \times \hat{\mathbb{R}}^d$  is formally defined as follows:

$$\mathcal{F}_s : L^2(\mathbb{R}^d \times \hat{\mathbb{R}}^d) \rightarrow L^2(\mathbb{R}^d \times \hat{\mathbb{R}}^d), \quad f \mapsto \mathcal{F}_s f : \lambda \mapsto \int_{\mathbb{R}^d \times \hat{\mathbb{R}}^d} f(\lambda') e^{-2\pi i [\lambda', \lambda]} d\lambda',$$

where

$$[\lambda', \lambda] = [(x', \xi'), (x, \xi)] = x' \cdot \xi - \xi' \cdot x, \quad \lambda, \lambda' \in \mathbb{R}^d \times \hat{\mathbb{R}}^d \tag{9}$$

is the standard symplectic form on  $\mathbb{R}^d \times \hat{\mathbb{R}}^d$ .

The dual lattice of  $\Lambda$  with respect to the standard symplectic form on  $\mathbb{R}^d \times \hat{\mathbb{R}}^d$  is the so-called *adjoint lattice*  $\Lambda^\circ \subset \mathbb{R}^d \times \hat{\mathbb{R}}^d$  of  $\Lambda$ ; and it is defined by the rule:  $\lambda' \in \Lambda^\circ$  if and only if  $[\lambda', \lambda] \in \mathbb{Z}$  for all  $\lambda \in \Lambda$ . Therefore, if  $\Lambda = a\mathbb{Z} \times b\mathbb{Z}$  then we have  $\Lambda^\circ = \frac{1}{b}\mathbb{Z} \times \frac{1}{a}\mathbb{Z}$ , and, in general, we have  $|\Lambda^\circ| = |\Lambda|^{-1}$  [29,30].

To illustrate the important role of the adjoint lattice and, consequently, the symplectic Fourier transformation, in time–frequency analysis, we mention the fact that  $(h, \Lambda)$  is a Riesz basis for its closed linear span in  $L^2(\mathbb{R}^d)$  if and only if  $(h, \Lambda^\circ)$  is a frame for  $L^2(\mathbb{R}^d)$  [30,31].

In the following,  $P_\Lambda$  denotes periodization by the lattice  $\Lambda$ , i.e.,  $P_\Lambda F(\lambda) = \sum_{\lambda' \in \Lambda} F(\lambda - \lambda')$ ,  $\lambda \in \mathbb{R}^d \times \hat{\mathbb{R}}^d / \Lambda$ .

**Theorem 2.1.** Given  $F \in L^2(\mathbb{R}^d \times \hat{\mathbb{R}}^d)$  and a full rank lattice  $\Lambda \subset \mathbb{R}^d \times \hat{\mathbb{R}}^d$ .

- (a) The family  $\{T_\lambda F\}_{\lambda \in \Lambda}$  is a Riesz basis for its closed linear span in  $L^2(\mathbb{R}^d \times \hat{\mathbb{R}}^d)$  if and only if  $P_{\Lambda^\circ} |\mathcal{F}_s F|^2 \asymp 1$  a.e. on  $\mathbb{R}^d \times \hat{\mathbb{R}}^d / \Lambda^\circ$ .
- (b) The family  $\{T_\lambda F\}_{\lambda \in \Lambda}$  is a frame for its closed linear span in  $L^2(\mathbb{R}^d \times \hat{\mathbb{R}}^d)$  if and only if  $P_{\Lambda^\circ} |\mathcal{F}_s F|^2 \asymp 1$  a.e. on  $(\mathbb{R}^d \times \hat{\mathbb{R}}^d / \Lambda^\circ) \setminus \{x: P_{\Lambda^\circ} |\mathcal{F}_s F|^2(x) = 0\}$ .

Theorem 2.1 and the material of Section 2.2 illustrate that the analysis of  $\{T_\lambda \sigma_{P_{g,h}}\}_{\lambda \in \Lambda}$  in  $L^2(\mathbb{R}^d \times \hat{\mathbb{R}}^d)$  and therefore of  $\{\rho(\lambda) P_{g,h}\}_{\lambda \in \Lambda}$  in  $HS(\mathbb{R}^d)$  requires only a thorough investigation of  $P_{\Lambda^\circ} |\mathcal{F}_s \sigma_{P_{g,h}}|^2$  on  $\mathbb{R}^d \times \hat{\mathbb{R}}^d / \Lambda^\circ$ .

To this end, note that for any rank one operator  $P_{g,h}$  we have

$$\sigma_{P_{g,h}}(\lambda) = \sigma_{P_{g,h}}(x, \xi) = \int_{\mathbb{R}^d} g(x) \bar{h}(x - y) e^{-2\pi i y \cdot \xi} dy = e^{-2\pi i x \cdot \xi} g(x) \bar{h}(\xi) \quad \text{a.e.}$$

and therefore

$$\begin{aligned} \mathcal{F}_s \sigma_{P_{g,h}}(\lambda) &= \mathcal{F}_s \sigma_{P_{g,h}}(x, \xi) = \int_{\mathbb{R}^d} \int_{\hat{\mathbb{R}}^d} g(x') \bar{h}(\xi') e^{-2\pi i (x' \cdot \xi' + x' \cdot \xi - x \cdot \xi')} dx' d\xi' \\ &= \int_{\mathbb{R}^d} g(x') \overline{h(x' - x)} e^{-2\pi i x' \cdot \xi} dx' = V_h g(x, \xi) = V_h g(\lambda). \end{aligned} \tag{10}$$

The results of Sections 2.2 and 2.3 allow us to prove Theorem 3.1.

### 3. Results

In [9], Feichtinger proved that if  $(g, \Lambda)$  is a Gabor frame for  $L^2(\mathbb{R}^d)$  generated by  $g \in S_0(\mathbb{R}^d)$ , then  $\{\rho(\lambda) P_g\}_{\lambda \in \Lambda}$  is a Riesz basis for its closed linear span in  $HS(\mathbb{R}^d)$  if and only if  $|\Psi| \asymp 1$ , where  $\Psi(\chi) = \sum_{\lambda \in \Lambda} |V_g g(\lambda)|^2 e^{2\pi i \lambda \cdot \chi}$ ,  $\chi \in \mathbb{R}^d \times \hat{\mathbb{R}}^d / \Lambda$ .

Theorem 3.1, part (a), is essentially Feichtinger’s theorem; and is, itself, the motivation for Theorem 3.1, parts (b) and (c). Theorem 3.1, part (b), and Theorem 3.2 are precise statements of our main theorem which was stated without hypotheses in Section 1. We emphasize that this is a *dichotomy theorem*, asserting that  $\{\rho(\lambda) P_{g,h}\}_{\lambda \in \Lambda}$  is either a Riesz basis or not a frame for its closed linear span in  $HS(\mathbb{R}^d)$ .

**Theorem 3.1.** Let  $g, h \in L^2(\mathbb{R}^d)$  and let  $\Lambda \subset \mathbb{R}^d \times \hat{\mathbb{R}}^d$  be a full rank lattice.

- (a) The family  $\{\rho(\lambda) P_{g,h}\}_{\lambda \in \Lambda}$  is a Riesz basis for its closed linear span in  $HS(\mathbb{R}^d)$  if and only if  $P_{\Lambda^\circ} |V_h g|^2 \asymp 1$  on  $\mathbb{R}^d \times \hat{\mathbb{R}}^d / \Lambda^\circ$ .
- (b) If  $(g, \Lambda)$  and  $(h, \Lambda)$  are Bessel sequences in  $L^2(\mathbb{R}^d)$ , then  $\{\rho(\lambda) P_{g,h}\}_{\lambda \in \Lambda}$  is either a Riesz basis or not a frame for its closed linear span in  $HS(\mathbb{R}^d)$ .
- (c) If  $g, h \in S_0(\mathbb{R}^d) \setminus \{0\}$ , then there exists  $r > 0$  such that, for all  $\alpha > r > 0$ ,  $\{\rho(\alpha \lambda) P_{g,h}\}_{\lambda \in \Lambda}$  is a Riesz basis for its closed linear span in  $HS(\mathbb{R}^d)$ .

In the case of  $g = h$ , we can drop the Bessel sequence condition in Theorem 3.1, part (b), and we obtain the following result.

**Theorem 3.2.** *Let  $g \in L^2(\mathbb{R}^d)$  and let  $\Lambda \subset \mathbb{R}^{2d}$  be a full rank lattice.  $\{\rho(\lambda)P_g\}_{\lambda \in \Lambda}$  is either a Riesz basis or not a frame for its closed linear span in  $HS(\mathbb{R}^d)$ .*

For the calculations in the proofs of Theorems 3.1 and 3.2, we need the following simple facts.

**Lemma 3.3.** *For  $g, h \in L^2(\mathbb{R}^d)$  we have  $\mathcal{F}_s|V_hg|^2 = V_hh\overline{V_gg}$ . For  $g = h$ , this is  $\mathcal{F}_s|V_gg|^2 = |V_gg|^2$ .*

Lemma 3.3 is proven in [18, p. 17].

**Lemma 3.4.** *Let  $F_n : \mathbb{R}^d \rightarrow \mathbb{R}^+$ ,  $n \in \mathbb{N}$ , be continuous functions with  $\sum_{n \in \mathbb{N}} F_n(x) \leq B$  a.e. Then  $\sum_{n \in \mathbb{N}} F_n(x) \leq B$  for all  $x \in \mathbb{R}^d$ .*

**Proof.** If there is  $x_0$  for which  $\infty \geq A = \sum_{n=1}^\infty F_n(x_0) > B$ , then there exists  $N \in \mathbb{N}$  such that

$$G_N(x_0) = \sum_{n=1}^N F_n(x_0) \geq \frac{1}{2}B + \frac{1}{2} \min\{A, B + 1\}.$$

Since  $G_N$  is continuous, there exists an open set  $V \subset \mathbb{R}^d$  such that

$$\sum_{n=1}^\infty F_n \geq G_N > \frac{3}{4}B + \frac{1}{4} \min\{A, B + 1\} > B$$

on  $V$ , and this is a contradiction.  $\square$

The crucial lemma to prove Theorem 3.1, part (b), and Theorem 3.2 is the following result.

**Lemma 3.5.** *Let  $\Lambda$  be a full rank lattice in  $\mathbb{R}^d \times \hat{\mathbb{R}}^d$ , and let  $g, h \in L^2(\mathbb{R}^d)$  with  $P_\Lambda|V_hg|^2 \in L^\infty(\mathbb{R}^d \times \hat{\mathbb{R}}^d / \Lambda)$ . If  $h = g$ , or if  $(g, \Lambda^\circ)$  and  $(h, \Lambda^\circ)$  are Bessel sequences, then  $P_\Lambda|V_hg|^2 = \Phi$  a.e. for some function  $\Phi$  which is continuous on  $\mathbb{R}^d \times \hat{\mathbb{R}}^d / \Lambda$ .*

**Proof.** We shall twice apply the Poisson Summation Formula for the symplectic Fourier transform. To this end, we define the symplectic Fourier transformation on  $L^2(\mathbb{R}^d \times \hat{\mathbb{R}}^d / \Lambda)$  as follows:

$$\mathcal{F}_s : L^2(\mathbb{R}^d \times \hat{\mathbb{R}}^d / \Lambda) \rightarrow l^2(\Lambda^\circ), \quad \mathcal{F}_s F(\lambda) = \int_{\mathbb{R}^d \times \hat{\mathbb{R}}^d / \Lambda} F(\lambda') e^{-2\pi i[\lambda', \lambda]} d\lambda'.$$

For  $F \in L^1(\mathbb{R}^d \times \hat{\mathbb{R}}^d)$  with  $P_\Lambda F \in L^2(\mathbb{R}^d \times \hat{\mathbb{R}}^d / \Lambda)$  and  $\lambda \in \Lambda^\circ$ , we have

$$\mathcal{F}_s P_\Lambda F(\lambda) = \int_{\mathbb{R}^d \times \hat{\mathbb{R}}^d / \Lambda} \left( \sum_{\lambda'' \in \Lambda} F(\lambda' - \lambda'') \right) e^{-2\pi i[\lambda', \lambda]} d\lambda' = \int_{\mathbb{R}^d \times \hat{\mathbb{R}}^d} F(\lambda') e^{-2\pi i[\lambda', \lambda]} d\lambda' = \mathcal{F}_s F(\lambda).$$

Therefore, the Poisson Summation Formula,

$$P_\Lambda F = |\Lambda^\circ| \sum_{\lambda \in \Lambda^\circ} \mathcal{F}_s F(\lambda) e^{2\pi i[\cdot, \lambda]}, \tag{11}$$

with convergence of the right-hand side in  $L^2(\mathbb{R}^d \times \hat{\mathbb{R}}^d / \Lambda)$ , is valid. We apply (11) and Lemma 3.3 to  $|V_h g|^2 \in L^1(\mathbb{R}^d \times \hat{\mathbb{R}}^d)$  with  $P_\Lambda |V_h g|^2 \in L^\infty(\mathbb{R}^d \times \hat{\mathbb{R}}^d / \Lambda) \subset L^2(\mathbb{R}^d \times \hat{\mathbb{R}}^d / \Lambda)$  to obtain

$$P_\Lambda |V_h g|^2(\lambda') = |\Lambda^\circ| \sum_{\lambda \in \Lambda^\circ} V_h h(\lambda) \overline{V_g g(\lambda)} e^{2\pi i[\lambda', \lambda]} \quad \text{a.e. } \lambda' \in \mathbb{R}^d \times \hat{\mathbb{R}}^d / \Lambda. \tag{12}$$

If  $(g, \Lambda^\circ)$  and  $(h, \Lambda^\circ)$  are Bessel sequences, then  $\{V_h h(\lambda)\}_{\lambda \in \Lambda^\circ}, \{V_g g(\lambda)\}_{\lambda \in \Lambda^\circ} \in l^2(\Lambda^\circ)$  and, consequently,  $\{V_h h(\lambda) \overline{V_g g(\lambda)}\}_{\lambda \in \Lambda^\circ} \in l^1(\Lambda^\circ)$ . Hence, the right-hand side of (12) is absolutely convergent and so it is continuous on  $\mathbb{R}^d \times \hat{\mathbb{R}}^d / \Lambda$ .

Let us now turn to the case  $h = g$  and  $P_\Lambda |V_g g|^2 \in L^\infty(\mathbb{R}^d \times \hat{\mathbb{R}}^d / \Lambda)$ .  $P_\Lambda |V_g g|^2$  bounded and  $|V_g g|^2$  continuous and positive imply  $P_\Lambda |V_g g|^2(\lambda) \leq \|P_\Lambda |V_g g|^2\|_{L^\infty}$  for all  $\lambda \in \mathbb{R}^d \times \hat{\mathbb{R}}^d / \Lambda$  by Lemma 3.4. In particular,  $P_\Lambda |V_g g|^2(0, 0) \in \mathbb{R}$ , i.e.,  $\{|V_g g|^2(\lambda)\}_{\lambda \in \Lambda} \in l^1(\Lambda)$ . Since  $\mathcal{F}_s |V_g g|^2 = |V_g g|^2$ , the adjoint version of Eq. (11) implies that  $\{|V_g g|^2(\lambda)\}_{\lambda \in \Lambda}$  is the symplectic Fourier transform of  $P_{\Lambda^\circ} |V_g g|^2$  and, therefore,

$$P_{\Lambda^\circ} |V_g g|^2(\lambda') = |\Lambda| \sum_{\lambda \in \Lambda} |V_g g|^2(\lambda) e^{2\pi i[\lambda', \lambda]} \quad \text{a.e. } \lambda' \in \mathbb{R}^d \times \hat{\mathbb{R}}^d / \Lambda^\circ. \tag{13}$$

The right-hand side of (13) is continuous and therefore bounded. Hence,  $P_{\Lambda^\circ} |V_g g|^2 \in L^\infty(\mathbb{R}^d \times \hat{\mathbb{R}}^d / \Lambda)$  and, applying Lemma 3.4 again, we conclude that  $P_{\Lambda^\circ} |V_g g|^2$  is bounded everywhere. In particular, we have  $\{|V_g g|^2(\lambda)\}_{\lambda \in \Lambda^\circ} \in l^1(\Lambda^\circ)$ .

Replacing  $\Lambda$  by  $\Lambda^\circ$  in (13) and repeating the argument above, we conclude that  $\{|V_g g|^2(\lambda)\}_{\lambda \in (\Lambda^\circ)^\circ} \in l^1((\Lambda^\circ)^\circ)$ .

The argument is completed by observing that  $(\Lambda^\circ)^\circ = \Lambda$  [30, p. 257], and therefore

$$P_\Lambda |V_g g|^2(\lambda') = |\Lambda^\circ| \sum_{\lambda \in \Lambda^\circ} |V_g g|^2(\lambda) e^{2\pi i[\lambda', \lambda]} \quad \text{a.e. } \lambda' \in \mathbb{R}^d \times \hat{\mathbb{R}}^d / \Lambda,$$

where the right-hand side is continuous on  $\mathbb{R}^d \times \hat{\mathbb{R}}^d / \Lambda$ .  $\square$

**Proof of Theorem 3.1.** (a) The equivalence in Theorem 3.1, part (a), follows directly from (8), (10), and Theorem 2.1, part (a). Alternatively it can be derived simply by using Feichtinger’s result in [9] which we mentioned at the beginning of Section 3, since Lemma 3.3 and (11) imply  $P_{\Lambda^\circ} |V_g g|^2 = \mathcal{F}_s^{-1} \{|V_g g(\lambda)|^2\}_{\lambda \in \Lambda}$ . For  $g \neq h$ , Feichtinger’s criterion requires an analysis of the lower bound of  $|\mathcal{F}_s^{-1} \{V_h h(\lambda) \overline{V_g g(\lambda)}\}_{\lambda \in \Lambda}|$ , which is made significantly easier by means of our observation that

$$P_{\Lambda^\circ} |V_h g|^2 = \mathcal{F}_s^{-1} \{V_h h(\lambda) \overline{V_g g(\lambda)}\}_{\lambda \in \Lambda}.$$

(b) In order for  $\{\rho(\lambda) P_{g,h}\}_{\lambda \in \Lambda}$  to form a non-exact frame,  $G = P_{\Lambda^\circ} |V_h g|^2$  would have to be bounded, vanish on a set of positive measure and be bounded away from zero off this set. Obviously, this criteria cannot be fulfilled for continuous  $G$ , and Lemma 3.5 therefore implies Theorem 3.1, part (b), for the case that  $(g, \Lambda)$  and  $(h, \Lambda)$  are Bessel sequences.

(c) To prove Theorem 3.1, part (c), let us observe that  $g, h \in S_0$  implies that  $P_{\Lambda^\circ} |V_g g|^2$  converges absolutely and uniformly on  $\mathbb{R}^d \times \hat{\mathbb{R}}^d / \Lambda$  [7, p. 255]. Since  $V_g g$  is uniformly continuous for any  $g \in L^2(\mathbb{R}^d)$ , we conclude that  $P_{\Lambda^\circ} |V_g g|^2$  is continuous and bounded.



Since  $g, h \neq 0$  there exists  $\lambda_0 \in \mathbb{R}^d \times \hat{\mathbb{R}}^d$  such that  $V_h g(\lambda_0) \neq 0$ . Using the continuity of  $V_h g$ , we can find an open, convex, and bounded set  $V$  such that  $|V_h g|^2 \geq \frac{1}{2}|V_h g|^2 > 0$  on  $V$ . For any full rank lattice  $\Lambda$  we can now choose  $r > 0$  so that  $V$  contains a fundamental domain of  $r^{-1}\Lambda^\circ$ , guaranteeing a lower bound of  $P_{\alpha^{-1}\Lambda^\circ}|V_g g|^2$  for any  $\alpha \geq r$ . The fact that  $(\alpha\Lambda)^\circ = \alpha^{-1}\Lambda^\circ$  completes the proof.  $\square$

**Proof of Theorem 3.2.** See the proof of Theorem 3.1, part (b).  $\square$

**Remark 3.6.** The hypotheses in Theorem 3.1, part (b), and Theorem 3.1, part (c), can be weakened considerably. For example, we could replace the Bessel sequence hypothesis on  $(g, \Lambda)$  and  $(h, \Lambda)$  in Theorem 3.1, part (b), with the hypothesis that  $\mathcal{F}_s^{-1}\{V_h h(\lambda)\overline{V_g g}\}_{\lambda \in \Lambda^\circ}$  be continuous on  $\mathbb{R}^d \times \hat{\mathbb{R}}^d / \Lambda$ ; and the hypothesis that  $g, h \in S_0(\mathbb{R}^d)$  in Theorem 3.1, part (c), can be replaced with the hypothesis  $|V_h g|^2 \in S_0(\mathbb{R}^d \times \hat{\mathbb{R}}^d)$ .

#### 4. Examples

Let us now provide examples illustrating our results for the case  $d = 1$ . The first example in the case of a Gaussian was pointed out to us by Hans Feichtinger.

**Example 4.1.** For  $g \in S_0(\mathbb{R})$  with  $V_g g(\lambda) \neq 0$  for all  $\lambda \in \mathbb{R} \times \hat{\mathbb{R}}$ , e.g., let  $g$  be a Gaussian, say  $g(x) = e^{-x^2}$ , we have that  $\{\rho(\lambda)P_{g,g}\}_{\lambda \in \Lambda}$  is a Riesz basis in  $HS(\mathbb{R})$  for any full rank lattice  $\Lambda$ .

**Example 4.2.** There exist non-exact frames in  $HS(\mathbb{R})$  of the form  $\{\rho(\lambda)H\}_{\lambda \in \Lambda}$ , where  $H$  is not rank one. For example, we may define  $H$  by means of its Kohn–Nirenberg symbol by choosing  $\mathcal{F}_s \sigma_H = \mathbf{1}_{[0,1]^2}$  and  $\Lambda = \frac{1}{2}(\mathbb{Z} \times \mathbb{Z})$ . Since  $\{M_\lambda \mathcal{F}_s \sigma_H\}_{\lambda \in \Lambda}$  forms a non-exact frame for its closed linear span in  $L^2(\mathbb{R} \times \hat{\mathbb{R}})$ , so does  $\{T_\lambda \sigma_H\}$ ; and, therefore,  $\{\rho(\lambda)H\}_{\lambda \in \Lambda}$  forms a non-exact frame for its closed linear span in  $HS(\mathbb{R})$ .

Note that any such example implies  $\sigma_H \notin S_0(\mathbb{R} \times \hat{\mathbb{R}})$ , since otherwise  $P_{\Lambda^\circ}|\sigma_H|^2$  is continuous.

**Example 4.3.** There also exist non-exact frames in  $HS(\mathbb{R})$  of rank one operators with smooth kernels, e.g., let  $g_0$  be a Gaussian and set  $\Lambda = \{(n, \frac{m}{2})\}_{m,n \in \mathbb{Z}}$ . Then  $(g_0 \otimes g_0, \Lambda \otimes \Lambda)$  is a frame for  $L^2(\mathbb{R} \times \hat{\mathbb{R}})$  and we obtain that  $\{\pi(\lambda)P_g \pi(\lambda')\}_{\lambda, \lambda' \in \Lambda}$  is a non-exact frame (for its closed linear span in  $HS(\mathbb{R})$ ) composed of rank one operators.

**Example 4.4.** There exist  $g \in S_0(\mathbb{R})$  and  $\Lambda \subset \mathbb{R} \times \hat{\mathbb{R}}$  such that  $\{\rho(\lambda)P_g\}_{\lambda \in \Lambda}$  is not a frame for its closed linear span in  $HS(\mathbb{R})$ . Consider any  $g \in S_0(\mathbb{R})$  such that  $\text{supp } g \subseteq [0, 1]$ ; then  $V_g g(x, \xi) = 0$  for  $|x| > 1$ . If  $\Lambda = \beta\mathbb{Z} \times \frac{1}{4}\mathbb{Z}$ ,  $\beta > 0$ , we have  $\Lambda^\circ = 4\mathbb{Z} \times \frac{1}{\beta}\mathbb{Z}$ . Since in this case  $P_{\Lambda^\circ}|V_g g|^2(\lambda) = 0$  whenever  $\lambda \in [2, 4] \times [0, \frac{1}{\beta}] \subset \mathbb{R} \times \hat{\mathbb{R}} / 4\mathbb{Z} \times \frac{1}{\beta}\mathbb{Z}$ , we have that  $\{\rho(\lambda)P_g\}_{\lambda \in \beta\mathbb{Z} \times \frac{1}{4}\mathbb{Z}}$ ,  $\beta > 0$  is not a frame for its closed linear span in  $HS(\mathbb{R})$ .

**Example 4.5.** We now illustrate that Theorem 3.1, part (c), does not hold for arbitrary  $g, h \in L^2(\mathbb{R})$ .

Let  $g = \sum_{k=1}^\infty T_k D_{k^{\frac{3}{2}}} \mathbf{1}_{[-\frac{1}{2}, \frac{1}{2}]} \in L^2(\mathbb{R})$ , where  $D_a f(x) = f(ax)$ . For example, we have  $T_4 D_{4^{\frac{3}{2}}} \mathbf{1}_{[-\frac{1}{2}, \frac{1}{2}]} = \mathbf{1}_{[-\frac{1}{8} + 5, \frac{1}{8} + 5]}$  and  $\|T_k D_{k^{\frac{3}{2}}} \mathbf{1}_{[-\frac{1}{2}, \frac{1}{2}]}\|_1 = \|T_k D_{k^{\frac{3}{2}}} \mathbf{1}_{[-\frac{1}{2}, \frac{1}{2}]}\|_2^2 = k^{-\frac{3}{2}}$ .

We shall prove that  $\{\rho(\lambda)P_g\}_{\lambda \in K(\mathbb{Z} \times \mathbb{Z})}$  is not a frame for its closed linear span in  $HS(\mathbb{R})$  for any  $K \in \mathbb{N}$  by showing that, for any  $K \in \mathbb{N}$ ,

$$P_{\frac{1}{K}(\mathbb{Z} \times \mathbb{Z})} |V_g g|^2 \notin L^\infty \left( \mathbb{R} \times \hat{\mathbb{R}} / \frac{1}{K}(\mathbb{Z} \times \mathbb{Z}) \right). \tag{14}$$

To this end observe that

$$\begin{aligned} V_g g(n, 0) &= \int g(x)g(x - n) dx = \int \sum_{k=n+1}^\infty T_k D_{k^{\frac{3}{2}}} \mathbf{1}_{[-\frac{1}{2}, \frac{1}{2}]}(x) dx = \sum_{k=n+1}^\infty k^{-\frac{3}{2}} \\ &> \int_{n+2}^\infty x^{-\frac{3}{2}} dx = 2(n + 2)^{-\frac{1}{2}}. \end{aligned}$$

Therefore

$$P_{\frac{1}{K}(\mathbb{Z} \times \mathbb{Z})} |V_g g|^2(0, 0) = \sum_{n,m \in \mathbb{Z}} \left| V_g g \left( 0 - \frac{n}{K}, 0 - \frac{m}{K} \right) \right|^2 \geq \sum_{n \in \mathbb{Z}} |V_g g(n, 0)|^2 \geq 2 \sum_{n=1}^\infty (n + 2)^{-1} = \infty.$$

An application of Lemma 3.4 gives (14).

**Remark 4.6.** In [32] it is shown that for  $g_1, g_2 \in S_0(\mathbb{R})$ , and  $f_1, f_2 \in L^2(\mathbb{R})$  we have  $V_{g_1} f_1 \cdot \overline{V_{g_2} f_2} \in S_0(\mathbb{R} \times \hat{\mathbb{R}})$ .

Example 4.5, on the other hand, shows that there exists  $g \in L^2(\mathbb{R})$  such that  $|V_g g|^2 \notin S_0(\mathbb{R} \times \hat{\mathbb{R}})$  since for  $g$  constructed in Example 4.5 we have

$$\infty = P_{\mathbb{Z} \times \mathbb{Z}} |V_g g|^2(0) = \left\langle |V_g g|^2, \sum_{n \in \mathbb{Z}^2} \delta_n \right\rangle$$

with  $\sum_{n \in \mathbb{Z}} \delta_n \in S'_0(\mathbb{R} \times \hat{\mathbb{R}})$ .

**Example 4.7.** We shall now consider a classical example, namely  $g_c = \mathbf{1}_{[0,c]}$  and  $\Lambda = a\mathbb{Z} \times b\mathbb{Z}$ ,  $a, b > 0$ . The question for which  $a, b, c$  the Gabor system  $(g_c, a\mathbb{Z} \times b\mathbb{Z})$  is a frame has been analyzed extensively by Janssen [33].

Note that  $\{\rho(an, bm)P_{g_c}\}_{n,m \in \mathbb{Z}}$  is a frame or Riesz basis for its closed linear span for  $g_c = \mathbf{1}_{[0,c]}$  if and only if  $\{\rho(\frac{a}{c}n, bcm)P_{g_1}\}_{n,m \in \mathbb{Z}}$  is the same. Hence, we shall analyze the function  $g = g_1 = \mathbf{1}_{[0,1]}$ , see Figs. 1 and 2. In this case,

$$V_g g(x, \xi) = \begin{cases} \int_0^{1+x} e^{-2\pi i t \xi} dt, & \text{for } -1 \leq x \leq 0, \\ \int_x^1 e^{-2\pi i t \xi} dt, & \text{for } 0 \leq x \leq 1, \\ 0, & \text{for } |x| \geq 1, \end{cases}$$

and therefore

$$|V_g g|^2(x, \xi) = \begin{cases} \frac{\sin^2 \pi(1-|x|)\xi}{\pi^2 \xi^2}, & \text{for } |x| \leq 1, \\ 0, & \text{for } |x| \geq 1. \end{cases}$$




Fig. 1. (A) Zero set of  $V_{\mathbf{1}_{[0,1]}} \mathbf{1}_{[0,1]}$  (dark). (B) Set containing all pairs  $(\alpha, \beta)$ ,  $\alpha, \beta > 0$  such that  $P_{\alpha\mathbb{Z} \times \beta\mathbb{Z}} |V_g g|^2$  has a root (dark). We include as reference the curve  $\alpha\beta = 1$  (dashed). (C) Set containing all pairs  $(a, b)$ ,  $a, b > 0$  such that  $\{\rho(an, bm) P_g\}_{n,m \in \mathbb{Z}}$  is not a Riesz basis in  $HS(\mathbb{R})$  (dark). The curve  $ab = 1$  is included (dashed).




Fig. 2. (A) Janssen tie, i.e., set containing pairs  $(a, b)$ ,  $a, b > 0$ , such that  $(\mathbf{1}_{[0,1]}, a, b)$  is not a frame (dark), set containing pairs  $(a, b)$ ,  $a, b > 0$ , such that  $(\mathbf{1}_{[0,1]}, a, b)$  is a frame (white). In the light area, it is known that  $(\mathbf{1}_{[0,1]}, a, b)$  is a frame if  $ab$  is irrational. (B) Superposition of Janssen tie and Fig. 1.B. (C) Superposition of Janssen tie and Fig. 1.C.

Thus,  $P_{\Lambda^\circ} |V_g g|^2 = P_{\frac{1}{b}\mathbb{Z} \times \frac{1}{a}\mathbb{Z}} |V_g g|^2$  is continuous and bounded and we can rule out the existence of non-exact frames for any  $a, b$ .

Elementary calculations show that, for  $\alpha = \frac{1}{b}$  and  $\beta = \frac{1}{a}$ ,  $P_{\alpha\mathbb{Z} \times \beta\mathbb{Z}} |V_g g|^2$  is not bounded below if and only if  $\alpha = \frac{2}{3}$  and  $\beta = \frac{3}{2}n$  for  $n \in \mathbb{N} \setminus \{1\}$ , or  $\frac{2}{3} < \alpha < 1$  and  $\beta = \frac{2m}{2-\alpha}$  for  $m \in \mathbb{N} \setminus \{1\}$ , or  $\frac{2k+1}{3k+1} < \alpha < 1$  and  $\beta = \frac{2k+1}{2-\alpha}$  for  $k \in \mathbb{N} \setminus \{1\}$ , or  $1 \leq \alpha < 2$  and  $\beta \in \bigcup_{l \geq 2}^\infty [l, \frac{l}{2-\alpha}]$ , for  $l \in \mathbb{N} \setminus \{1\}$ , or  $2 \leq \alpha$ .

Hence,  $\{\rho(an, bm) P_g\}_{n,m \in \mathbb{Z}}$  is not a Riesz basis for its closed linear span in  $HS(\mathbb{R})$ , and, therefore, by Theorem 3.2, it is not a frame for its closed linear span in  $HS(\mathbb{R})$  if and only if  $b \leq \frac{1}{2}$ , or  $\frac{1}{2} < b \leq 1$  and  $a \in \bigcup_{l \geq 2}^\infty [\frac{2b-1}{bl}, \frac{1}{l}]$ , or  $1 < b < \frac{3k+1}{2k+1}$  and  $a = \frac{2b-1}{(2k+1)b}$  for  $k \in \mathbb{N} \setminus \{1\}$ , or  $1 < b < \frac{3}{2}$  and  $a = \frac{2b-1}{2mb}$  for  $m \in \mathbb{N} \setminus \{1\}$ , or  $b = \frac{3}{2}$  and  $\beta = \frac{2}{3n}$  for  $n \in \mathbb{N} \setminus \{1\}$ .

### 5. Gabor multipliers

Multipliers play a central role in functional and harmonic analysis. The theory of multipliers is based on simple pointwise multiplication operators  $\mathcal{M}_s : L^2(X) \rightarrow L^2(X)$ ,  $f \mapsto s \cdot f$ , where  $X$  is a measure space and  $s$  is a bounded function defined on  $X$  [34].

In applied harmonic analysis, frequency domain multipliers, i.e., convolution operators,  $\hat{\mathcal{M}}_{\hat{s}} : f \mapsto s * f$  where  $\widehat{s * f} = \hat{s} \cdot \hat{f}$ , are widely used, e.g., to model time-invariant channels in signal processing. Here, we shall consider operators which are composed of an analysis operator  $C : L^2(\mathbb{R}^d) \rightarrow L^2(X)$ , whose range consists of real or complex valued functions or sequences, a pointwise multiplication by a fixed function (sequence)  $s$  on  $X$ , i.e., by the symbol  $s$  of the operator, and a synthesis operator  $T : L^2(X) \rightarrow L^2(\mathbb{R}^d)$ . For example, we have  $\hat{\mathcal{M}}_{\hat{s}} = \mathcal{F}^{-1} \circ \mathcal{M}_{\hat{s}} \circ \mathcal{F}$ .

Continuous Gabor multipliers are given by

$$V_h^* \circ \mathcal{M}_F \circ V_g : L^2(\mathbb{R}^d) \rightarrow L^2(\mathbb{R}^d), \quad f \mapsto \int_{\lambda} F(\lambda) \cdot V_g f(\lambda) \pi(\lambda) h \, d\lambda,$$

for  $g, h \in L^2(\mathbb{R}^d)$ , and they are widely discussed in the literature, e.g., in [35–37]. In the following, we shall discuss *discrete Gabor multipliers* which, as noted in the Introduction, are formally given by

$$G_m f = T_h \circ \mathcal{M}_m \circ C_g f = \sum_{\lambda \in \Lambda} m_{\lambda} \langle f, \pi(\lambda) h \rangle \pi(\lambda) g = \sum_{\lambda \in \Lambda} m_{\lambda} (\rho(\lambda) P_{g,h}) f, \tag{15}$$

for  $f \in L^2(\mathbb{R}^d)$ , where  $\Lambda$  is a full rank lattice in  $\mathbb{R}^d \times \hat{\mathbb{R}}^d$ ,  $g, h \in L^2(\mathbb{R}^d)$ , and the so-called *upper symbol*  $\{m_{\lambda}\}_{\lambda \in \Lambda}$  is a complex valued sequence [9,10]. The operator  $G_m$  in (15) is well defined and bounded on  $L^2(\mathbb{R}^d)$ , if, for example, the Gabor systems  $(g, \Lambda)$  and  $(h, \Lambda)$  are Bessel sequences and if  $\{m_{\lambda}\} \in l^{\infty}(\Lambda)$ . Thus,  $(g, \Lambda)$  and  $(h, \Lambda)$  are dual Gabor frames for  $L^2(\mathbb{R}^d)$  if and only if  $G_{\mathbf{1}} = Id_{L^2}$  where  $\mathbf{1}_{\lambda} = 1$  for all  $\lambda \in \Lambda$ .

Discrete Gabor multipliers on  $L^2(\mathbb{R})$  can be used to model time-varying filters in communications engineering. While a convolution operator represents a time-invariant filter which allows the removal of global frequency components in a signal, a Gabor multiplier allows for the decimation of a frequency band  $[\Omega_1, \Omega_2]$  during a time interval  $[T_1, T_2]$  by setting  $m_{\lambda} = 0$  for  $\lambda = (x, \xi) \in [T_1, T_2] \times [\Omega_1, \Omega_2] \cap \Lambda \subset \mathbb{R} \times \hat{\mathbb{R}}$ .

If  $(g, \Lambda)$  is an orthonormal basis of  $L^2(\mathbb{R}^d)$ , and, therefore  $|\Lambda| = 1$ , and if  $h = g$ , then, discrete Gabor multipliers associated to  $(g, \Lambda)$  are exactly those operators mapping  $L^2(\mathbb{R}^d)$  to  $L^2(\mathbb{R}^d)$  which are represented by bi-infinite diagonal matrices with respect to the orthonormal basis  $(g, \Lambda)$ . In this case, the operator  $G_m$  in (15) is bounded if and only if  $\{m_{\lambda}\}_{\lambda \in \Lambda}$  is bounded, and  $G_m$  is stable if and only if  $\{|m_{\lambda}|^{-1}\}_{\lambda \in \Lambda}$  is well defined and bounded. Nevertheless, families of Gabor multipliers associated to Gabor frames  $(g, \Lambda)$  and  $(h, \Lambda)$  are not simultaneously diagonalizable in general if  $|\Lambda| < 1$ .

A contribution to the study of Gabor multipliers in the case  $|\Lambda| \neq 1$  is given in terms of operator identification in Theorem 5.2. This result further illuminates the role of the critical density  $|\Lambda| = 1$  in the theory of Gabor multipliers. Recall that Figs. 1.A and 1.B show that  $\{\rho(\lambda) P_{g,h}\}_{\lambda \in \Lambda}$  may or may not be a Riesz basis for its closed linear span in the space  $HS(\mathbb{R}^d)$ , regardless if  $|\Lambda| < 1$ ,  $|\Lambda| = 1$ , or  $|\Lambda| > 1$ .

**Definition 5.1.** Let  $X$  and  $Y$  be normed linear spaces over  $\mathbb{C}$ ; and let  $\mathcal{L}(X, Y)$  be the space of bounded linear operators mapping  $X$  to  $Y$ . A normed space of linear operators  $\mathcal{Z} \subset \mathcal{L}(X, Y)$  is *identifiable* if there exists  $f \in X$  such that  $\|Zf\|_Y \asymp \|Z\|_{\mathcal{Z}}$  for all  $Z \in \mathcal{Z}$ .

The operator spaces  $\mathcal{Z}$  which are considered here are defined by fixing a full rank lattice  $\Lambda$  in  $\mathbb{R}^d \times \hat{\mathbb{R}}^d$  and  $g, h \in S_0(\mathbb{R}^d)$  with  $\{\rho(\lambda) P_{g,h}\}_{\lambda \in \Lambda}$  a Bessel sequence in  $HS(\mathbb{R}^d)$ . We set

$$\mathcal{Z} = \mathcal{G}(g, h, \Lambda) = \left\{ G_m = \sum_{\lambda \in \Lambda} m_\lambda \rho(\lambda) P_{g,h} : \{m_\lambda\} \in l^2(\Lambda) \right\} \subset HS(\mathbb{R}^d), \tag{16}$$

and choose as norm on  $\mathcal{Z}$  the Hilbert–Schmidt norm, i.e.,  $\|\cdot\|_{\mathcal{Z}} = \|\cdot\|_{HS}$ . The operators in  $\mathcal{G}(g, h, \Lambda) \subset \mathcal{L}(L^2(\mathbb{R}^d), L^2(\mathbb{R}^d))$  extend to  $S'_0(\mathbb{R}^d)$  since  $g, h \in S_0(\mathbb{R}^d)$ , i.e., we have  $\mathcal{G}(g, h, \Lambda) \subset \mathcal{L}(S'_0(\mathbb{R}^d), L^2(\mathbb{R}^d))$  with domain  $X = S'_0(\mathbb{R}^d)$  and range  $Y = L^2(\mathbb{R}^d)$ .

**Theorem 5.2.** *Given a full rank lattice  $\Lambda$  in  $\mathbb{R}^d \times \mathbb{R}^d$  and  $g, h \in S_0(\mathbb{R}^d)$  such that  $\{\rho(\lambda) P_{g,h}\}_{\lambda \in \Lambda}$  is a Riesz basis for its closed linear span in  $HS(\mathbb{R}^d)$ .*

- (a) *If  $|\Lambda| > 1$  and  $(g, \Lambda)$  and  $(h, \Lambda)$  are Riesz bases for their closed linear span in  $L^2(\mathbb{R}^d)$ , then  $\mathcal{G}(g, h, \Lambda)$  is identifiable.*
- (b) *If  $|\Lambda| < 1$ , then  $\mathcal{G}(g, h, \Lambda)$  is not identifiable.*

**Proof.** (a) Let  $|\Lambda| > 1$  and  $g, h \in S_0(\mathbb{R}^d)$  with  $(g, \Lambda)$  and  $(h, \Lambda)$  are Riesz bases for their closed linear span in  $L^2(\mathbb{R}^d)$ . In order to construct  $f \in S'_0(\mathbb{R}^d)$  which identifies  $\mathcal{G}(g, h, \Lambda)$ , we pick  $\tilde{g} \in S_0(\mathbb{R}^d)$  such that  $(\tilde{g}, \Lambda^\circ)$  is a dual frame of  $(g, \Lambda^\circ)$  for  $L^2(\mathbb{R}^d)$  [38]. Consequently we have  $V_g \tilde{g}(0) = 1$  and  $V_g \tilde{g}(\lambda) = 0$  if  $\lambda \in \Lambda \setminus \{0\}$  [7, p. 133], [39]. We have  $f = \sum \pi(\lambda) \tilde{g} \in S'_0(\mathbb{R}^d)$  with weak\*-convergence [22, p. 141], and, therefore,

$$\begin{aligned} \|G_m f\|_{L^2} &= \left\| \sum_{\lambda \in \Lambda} m_\lambda \langle f, \pi(\lambda)g \rangle \pi(\lambda)h \right\|_{L^2} = \left\| \sum_{\lambda \in \Lambda} m_\lambda \left\langle \sum_{\lambda' \in \Lambda} \pi(\lambda') \tilde{g}, \pi(\lambda)g \right\rangle \pi(\lambda)h \right\|_{L^2} \\ &= \left\| \sum_{\lambda \in \Lambda} m_\lambda \pi(\lambda)h \right\|_{L^2} \asymp \|\{m_\lambda\}\|_{l^2(\Lambda)} \asymp \|G_m\|_{HS}, \end{aligned}$$

since  $(h, \Lambda)$  is a Riesz basis for its closed linear span in  $L^2(\mathbb{R}^d)$  and  $\{\rho(\lambda) P_{g,h}\}_{\lambda \in \Lambda}$  is a Riesz basis for its closed linear span in  $HS(\mathbb{R}^d)$ . Hence,  $f$  identifies  $\mathcal{G}(g, h, \Lambda)$ .

(b) Let  $|\Lambda| < 1$  and  $g, h \in S_0(\mathbb{R}^d)$ , and suppose that  $f \in S'_0(\mathbb{R}^d)$  identifies  $\mathcal{G}(g, h, \Lambda)$ . Since  $\|\{m_\lambda\}\|_{l^2} \asymp \|G_m\|_{HS}$  by hypothesis, identification of  $\mathcal{G}(g, h, \Lambda)$  by  $f$  is equivalent to the fact that the operator  $\Phi_f : l^2(\Lambda) \rightarrow L^2(\mathbb{R}^d)$ ,  $\{m_\lambda\} \mapsto G_m f$  is bounded and stable.

Let  $\mathcal{M}$  be the multiplication operator given by  $\mathcal{M} : l^2(\Lambda) \rightarrow l^2(\Lambda)$ ,  $\{m_\lambda\} \mapsto \{m_\lambda \cdot \langle f, \pi(\lambda)g \rangle\}$  and observe that we have  $\Phi_f = T_h \circ \mathcal{M}$ . The multiplication operator  $\mathcal{M}$  is bounded since  $|\langle f, \pi(\lambda)g \rangle| \leq \|f\|_{S'_0} \|g\|_{S_0}$  for all  $\lambda \in \Lambda$  and, therefore,  $\|\mathcal{M}\{m_\lambda\}\|_{l^2} \leq \|f\|_{S'_0} \|g\|_{S_0} \|\{m_\lambda\}\|_{l^2}$ . By assumption, we have  $\Phi_f$  is stable and  $T_h$  is bounded, and, hence,  $\mathcal{M}$  is stable, i.e.,  $\{|\langle f, \pi(\lambda)g \rangle|^{-1}\}$  is bounded. This implies that  $\mathcal{M}$  is onto as well, and therefore  $\mathcal{M}$  is an homeomorphism.

Since  $|\Lambda| < 1$ ,  $T_h$  is not stable, and, since  $\mathcal{M}$  is bounded and onto, this contradicts the assumption that the operator  $\Phi_f$  is stable.  $\square$

Identifiability results such as Theorem 5.2 can be found in [12]. There, it is shown that classes of Hilbert–Schmidt operators which are characterized by a rectangular band limitation of their Kohn–Nirenberg symbols are identifiable if and only if the characterizing rectangle has area less than or equal to one. Similarly, it is shown that classes of Gabor frame operators are identifiable if and only if the generating lattice  $\Lambda$  of time–frequency shifts satisfies  $|\Lambda| \leq 1$ .

Additional applications of the time–frequency analysis of such operators are found in [30,40–49].

## Acknowledgment

We gratefully acknowledge valuable conversations about this material with Hans Feichtinger.

## References

- [1] J. von Neumann, *Mathematical Foundations of Quantum Mechanics*, Princeton Univ. Press, Princeton, NJ, 1932, 1949, and 1955.
- [2] D. Gabor, Theory of communication, *J. IEE London* 93 (3) (1946) 429–457.
- [3] I. Daubechies, *Ten Lectures on Wavelets*, CBMS-NSF Regional Conf. Ser. in Appl. Math., SIAM, Philadelphia, 1992.
- [4] G. Folland, *Harmonic Analysis in Phase Space*, Ann. of Math. Stud., Princeton Univ. Press, Princeton, NJ, 1989.
- [5] L. Cohen, *Time–Frequency Analysis*, Prentice–Hall, Englewood Cliffs, NJ, 1995.
- [6] H. Feichtinger, T. Strohmer (Eds.), *Gabor Analysis and Algorithms: Theory and Applications*, Birkhäuser, Boston, MA, 1998.
- [7] K. Gröchenig, *Foundations of Time–Frequency Analysis*, Appl. Numer. Harmon. Anal., Birkhäuser, Boston, MA, 2001.
- [8] H. Feichtinger, T. Strohmer (Eds.), *Advances in Gabor Analysis*, Appl. Numer. Harmon. Anal., Birkhäuser, Boston, MA, 2003.
- [9] H. Feichtinger, Spline-type spaces in Gabor analysis, in: *Wavelet Analysis*, Hong Kong, 2001, in: Ser. Anal., vol. 1, World Scientific, River Edge, NJ, 2002, pp. 100–122.
- [10] H. Feichtinger, K. Nowak, A first survey of Gabor multipliers, in: *Advances in Gabor Analysis*, in: Appl. Numer. Harmon. Anal., Birkhäuser, Boston, MA, 2003, pp. 99–128.
- [11] H.G. Feichtinger, M. Hampejs, G. Kracher, Best approximation of matrices by Gabor multipliers, Preprint, 2004.
- [12] W. Kozek, G. Pfander, Identification of operators with bandlimited symbols, Preprint, 2004.
- [13] C.L. Fefferman, The uncertainty principle, *Bull. Amer. Math. Soc. (N.S.)* 9 (2) (1983) 129–206.
- [14] I. Daubechies, The wavelet transform, time–frequency localization and signal analysis, *IEEE Trans. Inform. Theory* (1990) 961–1005.
- [15] H. Landau, On the density of phase-space expansions, *IEEE Trans. Inform. Theory* 39 (4) (1993) 1152–1156.
- [16] J. Benedetto, S. Li, The theory of multiresolution analysis frames and applications to filter banks, *Appl. Comput. Harmon. Anal.* 5 (4) (1998) 389–427.
- [17] J. Benedetto, O. Treiber, Wavelet frames: Multiresolution analysis and extension principles, in: *Wavelet Transforms and Time–Frequency Signal Analysis*, in: Appl. Numer. Harmon. Anal., Birkhäuser, Boston, MA, 2001, pp. 3–36.
- [18] K. Gröchenig, Uncertainty principles for time–frequency representations, in: H. Feichtinger, T. Strohmer (Eds.), *Advances in Gabor Analysis*, Birkhäuser, Boston, MA, 2003, pp. 11–30.
- [19] J. Benedetto, C. Heil, D. Walnut, Differentiation and the Balian–Low theorem, *J. Fourier Anal. Appl.* 1 (4) (1995) 355–403.
- [20] K. Gröchenig, D. Han, C. Heil, G. Kutyniok, The Balian–Low theorem for symplectic lattices in higher dimensions, *Appl. Comput. Harmon. Anal.* 13 (2) (2002) 169–176.
- [21] J. Benedetto, W. Czaja, A.Y. Maltsev, The Balian–Low theorem for the symplectic form on  $\mathbb{R}^{2d}$ , *J. Math. Phys.* 44 (4) (2003) 1735–1750.
- [22] H. Feichtinger, G. Zimmermann, A space of test functions for Gabor analysis, in: H. Feichtinger, T. Strohmer (Eds.), *Gabor Analysis and Algorithms: Theory and Applications*, Birkhäuser, Boston, MA, 1998, pp. 123–170.
- [23] J. Dieudonné, *Treatise on Analysis*, vol. II (translated from the French by I.G. MacDonald). Pure and Applied Mathematics, vol. 10-II, Academic Press, New York, 1970.
- [24] S. Gaal, *Linear Analysis and Representation Theory*, Springer-Verlag, New York, 1973.
- [25] J. Kohn, L. Nirenberg, An algebra of pseudo-differential operators, *Comm. Pure Appl. Math.* 18 (1965) 269–305.
- [26] J. Benedetto, S. Li, Narrow band frame multiresolution analysis with perfect reconstruction, in: *International Symposium on Time–Frequency and Time-Scale Analysis*, Philadelphia, 1994, pp. 36–39.
- [27] C. de Boor, R.A. DeVore, A. Ron, The structure of finitely generated shift-invariant spaces in  $L_2(\mathbb{R}^d)$ , *J. Funct. Anal.* 119 (1) (1994) 37–78.
- [28] A. Ron, Z. Shen, Frames and stable bases for shift-invariant subspaces  $L_2(\mathbb{R}^d)$ , *Canad. J. Math.* 47 (5) (1995) 1051–1094.
- [29] J. Conway, N. Sloane, E. Bannai, *Sphere-Packings, Lattices, and Groups*, Springer-Verlag, New York, 1987.

- [30] H. Feichtinger, W. Kozek, Quantization of TF-lattice invariant operators on elementary LCA groups, in: H. Feichtinger, T. Strohmer (Eds.), *Gabor Analysis and Algorithms: Theory and Applications*, Birkhäuser, Boston, MA, 1998, pp. 233–266.
- [31] A. Ron, Z. Shen, Weyl–Heisenberg frames and Riesz bases in  $L_b2(\mathbf{R}^d)$ , *Duke Math. J.* 89 (2) (1997) 237–282.
- [32] H. Feichtinger, F. Luef, Wiener amalgam spaces for the fundamental identity of Gabor analysis, Preprint, 2005.
- [33] A. Janssen, Zak transforms with few zeros and the tie, in: *Advances in Gabor Analysis*, in: *Appl. Numer. Harmon. Anal.*, Birkhäuser, Boston, MA, 2003, pp. 31–70.
- [34] R. Larsen, *An Introduction to the Theory of Multipliers*, Grundlehren Math. Wiss., Band 175, Springer-Verlag, New York, 1971.
- [35] P. Gibson, M. Lamoureux, G. Margrave, Representation of linear operators by Gabor multipliers, Preprint, 2004.
- [36] E. Cordero, K. Gröchenig, Time–frequency analysis of localization operators, *J. Funct. Anal.* 205 (1) (2003) 107–131.
- [37] P. Boggiatto, E. Cordero, K. Gröchenig, Generalized anti-Wick operators with symbols in distributional Sobolev spaces, *Integral Equations Operator Theory* 48 (4) (2004) 427–442.
- [38] K. Gröchenig, M. Leinert, Wiener’s lemma for twisted convolution and Gabor frames, *J. Amer. Math. Soc.* 17 (1) (2004) 1–18 (electronic).
- [39] J. Wexler, S. Raz, Discrete Gabor expansions, *Signal Process.* 21 (3) (1990) 207–221.
- [40] T. Kailath, Measurements on time-variant communication channels, *IEEE Trans. Inform. Theory* 8 (5) (1962) 229–236.
- [41] T. Kailath, Time-variant communication channels, *IEEE Trans. Inform. Theory: Inform. Theory. Progress Report* 1960–1963 (1963) 233–237.
- [42] W. Kozek, Matched generalized Gabor expansion of nonstationary processes, in: *Proc. IEEE Int. Conf. Signals, Systems, and Computers*, Pacific Grove, CA, 1993, pp. 499–503.
- [43] C. Heil, J. Ramanathan, P. Topiwala, Singular values of compact pseudodifferential operators, *J. Funct. Anal.* 150 (2) (1997) 426–452.
- [44] W. Kozek, Adaptation of Weyl–Heisenberg frames to underspread environments, in: H. Feichtinger, T. Strohmer (Eds.), *Gabor Analysis and Algorithms: Theory and Applications*, Birkhäuser, Boston, MA, 1998, pp. 323–352.
- [45] R. Rochberg, K. Tachizawa, Pseudodifferential operators, Gabor frames, and local trigonometric bases, in: H. Feichtinger, T. Strohmer (Eds.), *Gabor Analysis and Algorithms: Theory and Applications*, Birkhäuser, Boston, MA, 1998, pp. 171–192.
- [46] K. Gröchenig, C. Heil, Modulation spaces and pseudodifferential operators, *Integral Equations Operator Theory* 34 (4) (1999) 439–457.
- [47] D. Labate, Time–frequency analysis of pseudodifferential operators, *Monatsh. Math.* 133 (2) (2001) 143–156.
- [48] D. Labate, A unified characterization of reproducing systems generated by a finite family, *J. Geom. Anal.* 12 (3) (2002) 469–491.
- [49] C. Heil, Integral operators, pseudodifferential operators, and Gabor frames, in: *Advances in Gabor Analysis*, in: *Appl. Numer. Harmon. Anal.*, Birkhäuser, Boston, MA, 2003, pp. 153–169.